Stereocontrolled Synthesis of β -D-Mannopyranuronic Acid Esters: The First Synthesis of an Alginate Trisaccharide

Leendert J. van den Bos, Jasper Dinkelaar, Herman S. Overkleeft, Gijsbert A. van der Marel*

Leiden Institute of Chemistry, Leiden University, P.O. Box 9502, 2300 RA Leiden, The Netherlands,

marel_g@chem.leidenuniv.nl

General Procedures: ^1H and ^{13}C NMR spectra were recorded on a Brüker DMX-400 and a Brüker AV-400 (400/100 MHz), Brüker AV500 (500/125 MHz) and a Brüker DMX-600 (600/150 MHz) spectrometer. Chemical shifts (δ) are given in ppm relative to tetramethylsilane as internal standard. Coupling constants are given in Hz. All given ^{13}C spectra are proton decoupled. Mass spectra were recorded with PE/SCIEX API 165 with electrospray interface and Q-Star Applied Biosystems Q-TOF (TOF-section). Optical rotations were measured on a Propol automatic polarimeter. IR-spectra were recorded on a Shimadzu FTIR-8300. Traces of water in the donor and acceptor glycosides, diphenylsulfoxide and TTBP were removed by co-evaporation with toluene. TTBP was synthesized as described by Crich *et al.*¹ Dichloromethane (DCM, Baker p.a.) was boiled under reflux over P_2O_5 for 2h and distilled prior to use. Trifluoromethanesulfonic anhydride was distilled from P_2O_5 . Molecular sieves $3\text{\AA}$ were flame dried before use. Solvents used for flash chromatography were of pro analysi quality. Flash chromatography was performed on Fluka silica gel 60 (0.04 – 0.063 mm). TLC-analysis was conducted on DC-alufolien (Merck, Kieselgel60, F_{254}) with detection by UV-absorption (254 nm) were applicable and by spraying with 20% sulfuric acid in ethanol followed by charring at $\sim 150^\circ\text{C}$ or by spraying with a solution of $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}\cdot\text{H}_2\text{O}$ (25 g/l) and $(\text{NH}_4)_4\text{Ce}(\text{SO}_4)_4\cdot 2\text{H}_2\text{O}$ (10g/l) in 10% sulfuric acid in water followed by charring at $\sim 150^\circ\text{C}$. All reactions were performed under an inert atmosphere of Argon unless stated otherwise.

NB:

1. NMR-spectroscopy of the various uronates showed low intensity of the ^{13}C signals probably due to conformational equilibriums.

2. Compounds **16**, **17**, **18**, **19**, **20** contain azido groups, which are known to be explosive.

General Procedure for the TEMPO/BAIB Mediated Oxidation:

To a vigorously stirred solution of 0.3 mmol thioglycoside in 1 mL DCM and 0.5 mL H₂O was added 0.06 mmol TEMPO (0.2 equiv.) and 0.75 mmol BAIB (2.5 equiv.). Stirring was allowed until TLC indicated complete conversion of the starting material to a lower running spot (~ 45 min). The reaction mixture was quenched by the addition of 10 mL Na₂S₂O₃ solution (10% in H₂O). The mixture was then extracted twice with EtOAc (10 ml) and the combined organic layers were dried (MgSO₄), filtered and concentrated. Flash column chromatography using EtOAc/petroleum ether (for the uronic acids 2% (v/v) AcOH is used) afforded the pure oxidized products.

General Procedure for Glycosidations using Ph₂SO/Tf₂O:

A solution of 1-thio mannopyranoside uronic acid ester (1 equiv.), diphenyl sulfoxide (1.3 equiv.) and tri-*tert*-butylpyrimidine (2.5 equiv.) in DCM (0.05M) was stirred over activated MS3Å for 30 min. The mixture was cooled to -60°C before triflic acid anhydride (1.3 equiv.) was added. The mixture was allowed to warm to -45°C in 15 min followed by addition of acceptor (1.5 equiv.) in DCM (0.15M). Stirring was continued and the reaction mixture was allowed to warm to rT. The reaction mixture was diluted with EtOAc and washed with water. The water layer was extracted twice with EtOAc. The collected organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography (EtOAc/PE) and removal of the eluent afforded the corresponding disaccharides.

General Procedure for Glycosidations using NIS/TMSOTf:

Method A (pre-mixing donor and acceptor):

A solution of 1-thio mannopyranoside uronic acid ester (1 equiv.) and acceptor (1.5 equiv.) in DCM (0.05M) was stirred over activated MS3Å for 30 min before *N*-iodosuccinimide (1.3 equiv.) was added. The mixture was cooled to -40°C before trimethylsilyl trifluoromethanesulfonate (0.1 equiv.) was added.

Stirring was continued and the reaction mixture was allowed to warm to rT. When TLC-analysis indicated complete reaction, triethylamine (5 equiv.) was added. The reaction mixture was diluted with EtOAc and washed with water. The water layer was extracted twice with EtOAc. The collected organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography (EtOAc/PE) and removal of the eluent afforded the corresponding disaccharides.

Method B (pre-activation using NIS and equimolar TMSOTf):

A solution of 1-thio mannopyranoside uronic acid ester (1 equiv.) in DCM (0.05M) was stirred over activated $\text{MS3\AA}$ for 30 min before *N*-iodosuccinimide (1.3 equiv.) was added. The mixture was cooled to -40°C before trimethylsilyl trifluoromethanesulfonate (1.3 equiv.) was added. The mixture was allowed to warm to -30°C in 15 min followed by addition of acceptor (1.5 equiv.) in DCM (0.15M). Stirring was continued and the reaction mixture was allowed to warm to rT. When TLC-analysis indicated complete reaction triethylamine (5 equiv.) was added. The reaction mixture was diluted with EtOAc and washed with water. The water layer was extracted twice with EtOAc. The collected organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography (EtOAc/PE) and removal of the eluent afforded the corresponding disaccharides.

Synthesis of Mannuronic Acid Ester Building Blocks:

Known phenyl 1-thio- α -D-mannopyranoside **22** was converted by standard procedures into benzylated derivatives **23**² and **24**³. Compound **25** was readily obtained according to a procedure described by Kong and co-workers.⁴ Thus, regioselective protection of the C-6 and C-3 position as a silyl-ether was followed by benzylation of the residual C-2 and C-4 positions using BnBr and NaH. Removal of the TBS-ether functionalities gave the desired phenyl 2,4-di-*O*-benzyl-1-thio- α -D-mannopyranoside **25** in 30% yield. Contrary to what is reported by Kong and co-workers also some 3,4-di-*O*-benzylated product was isolated. The next step was oxidation of these substrates using the 2,2,6,6-tetramethyl-1-piperidinyloxy free radical (TEMPO)/[bis(acetoxy)-iodo]benzene (BAIB) reagent system.⁵ In the case of compounds **23** and **24** the corresponding open form uronic acids were obtained (**26** and **28**, respectively), which were methylated using MeI/K₂CO₃ giving uronic ester donors **14** and **3**. Standard acetylation of compound **14** using Ac₂O/pyridine afforded donor **1**. Contrary, the 3,6-unprotected diol system **25** underwent tandem oxidation/lactonization affording lactone **27**.⁶ Acidic cleavage of the lactone **27** using Amberlite H⁺/MeOH followed by standard acetylation yielded donor **2**.

Scheme 1.

Reagents and conditions: (a) for compound **23**: ref (2), for compound **24**: ref (3), for compound **25**: TBSCl, Im., DMF, 0°C – rT, 24h then BnBr, NaH, DMF, 0°C – rT, 1h then cat. H⁺, MeOH, reflux, 10h, 30% (over 3 steps) (b) TEMPO, BAIB, DCM/H₂O (2/1), 0°C–rT, 15–30 min (87%, **26**), (63%, **27**), (89%, **28**) (c) MeI, K₂CO₃, DMF, rT, 10h (95%, **14**), (93%, **3**) (d) Ac₂O, pyridine, rT, 1h, quant. (e) Amberlite IR120 H⁺ resin, MeOH, reflux, 2h then Ac₂O, pyridine, rT, 1h, 81% (over 2 steps).

Phenyl 2,3-di-*O*-benzyl-1-thio- α -D-mannopyranosiduronic acid **26**: Diol **23** was converted according to the general procedure delivering 1-thio mannuronic acid **26** in 87% yield as a colorless oil. TLC: 50% EtOAc/PE (5% AcOH); $[\alpha]_D^{22}$: +16° ($c = 0.4$, CHCl₃); ¹H NMR (500 MHz, MeOD) δ = 3.72 (dd, 1H, $J_{3,4} = 8.0$ Hz, $J_{3,2} = 3.0$ Hz, H-3), 3.94 (t, 1H, $J_{2,3} = 3.5$ Hz, H-2), 4.26 (dd, 1H, $J_{4,3} = 8.0$ Hz, $J_{4,5} = 7.5$ Hz, H-4), 4.43 (d, 1H, $J_{5,4} = 7.0$ Hz, H-5), 4.56 (d, 1H, $J = 10.5$ Hz, CHHPh), 4.60 (d, 1H, $J = 10.5$ Hz, CHHPh), 4.63 (d, 1H, $J = 10.5$ Hz, CHHPh), 4.71 (d, 1H, $J = 10.5$ Hz, CHHPh), 5.61 (d, 1H, $J_{1,2} = 3.5$ Hz, H-1), 7.27 – 7.59 (m, 15H, H Arom); ¹³C NMR (100 MHz, MeOD) δ = 69.3 (C-4), 73.2 (CH₂ Bn), 73.5 (CH₂ Bn), 75.5 (C-5), 76.8 (C-2), 79.0 (C-3), 86.2 (C-1), 128.4 – 135.2 (CH Arom), 139.3 (C_q Bn), 139.5 (C_q Bn), 172.5 (C=O COOMe); ¹³C-GATED (125 MHz, MeOD): 86.2 ($J_{C1,H1} = 167$ Hz, C-1); ESI-MS: 489.2 (M+Na⁺); HRMS: (M+Na⁺) calcd for C₂₆H₂₆O₆SNa 489.13423, found 489.13446.

Methyl (phenyl 2,3-di-*O*-benzyl-1-thio- α -D-mannopyranoside) uronate **14**: To a stirred solution of 965 mg compound **26** (2.1 mmol) in 10 mL DMF was added 0.3 mL MeI (5 mmol, 2.5 equiv.) and 200 mg K₂CO₃. The solution was stirred overnight and quenched by the addition of 2 mL MeOH. The reaction mixture was taken up in EtOAc and washed with H₂O. The water layer was extracted with EtOAc. The collected organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. Flash column chromatography (EtOAc/PE) afforded 0.96 g of the title compound **14** (1.9 mmol, 95%) as a colorless oil. TLC: 35% EtOAc/PE; $[\alpha]_D^{22}$: +40° ($c = 1$, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ = 2.98 (s, 1H, OH-4), 3.71 (dd, 1H, $J_{3,4} = 7.2$ Hz, $J_{2,3} = 2.4$ Hz, H-3), 3.78 (s, 3H, CH₃ COOMe), 3.95 (t, 1H, $J_{2,3} = 2.4$ Hz, $J_{2,1} = 1.6$ Hz, H-2), 4.38 (t, 1H, $J_{4,3} = 7.2$ Hz, H-4), 4.57 (d, 1H, $J = 9.6$ Hz, CHHPh), 4.60 (d, 1H, $J = 9.6$ Hz, CHHPh), 4.61 (s, 1H, H-5), 4.66 (d, 1H, $J = 9.6$ Hz, CHHPh), 4.69 (d, 1H, $J = 9.6$ Hz, CHHPh), 5.61 (d, 1H, $J_{1,2} = 1.6$ Hz, H-1), 7.24 – 7.45 (m, 15H, H Arom); ¹³C NMR (100 MHz, CDCl₃) δ = 52.6 (CH₃ COOMe), 68.5 (C-4), 72.3 (CH₂ Bn), 72.5 (CH₂ Bn), 72.6 (C-5), 75.7 (C-2), 78.1 (C-3), 86.2 (C-1), 127.6 – 131.5 (CH Arom), 133.7 (C_q SPh), 137.6 (C_q Bn), 137.9 (C_q Bn), 170.3 (C=O COOMe); ESI-MS: 481.2 (M+H⁺), 498.4 (M+NH₄⁺), 503.3 (M+Na⁺), 519.0 (M+K⁺); HRMS: (M+Na⁺) calcd for C₂₇H₂₈O₆SNa 503.14988, found 503.14992.

Methyl (phenyl 4-*O*-acetyl-2,3-di-*O*-benzyl-1-thio- α -D-mannopyranoside) uronate **1**:

Compound **14** (0.96 g, 1.9 mmol) was treated with 3 mL pyridine/ Ac_2O (3:1) solution for 8h followed by the addition 2 mL MeOH. The reaction mixture was taken up in EtOAc and washed with 2M HCl and saturated NaHCO_3 . The organic layer is dried (MgSO_4), filtered and concentrated under reduced pressure. Flash column chromatography (EtOAc/PE) afforded 1.04 g of the title compound **1** (1.9 mmol, quant.) as a colorless oil. TLC: 25% EtOAc/PE; $[\alpha]_D^{22}$: $+28^\circ$ ($c = 1$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) $\delta = 2.03$ (s, 3H, CH_3 Ac), 3.60 (s, 3H, CH_3 COOMe), 3.75 (bd, 1H, $J_{2,1} = 5.0$ Hz, H-2), 3.80 (dd, 1H, $J_{3,4} = 6.0$ Hz, $J_{3,2} = 2.8$ Hz, H-3), 4.49 (d, 1H, $J = 11.9$ Hz, CHHPh), 4.54 (d, 1H, $J_{5,4} = 4.5$ Hz, H-5), 4.57 (d, 1H, $J = 12.1$ Hz, CHHPh), 4.61 (d, 1H, $J = 12.1$ Hz, CHHPh), 4.63 (d, 1H, $J = 11.9$ Hz, CHHPh), 5.55 (t, 1H, $J_{4,3} = 6.0$ Hz, $J_{4,5} = 4.5$ Hz, H-4), 5.77 (d, 1H, $J_{1,2} = 7.0$ Hz, H-1), 7.22 – 7.59 (m, 15H, H Arom); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 20.8$ (CH_3 Ac), 52.4 (CH_3 COOMe), 69.4 (C-4), 72.4 (CH_2 Bn), 72.7 (C-5), 73.1 (C-2, C-3), 82.9 (C-1), 126.7 – 131.3 (CH Arom), 133.6 (C_q SPh), 137.3 (C_q Bn), 137.4 (C_q Bn), 168.4 (C=O Ac or COOMe), 169.6 (C=O COOMe or Ac); ^{13}C -GATED (100 MHz, CDCl_3): 82.9 ($J_{\text{C1,H1}} = 168$ Hz, C-1); ESI-MS: 624.3 ($\text{M} + \text{Et}_3\text{N}^+\text{H}$); HRMS: ($\text{M} + \text{NH}_4^+$) calcd for $\text{C}_{29}\text{H}_{34}\text{O}_7\text{SN}$ 540.20560, found 540.20614.

Methyl (phenyl 4-*O*-levulinoyl-2,3-di-*O*-benzyl-1-thio- α -D-mannopyranoside) uronate

15: Compound **14** (276 mg, 0.57 mmol) was dissolved in 5.7 mL pyridine and levulinoyl anhydride (0.5M in dioxane, 3.4 mL, 1.7 mmol) was added. The reaction was stirred overnight, after which the mixture was diluted with EtOAc, washed with 1M HCl, saturated aqueous NaHCO_3 and brine. The organic layer was dried (MgSO_4), filtered and concentrated under reduced pressure. The crude product was purified by flash column chromatography (20% EtOAc/PE) to yield 330 mg of the title compound **15** (0.57 mmol, 86%) as a colorless oil. TLC: 30% EtOAc/PE; ^1H NMR (400 MHz, CDCl_3) $\delta = 2.17$ (s, 3H, CH_3 Lev), 2.52 (q, 2H, $J = 6.4$ Hz, $J = 6.0$ Hz, CH_2 Lev), 2.69 – 2.72 (m, 2H, $J = 6.0$ Hz, $J = 3.2$ Hz, CH_2 Lev), 3.58 (s, 3H, CH_3 COOMe), 3.74 (bd, 1H, $J_{2,1} = 6.0$ Hz, H-2), 3.86 (dd, 1H, $J_{3,4} = 6.0$ Hz, $J_{3,2} = 2.8$ Hz, H-3), 4.53 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.54 (d, 1H, $J_{5,4} = 5.6$ Hz, H-5), 4.56 (d, 1H, $J = 12.8$ Hz, CHHPh), 4.61 (d, 1H, $J = 12.8$ Hz, CHHPh), 4.64 (d, 1H, $J = 12.0$ Hz, CHHPh), 5.57 (t, 1H, $J_{4,3} = 5.6$ Hz, H-4), 5.78 (d, 1H, $J_{1,2} = 7.2$ Hz, H-1), 7.21 – 7.60 (m, 15H, H Arom); ^{13}C NMR (100 MHz,

CDCl₃) δ = 27.8 (CH₂ Lev), 29.7 (CH₃ Lev), 37.7 (CH₂ Lev), 52.2 (CH₃ COOMe), 69.6 (C-4), 72.9 (C-5), 73.8 (C-3), 74.3 (C-2), 82.9 (C-1), 168.4 (C=O Lev or COOMe), 171.5 (C=O COOMe or Lev), 206.0 (C=O Lev ketone); ESI-MS: 601.2 (M+Na⁺). HRMS: (M+NH₄⁺) calcd for C₃₂H₃₈O₈SN 596.23126, found 596.23024.

Phenyl 2,4-di-O-benzyl- α -D-mannopyranoside **25**: To a cooled (0°C) solution of 2.71 g compound **22** (10.0 mmol) in 50 mL DMF was added 3.24 g TBSCl (21.5 mmol, 2.15 equiv.) and 3.18 g imidazole (21.5 mmol, 2.15 equiv.). The mixture was allowed to warm to room temperature. After stirring for 8 h the reaction was quenched with MeOH, taken up in EtOAc and washed with H₂O. The organic layer were dried (MgSO₄), filtered and concentrated under reduced pressure. The crude product was filtered through a plug of silica gel using (10% EtOAc/PE) as the eluent. After evaporation the product was dissolved in 50 mL DMF and cooled to 0°C before 3 ml BnBr (25 mmol, 2.5 equiv.) and 1 g NaH (60% in min. oil, 25 mmol, 2.5 equiv.) were added. The mixture was allowed to stir for 12 h before the reaction was quenched by addition of MeOH, taken up in Et₂O and washed with H₂O. The organic layers was dried (MgSO₄), filtered and concentrated under reduced pressure. The oily substance was suspended in 50 mL MeOH and a catalytic amount of *p*-TsOH was added. The mixture was refluxed until TLC indicated complete conversion. After addition of 5 mL NEt₃ the reaction mixture was concentrated under reduced pressure. Flash chromatography afforded 1.35 g of the title compound **25** (2.98 mmol, 30%) as a colorless oil. TLC: 50% EtOAc/PE; $[\alpha]_D^{22}$: +59° (c = 1, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ = 1.84 (bs, 1H, OH-6), 2.42 (d, 1H, $J_{OH3,H3}$ = 7.6 Hz, OH-3), 3.74 (t, 1H, $J_{4,3}$ = $J_{4,5}$ = 9.6 Hz, H-4), 3.78 – 3.85 (m, 2H, H-6), 4.0 (bs, 2H, H-2, H-3), 4.12 (dt, 1H, $J_{5,6}$ = 4.0 Hz, $J_{5,6}$ = 3.2 Hz, H-5), 4.56 (d, 1H, J = 11.6 Hz, CHHPh), 4.67 (d, 1H, J = 11.6 Hz, CHHPh), 4.75 (d, 1H, J = 11.6 Hz, CHHPh), 4.93 (d, 1H, J = 11.6 Hz, CHHPh), 5.57 (s, 1H, H-1), 7.24 – 7.48 (m, 15H, H Arom); ¹³C NMR (100 MHz, CDCl₃) δ = 62.1 (C-6), 72.2 (C-2 or C-3), 72.4 (C-5), 72.5 (CH₂ Bn), 74.9 (CH₂ Bn), 76.4 (C-4), 79.7 (C-3 or C-2), 85.1 (C-1), 127.4 – 131.9 (CH Arom), 133.6 (C_q SPh), 137.2 (C_q Bn), 138.1 (C_q Bn); ¹³C-GATED (100 MHz, CDCl₃): 85.1 ($J_{C1,H1}$ = 165 Hz, C-1); ESI-MS: 475.1 (M+Na⁺); HRMS: (M+Na⁺) calcd for C₂₆H₂₈O₅SN 475.15497, found 475.15565.

Phenyl 2,4-di-*O*-benzyl-1-thio- α -D-mannopyranosidurono-6,3-lactone **27**: Diol **25** was

converted according to the general procedure delivering 1-thio mannuronic acid lactone

27 in 63% yield as a colorless oil. TLC: 25% EtOAc/PE; ^1H NMR (500 MHz, CDCl_3) δ

= 3.94 (dd, 1H, $J_{2,1} = 9.0$ Hz, $J_{2,3} = 1.0$ Hz, H-2), 4.03 (dd, 1H, $J_{4,3} = 6.0$ Hz, $J_{4,5} = 3.0$ Hz, H-4), 4.35 (d, 1H, $J_{5,4} = 2.5$ Hz, H-5), 4.44 (d, 1H, $J = 11.5$ Hz, CHHPh), 4.58 (d, 1H, $J = 11.5$ Hz, CHHPh), 4.66 (d, 1H, $J = 11.5$ Hz, CHHPh), 4.70 (d, 1H, $J_{3,4} = 6.0$ Hz, H-3), 4.78 (d, 1H, $J = 11.5$ Hz, CHHPh), 5.00 (d, 1H, $J_{1,2} = 9.0$ Hz, H-1), 7.24 – 7.65 (m, 15H, H Arom); ^{13}C NMR (125 MHz, CDCl_3) δ = 71.7 (CH_2 Bn), 72.0 (C-5), 72.9 (CH_2 Bn), 73.1 (C-2), 74.9 (C-4), 77.8 (C-3), 84.3 (C-1), 127.8 – 132.4 (CH Arom), 132.1 (C_q SPh), 136.3 (C_q Bn), 137.2 (C_q Bn), 169.6 (C=O lactone); ^{13}C -GATED (125 MHz, CDCl_3): 84.3 ($J_{\text{C1,H1}} = 158$ Hz, C-1); ESI-MS: 448.2 ($\text{M}+\text{H}^+$); HRMS: ($\text{M}+\text{H}^+$) calcd for $\text{C}_{26}\text{H}_{25}\text{O}_5\text{S}$ 448.13444, found 448.19671.

Methyl (phenyl 3-*O*-acetyl-2,4-di-*O*-benzyl-1-thio- α -D-mannopyranoside) uronate **2**:

Compound **27** (170 mg, 0.37 mmol) was dissolved in MeOH and, after addition of 200

mg Amberlite IR120 H^+ resin, heated under reflux. After 4h, TLC analysis indicated complete conversion of starting material into a lower running spot. The reaction mixture was filtered and concentrated under reduced pressure. The product was treated with 5 ml pyridine/ Ac_2O (3:1) for 8h followed by addition of MeOH. The reaction mixture was diluted with EtOAc and washed with 2M HCl and sat. NaHCO_3 . The organic layer was dried (MgSO_4), filtered and concentrated under reduced pressure. Flash column chromatography (EtOAc/PE) afforded 156 mg of the title compound **2** (0.3 mmol, 81%) as a colorless oil. TLC: 25% EtOAc/PE; $[\alpha]_D^{22}$: $+29^\circ$ ($c = 1.5$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ = 1.97 (s, 3H, CH_3 Ac), 3.72 (CH_3 COOMe), 3.99 (dd, 1H, $J_{2,1} = 5.1$ Hz, $J_{2,3} = 3.1$ Hz, H-2), 4.25 (t, 1H, $J_{4,3} = J_{4,5} = 6.9$ Hz, H-4), 4.48 (d, 1H, $J = 11.8$ Hz, CHHPh), 4.59 (d, 1H, $J = 11.8$ Hz, CHHPh), 4.61 (d, 1H, $J = 11.4$ Hz, CHHPh), 4.67 (d, 1H, $J = 11.4$ Hz, CHHPh), 4.68 (d, 1H, $J_{5,4} = 6.9$ Hz, H-5), 5.25 (dd, 1H, $J_{3,4} = 7.3$ Hz, $J_{3,2} = 3.1$ Hz, H-3), 5.58 (d, 1H, $J_{1,2} = 5.1$ Hz, H-1), 7.24 – 7.56 (m, 15H, H Arom); ^{13}C NMR (100 MHz, CDCl_3) δ = 20.8 (CH_3 Ac), 52.4 (CH_3 COOMe), 70.9 (C-3), 72.3 (CH_2 Bn), 72.7 (C-5), 73.8 (CH_2 Bn), 74.6 (C-2), 75.0 (C-4), 84.1 (C-1), 127.5 – 131.7 (CH Arom), 133.3 (C_q SPh), 137.3 (C_q Bn), 137.6 (C_q Bn), 169.4 (C=O Ac or COOMe), 169.7 (C=O COOMe or Ac); ^{13}C -GATED (100 MHz, CDCl_3): 84.1

($J_{C1,H1}$ = 165 Hz, C-1); ESI-MS: 540.4 ($M+NH_4^+$); HRMS: ($M+NH_4^+$) calcd for $C_{29}H_{34}O_7SN$ 540.20560, found 540.20607.

Phenyl 2,3,4-tri-*O*-benzyl-1-thio- α -D-mannopyranosiduronic acid **28**: Compound **24**

was converted according to the general procedure delivering 1-thio mannuronic acid **28** in 68% yield as a colorless oil. TLC: 25% EtOAc/PE; $[\alpha]_D^{22}$: +34° (c = 1, $CHCl_3$); 1H NMR (400 MHz, $CDCl_3$) δ = 3.81 (dd, 1H, $J_{3,4}$ = 6.8 Hz, $J_{3,2}$ = 2.8 Hz, H-3), 3.91 (dd, 1H, $J_{2,1}$ = 5.2 Hz, $J_{2,3}$ = 2.8 Hz, H-2), 4.23 (t, 1H, $J_{4,3}$ = $J_{4,5}$ = 6.8 Hz, H-4), 4.52 (d, 1H, J = 12.0 Hz, $CHHPh$), 4.54 (d, 1H, J = 12.0 Hz, $CHHPh$), 4.57 (d, 1H, J = 12.0 Hz, $CHHPh$), 4.64 (d, 1H, $J_{5,4}$ = 6.8 Hz, H-5), 4.65 – 4.73 (m, 3H, 2x $CHHPh$, $CHHPh$), 5.67 (d, 1H, $J_{1,2}$ = 5.2 Hz, H-1), 7.22 – 7.51 (m, 20H, H Arom); ^{13}C NMR (100 MHz, $CDCl_3$) δ = 72.0, 72.3 (CH_2 Bn), 72.3 (CH_2 Bn), 74.9, 75.7 127.4 – 131.5 (CH Arom), 133.5 (C_q SPh), 137.5 (C_q Bn), 137.7 (C_q Bn), 172.4 (C=O COOH). ESI-MS: 579.3 ($M+Na^+$); HRMS: ($M+Na^+$) calcd for $C_{33}H_{32}O_6SNa$ 579.18118 found 579.18222.

Methyl (phenyl 2,3,4-tri-*O*-benzyl-1-thio- α -D-mannopyranoside) uronate **3**: To a stirred

solution of 373 mg compound **28** (0.67 mmol) in 2.2 mL DMF was added 92 μ L MeI (1.5 mmol, 2 equiv.) and 200 mg K_2CO_3 . The solution was stirred overnight and quenched by the addition of 2 mL MeOH. The reaction mixture was diluted with EtOAc and washed with H_2O . The water layer was extracted with EtOAc. The collected organic layers were dried ($MgSO_4$), filtered and concentrated under reduced pressure. Flash column chromatography (EtOAc/PE) afforded 355 mg of the title compound **3** (0.62 mmol, 92%) as a colorless oil. TLC: 25% EtOAc/PE; 1H NMR (400 MHz, $CDCl_3$) δ = 3.46 (s, 3H, CH_3 COOMe), 3.81 (dd, 1H, $J_{3,4}$ = 7.0 Hz, $J_{3,2}$ = 2.8 Hz, H-3), 3.89 (dd, 1H, $J_{2,1}$ = 5.2 Hz, $J_{2,3}$ = 2.8 Hz, H-2), 4.26 (t, 1H, $J_{4,3}$ = 7.0 Hz, $J_{4,5}$ = 6.2 Hz, H-4), 4.52 (d, 1H, J = 12.0 Hz, $CHHPh$), 4.55 (s, 2H, CH_2 Bn), 4.63 (d, 1H, $J_{5,4}$ = 6.2 Hz, H-5), 4.64 – 4.69 (m, 3H, 2x $CHHPh$, $CHHPh$), 5.67 (d, 1H, $J_{1,2}$ = 5.2 Hz, H-1), 7.23 – 7.53 (m, 20H, CH Arom); ^{13}C NMR (100 MHz, $CDCl_3$) δ = 52.2 (CH_3 COOMe), 72.1 (CH_2 Bn), 72.4 (CH_2 Bn), 72.9 (C-5), 73.7 (CH_2 Bn), 74.9 (C-2), 75.6 (C-4), 77.1 (C-3), 84.7 (C-1), 127.2 – 131.4 (CH Arom), 134.9 (C_q SPh), 137.8 (2x C_q Bn), 137.9 (C_q Bn), 169.2 (C=O COOMe); ^{13}C -GATED (100

MHz, CDCl₃): 84.7 ($J_{C1,H1}$ = 163 Hz, C-1); ESI-MS: 593.2 (M+Na⁺); HRMS: (M+Na⁺) calcd for C₃₄H₃₄O₆SNa 593.19683 found 593.19780.

Methyl (2,3,4-tri-*O*-benzyl-6-*O*-(methyl (4-*O*-acetyl-2,3-di-*O*-benzyl-β-D-mannopyranosyl) uronate)-α-D-glucopyranoside **8**: Colorless oil. TLC: 30% EtOAc/PE; $[\alpha]_D^{22}$: -11° (c = 0.6, CHCl₃); ¹H NMR (500 MHz,

CDCl₃) δ = 2.02 (s, 3H, CH₃ Ac), 3.31 (s, 3H, CH₃ OMe), 3.37 (dd, 1H, $J_{3',4'} = 9.5$ Hz, $J_{3',2'} = 2.5$ Hz, H-3'), 3.41 (m, 2H, $J_{4,5} = J_{4,3} = 9.5$ Hz, H-4 and H-6), 3.50 (dd, 1H, $J_{2,3} = 9.5$ Hz, $J_{2,1} = 3.5$ Hz, H-2), 3.71 (s, 4H, H-2' and CH₃ COOMe), 3.75 (d, 1H, $J_{5',4'} = 9.5$ Hz, H-5'), 3.79 (bt, 1H, $J_{5,4} = 8.0$ Hz, $J_{5,6} = 6.0$ Hz, H-5), 4.01 (t, 1H, $J_{3,2} = J_{3,4} = 9.5$ Hz, H-3), 4.12 (bs, 2H, H-1' and H-6), 4.36 (d, 1H, $J = 12.5$ Hz, CHHPh), 4.47 (d, 1H, $J = 12.5$ Hz, CHHPh), 4.49 (d, 1H, $J = 12.5$ Hz, CHHPh), 4.56 (d, 1H, $J_{1,2} = 3.5$ Hz, H-1), 4.66 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.75 (d, 1H, $J = 11.5$ Hz, CHHPh), 4.79 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.81 (d, 1H, $J = 11.5$ Hz, CHHPh), 4.84 (d, 1H, $J = 10.5$ Hz, CHHPh), 4.90 (d, 1H, $J = 12.5$ Hz, CHHPh), 5.01 (d, 1H, $J = 10.5$ Hz, CHHPh), 4.87 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.5$ Hz, H-4'), 7.18 – 7.37 (m, 25H H Arom); ¹³C NMR (125 MHz, CDCl₃) δ = 20.8 (CH₃ Ac), 48.5 (CH₃ COOMe), 52.5 (CH₃ OMe), 68.8 (C-6), 68.9 (C-4'), 69.7 (C-5), 71.5 (CH₂ OBn), 72.9 (C-2'), 73.3 (CH₂ Bn), 73.6 (CH₂ Bn), 73.7 (C-5'), 74.7 (CH₂ Bn), 75.7 (CH₂ Bn), 77.6 (C-4), 78.2 (C-3'), 79.8 (C-2), 82.1 (C-3), 97.8 (C-1), 101.6 (C-1'), 127.4 – 128.5 (CH Arom), 137.6 (C_q Bn), 138.0 (C_q Bn), 138.3 (2xC_q Bn), 138.7 (C_q Bn), 167.9 (C=O Ac or COOMe), 169.5 (C=O Ac or COOMe); ¹³C-GATED (125 MHz, CDCl₃): 97.8 ($J_{C1,H1}$ = 169 Hz, C-1), 101.6 ($J_{C1',H1'}$ = 156 Hz, C-1'); ESI-MS: 894.5 (M+NH₄⁺), 900.5 (M+Na⁺), 915.4 (M+K⁺); HRMS: (M+NH₄⁺) calcd for C₅₁H₆₀O₁₃N 894.40647, found 894.40531.

N.B.: Conclusive evidence for the β-orientation was gained by reducing both ester functions after which the heteronuclear coupling constant was determined at ¹ J_{CH} = 153 Hz.

Methyl (2,3,4-tri-*O*-benzyl-6-*O*-(methyl (4-*O*-acetyl-2,3-di-*O*-benzyl-α-D-mannopyranosyl) uronate)-α-D-glucopyranoside **9**: Colorless oil. TLC: 30%

EtOAc/PE; ^1H NMR (400 MHz, CDCl_3) δ = 1.96 (s, 3H, CH_3 Ac), 3.36 (s, 3H, CH_3 OMe), 3.48 (t, 1H, $J_{4,3}$ = $J_{4,5}$ = 9.2 Hz, H-4), 3.50 (dd, 1H, $J_{2,3}$ = 9.6 Hz, $J_{2,1}$ = 3.6 Hz, H-2), 3.68 (s, 3H, CH_3 COOMe), 3.71 – 3.77 (m, 2H, H-5, H-6), 3.86 (t, 1H, $J_{2',1'} = J_{2',3'} = 3.0$ Hz, H-2'), 3.90 (dd, 1H, $J_{6,5}$ = 11.1 Hz, $J_{6,5}$ = 4.8 Hz, H-6), 3.99 (t, 1H, $J_{3,2}$ = 9.6 Hz, $J_{3,4}$ = 9.2 Hz, H-3), 4.19 (t, 1H, $J_{4',3'} = J_{4',5'} = 8.7$ Hz, H-4'), 4.29 (d, 1H, $J_{5',4'} = 8.7$ Hz, H-5'), 4.53 (d, 1H, J = 12.3 Hz, CHHPh), 4.56 (d, 1H, $J_{1,2}$ = 3.6 Hz, H-1), 4.61 (d, 1H, J = 10.9 Hz, CHHPh), 4.63 (d, 1H, J = 12.0 Hz, CHHPh), 4.65 (s, 2H, CH_2 Bn), 4.67 (d, 1H, J = 12.3 CHHPh), 4.78 (d, 1H, J = 12.0 Hz, CHHPh), 4.82 (d, 1H, J = 10.9 Hz, CHHPh), 4.92 (d, 1H, J = 10.9 Hz, CHHPh), 4.99 (d, 1H, J = 11.1 Hz, CHHPh), 5.00 (s, 1H, H-1'), 5.24 (dd, 1H, $J_{3',4'} = 8.6$ Hz, $J_{3',2'} = 3.2$ Hz, H-3'), 7.23 – 7.38 (m, 25H, H Arom); ^{13}C NMR (100 MHz, CDCl_3) δ = 20.9 (CH_3 Ac), 52.3 (CH_3 OMe), 55.1 (CH_3 COOMe), 66.7 (C-6), 69.8 (C-5), 71.5 (C-5'), 72.5 (C-3'), 72.9 (CH_2 Bn), 73.3 (CH_2 Bn), 74.3 (CH_2 Bn), 74.8 (C-4'), 75.0 (CH_2 Bn), 75.0 (C-2'), 75.7 (CH_2 Bn), 77.6 (C-4), 79.9 (C-2), 82.1 (C-3), 97.8 (C-1), 98.5 (C-1'); 127.5 – 129.1 (CH Arom), 137.7 (C_q Bn), 137.9 (C_q Bn), 138.1 (C_q Bn), 138.6 (C_q Bn), 169.6 (C=O COOMe or Ac), 169.9 (C=O Ac or COOMe); ^{13}C -GATED (100 MHz, CDCl_3): 97.8 ($J_{\text{Cl},\text{H}1}$ = 166 Hz, C-1), 98.5 ($J_{\text{Cl},\text{H}1'} = 170$ Hz, C-1'); ESI-MS: 900.6 ($\text{M}+\text{Na}^+$); HRMS: ($\text{M}+\text{NH}_4^+$) calcd for $\text{C}_{51}\text{H}_{60}\text{O}_{13}\text{N}$ 894.40647, found 894.40545.

para-Methoxyphenyl (2-*O*-benzyl-4,6-*O*-benzylidene-3-*O*-(methyl (2,3,4-tri-*O*-benzyl-β-D-mannopyranosyl) uronate)-β-D-galactopyranoside **10**: Colorless oil. TLC: 30% EtOAc/PE; $[\alpha]_{\text{D}}^{22}$: -4° (c

= 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ = 3.16 (dd, 1H, $J_{3',4'} = 9.6$ Hz, $J_{3',2'} = 2.9$ Hz, H-3'), 3.56 (s, 1H, H-5), 3.63 (d, 1H, $J_{2',1'} = J_{2',3'} = 2.8$ Hz, H-2'), 3.72 (s, 3H, CH_3 COOMe or pMP), 3.73 (d, 1H, $J_{5',4'} = 9.2$ Hz, H-5'), 3.79 (s, 3H, CH_3 COOMe or pMP), 3.84 (dd, 1H, $J_{3,2}$ = 9.9 Hz, $J_{3,4}$ = 3.4 Hz, H-3), 4.08 (dd, 1H, $J_{2,3}$ = 9.9 Hz, $J_{2,1}$ = 7.6 Hz, H-2), 4.09 (dd, 1H, $J_{6,5}$ = 12.3 Hz, $J_{6,6}$ = 1.7 Hz, H-6), 4.18 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.7$ Hz, H-4'), 4.28 (d, 1H, J = 11.6 Hz, CHHPh), 4.30 (d, 1H, J = 11.3 Hz, CHHPh), 4.36 (d, 1H, $J_{6,5}$ = 12.3 Hz, H-6), 4.37 (d, 1H, J = 11.6 Hz, CHHPh), 4.39 (d, 1H, $J_{4,3}$ = 3.4 Hz, H-4), 4.61 (d, 1H, J = 10.7 Hz, CHHPh), 4.71 (s, 1H, H-1'), 4.85 (d, 1H, J = 10.7 CHHPh), 4.91 (d, 1H, $J_{1,2}$ = 7.6 Hz, H-1), 4.93 (s, 2H, CH_2 Bn), 4.96 (d, 1H, J = 11.6 Hz, CHHPh), 5.62 (s, 1H, CHPh), 6.84 (d, 1H, J = 9.0 Hz, H Arom pMP), 7.05 (d, 1H, J = 9.0 Hz, H Arom pMP), 7.10 – 7.60 (m, 25H, H Arom); ^{13}C NMR (100 MHz,

CDCl₃) δ = 52.3 (CH₃ COOMe or pMP), 55.6 (CH₃ COOMe or pMP), 66.7 (C-5), 68.9 (C-6), 71.8 (CH₂ Bn), 71.9 (C-2'), 73.4 (CH₂ Bn), 75.2 (2xCH₂ Bn), 75.3 (C-5'), 75.7 (C-4'), 75.8 (C-4), 77.7 (C-3), 79.0 (C-2), 82.0 (C-3'), 100.8 (CHPh), 103.2 (C-1 and C-1'), 114.5 (CH Arom pMP), 118.9 (CH Arom pMP), 126.3 – 128.7 (CH Arom), 137.9 (C_q Bn), 138.1 (C_q Bn), 138.3 (C_q Bn), 138.4 (C_q Bn), 151.5 (C_q pMP), 155.4 (C_q pMP), 168.7 (C=O COOMe); ¹³C-GATED (100 MHz, CDCl₃): 103.2 ($J_{Cl',HI'} = 159$ Hz, C-1'), 103.2 ($J_{Cl,HI} = 159$ Hz, C-1); ESI-MS: 947.3 (M+Na⁺); HRMS: (M+Na⁺) calcd for C₅₅H₅₆O₁₃Na 947.36131, found 947.36288.

para-Methoxyphenyl (2-*O*-benzyl-4,6-*O*-benzylidene-3-*O*-(methyl 4-*O*-acetyl-2,3-di-*O*-benzyl- β -D-mannopyranosyl)uronate)- β -D-galactopyranoside **11**: Colorless oil. TLC: 30% EtOAc/PE; $[\alpha]_D^{22}$: -28°

($c = 0.8$, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ = 1.97 (s, 3H, CH₃ Ac), 3.12 (dd, 1H, $J_{3',4'} = 9.8$ Hz, $J_{3',2'} = 2.8$ Hz, H-3'), 3.53 (s, 1H, H-5), 3.65 (d, 1H, $J_{2',1'} = J_{2',3'} = 2.7$ Hz, H-2'), 3.70 (s, 1H, $J_{5',4'} = 9.7$ Hz, H-5'), 3.72 (s, 3H, CH₃ COOMe or OMe), 3.78 (s, 3H, CH₃ OMe or COOMe), 3.83 (dd, 1H, $J_{3,2} = 9.9$ Hz, $J_{3,4} = 3.4$ Hz, H-3), 4.07 – 4.10 (m, 2H, H-2, H-6), 4.18 (d, 1H, $J = 12.1$ Hz, CHHPh), 4.30 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.37 (m, 3H, H-6, 2xCHHPh), 4.42 (d, 1H, $J_{4,3} = 3.4$ Hz, H-4), 4.70 (s, 1H, H-1'), 4.88 (d, 1H, $J = 11.3$ Hz, CHHPh), 4.90 (s, 1H, H-1), 4.95 (d, 1H, $J = 11.3$ Hz, CHHPh), 5.45 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.7$ Hz, H-4'), 5.64 (s, 1H, CHPh), 6.84 (d, 1H, $J = 9.0$ Hz, H Arom pMP), 7.07 (d, 1H, $J = 9.0$ Hz, H Arom pMP), 7.10 – 7.60 (m, 20H, H Arom); ¹³C NMR (125 MHz, CDCl₃) δ = 20.7 (CH₃ Ac), 52.5 (CH₃ OMe), 55.6 (CH₃ COOMe), 66.7 (C-5), 68.7 (C-4'), 68.9 (C-6), 71.6 (C-2'), 68.9 (CH₂ Bn), 73.4 (CH₂ Bn), 73.5 (C-5'), 75.2 (CH₂ Bn), 75.7 (C-4), 77.7 (C-3), 79.0 (C-2, C-3'), 100.8 (CHPh), 102.5 (C-1'), 103.1 (C-1), 114.5 (CH Arom pMP), 118.8 (CH Arom pMP), 126.3 – 128.8 (CH Arom), 137.7 (C_q Bn), 137.9 (C_q Bn), 138.2 (C_q Bn), 138.5 (C_q Bn), 151.4 (C_q pMP), 155.4 (C_q pMP), 167.9 (C=O Ac or COOMe), 169.5 (C=O COOMe or Ac); ¹³C-GATED (125 MHz, CDCl₃): 102.5 ($J_{Cl',HI'} = 164$ Hz, C-1'), 103.1 ($J_{Cl,HI} = 158$ Hz, C-1); ESI-MS: 894.5 (M+NH₄⁺), 900.3 (M+Na⁺); HRMS: (M+NH₄⁺) calcd for C₅₀H₅₆O₁₄N 894.37008, found 894.36879.

3-*O*-(methyl (4-*O*-acetyl-2,3-di-*O*-benzyl-β-*D*-mannopyranosyl) uronate)-

1,2;5,6-di-*O*-isopropylidene-α-*D*-glucopyranoside **12**: Colorless oil. TLC:

30% EtOAc/PE; $[\alpha]_D^{22}$: -34° ($c = 0.7$, CHCl_3); ^1H NMR (400 MHz,

CDCl_3) $\delta = 1.22$ (s, 3H, CH_3 isoprop), 1.24 (s, 3H, CH_3 isoprop), 1.35 (s,

3H, CH_3 isoprop), 1.49 (s, 3H, CH_3 isoprop), 1.96 (s, 3H, CH_3 Ac), 3.42 (dd, 1H, $J_{3',4'} = 9.6$ Hz, $J_{3',2'} = 3.2$ Hz, H-3'), 3.65 (s, 3H, CH_3 COOMe), 3.76 (d, 1H, $J_{5',4'} = 9.2$ Hz, H-5'), 3.77 (d, 1H, $J_{2',1'} = J_{2',3'} = 3.2$ Hz, H-2'), 3.97 (dd, 1H, $J_{6,5} = 8.4$ Hz, $J_{6,6} = 6.0$ Hz, H-6), 4.09 (dd, 1H, $J_{6,5} = 8.4$ Hz, $J_{6,6} = 6.8$ Hz, H-6), 4.18 (d, 1H, $J_{3,2} = J_{3,4} = 3.2$ Hz, H-3), 4.25 (dd, 1H, $J_{4,5} = 4.9$ Hz, $J_{4,3} = 3.2$ Hz, H-4), 4.33 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.34 (d, 1H, $J_{3,2} = J_{2,1} = 4.0$ Hz, H-2), 4.39 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.45 (dd, 1H, $J_{5,6} = 11.2$ Hz, $J_{5,4} = 1.6$ Hz, H-5), 4.46 (s, 1H, H-1'), 4.67 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.76 (d, 1H, $J = 12.4$ Hz, CHHPh), 5.43 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.6$ Hz, H-4'), 5.82 (d, 1H, $J_{1,2} = 3.6$ Hz, H-1), 7.15 – 7.32 (m, 10H, H Arom); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 20.8$ (CH_3 Ac), 20.0 (CH_3 isoprop), 26.3 (CH_3 isoprop), 26.5 (CH_3 isoprop), 26.7 (CH_3 isoprop), 52.6 (CH_3 COOMe), 65.9 (C-6), 68.6 (C-3'), 71.8 (CH_2 Bn), 73.3 (C-5'), 73.5 (C-5 or C-2'), 73.6 (C-2' or C-5), 73.9 (CH_2 Bn), 78.2 (C-4'), 80.5 (C-4), 81.8 (C-3), 83.0 (C-2), 99.8 (C-1'), 104.9 (C-1), 108.4 (C_q isoprop), 111.9 (C_q isoprop), 127.5 – 128.4 (CH Arom), 137.6 (C_q Bn), 137.9 (C_q Bn), 167.7 (C=O Ac or COOMe), 169.5 (C=O Ac or COOMe); ^{13}C -GATED (125 MHz, CDCl_3): 99.8 ($J_{\text{Cl,H1}} = 157$ Hz, C-1'), 104.9 ($J_{\text{Cl,H1}} = 181$ Hz, C-1); ESI-MS: 673.5 ($\text{M}+\text{H}^+$); HRMS: ($\text{M}+\text{H}^+$) calcd for $\text{C}_{35}\text{H}_{45}\text{O}_{13}$ 673.28547 found 673.28598.

para-Methoxyphenyl (methyl 2,3-di-*O*-benzyl-4-*O*-(methyl (4-*O*-acetyl-2,3-di-*O*-benzyl-α-*D*-mannopyranosyl) uronate)-β-*D*-galactopyranoside) uronate

13α: Colorless oil. TLC: 30% EtOAc/PE; $[\alpha]_D^{22}$: $+2^\circ$ ($c = 0.3$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) $\delta = 2.03$ (s, 3H, CH_3 Ac), 3.53 (dd, 1H, $J_{3,2} = 7.3$

Hz, $J = 3.1$ Hz, H-3), 3.54 (s, 3H, CH_3 COOMe), 3.57 (s, 3H, CH_3 COOMe), 3.77 (s, 4H, H-2', CH_3 OMe), 3.86 (dd, 1H, $J_{3',4'} = 8.3$ Hz, $J_{3',2'} = 3.0$ Hz, H-3'), 3.91 (dd, 1H, $J_{2,1} = 9.8$ Hz, $J_{2,3} = 7.8$ Hz, H-2), 4.09 (s, 1H, H-4), 4.53 (d, 1H, $J = 12.3$ Hz, CHHPh), 4.54 (d, 1H, $J_{5,4} = 1.1$ Hz, H-5), 4.61 (d, 1H, $J = 12.3$ Hz, CHHPh), 4.62 (d, 1H, $J = 11.7$ Hz, CHHPh), 4.65 (d, 1H, $J = 10.9$ Hz, CHHPh), 4.68 (d, 1H, $J_{5',4'} = 8.1$ Hz, H-5'), 4.76 (d, 1H, $J = 12.6$ Hz, CHHPh), 4.80 (d, 1H, $J_{1,2} = 7.7$ Hz, H-1), 4.83 (d, 1H, $J = 12.6$ Hz,

CHHPh), 4.91 (d, 1H, $J = 10.9$ Hz, CHHPh), 5.01 (d, 1H, $J = 10.9$ Hz, CHHPh), 5.15 (d, 1H, $J_{1',2'} = 2.5$ Hz, H-1'), (t, 1H, $J_{4',3'} = J_{4',5'} = 8.1$ Hz, H-4'), 6.81 (d, 1H, $J = 9.1$ Hz, CH Arom pMP), 7.07 (d, 1H, $J = 9.1$ Hz, CH Arom pMP), 7.24 – 7.40 (m, 20H, CH Arom); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 20.9$ (CH_3 Ac), 52.3 (CH_3 COOMe), 52.4 (CH_3 COOMe), 55.6 (CH_3 OMe), 69.3 (C-4'), 71.2 (C-5'), 71.8 (CH_2 Bn), 72.6 (CH_2 Bn), 72.8 (CH_2 Bn), 73.3 (C-4), 73.6 (C-2'), 75.2 (CH_2 Bn), 75.3 (C-5), 76.4 (C-3'), 77.6 (C-2), 78.7 (C-3), 99.8 (C-1'), 103.1 (C-1), 114.5 (CH Arom pMP), 118.8 (CH Arom pMP), 127.4 – 128.3 (CH Arom), 137.7 (C_q Bn), 138.0 (C_q Bn), 138.3 (C_q Bn), 151.6 (C_q pMP), 155.5 (C_q pMP), 167.7 (C=O COOMe or Ac), 169.0 (C=O COOMe or Ac), 169.9 (C=O COOMe or Ac); ^{13}C -GATED (125 MHz, CDCl_3): 99.8 ($J_{\text{Cl}',\text{H}1'} = 168$ Hz, C-1'), 103.1 ($J_{\text{Cl},\text{H}1} = 160$ Hz, C-1); ESI-MS: 924.4 ($\text{M}+\text{NH}_4^+$), 930.3 ($\text{M}+\text{Na}^+$); HRMS: ($\text{M}+\text{NH}_4^+$) calcd for $\text{C}_{51}\text{H}_{58}\text{O}_{15}\text{N}$ 924.38010, found 924.38025.

para-Methoxyphenyl (methyl 2,3-di-*O*-benzyl-4-*O*-(methyl (4-*O*-acetyl-2,3-di-*O*-benzyl-β-D-mannopyranosyl) uronate)-β-D-galactopyranoside uronate **13β**: Colorless oil. TLC: 30% EtOAc/PE; $[\alpha]_{\text{D}}^{22}$: -34° ($c = 0.8$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) $\delta = 1.99$ (s, 3H, CH_3 Ac), 3.24 (dd, 1H, $J_{3',4'} = 9.8$ Hz, $J_{3',2'} = 2.8$ Hz, H-3'), 3.63 (dd, 1H, $J_{3,2} = 9.6$ Hz, $J_{3,4} = 3.0$ Hz, H-3), 3.67 (d, 1H, $J_{5',4'} = 9.9$ Hz, H-5'), 3.72 (s, 3H, CH_3 COOMe), 3.77 (s, 3H, CH_3 OMe), 3.79 (s, 3H, CH_3 COOMe), 3.93 (d, 1H, $J_{2',1'} = J_{2',3'} = 2.9$ Hz, H-2'), 3.95 (dd, 1H, $J_{2,3} = 9.6$ Hz, $J_{2,1} = 7.7$ Hz, H-2), 4.15 (d, 1H, $J_{5,4} = 1.1$ Hz, H-5), 4.25 (d, 1H, $J = 12.3$ Hz, CHHPh), 4.86 (d, 1H, $J = 12.3$ Hz, CHHPh), 4.53 (dd, 1H, $J_{4,3} = 3.0$ Hz, $J_{4,5} = 1.1$ Hz, H-4), 4.62 (d, 1H, $J = 11.7$ Hz, CHHPh), 4.76 (d, 1H, $J = 11.7$ Hz, CHHPh), 4.77 (d, 1H, $J_{1',2'} = 2.9$ Hz, H-1'), 4.78 (d, 1H, $J = 10.9$ Hz, CHHPh), 4.84 (d, 1H, $J_{1,2} = 7.7$ Hz, H-1), 4.86 (d, 1H, $J = 12.6$ Hz, CHHPh), 4.95 (d, 1H, $J = 12.6$ Hz, CHHPh), 5.05 (d, 1H, $J = 10.9$ Hz, CHHPh), 5.41 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.8$ Hz, H-4'), 6.82 (d, 1H, $J = 9.1$ Hz, CH Arom pMP), 7.10 (d, 1H, $J = 9.1$ Hz, CH Arom pMP), 7.23 – 7.54 (m, 20H, CH Arom); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 20.7$ (CH_3 Ac), 52.4 (CH_3 COOMe), 52.6 (CH_3 COOMe), 55.6 (CH_3 OMe), 68.6 (C-4'), 71.1 (CH_2 Bn), 71.7 (C-2'), 73.1 (C-4), 73.3 (CH_2 Bn), 73.5 (CH_2 Bn), 73.7 (C-5'), 73.7 (C-5), 75.3 (CH_2 Bn), 78.5 (C-2), 78.6 (C-3'), 80.9 (C-3), 101.4 (C-1'), 103.2 (C-1), 114.5 (CH Arom pMP), 119.3 (CH Arom pMP), 127.3 – 128.8 (CH Arom), 137.7 (C_q Bn), 137.9 (C_q Bn), 138.3 (C_q Bn), 138.6 (C_q Bn), 151.4 (C_q pMP), 155.6 (C_q pMP), 167.1 (C=O COOMe or Ac), 167.9 (C=O COOMe or Ac), 169.6

(C=O COOMe or Ac); ^{13}C -GATED (100 MHz, CDCl_3): 101.4 ($J_{\text{Cl}',\text{Hl}'} = 160$ Hz, C-1'), 103.2 ($J_{\text{Cl},\text{Hl}} = 160$ Hz, C-1); ESI-MS: 924.5 ($\text{M}+\text{NH}_4^+$), 930.4 ($\text{M}+\text{Na}^+$); HRMS: ($\text{M}+\text{Na}^+$) calcd for $\text{C}_{51}\text{H}_{54}\text{O}_{15}\text{Na}$ 929.33549, found 929.33706.

3-Azidopropyl (methyl (4-*O*-acetyl-2,3-di-*O*-benzyl- β -D-mannopyranoside)uronate) **16**: A solution of 102 mg donor **1** (0.2 mmol 1 equiv.), 51 mg diphenyl

sulfoxide (0.25 mmol, 1.3 equiv.) and 124 mg tri-*tert*-butylpyrimidine (0.5 mmol, 2.5 equiv.) in 5 mL DCM was stirred over activated $\text{MS3\AA}$ for 30 min. The mixture was cooled to -60°C before 41 μL triflic acid anhydride (0.25 mmol, 1.3 equiv.) was added. The mixture was allowed to warm to -40°C in 15 min followed by cooling to -80°C . At that temperature 30 mg acceptor (0.3 mmol, 1.5 equiv.) in 1 mL DCM was added. Stirring was continued and the reaction mixture was allowed to warm to rT followed by addition of 2 mL pyridine and 1 mL Ac_2O . Stirring was continued for a further 10h. After addition of MeOH, the reaction mixture was diluted with EtOAc and washed with 1M HCl and NaHCO_3 (sat.). The organic layer was dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography (EtOAc/PE) and removal of the eluent afforded 80 mg of title compound **16** ($\alpha/\beta = 1/5$, 80%) as a colorless oil. TLC: 50% EtOAc/PE; $[\alpha]_{\text{D}}^{22}$: -2° ($c = 0.4$, CHCl_3); IR (neat, cm^{-1}): 1091, 1211, 1265, 1450, 1732, 2110; ^1H NMR (400 MHz, CDCl_3) $\delta = 1.84 - 1.96$ (m, 2H, CH_2 azidopropyl), 2.02 (s, 3H, CH_3 Ac), 3.39 (t, 2H, CH_2 azidopropyl), 3.48 – 3.56 (m, 2H, $J = 2.4$ Hz, H-3, CH azidopropyl), 3.72 (s, 3H, CH_3 COOMe), 3.85 (d, 1H, $J_{5,4} = 9.6$ Hz, H-5), 3.89 (d, 1H, $J_{2,1} = J_{2,3} = 2.0$ Hz, H-2), 4.01 – 4.07 (m, 1H, CH_2 azidopropyl), 4.38 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.44 (s, 1H, H-1), 4.50 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.80 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.92 (d, 1H, $J = 12.4$ Hz, CHHPh), 5.52 (t, 1H, $J_{4,3} = J_{4,5} = 9.6$ Hz, H-4), 7.21 – 7.42 (m, 10H, H Arom); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 20.8$ (CH_3 Ac), 29.0 (CH_2 azidopropyl), 48.3 (CH_2 azidopropyl), 52.6 (CH_3 COOMe), 66.9 (CH_2 azidopropyl), 68.9 (C-4), 71.5 (CH_2 Bn), 73.0 (C-2), 73.7 (C-5), 73.8 (CH_2 Bn), 78.1 (C-3), 101.5 (C-1), 127.4 – 128.3 (CH Arom), 137.7 (C_q Bn), 138.2 (C_q Bn), 167.9 (C=O Ac), 169.6 (C=O COOMe); ^{13}C -GATED (100 MHz, CDCl_3): 101.5 ($J_{\text{Cl},\text{Hl}} = 154$ Hz, C-1); ESI-MS: 536.2 ($\text{M}+\text{Na}^+$); HRMS: ($\text{M}+\text{Na}^+$) calcd for $\text{C}_{26}\text{H}_{31}\text{O}_8\text{N}_3\text{Na}$ 536.20034, found 536.20156.

3-Azidopropyl (methyl 2,3-di-*O*-benzyl-β-D-mannopyranoside) uronate **17**: To a stirred solution of 80 mg compound **16** (1.6 mmol) in MeOH was added a catalytic amount of sodium methoxide (30% wt in MeOH). The mixture was stirred for 6h before Amberlite IR120+ resin was added to acidify the solution to pH = 6. Subsequent filtration and concentration under reduced pressure afforded the crude product. Flash chromatography (EtOAc/PE) and removal of the eluent afforded 44 mg of title compound **17** (β-isomer was selectively hydrolyzed, 60%) as a colorless oil. TLC: 50% EtOAc/PE; $[\alpha]_D^{22}$: -21° ($c = 0.2$, CHCl_3); IR (neat, cm^{-1}): 1091, 1211, 1265, 1450, 1732, 2110; ^1H NMR (400 MHz, CDCl_3) $\delta = 1.89 - 2.02$ (m, 2H, CH_2 azidopropyl), 3.02 (s, 1H, OH-4), 3.39 (dd, 1H, $J_{3,4} = 9.5$ Hz, $J_{3,2} = 2.9$ Hz, H-3), 3.44 (t, 2H, $J = 6.7$ Hz, CH_2 azidopropyl), 3.57 – 3.66 (m, 1H, CH azidopropyl), 3.80 (d, 1H, $J_{5,4} = 9.6$ Hz, H-5), 3.85 (s, 3H, CH_3 COOMe), 3.92 (d, 1H, $J_{2,1} = J_{2,3} = 2.7$ Hz, H-2), 4.06 – 4.11 (m, 1H, CH azidopropyl), 4.34 (t, 1H, $J_{4,3} = J_{4,5} = 9.6$ Hz, H-4), 4.47 (s, 1H, H-1), 4.55 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.61 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.81 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.97 (d, 1H, $J = 12.4$ Hz, CHHPh), 7.30 – 7.47 (m, 10H H Arom); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 28.9$ (CH_2 azidopropyl), 48.2 (CH_2 azidopropyl), 52.5 (CH_3 COOMe), 66.8 (CH_2 azidopropyl), 68.2 (C-4), 71.7 (CH_2 Bn), 73.4 (C-2), 74.1 (CH_2 Bn), 74.9 (C-5), 80.2 (C-3), 101.9 (C-1), 127.4 – 128.3 (CH Arom), 137.7 (C_q Bn), 138.2 (C_q Bn), 169.6 (C=O COOMe); ^{13}C -GATED (100 MHz, CDCl_3): 101.9 ($J_{\text{C1,H1}} = 153$ Hz, C-1); ESI-MS: 472.3 ($\text{M}+\text{H}^+$), 494.2 ($\text{M}+\text{Na}^+$), 489.3 ($\text{M}+\text{K}^+$); HRMS: ($\text{M}+\text{Na}^+$) calcd for $\text{C}_{24}\text{H}_{29}\text{O}_7\text{N}_3\text{Na}$ 494.18977, found 494.19008.

Phenyl ((methyl 2,3-di-*O*-benzyl-4-*O*-(methyl levulinoyl)-2,3-di-*O*-benzyl-β-D-mannopyransyl) uronate)-α-D-mannopyranoside) uronate **18**: Disaccharide **18** was obtained from donor **15** and acceptor **17** according to the method A of the general procedure for glycosidations using NIS/TMSOTf giving 73 mg title compound **18** (0.078 mmol, 78%) as a colorless oil. TLC: 30% EtOAc/PE; $[\alpha]_D^{22}$: -26° ($c = 1.0$, CHCl_3); IR (neat, cm^{-1}): 1024, 1053, 1361, 1454, 1496, 1747, 2110, 2869; ^1H NMR (400 MHz, CDCl_3) $\delta = 1.83 - 1.89$ (m, 2H, CH_2 azidopropyl), 2.14 (CH_3 Lev), 2.50 (q, 2H, $J = 6.0$ Hz, CH_2 Lev), 2.67 (t, 2H, $J = 6.0$ Hz, CH_2 Lev), 3.34 (t, 1H, $J = 6.8$ Hz, CH_2 azidopropyl), 3.45 (dd, 1H, $J_{3',4'} = 9.6$ Hz, $J_{3',2'} = 2.8$ Hz, H-3'), 3.49 – 3.54 (m, 1H, CH azidopropyl), 3.54 (s, 3H, CH_3 COOMe), 3.60 (dd, 1H, $J_{3,4} = 8.8$ Hz, $J_{3,2} = 3.2$ Hz, H-3), 3.62

(s, 3H, CH₃ COOMe), 3.68 (d, 1H, $J_{5',4'} = 10.0$ Hz, H-5'), 3.82 (bs, 2H, H-2, H-2'), 3.87 (d, 1H, $J_{5,4} = 8.8$ Hz, H-5), 4.98 – 4.02 (m, 1H, CH azidopropyl), 4.39 (t, 1H, $J_{4,3} = J_{4,5} = 8.8$ Hz, H-4), 4.42 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.46 (s, 1H, H-1), 4.52 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.56 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.69 (s, 1H, H-1'), 4.70 (d, 1H, $J = 12.8$ Hz, CHHPh), 4.83 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.78 (d, 1H, $J = 12.8$ Hz, CHHPh), 4.83 (d, 1H, $J = 12.0$ Hz, CHHPh), 5.42 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.6$ Hz, H-4'), 7.22 – 7.38 (m, 20H, H Arom); ¹³C NMR (100 MHz, CDCl₃) δ = 27.8 (CH₂ Lev), 29.0 (CH₂ azidopropyl), 29.8 (CH₃ Lev), 37.7 (CH₂ Lev), 48.2 (CH₂ azidopropyl), 52.3 (CH₃ COOMe), 66.8 (CH₂ azidopropyl), 68.9 (C-4'), 71.7 (CH₂ Bn), 72.2 (CH₂ Bn), 73.3 (C-5'), 73.9 (CH₂ Bn), 73.9 (C-2 or C-2'), 74.3 (CH₂ Bn), 74.3 (C-5), 74.9 (C-2' or C-2), 77.5 (C-4), 78.5 (C-3'), 79.3 (C-3), 101.8 (C-1), 102.3 (C-1'), 127.1 – 128.3 (CH Arom), 137.8 (C_q Bn), 138.3 (C_q Bn), 138.6 (C_q Bn), 138.6 (C_q Bn), 167.7 (C=O COOMe), 168.6 (C=O COOMe), 171.6 (C=O Lev ester), 206.1 (C=O Lev ketone); ¹³C-GATED (100 MHz, CDCl₃): 101.8 ($J_{C1,H1} = 155$ Hz, C-1), 102.3 ($J_{C1',H1'} = 156$ Hz, C-1'); ESI-MS: 962.5 (M+Na⁺); HRMS: (M+Na⁺) calcd for C₅₀H₅₇O₁₅N₃Na 962.36819, found 962.36725.

Azidopropyl 3-((methyl 2,3-di-*O*-benzyl-4-*O*-(methyl (2,3-di-*O*-benzyl- β -D-mannopyransyl)uronate)- α -D-mannopyranoside) uronate **19**: Disaccharide **18** (65 mg, 0.069 mmol) was dissolved in a mixture of pyridine and acetic acid (1 mL, 4/1 v/v), after which 17 μ L hydrazine monohydrate (0.35 mmol) was added. The mixture was stirred for 30 min and then diluted with EtOAc, washed with 2M HCl and saturated NaHCO₃. The organic layer was dried (MgSO₄), filtered and concentrated under reduced pressure. Flash column chromatography (EtOAc/hexane) afforded 55 mg of the title compound **19** (0.66 mmol, 95%) as a colorless oil. TLC: 20% EtOAc/DCM; $[\alpha]_D^{22}$: +35.6° (c = 0.7, CHCl₃); IR (neat, cm⁻¹): 1026, 1053, 1361, 1454, 1496, 1747, 2094, 2869; ¹H NMR (400 MHz, CDCl₃) δ = 1.82 – 1.94 (m, 2H, CH₂ azidopropyl), 2.94 (bs, 1H, OH-4'), 3.31 (dd, 1H, $J_{3',4'} = 9.6$ Hz, $J_{3',2'} = 3.8$ Hz, H-3'), 3.35 (t, 2H, $J = 7.0$ Hz, CH₂ azidopropyl), 3.51 – 3.55 (m, 1H, CH azidopropyl), 3.59 (d, 1H, $J_{5',4'} = 9.6$ Hz, H-5'), 3.62 (s, 3H, CH₃ COOMe), 3.64 (s, 3H, CH₃ COOMe), 3.65 (dd, 1H, $J_{3,4} = 8.8$ Hz, $J_{3,2} = 2.8$ Hz, H-3), 3.83 (d, 1H, $J_{2',3'} = 2.8$ Hz, H-2'), 3.86 (d, 1H, $J_{2,3} = 2.8$ Hz, H-2), 3.90 (d, 1H, $J_{5,4} = 8.8$ Hz, H-5), 4.00 – 4.06 (m, 1H, CH azidopropyl), 4.19 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.6$ Hz, H-4'), 4.45 (t, 1H, $J_{4,3} = J_{4,5} = 8.6$ Hz, H-4), 4.48 (s, 1H, H-

1), 4.56 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.57 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.60 (d, 1H, $J = 12.4$ Hz, CHHPh), 4.68 (d, 1H, $J = 12.8$ Hz, CHHPh), 4.73 (s, 1H, H-1'), 4.74 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.76 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.78 (d, 1H, $J = 12.8$ Hz, CHHPh), 4.83 (d, 1H, $J = 12.0$ Hz, CHHPh), 7.23 – 7.38 (m, 20H, H Arom); ^{13}C NMR (100 MHz, CDCl_3) $\delta = 29.0$ (CH_2 azidopropyl), 48.2 (CH_2 azidopropyl), 52.3 (CH_3 COOMe), 52.4 (CH_3 COOMe), 66.8 (CH_2 azidopropyl), 68.1 (C-4'), 71.8 (CH_2 Bn), 72.1 (CH_2 Bn), 73.9 (CH_2 Bn), 73.9 (C-2), 74.5 (C-5), 74.6 (CH_2 Bn), 74.8 (C-5'), 75.2 (C-2'), 77.2 (C-4), 79.3 (C-5'), 80.4 (C-3'), 101.7 (C-1), 102.4 (C-1'), 127.0 – 128.4 (CH Arom), 137.9 (C_q Bn), 138.3 (C_q Bn), 138.6 (C_q Bn), 138.8 (C_q Bn), 168.6 (C=O COOMe), 169.8 (C=O COOMe); ^{13}C -GATED (100 MHz, CDCl_3): 101.7 ($J_{\text{C1,H1}} = 154$ Hz, C-1), 102.4 ($J_{\text{C1',H1'}} = 155$ Hz, C-1'); ESI-MS: 864.3 ($\text{M}+\text{Na}^+$).

3-Azidopropyl (methyl 2,3-di-O-benzyl-4-O-(methyl 2,3-di-O-benzyl-4-O-(methyl 3-azidopropyl)uronate)-β-D-mannopyranosyl)uronate)-β-D-mannopyranoside)uronate

20: Trisaccharide **20** was obtained from donor **15** and acceptor **19** according to the method A of the general procedure for glycosidations using NIS/TMSOTf giving 33 mg title compound **20** ($\alpha/\beta = 1/8$, 0.026 mmol, 70%) as a colorless oil. TLC: 30% EtOAc/PE; IR (neat, cm^{-1}): 1091, 1211, 1265, 1450, 1732, 2110; ^1H NMR (600 MHz, CDCl_3) $\delta = 1.82 - 1.90$ (m, 2H, CH_2 azidopropyl), 2.14 (s, 3H, CH_3 Lev), 2.52 (q, 2H, $J = 7.2$ Hz, $J = 6.6$ Hz, CH_2 Lev), 2.67 (t, 2H, $J = 6.6$ Hz, CH_2 Lev), 3.34 (t, 2H, CH_2 azidopropyl), 3.41 (dd, 1H, $J_{3'',4''} = 9.6$ Hz, $J_{3'',2''} = 2.4$ Hz, H-3''), 3.49 (dd, 1H, $J_{3,4} = 9.0$ Hz, $J_{3,2} = 3.3$ Hz, H-3), 3.45 (s, 4H, CH azidopropyl and CH_3 COOMe), 3.56-3.60 (m, 4H, H-3' and CH_3 COOMe), 3.65 (d, 1H, $J_{5'',4''} = 9.6$ Hz, H-5''), 3.74 (d, 1H, $J_{2,1} = J_{2,3} = 2.4$ Hz, H-2), 3.76 (d, 1H, $J_{5,4} = 9.6$ Hz, H-5), 3.79 (d, 1H, $J_{2'',1''} = J_{2'',3''} = 2.4$ Hz, H-2''), 3.81 (d, 1H, $J_{2',1'} = J_{2',3'} = 1.8$ Hz, H-2'), 3.82 (d, 1H, $J_{5',4'} = 8.4$ Hz, H-5'), 4.00 – 4.03 (m, 1H, $J = 6.1$ Hz, $J = 5.4$ Hz, CH azidopropyl), 4.30 (t, 1H, $J_{4,3} = J_{4,5} = 9.6$ Hz, H-4), 4.41 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.42 (t, 1H, $J_{4',3'} = J_{4',5'} = 9.4$ Hz, H-4'), 4.44 (s, 1H, H-1), 4.47 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.53 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.56 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.59 (s, 1H, H-1''), 4.60 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.65 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.71 (s, 1H, H-1'), 4.75 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.77 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.79 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.79 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.81 (d, 1H, $J = 12.0$ Hz, CHHPh), 4.83 (d, 1H, $J = 12.0$ Hz, CHHPh), 5.40 (t, 1H, $J_{4'',3''} = J_{4'',5''}$,

= 9.6 Hz, H-4''), 7.16 – 7.36 (m, 30H, H Arom); ^{13}C NMR (150 MHz, CDCl_3) δ = 27.8 (CH_2 Lev), 29.0 (CH_2 azidopropyl), 29.8 (CH_3 Lev), 37.8 (CH_2 Lev), 48.2 (CH_2 azidopropyl), 52.1 (CH_3 COOMe), 52.3 (CH_3 COOMe), 52.4 (CH_3 COOMe), 66.8 (CH_2 azidopropyl), 68.9 (C-4''), 71.6 (CH_2 Bn), 72.3 (CH_2 Bn), 72.5 (CH_2 Bn), 73.3 (C-5''), 73.8 (CH_2 Bn), 74.1 (C-2''), 74.3 (CH_2 Bn), 74.4 (C-5), 74.5 (C-5'), 74.5 (CH_2 Bn), 74.9 (C-2''), 75.9 (C-2), 77.1 (C-4'), 77.7 (C-4''), 78.4 (C-3''), 79.3 (C-3'), 79.6 (C-3), 101.6 (C-1), 102.4 (C-1''), 102.5 (C-1'), 127.2 – 128.4 (CH Arom), 137.8 (C_q Bn), 138.4 (C_q Bn), 138.7 (C_q Bn), 138.7 (C_q Bn), 138.8 (C_q Bn), 167.8 (C=O COOMe or Lev), 168.6 (C=O COOMe or Lev), 168.6 (C=O COOMe or Lev), 171.6 (C=O COOMe or Lev), 206.2 (C=O Lev ketone); ^{13}C -GATED (150 MHz, CDCl_3): 101.6 ($J_{\text{C1,H1}} = 155$ Hz, C-1), 102.4 ($J_{\text{C1'',H1''}} = 158$ Hz, C-1''), 102.5 ($J_{\text{C1',H1'}} = 156$ Hz, C-1'); ESI-MS: 1332.5 ($\text{M}+\text{Na}^+$); HRMS: ($\text{M}+\text{Na}^+$) calcd for $\text{C}_{71}\text{H}_{79}\text{O}_{21}\text{N}_3\text{Na}$ 1332.50983, found 1332.51029.

Trisaccharide **20** (35mg, 0.027mmol) was dissolved in 0.5 ml pyridine/AcOH (4/1) before 5 μL $\text{N}_2\text{H}_4\cdot\text{H}_2\text{O}$ (0.1mmol) was added. The mixture was allowed to stir for 10 min, diluted with EtOAc and subsequently washed with 2M HCl and sat NaHCO_3 . The organic layer is dried (MgSO_4), filtered and concentrated under reduced pressure. The crude intermediate was dissolved in 1 ml THF/ H_2O and treated with 0.15 ml KOH (0.45M) for 1h. The reaction mixture was neutralized with Amberlite IR120 to pH=3, filtered and concentrated under reduced pressure. The residual oil is dissolved in 1 mL MeOH/AcOH (10/1) before 25 mg Pd/C (10%) is added. The reaction mixture is stirred overnight under an H_2 -atmosphere and filtered. Gel filtration (HW-40, 0.15M Et_4NOAc in H_2O) of the residual afforded the desired trisaccharide **21** (5 mg, 8.3 μmol , 35% over 3 steps) as a white foam. TLC: 50% *i*PrOH/ H_2O ; ^1H NMR (400 MHz, CDCl_3) δ = 1.99-2.03 (m, 2H, CH_2 spacer), 3.14-3.21 (m, 2H, CH_2 spacer), 3.38 (s, 1H), 3.67-3.70 (m, 2H), 3.74-3.78 (m, 1H), 3.79-3.84 (m, 3H), 3.86-3.90 (m, 1H), 3.89-3.93 (m, 2H), 3.97-4.03 (m, 2H), 4.05-4.07 (m, 2H), 4.66 (s, 1H), 4.69 (s, 1H), 4.77 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ = 27.5, 38.5, 63.6, 67.8, 68.4, 69.3, 70.1, 70.6, 70.8, 71.2, 72.3, 73.4, 75.7, 76.7, 76.8, 100.7, 100.8, 101.0, 176.1, 176.4, 177.0; ESI-MS: 629.7 ($\text{M}+\text{H}^+$).

References

- 1 Crich, D.; Smith, M.; Yao, Q. Picione, J. *Synthesis*, **2001**, 323–326.
- 2 Crich, D.; Smith, M. *J. Am. Chem. Soc.* **2002**, *124*, 8867–8869.
- 3 Zuurmond, H.M.; Van der Klein, P.A.M.; Van der Marel, G.A.; Van Boom, J.H. *Tetrahedron* **1993**, *49*, 6501–6514.
- 4 Du, Y.; Zhang, M.; Kong, F. *Org. Lett.* **2000**, *2*, 3797–3800.
- 5 Van den Bos, L.J.; Codée, J.D.C.; Van der Toorn, J.C., Boltje, T.J.; Van Boom, J.H., Overkleeft, H.S.; Van der Marel, G.A. *Org. Lett.* **2004**, *6*, 2165–2168.
- 6 Van den Bos, L.J.; Litjens, R.E.J.N.; Van den Berg, R.J.B.H.N.; Overkleeft, H.S.; Van der Marel, G.A. *Org. Lett.* **2005**, *7*, 2007–2010.