Supporting Information

Stereoselective Hydrothiolation of Alkynes Catalyzed by Cesium Base: Facile Access to (Z)-1-Alkenyl
Sulfides

Azusa Kondoh, Kazuaki Takami, Hideki Yorimitsu*, and Koichiro Oshima*

*Department of Material Chemistry, Graduate School of Engineering, Kyoto University, Kyoto-daigaku
Katsura, Nishikyo-ku, Kyoto 615-8510, Japan*

Contents

Chart 1. ^1H NMR Spectrum of 1k	S2
Chart 2. ^{13}C NMR Spectrum of 1k	S3
Chart 3. ^1H NMR Spectrum of 1m	S4
Chart 4. ^{13}C NMR Spectrum of 1m	S5
Chart 5. ^1H NMR Spectrum of 1n	S6
Chart 6. ^{13}C NMR Spectrum of 1n	S7
Chart 7. ^1H NMR Spectrum of 1o	S8
Chart 8. ^{13}C NMR Spectrum of 1o	S9

Chart 1. ^1H NMR Spectrum of **1k**

Chart 2. ^{13}C NMR Spectrum of **1k**

Chart 3. ^1H NMR Spectrum of **1m**

Chart 4. ^{13}C NMR Spectrum of **1m**

1m

Chart 5. ^1H NMR Spectrum of **1n**

1n
1.1
1.5
3.9
6.5-7.5
7.5-8.5

Chart 6. ^{13}C NMR Spectrum of **1n**

1n
S
1

Chart 7. ^1H NMR Spectrum of **1o**

Chart 8. ^{13}C NMR Spectrum of **1o**