Supporting information for:**Stereoselective *N*-Glycosylation By Staudinger Ligation**

Yi He, Ronald J. Hinklin, Jiyoung Chang and Laura L. Kiessling

General Procedures and Materials:

All moisture- and oxygen-sensitive reactions were carried out in flame-dried glassware under nitrogen atmosphere unless otherwise specified. All purchased materials were used without further purification, unless otherwise noted. Methylene chloride was distilled from calcium hydride, methanol was distilled from magnesium and pyridine was distilled from sodium and anthracene. Tetrahydrofuran and toluene were distilled over sodium and benzophenone.

The glycosyl azides employed were synthesized using known procedures.^{1, 2} Diethylphosphine and diphenylphosphine were purchased from Strem Chemicals. Dibutylphosphine was synthesized by reduction of tributylphosphine oxide,³ and the crude product was used in subsequent reactions.

Analytical thin layer chromatography (TLC) was carried out on E. Merck (Darmstadt) TLC plates pre-coated with silica gel 60 F₂₅₄ (250 Å layer thickness). Analyte visualization was accomplished using an UV lamp and/or by charring with at least one of the following solutions: a *p*-anisaldehyde stain (18 mL *p*-anisaldehyde, 7.5 mL glacial acetic acid, 25 mL concentrated H₂SO₄, 675 mL absolute EtOH), ninhydrin stain (200 mg ninhydrin, 95 mL *n*-butanol, 5 mL 10% AcOH), or potassium permanganate stain (3 g KMnO₄, 20 g K₂CO₃, 5 mL 5% aqueous NaOH, 300 mL H₂O). Flash chromatography was performed on Scientific Adsorbents Incorporated silica gel (32-63 μM, 60 Å pore size) using distilled reagent grade hexanes and ACS grade ethyl acetate or methanol and chloroform. The term, “concentrated under reduced pressure” refers to the removal of solvents and other volatile materials using a rotary evaporator at water aspirator pressure (< 20 torr) while maintaining the water bath temperature below

40 °C, and then removing residual solvent at high vacuum (< 0.1 torr). The term, “high vacuum,” refers to vacuum achieved by a mechanical belt-drive oil pump.

^1H , ^{13}C , and ^{31}P nuclear magnetic resonance (NMR) spectra were recorded on Bruker AC-300 and chemical shifts are reported relative to tetramethylsilane or residual solvent peaks in parts per million (CHCl_3 : ^1H : δ 7.26, ^{13}C : δ 77.0; ^{31}P chemical shifts are relative to an external reference standard of phosphoric acid 0.0). ^1H NMR data are assumed to be first order. Peak multiplicity is reported as: singlet (s), doublet (d), triplet (t), quartet (q), multiplet (m), and broad (br). Mass spectrometry was performed on a Micromass LCT instrument.

Synthetic Procedures:

Diethylphosphino(borane)methane alcohol (**2a**)

Borane-dimethylsulfide (11 M solution in tetrahydrofuran, 2.5 mL, 27.5 mmol) was added to a round-bottom flask with freshly distilled tetrahydrofuran (10 mL). To this solution, diethylphosphine **1a** (2.0 mL, 17.5 mmol) was added in one portion. The solution was stirred under nitrogen for 2 h, and was quenched slowly with ice (~10 g). The resulting solution was diluted with brine (20 mL), and extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with brine (2 x 10 mL), dried over sodium sulfate, filtered, and the solvent was removed under reduced pressure. The crude phosphine-borane was used without purification. The crude phosphine-borane was dissolved in a mixture of aqueous formaldehyde (37% w/w, 5 mL) and tetrahydrofuran (10 mL). To this solution, potassium hydroxide (1.0 g, 17.9 mmol) was added. The solution was stirred for 4 h, and the volatile solvent was removed under reduced pressure. The resulting aqueous solution was extracted with ethyl acetate (3 x 10 mL), washed with brine (1 x 10 mL), dried over sodium sulfate, filtered, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield **2a** as a colorless liquid (2.3 g, 17.5 mmol, 100%). ^1H

NMR (CDCl₃) δ 3.96 (s, 2H), 1.76-1.63 (m, 4H), 1.18-1.10 (m, 6H), 0.77- -0.13 (m, 3H); ¹³C NMR (CDCl₃) δ 55.7 (d, J =40.2 Hz), 13.0 (d, J =35.3 Hz), 6.3 (d, J =2.3 Hz); ³¹P NMR (CDCl₃) δ 23.0 (q, J =56.7 Hz).

Dibutylphosphino(borane)methane alcohol (2b)

Borane-dimethylsulfide (11 M solution in tetrahydrofuran, 2.0 mL, 22.0 mmol) was added to a round-bottom flask with freshly distilled tetrahydrofuran (10 mL). To this solution, crude dibutylphosphine **1b** (2.6 g, 17.9 mmol) was added in one portion. The solution was stirred under nitrogen for 2 h, and was quenched slowly with ice (~10 g). The resulting solution was diluted with brine (20 mL), and extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with brine (2 x 10 mL), dried over sodium sulfate, filtered, and the solvent was removed under reduced pressure. The crude phosphine-borane was used without purification. The crude phosphine-borane was dissolved in a mixture of aqueous formaldehyde (37% w/w, 5 mL) and tetrahydrofuran (10 mL). To this solution, potassium hydroxide (1.0 g, 17.9 mmol) was added. The solution was stirred for 4 h, and the volatile solvent was removed under reduced pressure. The resulting aqueous solution was extracted with ethyl acetate (3 x 10 mL), washed with brine (1 x 10 mL), dried over sodium sulfate, filtered, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield **2b** as a colorless liquid (1.7 g, 9.3 mmol, 51%). ¹H NMR (CDCl₃) δ 3.94 (s, 2H), 2.22 (s, 1H), 1.64-1.43 (m, 12H), 0.91 (t, J =7.1 Hz, 6H); ¹³C NMR (CDCl₃) δ 57.0 (d, J =40.8 Hz), 24.5, 24.2 (d, J =12.6 Hz), 20.3 (d, J =34.2 Hz), 13.4; ³¹P NMR (CDCl₃) δ 18.7 (q, J =54.3 Hz); ESI (m/z): [M+Na]⁺ calcd for C₆H₁₈BO₃PSNa, 235.1; found 235.0.

Diphenylphosphino(borane)methane alcohol (2c)

Borane-dimethylsulfide (11 M solution in tetrahydrofuran, 5.0 mL, 55.5 mmol) was added to a round-bottom flask with freshly distilled tetrahydrofuran (20 mL). To this solution, diphenylphosphine **1c** (5 mL, 28.7 mmol) was added in one portion. The solution was stirred under nitrogen for 2 h, and was quenched slowly with ice (~10 g). The resulting solution was diluted with brine (20 mL), and extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with brine (2 x 10 mL), dried over sodium sulfate, filtered, and the solvent was removed under reduced pressure to yield pure diphenylphosphine-borane (5.7 g, 100% yield). The crude phosphine-borane was used without purification. Part of the phosphine-borane (1.3 g, 6.5 mmol) was dissolved in a mixture of aqueous formaldehyde (37% w/w, 5 mL) and tetrahydrofuran (10 mL). To this solution, potassium hydroxide (750 mg, 13.4 mmol) was added. The solution was stirred for 2 h, and the volatile solvent was removed under reduced pressure. The resulting aqueous solution was extracted with ethyl acetate (3 x 10 mL), washed with brine (1 x mL), dried over sodium sulfate, filtered, and the solvent was removed under reduced pressure to yield pure alcohol **2c** as a colorless liquid (1.6 g, 28.7 mmol, 100%). ^1H NMR (CDCl_3) δ 7.78-7.43 (m, 10H), 4.42 (s, 3H), 2.76 (br, 1H), 1.46-0.52 (m, 3H); ^{13}C NMR (CDCl_3) δ 132.6 (d, $J=8.3$ Hz), 131.5, 128.8 (d, $J=9.3$ Hz), 126.6 (d, $J=54.6$ Hz), 60.0 (d, $J=42.0$ Hz); ^{31}P NMR (CDCl_3) δ 27.7 (d, $J=39.0$ Hz); ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{13}\text{H}_{16}\text{BONa}$ 253.1; found 253.1.

Diethylphosphino(borane)methyl methanesulfonate

Alcohol **2a** (120.0 g, 0.9 mmol) was dissolved in freshly distilled methylene chloride (1 mL), followed by addition of triethylamine (260.0 μL , 1.9 mmol). The solution was cooled to 0 $^{\circ}\text{C}$ with ice-water bath. To this solution, methanesulfonyl chloride (85.0 μL , 1.1 mmol) was added dropwise. The solution was stirred under nitrogen for 15 h. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield the mesylate as a slightly yellow liquid (130.0 mg, 0.6 mmol, 67%). ^1H NMR (CDCl_3) δ 4.46 (d, $J=1.8$ Hz, 2H), 3.07 (s,

3H), 1.84-1.49 (m, 4H), 1.22-1.11 (m, 6 H), 0.75- -0.13 (m, 3H); ^{13}C NMR (CDCl_3) δ 61.5 (d, $J=30.0$ Hz), 37.3, 14.0 (d, $J=35.2$ Hz), 6.5; ^{31}P NMR (CDCl_3) δ 26.1 (q, $J=42.8$ Hz); ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_6\text{H}_{18}\text{BO}_3\text{PSNa}$, 235.1; found 235.0.

Dibutylphosphino(borane)methyl methanesulfonate

Alcohol **2b** (1.2 g, 6.1 mmol) was dissolved in freshly distilled methylene chloride (10 mL), followed by addition of triethylamine (1.0 mL, 7.2 mmol). The solution was cooled to 0 °C with ice-water bath. To this solution, methanesulfonyl chloride (570.0 μL , 7.4 mmol) was added dropwise. The solution was stirred under nitrogen for 15 h. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield the mesylate as a colorless liquid (1.7 g, 6.1 mmol, 100%). ^1H NMR (CDCl_3) δ 4.42 (br s, 2H), 3.05 (s, 3H), 1.99-1.79 (m, 4H), 1.59-1.55 (m, 8H), 0.89 (t, $J=6.7$ Hz, 6 H), 0.68- -0.13 (m, 3H); ^{13}C NMR (CDCl_3) δ 62.2 (d, $J=30.2$ Hz), 37.1, 24.2, 23.9 (d, $J=13.1$ Hz), 20.5 (d, $J=34.3$ Hz), 13.3; ^{31}P NMR (CDCl_3) δ 21.5 (d, $J=56.4$ Hz); ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{11}\text{H}_{26}\text{BO}_3\text{PSNa}$, 293.1; found 293.1.

Diphenylphosphino(borane)methyl methanesulfonate

Alcohol **2c** (1.1 g, 5.0 mmol) was dissolved in freshly distilled methylene chloride (10 mL), followed by addition of triethylamine (1.4 mL, 10.1 mmol). The solution was cooled to 0 °C with ice-water bath. To this solution, methanesulfonyl chloride (580.0 μL , 7.4 mmol) was added dropwise. The solution was stirred under nitrogen for 15 h. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield the mesylate as a white solid (1.2 g, 3.9 mmol, 78%). ^1H NMR (CDCl_3) δ 7.45-7.77 (m, 10H), 4.92 (d, $J=2.1$ Hz, 2H), 2.86 (s,

3H), 1.54-0.87 (m, 3H); ^{13}C NMR (CDCl_3) δ 132.9 (d, $J=9.3$ Hz), 132.3 (d, $J=3$ Hz), 129.1 (d, $J=10.9$ Hz), 125.0 (d, $J=57.4$ Hz), 64.5 (d, $J=38.7$ Hz), 37.4; ^{31}P NMR (CDCl_3) δ 21.9 (d, $J=59.2$ Hz); ESI (m/z): $[2\text{M}+\text{Na}+\text{MeOH}]^+$ calcd for $\text{C}_{29}\text{H}_{40}\text{B}_2\text{O}_7\text{P}_2\text{S}_2\text{Na}$ 671.2; found 671.1.

Diethylphosphino(borane)methanethiol acetate (**3a**)

The mesylate (228.0 mg, 1.0 mmol) was dissolved in anhydrous N,N' -dimethylformamide (2 mL), followed by addition of thiolacetic acid (85.0 μL , 1.2 mmol). To this solution cesium carbonate (340.0 mg, 1.0 mmol) was added portionwise. The color of the solution darkened. The solution was stirred under nitrogen for 12 h. The resulting dark solution was diluted with water (10 mL), extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with brine (3 x 5 mL), dried over magnesium sulfate, filtered, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (9:1 hexanes:ethyl acetate) to yield thioacetate **3a** as a pale yellow liquid (174.0, 0.9 mmol, mg, 87%). ^1H NMR (CDCl_3) δ 6.31 (d, $J=6.0$ Hz, 2H), 2.40 (s, 3H), 1.66 (dq, $J=10.2$, 7.6 Hz, 4H), 1.14 (dt, $J=16.1$, 7.9 Hz, 6H), 0.86-0.04 (m, 3H); ^{13}C NMR (CDCl_3) δ 193.4, 30.2, 20.3 (d, $J=112.3$ Hz), 15.5 (d, $J=141.0$ Hz), 3.7 (d, $J=9.6$ Hz); ^{31}P NMR (CDCl_3) δ 25.9 (q, $J=47.6$ Hz); ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_7\text{H}_{18}\text{BOPSNa}$, 215.1; found 215.0.

Dibutylphosphino(borane)methanethiol acetate (**3b**)

The mesylate (976.0 mg, 3.6 mmol) was dissolved in anhydrous N,N' -dimethylformamide (10 mL), followed by addition of thiolacetic acid (520.0 μL , 7.3 mmol). The solution was cooled to -40 $^{\circ}\text{C}$ with dry ice-isopropanol bath. To this solution cesium carbonate (2.4 g, 7.3 mmol) was added portionwise. The color of the solution

darkened. The solution was stirred under nitrogen, while the cold bath warmed up to room temperature over 8 h. The resulting dark solution was diluted with water (40 mL), extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with brine (3 x 10 mL), dried over magnesium sulfate, filtered, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (9:1 hexanes:ethyl acetate) to yield thioacetate **3b** as a colorless liquid (697.0 mg, 2.8 mmol, 77%). ^1H NMR (CDCl_3) δ 6.21 (d, $J=5.8$, 2H), 2.36 (s, 3H), 1.61-1.19 (m, 12H), 0.87 (t, $J=7.1$ Hz, 6H), 0.66- -0.05 (m, 3H); ^{13}C NMR (CDCl_3) δ 193.2, 30.0, 24.4, 24.1 (d, $J=12.7$ Hz), 22.3 (d, $J=34.4$ Hz), 21.0 (d, $J=28.4$ Hz), 13.3; ^{31}P NMR (CDCl_3) δ 21.5 (d, $J=72.6$ Hz); ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{11}\text{H}_{26}\text{BOPSNa}$, 271.1; found 271.1.

Diphenylphosphino(borane)methanethiol acetate (**3c**)

The mesylate (872.0 mg, 2.8 mmol) was dissolved in anhydrous N,N' -dimethylformamide (5 mL), followed by addition of thiolacetic acid (306.0 μL , 4.3 mmol). The solution was cooled to $-20\text{ }^\circ\text{C}$ with dry ice-isopropanol bath. To this solution cesium carbonate (1.8 g, 5.5 mmol) was added portionwise. The color of the solution darkened. The solution was stirred under nitrogen, while the cold bath warmed up to room temperature over 8 h. The resulting dark solution was diluted with water (20 mL), extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed with brine (3 x 10 mL), dried over magnesium sulfate, filtered, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (4:1 hexanes:ethyl acetate) to yield thioacetate **3c** as a white solid (531.0 mg, 1.8 mmol, 65%). ^1H NMR (CDCl_3) δ 7.44-7.75 (m, 10H), 3.72 (d, $J=7.2$ Hz, 2H), 2.23 (s, 3H), 1.54-0.87 (m, 3H); ^{13}C NMR (CDCl_3) δ 193.2, 132.2 (d, $J=9.2$ Hz), 131.9 (d, $J=1.5$ Hz), 128.6 (d, $J=10.3$ Hz), 127.3 (d, $J=56.0$ Hz), 29.8, 23.5 (d, $J=35.6$ Hz); ^{31}P NMR (CDCl_3) δ 22.4 (d, $J=69.3$ Hz); ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{15}\text{H}_{18}\text{BOPSNa}$, 311.1; found 311.1.

Boc-*N*-β-diethylphosphino(borane)methanethiol-*L*-asparagine benzyl ester (4a)

Thioacetate **3a** (410.0 mg, 2.1 mmol) was dissolved in freshly distilled methanol (5 mL), followed by addition of sodium methoxide (1 N solution in methanol, 2.5 mL). After stirring under nitrogen for 10 min, the solution was neutralized by aqueous phosphate buffer (pH = 6.5), and was extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with brine (1 x 10 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. The crude thiol was dissolved in freshly distilled methylene chloride (10 mL), followed by addition of Boc-Asp-OBz (760 mg, 2.35 mmol), 1,3-dicyclohexylcarbodiimide (485.0 mg, 2.4 mmol), and cat. *N,N'*-dimethyl-4-aminopyridine. The solution was stirred under nitrogen for 4 h, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield the thioester **4a** as a colorless oil (946.0 mg, 2.1 mmol, 98%). ¹H NMR (CDCl₃) δ 7.31 (m, 5H), 5.44 (d, *J*=7.5 Hz, 1H), 5.14 (s, 2H), 4.57 (m, 1H), 3.20 (br s, 2H), 3.08 (d, *J*=6 Hz), 1.59 (qt, *J*=9.2, 6.9 Hz, 4H), 1.40 (s, 9H), 1.07 (dt, *J*=16.2, 8.4 Hz, 6H), 0.51–0.26 (m, 3H); ¹³C NMR (CDCl₃) δ 234.9, 194.7, 170.2, 134.9, 128.5, 128.4, 129.2, 80.2, 67.6, 50.5, 28.2, 20.1 (d, *J*=28.0 Hz), 15.3 (d, *J*=35.6 Hz), 6.6 (d, *J*=2.4 Hz); ³¹P NMR (CDCl₃) δ 26.2 (d, *J*=142.8 Hz); ESI (m/z): [M+Na]⁺ calcd for C₂₁H₃₅BNO₅PSNa, 478.2; found 478.1.

Boc-*N*-β-dibutylphosphino(borane)methanethiol-*L*-asparagine benzyl ester (4b)

Thioacetate **3b** (653.0 mg, 2.6 mmol) was dissolved in freshly distilled methanol (5 mL), followed by addition of sodium methoxide (1 N solution in methanol, 3.0 mL). After stirring under nitrogen for 10 min, the solution was neutralized by aqueous phosphate

buffer (pH = 6.5), and was extracted with ethyl acetate (3 x 10 mL). The combined organic layers were washed with brine (1 x 10 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. The crude thiol was dissolved in freshly distilled methylene chloride (10 mL), followed by addition of Boc-Asp-OBz (940.0 mg, 2.9 mmol), 1,3-dicyclohexylcarbodiimide (600.0 mg, 2.9 mmol), and cat. *N,N'*-dimethyl-4-aminopyridine. The solution was stirred under nitrogen for 4 h, and the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield thioester **4b** as a colorless oil (1.4 g, 2.6 mmol, 100%). ¹H NMR (CDCl₃) δ 7.40-7.30 (m, 5H), 5.41 (d, *J*=7.6 Hz), 5.18 (s, 3H), 4.63-4.57 (m, 1H), 3.30-3.10 (m, 3H), 1.60-1.37 (m, 21H), 0.92 (t, *J*=7.2 Hz, 6H), 0.1 (br s, 3H); ¹³C NMR (CDCl₃) δ 194.4, 170.0, 154.8, 134.8, 128.2, 128.14, 127.9, 79.9, 67.2, 53.3, 50.2, 44.8, 34.6, 27.9, 24.7, 23.9 (d, *J*=13.1 Hz), 22.0 (d, *J*=33.3 Hz), 20.7 (d, *J*=27.0 Hz), 13.2; ³¹P NMR (CDCl₃) δ 21.8; ESI (m/z): [M+Na]⁺ calcd for C₂₅H₄₃BNO₅PSNa, 534.3; found 534.1.

Boc-*N*-β-(diphenylphosphino(borane))methanethiol-*L*-asparagine benzyl ester (4c)

Thioacetate **3c** (1.8 g, 6.3 mmol) was dissolved in freshly distilled methanol (30 mL), followed by addition of sodium methoxide (439.0 mg, 8.1 mL). After stirring under nitrogen for 10 min, the solution was neutralized by 1 N hydrochloric acid, and was extracted with ethyl acetate (2 x 30 mL). The combined organic layers were washed by brine (1 x 20 mL), dried over sodium sulfate, filtered, and concentrated under reduced pressure. The crude thiol was dissolved in freshly distilled methylene chloride (40 mL), followed by addition of Boc-Asp-OBz (2.0 g, 6.3 mmol), 1,3-dicyclohexylcarbodiimide (206.0 mg, 6.9 mmol), and cat. *N,N'*-dimethyl-4-aminopyridine. The solution was stirred under nitrogen for 4 h, and the solution was filtered through a pad of celite. The solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield the thioester **4c** as a colorless oil (3.3

g, 5.8 mmol, 93%). ^1H NMR (CDCl_3) δ 7.73-7.27 (m, 15H), 5.25-5.13 (m, 2H), 4.55-4.49 (m, 1H), 3.69 (qd, $J=14.3$, 7.3 Hz, 2H), 3.04 (qd, $J=16.5$, 5.1 Hz, 2H), 1.44 (s, 3H), 1.54-0.87 (m, 3H); ^{13}C NMR (CDCl_3) δ 194.6, 170.2, 155.0, 35.0, 132.4 (d, $J=15.9$ Hz), 132.4 (d, $J=2.3$ Hz), 131.8, 128.9 (d, $J=3.1$ Hz), 128.8 (d, $J=2.9$ Hz), 128.5, 128.3, 128.0, 127.8, 127.6, 127.0 (d, $J=13.5$ Hz), 80.2, 67.5, 50.4, 44.9, 28.2, 23.6 (d, $J=34.2$ Hz); ^{31}P NMR (CDCl_3) δ 22.7; ESI (m/z): $[\text{M}+\text{Na}]^+$ $\text{C}_{29}\text{H}_{35}\text{BNO}_5\text{PSNa}$, 574.1; found 574.1.

Boc-*N*- β -diethylphosphinomethanethiol-*L*-asparagine benzyl ester (**4a**)

Phosphine-borane **4a** (0.10 g, 0.22 mmol) and 1,4-diazabicyclo[2.2.2]octane (DABCO) (50.00 mg, 0.44 mmol) was dissolved in freshly distilled toluene (1 mL). The solution was submerged in an oil-bath pre-heated to 80 °C. The solution was stirred under argon for 1 h. After cooling to room temperature, the solvent was removed under reduced pressure. The crude product was suspended in minimum amount of hexanes and ethyl acetate mixture (10:1 hexanes:ethyl acetate). The suspension was passed through a short column of silica gel under nitrogen. The column was washed with the same solvent mixture until no addition product was observed in the eluent. Removal of the solvent under reduced pressure yielded pure **5a** as a colorless oil (92.00 mg, 0.21 mmol, 95%). ^1H NMR (CDCl_3) δ 7.35-7.30 (m, 5H), 5.44 (d, $J=8.5$ Hz, 1H), 1.92 (s, 2H), 4.65-4.55 (m, 1H), 3.31-3.11 (m, 2H), 3.01 (d, $J=4.5$ Hz, 2H), 1.58-1.36 (m, 13H), 1.05 (dt, $J=15.5$, 7.6 Hz, 6H); ^{13}C (CDCl_3) δ 196.5, 170.6, 155.2, 135.2, 128.6, 128.4, 128.2, 80.2, 67.5, 50.6, 45.2, 29.7, 28.3, 23.0 (d, $J=26.9$ Hz), 18.1 (d, $J=9.4$ Hz), 19.0 (d, $J=13.4$ Hz); ^{31}P NMR (CDCl_3) δ -18.5.

Boc-*N*- β -dibutylphosphinomethanethiol-*L*-asparagine benzyl ester (**5b**)

Phosphine-borane **4b** (0.18 mg, 0.35 mmol) and 1,4-diazabicyclo[2.2.2]octane (DABCO) (80.00 mg, 0.71 mmol) was dissolved in freshly distilled toluene (1 mL). The solution was submerged in an oil-bath pre-heated to 80 °C. The solution was stirred under argon for 30 min. After cooling to room temperature, the solvent was removed under reduced pressure. The crude product was suspended in minimum amount of hexanes and ethyl acetate mixture (10:1 hexanes:ethyl acetate). The suspension was passed through a short column of silica gel under nitrogen. The column was washed with the same solvent mixture until no additional product was observed in the eluent. Removal of the solvent under reduced pressure yielded pure **5b** as a colorless oil (0.18 mg, 0.35 mmol, 100%). ¹H NMR (CDCl₃) δ 7.35 (br s, 5H), 5.45 (d, *J*=8.5 Hz, 1H), 5.18 (s, 2H), 4.59 (dt, *J*=8.1, 4.9 Hz, 1H), 3.26 (qd, *J*=16.1 Hz, 4.9 Hz, 2H), 3.01 (qd, *J*=15.0, 5.0 Hz, 2H), 1.44-1.38 (m, 21H), 0.91 (t, *J*=6.3 Hz, 6H); ¹³C (CDCl₃) δ 196.5, 170.6, 155.2, 135.1, 128.5, 128.3, 128.2, 80.1, 67.4, 50.5, 45.1, 28.2, 28.8 (d, *J*=13.5 Hz), 25.8 (d, *J*=11.5 Hz), 24.3 (d, *J*=11.4), 23.9 (d, *J*=29.7 Hz), 13.7; ³¹P NMR (CDCl₃) δ -26.8; ESI (m/z): [M+H]⁺ calcd for C₂₅H₄₀O₃PS, 498.2; found 498.1.

Boc-*N*-β-diphenylphosphinomethanethiol-*L*-asparagine benzyl ester (**5c**)

Phosphine-borane **4c** (0.17 mg, 0.31 mmol) and 1,4-diazabicyclo[2.2.2]octane (DABCO) (71.00 mg, 0.62 mmol) was dissolved in freshly distilled toluene (1 mL). The solution was submerged in an oil-bath pre-heated to 60 °C. The solution was stirred under argon for 1 h. After cooling to room temperature, the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) under nitrogen to yield **5c** as a colorless oil (0.15 mg, 0.28 mmol, 91%). ¹H NMR (CDCl₃) δ 7.45-7.35 (m, 15H), 5.44 (d, *J*=9.7 Hz, 1H), 5.18 (s, 2H), 4.62-4.56 (m, 1H), 3.52 (qd, *J*=13.5, 4.2 Hz, 2H), 3.22 (qd, *J*=16.5, 4.8 Hz, 2H), 1.46 (s, 9H); ¹³C (CDCl₃) δ 195.7, 170.4, 155.1, 136.4 (d, *J*=13.7 Hz), 135.1, 132.7 (d, *J*=1.3 Hz), 132.5 (d, *J*=1.3 Hz), 129.1, 128.5, 128.4, 128.3, 128.2, 80.0, 67.4, 50.4, 45.0, 29.2, 25.6 (d,

$J=24.0$ Hz); ^{31}P NMR (CDCl_3) δ -11.8 ; ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd $\text{C}_{29}\text{H}_{32}\text{NO}_5\text{PSNa}$, 560.2; found 560.2.

**Boc-*N*- β -2,3,4,6-tetra-*O*-benzyl- β -*D*-glucopyranosyl-*L*-asparagine benzyl ester
(ligation product from glycosyl azide 6)**

Glycosyl azide **6** (93 mg, 165 μmol) was dried by azeotropic removal of water with freshly distilled toluene (3 x) in a round-bottom flask, and the flask was purged with argon. Anhydrous *N,N'*-dimethylformamide (2 mL) was added the flask. The resulting solution of the **6** was transferred via cannula under argon to another flask, previously purged with argon, containing freshly prepared diethylphosphinothioester **5a** (44 mg, 100 μmol). The solution was stirred under argon for 12 h, and water (200 μL) was added. The solution was stirred for additional 30 min, and the solvent was removed under high vacuum. The crude product was purified by flash chromatography (3:1 hexanes:ethyl acetate) to yield a white solid (46 mg, 54 μmol , 54%). The ligations of **6** with phosphinothioesters **5b**, **5c** were carried out in a similar fashion. ^1H NMR (CDCl_3) δ 7.36-7.23 (m, 23H), 7.13-7.09 (m, 2H), 5.68 (d, $J=9.4$ Hz, 1H), 5.35 (d, $J=8.5$, 1H), 5.15 (s, 2H), 4.96 (t, $J=8.8$ Hz, 1H), 4.91 (s, 2H), 4.79-4.60 (AB, $J_{\text{AB}}=11.9$, 2H), 4.5 (m, 1H), 4.81-4.47 (AB, $J_{\text{AB}}=10.8$, 2H), 4.63-4.41 (AB, $J_{\text{AB}}=12.1$, 2H), 3.74-3.63 (m, 4H), 3.44 (m, 1H), 3.32-3.26 (m, 1H), 2.76-2.39 (ABX, $J_{\text{AB}}=17.3$, $J_{\text{AX}}=4.2$, $J_{\text{BX}}=3.8$ Hz, 2H), 1.4 (s, 9H); ^{13}C NMR (CDCl_3) δ 171.1, 170.2, 155.6, 138.2, 137.9, 137.7, 137.6, 135.4, 129.0, 128.7, 128.4, 128.3, 128.2, 128.1, 128.0, 127.9, 127.8, 127.5, 127.4, 86.1, 79.9, 78.9, 78.6, 77.4, 76.2, 75.7, 74.9, 74.4, 73.4, 67.9, 67.2, 49.9, 37.7, 28.2; ESI (m/z): $[\text{M}+\text{Na}]^+$ calcd for $\text{C}_{50}\text{H}_{56}\text{N}_2\text{O}_{10}\text{Na}$, 867.4; found 867.3.

Boc-*N*-β-hepta-O-benzyl-β-*D*-lactosyl-*L*-asparagine benzyl ester (ligation product from glycosyl azide **7)**

Glycosyl azide **7** (188 mg, 188 μmol) was dried by azeotropic removal of water with freshly distilled toluene (3 x) in a round-bottom flask, and the flask was purged with argon. Anhydrous *N,N*-dimethylformamide (2 mL) was added the flask. The resulting solution of the **7** was transferred via cannula under argon to another flask, previously purged with argon, containing freshly prepared diethylphosphinothioester **5a** (66 mg, 150 μmol). The solution was stirred under argon for 12 h, and water (200 μL) was added. The solution was stirred for additional 30 min, and the solvent was removed under high vacuum. The crude product was purified by flash chromatography (2:1 hexanes:ethyl acetate) to yield a white solid (98 mg, 77 μmol, 51%). The ligations of **7** with phosphinothioesters **5b**, **5c** were carried out in a similar fashion. ¹H NMR (CDCl₃) δ 7.40-7.20 (m, 40H), 5.67 (d, *J*=9.0 Hz, 1H), 5.29 (d, *J*=9.0 Hz, 1H), 5.15 (s, 2H), 5.13 (d, *J*=10.3 Hz, 1H), 4.98 (d, *J*=11.4 Hz, 1H), 4.92 (t, *J*=9.1 Hz, 1H), 4.77 (t, *J*=10.3 Hz, 2H), 4.72-4.68 (m, 3H), 4.62 (d, *J*=11.9 Hz, 1H), 4.57-4.44 (m, 3H), 4.39 (d, *J*=11.8 Hz, 1H), 4.30-4.24 (m, 3H), 4.00 (t, *J*=9.6 Hz, 1H), 3.88 (d, *J*=2.6 Hz, 1H), 3.82 (dd, *J*=10.7, 2.0 Hz, 1H), 3.73 (dd, *J*=9.8, 7.8 Hz, 1H), 3.63 (t, *J*=9.0 Hz, 1H), 3.55-3.49 (m, 2H), 3.40 (t, *J*=5.2 Hz, 1H), 3.37-3.26 (m, 3H), 3.21 (t, *J*=9.0 Hz, 1H), 2.73-2.35 (ABX, *J*_{AB}=16.8, *J*_{AX}=4.8, *J*_{BX}=3.5 Hz, 2H), 1.39 (s, 9H); ¹³C NMR (CDCl₃) δ 171.1, 170.2, 139.0, 138.9, 138.6, 138.5, 138.2, 138.0, 137.9, 135.5, 129.0, 128.7, 128.5, 128.4, 128.3, 128.2, 128.1, 128.0, 127.9, 127.8, 127.7, 127.6, 127.5, 127.3, 102.6, 84.3, 82.3, 79.8, 78.6, 78.1, 76.5, 75.9, 75.5, 75.3, 74.3, 73.7, 73.4, 73.0, 72.5, 68.2, 67.5, 67.3, 49.9, 37.7, 28.2; ESI (*m/z*): [M+Na]⁺ calcd for C₇₇H₈₄N₂O₁₅Na, 1299.5; found 1299.5.

Boc-*N*-β-2,3,4,6-tetra-O-acetyl-β-*D*-glucopyranosyl-*L*-asparagine benzyl ester
(ligation product from glycosyl azide **8, **10**)**

Glycosyl azide **8** (64 mg, 171 μmol) was dried by azeotropic removal of water with freshly distilled toluene (3 x) in a round-bottom flask, and the flask was purged with argon. Anhydrous *N,N'*-dimethylformamide (2 mL) was added the flask. The resulting solution of the **7** was transferred via cannula under argon to another flask, previously purged with argon, containing freshly prepared diethylphosphinothioester **5a** (63 mg, 143 μmol). The solution was stirred under argon for 12 h, and water (200 μL) was added. The solution was stirred for additional 30 min, and the solvent was removed under high vacuum. The crude product was purified by flash chromatography (1:1 hexanes:ethyl acetate) to yield a white solid (40 mg, 61 μmol, 43%). The ligations of **8** and **10** with phosphinothioesters **5b**, **5c** were carried out in a similar fashion. The product was purified by flash chromatography (1:1 hexanes:ethyl acetate) to yield a white solid. ¹H NMR (CDCl₃) δ 7.40-7.33 (m, 4H), 6.38 (d, *J*=9.1 Hz), 5.70 (d, *J*=8.0 Hz), 5.31 (t, *J*=9.4 Hz, 1H), 5.24-5.18 (m, 2H), 5.08 (t, *J*=9.7 Hz, 1H), 4.91 (t, *J*=9.6 Hz, 1H), 4.61-4.58 (m, 1H), 4.20 (ABX, *J*_{AB}=12.5, *J*_{AX}=4.0, *J*_{BX}=1.9 Hz, 2H), 3.82-3.77 (m, 1H), 2.81 (ABX, *J*_{AB}=16.5 *J*_{AX}=3.9, *J*_{BX}=4.1 Hz, 2H), 2.09 (s, 3H), 2.05 (s, 3H), 2.03 (s, 3H), 2.00 (s, 3H), 1.43 (s, 9H); ¹³C NMR (CDCl₃) δ 171.1, 170.6, 170.5, 169.8, 169.5, 155.5, 135.3, 128.5, 128.2, 127.9, 80.1, 78.1, 77.2, 73.6, 72.5, 70.4, 68.0, 67.24, 61.57, 50.1, 37.8, 28.2, 20.7, 20.5, 20.5; ESI (m/z): [M+Na]⁺ calcd for C₃₀H₄₀N₂O₁₄Na, 675.2; found 675.1.

Boc-*N*-β-2-acetamido-3,4,6-tri-*O*-acetyl-β-*D*-glucopyranosyl-*L*-asparagine benzyl ester (ligation product from glycosyl azide **9)**

Glycosyl azide **9** (55 mg, 148 μmol) was dried by azeotropic removal of water with freshly distilled toluene (3 x) in a round-bottom flask, and the flask was purged with argon. Anhydrous *N,N'*-dimethylformamide (2 mL) was added the flask. The resulting solution of the **7** was transferred via cannula under argon to another flask, previously purged with argon, containing freshly prepared diethylphosphinothioester **5a** (60 mg, 136 μmol). The solution was stirred under argon for 12 h, and water (200 μL) was added. The solution was stirred for additional 30 min, and the solvent was removed under high vacuum. The crude product was purified by flash chromatography (1:9 hexanes:ethyl acetate) to yield a white solid (28 mg, 43 μmol, 32%). The ligations of **9** with phosphinothioesters **5b**, **5c** were carried out in a similar fashion. ¹H NMR (CDCl₃) δ 7.37-7.32 (m, 4H), 7.17 (d, *J*=8.1 Hz), 5.88 (d, *J*=7.9 Hz, 1H), 5.74 (d, *J*=8.9, 1H), 5.22-4.96 (m, 4H), 4.61-4.55 (m, 1H), 4.28 (dd, *J*=4.2, 12.6 Hz, 1H), 4.08-4.05 (m, 1H), 3.73-3.70 (m, 1H), 2.77 (ABX, *J*_{AB}=16.5, *J*_{AX}=3.9, *J*_{BX}=4.1 Hz, 2H), 2.10 (s, 3H), 2.08 (s, 3H), 2.04 (s, 3H), 1.85 (s, 3H), 1.41 (s, 9H); ¹³C NMR (CDCl₃) δ 171.2, 171.1, 170.6, 170.5, 169.8, 169.5, 155.5, 135.3, 128.5, 128.3, 127.9, 80.1, 78.1, 73.7, 72.5, 70.4, 68.0, 67.3, 61.6, 50.1, 37.8, 28.2, 20.7, 20.5, 20.5; ESI (m/z): [M+Na]⁺ calcd for C₃₀H₄₀N₃O₁₃Na, 674.3; found 674.0.

- (1) Matsubara, K.; Mukaiyama, T. *Chem. Lett.* **1994**, 247-250.
- (2) Lehnhoff, S.; Goebel, M.; Karl, R. M.; Klosel, R.; Ugi, I. *Angew. Chem. Int. Ed. Engl.* **1995**, 34, 1104-1107.
- (3) Horner, L.; Beck, P.; Hoffmann, H. *Chem. Ber.* **1959**, 92, 2088-2094.

^1H NMR of compound **1a**

^{13}C NMR of compound **1a**

^{31}P NMR of compound **1a**

^1H NMR of compound **1b**

^{13}C NMR of compound **1b**

^{31}P NMR of compound **1b**

^1H NMR of compound **1c**

^{13}C NMR of compound **1c**

^{31}P NMR of compound **1c**

^1H NMR of diethylphosphino(borane)methyl
methansulfonate

^{13}C NMR of diethylphosphino(borane)methyl
methansulfonate

^{31}P NMR of diethylphosphino(borane)methyl
methanesulfonate

^1H NMR of dibutylphosphino(borane)methyl
methansulfonate

^{13}C NMR of dibutylphosphino(borane)methyl
methansulfonate

^{31}P NMR of dibutylphosphino(borane)methyl
methanesulfonate

^1H NMR of diphenylphosphino(borane)methyl
methansulfonate

^{13}C NMR of diphenylphosphino(borane)methyl
methansulfonate

^{31}P NMR of diphenylphosphino(borane)methyl
methanesulfonate

^1H NMR of compound **3a**

^{31}P NMR of compound **3a**

^{13}H NMR of compound **3a**

^1H NMR of compound **3b**

^{13}C NMR of compound **3b**

^{31}P NMR of compound **3b**

^1H NMR of compound **3c**

^{13}C NMR of compound **3c**

^{31}P NMR of compound **3c**

^1H NMR of compound **4a**

^{13}C NMR of compound **4a**

^{31}P NMR of compound **4a**

^1H NMR of compound **4b**

^{13}C NMR of compound **4b**

^{31}P NMR of compound **4b**

^1H NMR of compound **4c**

^{13}C NMR of compound **4c**

^{31}P NMR of compound **4c**

^1H NMR of compound **5a**

^{13}C NMR of compound **5a**

^{31}P NMR of compound **5a**

^1H NMR of compound **5b**

^{13}C NMR of compound **5b**

^{31}P NMR of compound **5b**

^1H NMR of compound **5c**

^{13}C NMR of compound **5c**

^{31}P NMR of compound **5c**

¹³C NMR