Stereoselective Suzuki-Miyaura Cross-Coupling Reactions of Potassium Alkenyltrifluoroborates with Alkenyl Bromides

Gary A. Molander and Luciana A. Felix*

Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania,
Philadelphia, Pennsylvania 19104-6323

gmolandr@sas.upenn.edu

Table of Contents

General.....	S2
Potassium alkenyltrifluoroborate synthesis.....	S2
Representative procedure for the cross-coupling reaction of potassium alkenyltrifluoroborates.....	S6
References.....	S16
NMR spectra.....	S17

Supporting Information

General: ^1H -, ^{13}C - and ^{19}F -NMR spectra were measured on spectrometers at 500.39 or 300.13, 125.75, and 470.55 MHz, respectively. ^{19}F -NMR chemical shifts were referenced to external CFCl_3 (0.0 ppm). ^{11}B NMR spectra at 64.2 MHz were obtained on a spectrometer equipped with the appropriate decoupling accessories. All ^{11}B chemical shifts were referenced to external $\text{BF}_3\cdot\text{OEt}_2$ (0.0 ppm) with a negative sign indicating an upfield shift. THF was distilled from sodium. All of the cross-coupling reactions were carried out under a nitrogen atmosphere. H_2O used for cross-coupling reactions was degassed using nitrogen.

Potassium (*E*)-1-Dec-1-enyl Trifluoroborate (1a).¹ To a solution of *trans*-2-(decenyl) vinylboronic acid (1.83 g, 10 mmol) in Et_2O (20 mL) was added KHF_2 (2.18 g, 28 mmol), followed by H_2O (9 mL) over a period of 30 min. After stirring at rt for 3 h, the solution was concentrated and the crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et_2O , resulting in a white solid (2.04 g, 8.50 mmol, 85%). The spectral data obtained were in accordance with those described in the literature.

Potassium *trans*-4-Phenyl-1-but-1-enyl Trifluoroborate (1b). To a solution of *trans*-4-phenyl-1-but-1-enylboronic acid² (2.6 g, 20 mmol) in Et_2O (40 mL) was added KHF_2 (4.7 g, 60 mmol), followed by H_2O (20 mL). After stirring at rt for 2 h, the solution was concentrated and the solid crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et_2O , resulting in a white solid (4.2 g, 17.6 mmol, 88%), mp > 260 °C. ^1H NMR (500 MHz, acetone- d_6) δ 7.26-7.12 (m, 5H), 5.79 (dt, J = 17.6, 6.7 Hz, 1H), 5.51-5.41 (m, 1H), 2.65-2.59 (m, 2H), 2.27-2.20 (m, 2H); ^{13}C NMR (125.8 MHz, DMSO- d_6) δ 142.5, 132.8, 132.7, 128.2 (2C), 128.0 (2C), 125.4, 37.3, 35.6; ^{19}F NMR (471 MHz, acetone- d_6) δ -

137.8. ^{11}B NMR (64.2 MHz, acetone- d^6) δ 3.22. HRMS (CI): m/z calcd. For $\text{C}_{10}\text{H}_{11}\text{BF}_3$ (M-K^+) 199.0905, found 199.0901. IR (neat): 1640, 1451, 952, 918 cm^{-1} .

Potassium (Z)-4-Phenyl-1-but-1-enyl Trifluoroborate (1c). To a solution of 4,4,5,5-tetramethyl-2-[(Z)-4-phenylbut-1-enyl]-1,3,2-dioxaborolane³ (2.06 g, 8 mmol) in acetonitrile (15 mL) was added KHF_2 (1.87 g, 24 mmol), followed by H_2O (5 mL). After stirring at rt for 2 h, the solution was concentrated and the solid crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et_2O , resulting in a white solid (1.24 g, 5.2 mmol, 65%), mp $> 260^\circ\text{C}$. ^1H NMR (500 MHz, acetone- d^6) δ 7.22-7.07 (m, 5H), 5.64 (brs, 1H), 5.41-5.31 (m, 1H), 2.62-2.57 (m, 2H), 2.53-2.48 (m, 2H); ^{13}C NMR (125.8 MHz, DMSO- d^6) δ 142.5, 132.8, 132.7, 128.2 (2C), 128.0 (2C), 125.4, 37.3, 35.6; ^{19}F NMR (471 MHz, acetone- d^6) δ -133.0. ^{11}B NMR (64.2 MHz, acetone- d^6) δ 3.07 (q, $J = 57.7\text{ Hz}$). HRMS (CI): m/z calcd. For $\text{C}_{10}\text{H}_{11}\text{BF}_3$ (M-K^+) 199.0905, found 199.0983. IR (neat): 1630, 1453, 935, 911 cm^{-1} .

Potassium Vinyl Trifluoroborate (1d).⁴ To a solution of trimethyl borate (2.32 g, 22.3 mmol) in THF (15 mL) was added dropwise vinylmagnesium bromide (1.0 M THF solution, 18.0 mL, 18.0 mmol) at -78°C . The resulting suspension was stirred for 20 min at -78°C and then allowed to warm to rt for 1 h. The mixture was then cooled to 0°C and KHF_2 (6.97 g, 89.2 mmol) was added followed by H_2O (12 mL) over a period of 30 min. After stirring at rt for 20 min, the solution was concentrated and the crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et_2O , obtaining a white solid (2.25 g, 16.8 mmol, 93%). The spectral data obtained were in accordance with those described in the literature.

Potassium *trans*-1-Ethylbut-1-enyl Trifluoroborate (1e).¹ To a solution of 1-ethenylbut-1-enyl boronic acid (1.28 g, 10 mmol) in Et_2O (20 mL) was added KHF_2 (2.18 g, 28 mmol), followed by H_2O

(9 mL) over a period of 30 min. After stirring at rt for 3 h, the solution was concentrated and the crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O, resulting in a white solid (1.19 g, 6.3 mmol, 65%). The spectral data obtained were in accordance with those described in the literature.

Potassium α -Styryl Trifluoroborate (1f). To a solution of α -phenylethenyl boronic acid⁵ (1.48 g, 10 mmol) in MeOH (20 mL) was added KHF₂ (2.18 g, 30 mmol), followed by H₂O (10 mL). After stirring at rt for 3 h, the solution was concentrated and the crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O, resulting in a white solid (0.84 g, 4.0 mmol, 40%), mp > 260 °C. ¹H NMR (300 MHz, acetone-d₆) δ 7.51 (d, J = 9.0 Hz, 2H), 7.14 (t, J = 9.0 Hz, 2H), 7.03 (t, J = 7.2 Hz, 1H), 5.33 (s, 2H); ¹³C NMR (125.8 MHz, DMSO-d₆) δ 143.5, 127.2 (2C), 126.7, 124.8 (2C), 115.8, 115.9; ¹⁹F NMR (282.4 MHz, acetone-d₆) δ -141.6. ¹¹B NMR (64.2 MHz, acetone-d₆) δ 3.06 (d, J = 49.4). Anal. Calcd for C₈H₇BF₃K: C, 41.33; H, 3.21. found: C, 41.63; H, 3.07. IR (neat): 1702, 1609, 925 cm⁻¹.

Potassium *trans*-Styryl Trifluoroborate (1g).⁴ To a solution of *trans*-2-phenylethenyl boronic acid (1.48 g, 10 mmol) in Et₂O (20 mL) was added KHF₂ (2.18 g, 28 mmol), followed by H₂O (9 mL) over a period of 30 min. After stirring at rt for 3 h, the solution was concentrated and the crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O, resulting in a white solid (1.89 g, 9.0 mmol, 90%). The spectral data obtained were in accordance with those described in the literature.

Potassium *trans*-6-Chlorohex-1-enyl Trifluoroborate (1h). To a solution of *trans*-6-chlorohex-1-enyl boronic acid² (1.79 g, 10 mmol) in Et₂O (20 mL) was added KHF₂ (2.34 g, 30 mmol), followed by H₂O (10 mL). After stirring at rt for 2 h, the solution was concentrated and the crude material was

dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O, resulting in a white solid (1.79 g, 80%), mp > 260 °C. ¹H NMR (500 MHz, acetone-d₆) δ 5.71 (dt, *J* = 17.6, 5.5 Hz, 1 H), 5.30 (m, 1H), 3.58 (t, *J* = 6.8 Hz, 2H), 1.97 (q, *J* = 7.3 Hz, 2H), 1.75 (dt, *J* = 15, 6.8 Hz, 2H), 1.48 (dt, *J* = 15, 7.3 Hz, 2H); ¹³C NMR (125.8 MHz, DMSO-d₆) δ 133.2, 133.1, 45.4, 34.4, 31.7, 26.2; ¹⁹F NMR (282.4 MHz, acetone-d₆): δ -137.8. ¹¹B NMR (64.2 MHz, acetone-d₆) δ 1.81. Anal. Calcd for C₆H₁₀ClBF₃K: C, 31.46; H, 3.54. found: C, 31.51; H, 3.41. IR (neat): 1704, 1644, 939 cm⁻¹.

Potassium *trans*-1-Hex-5-enoic Acid Trifluoroborate, Methyl Ester (1i).¹ To a solution of *trans*-1-hex-5-enoic boronic acid, methyl ester (1.72 g, 10 mmol) in Et₂O (20 mL) was added KHF₂ (2.18 g, 28 mmol), followed by H₂O (9 mL). After stirring at rt for 1.5 h, the solution was concentrated and the crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O, resulting in a white solid (1.16 g, 5 mmol, 50%). The spectral data obtained were in accordance with those described in the literature.

Potassium 2-Trimethylsilylanyl-Vinyl Trifluoroborate (1j). To a solution of 4,4,5,5-tetramethyl-2-((*E*)-2-(trimethylsilyl)vinyl)-1,3,2-dioxaborolane⁶ (1.13 g, 5 mmol) in MeCN (15 mL) was added KHF₂ (1.17 g, 15 mmol), followed by H₂O (5 mL). After stirring at rt for 2 h, the solution was concentrated and the solid crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O, resulting in a white solid (0.72 g, 3.5 mmol, 70%), mp > 260 °C. ¹H NMR (500 MHz, acetone-d₆) δ 6.41 (dq, *J* = 21.6, 3.3 Hz, 1H), 6.08 (d, *J* = 21.7 Hz, 1H), -0.03 (s, 9H); ¹³C NMR (125.8 MHz, DMSO-d₆) δ 133.7, -0.9 (3C); ¹⁹F NMR (471 MHz, acetone-d₆) δ -143.0. ¹¹B NMR (64.2 MHz, acetone-d₆) δ 3.55. Anal. Calcd for C₅H₁₁SiBF₃K: C, 26.07; H, 3.90. found: C, 26.12; H, 3.99. IR (neat): 1246, 1705 cm⁻¹.

Potassium 4-Cyano-But-1-enyl Trifluoroborate (1k). To a solution of (*E*)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pent-4-enenitrile⁶ (1.03 g, 5 mmol) in MeCN (15 ml) was added KHF₂ (1.17 g, 15 mmol), followed by H₂O (5 mL). After stirring at rt for 2 h, the solution was concentrated and the solid crude material was dissolved in acetone, filtered, and concentrated. The resulting white solid was purified by dissolving in hot acetone and precipitating with Et₂O, resulting in a white solid (0.69 g, 3.7 mmol, 75%), mp = 146-148 °C. ¹H NMR (500 MHz, DMSO-*d*⁶) δ 5.68 (dt, *J* = 17.5, 5.6 Hz, 1H), 5.59-5.55 (m, 1H), 2.64 (t, *J* = 7.1 Hz, 2H), 2.35 (q, *J* = 6.5 Hz, 2H); ¹³C NMR (125.8 MHz, DMSO-*d*⁶) δ 130.0, 129.9, 120.8, 30.6, 16.4; ¹⁹F NMR (471 MHz, acetone-*d*⁶) δ -142.0. ¹¹B NMR (64.2 MHz, acetone-*d*⁶) δ 3.07 (d, *J* = 57.7 Hz). Anal. Calcd for C₅H₆NBF₃K: C, 32.09; H, 2.19. found: C, 32.17; H, 2.04. IR (neat): 2256, 1645 cm⁻¹.

Representative Procedure for the Cross-coupling Reaction of Alkenyl Halides with Potassium Alkenyltrifluoroborates. (*E,Z*)-1-Chloro-pentadeca-4,6-diene (2a). To a mixture of potassium (*E*)-1-dec-1-enyl trifluoroborate (**1a**) (370.4 mg, 1.1 mmol), (*Z*)-1-bromo-5-chloro-pent-1-ene⁷ (183.5 mg, 1 mmol), Cs₂CO₃ (977 mg, 3 mmol), Pd(OAc)₂ (11 mg, 0.05 mmol), and PPh₃ (26 mg, 0.1 mmol) was added THF-H₂O (10:1, 4 mL). The reaction was heated at 70 °C with stirring under a nitrogen atmosphere for 2 h, then cooled to rt and diluted with water (3 mL). The resulting mixture was extracted with Et₂O. The organic layers were combined and washed with 1 N HCl and brine, dried over MgSO₄, and then filtered. The solvent was removed under vacuum, and the crude product was purified by silica gel chromatography (eluting with hexane) to afford a colorless oil (200 mg, 85%): ¹H NMR (500 MHz, CDCl₃) δ 6.30 (ddt, *J* = 15, 10.9, 1.2 Hz, 1H), 6.01 (dd, *J* = 10.9, 10.7 Hz, 1H), 5.69 (dt, *J* = 15, 6.9 Hz, 1H), 5.24 (dt, *J* = 10.7, 7.7 Hz, 1H), 3.54 (t, *J* = 6.6 Hz, 2H), 2.32 (q, *J* = 7.7 Hz, 2H), 2.10 (q, *J* = 7.0 Hz, 2H), 1.86 (dt, *J* = 14.2, 6.7 Hz, 2H), 1.39 (dt, *J* = 14.2, 7.2 Hz, 2H), 1.30-1.27 (m, 10H), 0.88 (t, *J* = 7.0 Hz, 3H); ¹³C NMR (125.8 MHz, CDCl₃) δ 135.7, 130.2, 127.2, 125.2, 44.4, 32.9, 32.5, 31.9, 29.5,

29.3, 29.3, 29.2, 24.8, 22.7, 14.1; HRMS (CI): m/z calcd. For $C_{15}H_{27}Cl$ (M^+) 242.1800, found 242.1805; IR (neat): 1325, 1265, 1167, 1128, 1067 cm^{-1} .

(*E,E*)-(9-Chloro-nona-3,5-dienyl)-benzene (2b). This product was obtained by the general method from the reaction of potassium (*E*)-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and (*E*)-1-bromo-5-chloro-pent-1-ene⁸ to afford the title compound in 86% yield. 1H NMR (500 MHz, $CDCl_3$) δ 7.27 (t, J = 7.6 Hz, 2H), 7.18-7.16 (m, 3H), 6.07-6.00 (m, 2H), 5.69 (dt, J = 14.0, 7.0 Hz, 1H), 5.52 (dt, J = 14.0, 7.0 Hz, 1H), 3.52 (t, J = 6.6 Hz, 2H), 2.69 (t, J = 8.0 Hz, 2H), 2.38 (q, J = 6.9 Hz, 2H), 2.21 (q, J = 7.0 Hz, 2H), 1.85 (dt, J = 14.2, 6.9 Hz, 2H); ^{13}C NMR (125.8 MHz, $CDCl_3$) δ 141.8, 132.0, 131.6, 130.5, 130.2, 128.4 (2C), 128.3 (2C), 125.8, 44.3, 35.8, 34.4, 32.1, 29.6; HRMS (CI): m/z calcd for $C_{15}H_{19}Cl$ (M^+) 234.1170, found 234.1177; IR (neat) 3026, 2929, 1720, 1693, 971 cm^{-1} .

(*E,Z*)-(9-Chloro-nona-3,5-dienyl)-benzene (2c). This product was obtained by the general method from the reaction of potassium (*E*)-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and (*Z*)-1-bromo-5-chloro-pent-1-ene⁷ to afford the title compound in 88% yield. 1H NMR (500 MHz, $CDCl_3$) δ 7.30 (t, J = 7.5 Hz, 2H), 7.22-7.19 (m, 3H), 6.37 (ddt, J = 15.1, 10.9, 1.2 Hz, 1H), 6.03 (dd, J = 10.9, 10.7 Hz, 1H), 5.75 (dt, J = 15.1, 6.9 Hz, 1H), 5.30 (dt, J = 10.7, 7.7 Hz, 1H), 3.55 (t, J = 6.6 Hz, 2H), 2.75 (t, J = 8.1 Hz, 2H), 2.46 (q, J = 7.0 Hz, 2H), 2.34 (q, J = 7.7 Hz, 2H), 1.87 (dt, J = 14.2, 6.7 Hz, 2H); ^{13}C NMR (125.8 MHz, $CDCl_3$) δ 141.7, 134.2, 130.0, 128.4 (2C), 128.3 (2C), 127.8, 125.9, 125.8, 44.3, 35.7, 34.6, 32.4, 24.8; HRMS (CI): m/z calcd. For $C_{15}H_{18}Cl$ ($M-H^+$) 233.1092, found 233.1081; IR (neat): 3026, 2927, 1714, 971 cm^{-1} .

(*Z,E*)-(9-Chloro-nona-3,5-dienyl)-benzene (2d). This product was obtained by the general method from the reaction of potassium (*Z*)-4-phenyl-1-but-1-enyl trifluoroborate (**1c**) and (*E*)-1-bromo-5-chloro-pent-1-ene⁸ to afford the title compound in 85% yield. 1H NMR (500 MHz, $CDCl_3$) δ 7.29-7.16

(m, 5H), 6.29 (dddd, $J = 14.9, 10.9, 2.5, 1.2$ Hz, 1H), 5.95 (dd, $J = 10.9, 10.7$ Hz, 1H), 5.59 (dt, $J = 15.0, 7.0$ Hz, 1H), 5.35 (dt, $J = 10.7, 7.5$ Hz, 1H), 3.52 (t, $J = 6.6$ Hz, 2H), 2.69 (t, $J = 7.4$ Hz, 2H), 2.47 (q, $J = 7.9$ Hz, 2H), 2.23 (q, $J = 7.1$ Hz, 2H), 1.84 (dt, $J = 14.3, 6.6$ Hz, 2H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 141.7, 132.4, 129.4, 128.8 (2C), 128.4 (2C), 128.2, 126.8, 125.8, 44.3, 35.8, 32.1, 29.9, 29.5; HRMS (CI): m/z calcd. For $\text{C}_{15}\text{H}_{18}\text{Cl}$ ($\text{M}-\text{H}^+$) 233.1092, found 233.1083; IR (neat): 3026, 2927, 1716, 971 cm^{-1} .

(Z,Z)-(9-Chloro-nona-3,5-dienyl)-benzene (2e). This product was obtained by the general method from the reaction of potassium (*Z*)-4-phenyl-1-but-1-enyl trifluoroborate (**1c**) and (*Z*)-1-bromo-5-chloro-pent-1-ene⁷ to afford the title compound in 86% yield. ^1H NMR (500 MHz, CDCl_3) δ 7.20 (t, $J = 7.0$ Hz, 2H), 7.10-7.09 (m, 3H), 6.23-6.17 (m, 2H), 5.44 (dt, $J = 9.0, 7.5$ Hz, 1H), 5.31 (dt, $J = 8.0, 7.5$ Hz, 1H), 3.42 (t, $J = 6.5$ Hz, 2H), 2.61 (t, $J = 7.5$ Hz, 2H), 2.42 (q, $J = 8.0$ Hz, 2H), 2.24 (q, $J = 7.0$ Hz, 2H), 1.75 (dt, $J = 14.2, 7.0$ Hz, 2H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 141.7, 131.5, 129.7, 128.4 (2C), 128.3 (2C), 125.8, 124.8, 123.7, 44.4, 35.8, 32.3, 29.3, 24.6; HRMS (CI): m/z calcd. For $\text{C}_{15}\text{H}_{18}\text{Cl}$ ($\text{M}-\text{H}^+$) 233.1092, found 233.1091; IR (neat): 3026, 2927, 1720 cm^{-1} .

Trans-(6-Methyl-hepta-3,5-dienyl)-benzene (2f). This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and 1-bromo-2-methylpropene to afford the title compound in 70% yield. ^1H NMR (500 MHz, CDCl_3) δ 7.27 (t, $J = 7.7$ Hz, 2H), 7.19-7.16 (m, 3H), 6.26 (dd, $J = 15.0, 10.8$ Hz, 1H), 5.78 (d, $J = 10.8$ Hz, 1H), 5.58 (dt, $J = 15.0, 6.9$ Hz, 1H), 2.70 (t, $J = 8.3$ Hz, 2H), 2.40 (q, $J = 7.2$ Hz, 2H), 1.75 (s, 3H), 1.72 (s, 3H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 142.0, 133.3, 130.7, 128.4 (2C), 128.3 (2C), 127.2, 125.7, 124.9, 36.11, 34.8, 25.9, 18.2; HRMS (CI): m/z calcd. For $\text{C}_{14}\text{H}_{18}$ (M^+) 186.1408, found 186.1401; IR (neat): 3025, 2922, 1452, 1377, 957, 745 cm^{-1} .

***Trans*-(5,6-Dimethyl-hepta-3,5-dienyl)-benzene (2g).** This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and 2-bromo-3-methyl-but-2-ene to afford the title compound in 95% yield. ¹H NMR (500 MHz, CDCl₃) δ 7.28 (t, *J* = 7.7 Hz, 2H), 7.20-7.16 (m, 3H), 6.54 (d, *J* = 15.4 Hz, 1H), 5.61 (dt, *J* = 15.4, 6.9 Hz, 1H), 2.73 (t, *J* = 8.4 Hz, 2H), 2.44 (q, *J* = 7.2 Hz, 2H), 1.77 (t, *J* = 12 Hz, 9H); ¹³C NMR (125.8 MHz, CDCl₃) δ 142.1, 129.8, 129.1, 128.4 (2C), 128.2 (2C), 127.0, 125.9, 125.7, 36.5, 35.3, 21.6, 20.2, 14.4; HRMS (CI): *m/z* calcd. For C₁₅H₂₀ (M⁺) 200.1565, found 200.1557; IR (neat): 3026, 2919, 1450, 1373, 959, 698 cm⁻¹.

***Trans*-3-Methylene-7-phenyl-hept-4-en-1-ol (2h).** This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and 3-bromo-but-3-en-1-ol to afford the title compound in 79% yield. ¹H NMR (500 MHz, CDCl₃) δ 7.31-7.27 (m, 2H), 7.21-7.18 (m, 3H), 6.11 (d, *J* = 15.8 Hz, 1H), 5.78 (dt, *J* = 15.8 and 6.9 Hz, 1H), 5.03 (s, 1H), 4.96 (s, 1H), 3.73 (t, *J* = 6.9 Hz, 2H), 2.73 (t, *J* = 6.8 Hz, 2H), 2.49 (t, *J* = 6.4 Hz, 2H), 2.44 (q, *J* = 7.0 Hz, 2H), 1.47 (s, 1H); ¹³C NMR (125.8 MHz, CDCl₃) δ 142.3, 141.6, 132.0, 130.0, 128.4 (2C), 128.3 (2C), 125.9, 115.7, 61.1, 35.8, 35.6, 34.6; HRMS (CI): *m/z* calcd. For C₁₄H₁₈O (M⁺) 202.1357, found 202.1350; IR (neat): 3400, 3027, 2936 cm⁻¹.

***Trans*-3-Methyl-2-(4-phenyl-but-1-enyl)-cyclopent-2-enone (2i).** This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and 2-bromo-3-methyl-cyclopent-2-enone to afford the title compound in 95% yield. ¹H NMR (500 MHz, CDCl₃) δ 7.27 (t, *J* = 7.5 Hz, 2H), 7.20-7.15 (m, 3H), 6.75 (dt, *J* = 15.8, 6.7 Hz, 1H), 6.08 (d, *J* = 15.8 Hz, 1H), 2.75 (t, *J* = 7.3 Hz, 2H), 2.49-2.44 (m, 4H), 2.39-2.38 (m, 2H), 2.08 (s, 3H); ¹³C NMR (125.8 MHz, CDCl₃) δ 208.2, 169.6, 141.9, 135.3, 134.8, 128.4 (2C), 128.3 (2C), 125.8, 119.3, 35.9, 35.7, 34.8, 31.5, 17.5; HRMS (CI): *m/z* calcd. For C₁₆H₁₈O (M⁺) 226.1357, found 226.1348; IR (neat): 3025, 2918, 2852, 1625 cm⁻¹.

2-Benzylidene-6-phenyl-hex-3-enal (2j). This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and *trans*- α -bromocinnamaldehyde to afford the title compound in 88% yield. ^1H NMR (500 MHz, CDCl_3) δ 9.64 (s, 1H), 7.36 (m, 5H), 7.29 (t, $J = 7.4$, 2H), 7.24-7.19 (m, 3H), 7.07 (s, 1H), 6.70 (dt, $J = 16.0$, 6.9 Hz, 1H), 6.17 (d, $J = 16.0$, 1H), 2.79 (t, $J = 7.5$ Hz, 2H), 2.53 (q, $J = 7.3$ Hz, 2H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 194.2, 148.3, 141.6, 138.3, 136.5, 134.9, 130.1 (2C), 129.5, 128.6 (2C), 128.5 (2C), 128.3 (2C), 125.9, 121.4, 35.8, 35.2; HRMS (CI): m/z calcd. For $\text{C}_{19}\text{H}_{18}\text{O}$ (M^+) 262.1357, found 262.1356; IR (neat) 3026, 2925, 1687 cm^{-1} .

2,3-Bis-(4-phenyl-but-1-enyl)-but-2-enedioic Acid, Dimethyl Ester (2k). This product was obtained by the following procedure: To a mixture of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) (285 mg, 1.2 mmol), 2,3-dibromo-but-2-ene-dioic acid, dimethyl ester⁹ (151 mg, 0.5 mmol), Cs_2CO_3 (1955 mg, 6 mmol), and $\text{PdCl}_2\text{dppf}\cdot\text{CH}_2\text{Cl}_2$ (41 mg, 0.05 mmol) was added THF- H_2O (10:1, 4 mL). The reaction was heated at 70 $^\circ\text{C}$ with stirring under a nitrogen atmosphere for 4 h, then cooled to rt and diluted with H_2O (3 mL). The resulting mixture was extracted with Et_2O . The organic layers were combined and washed with a saturated solution of NH_4Cl and brine, dried over MgSO_4 , and then filtered. The solvent was removed under vacuum and the crude product was purified by silica gel chromatography (eluting with hexane/ EtOAc) to afford a yellow oil (158 mg, 78%). ^1H NMR (500 MHz, CDCl_3) δ 7.24-7.20 (m, 4H), 7.12-7.08 (m, 6H), 6.13 (d, $J = 15.8$ Hz, 2H), 5.85 (dt, $J = 15.8$, 6.9 Hz, 2H), 3.68 (s, 6H), 2.65 (t, $J = 7.7$ Hz, 4H), 2.41 (q, $J = 7.5$ Hz, 4H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 168.0 (2C), 141.1 (2C), 139.3 (2C), 133.3 (2C), 128.4 (4C), 128.3 (4C), 126.0 (2C), 123.6 (2C), 52.2 (4C), 35.14 (2C); HRMS (CI): m/z calcd. For $\text{C}_{26}\text{H}_{28}\text{O}_4\text{Na}$ ($\text{M} + \text{Na}^+$) 427.1885, found 427.1900; IR (neat) 3026, 2925, 1731 cm^{-1} .

10-Phenyl-deca-5,7-dienenitrile (2l). This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and (Z)-5-bromo-pent-4-enenitrile¹⁰ to afford the title compound in 78% yield. ¹H NMR (500 MHz, CDCl₃) δ 7.29 (t, *J* = 7.2 Hz, 2H), 7.19-7.17 (m, 3H), 6.29 (ddt, *J* = 15.0, 10.8, 1.2 Hz, 1H), 6.03 (dd, *J* = 10.8, 10.7 Hz, 1H), 5.74 (dt, *J* = 15.0, 6.9 Hz, 1H), 5.21 (dt, *J* = 10.7, 7.7 Hz, 1H), 2.72 (t, *J* = 7.5 Hz, 2H), 2.44 (q, *J* = 7.5 Hz, 2H), 2.31-2.27 (m, 4H), 1.72 (dt, *J* = 14.4, 7.2 Hz, 2H); ¹³C NMR (125.8 MHz, CDCl₃): δ 141.6, 134.9, 130.8, 128.4 (2C), 128.3 (2C), 126.7, 125.8, 125.6, 119.6, 35.6, 34.5, 26.3, 26.3, 16.3; HRMS (CI): *m/z* calcd. For C₁₆H₁₉N (M⁺) 225.1517, found 225.1514; IR (neat) 3023, 2931, 2250 cm⁻¹.

(2-(4-Phenyl-but-1-enyl)-1*H*-indol-3-yl)-acetonitrile (2m): This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and 2-bromo-1*H*-indol-3-yl)-acetonitrile to afford the title compound in 90% yield. ¹H NMR (500 MHz, CDCl₃) δ 8.12 (s, 1H), 7.44 (d, *J* = 7.5 Hz, 1H), 7.20 (t, *J* = 8.0 Hz, 2H), 7.14-7.05 (m, 5H), 7.01 (t, *J* = 8.0 Hz, 1H), 6.30 (d, *J* = 16.0 Hz, 1H), 5.93 (dt, *J* = 16.0 and 6.5 Hz, 1H), 3.57 (s, 2H), 2.68 (t, *J* = 7.9 Hz, 2H), 2.45 (q, *J* = 6.5 Hz, 2H); ¹³C NMR (125.8 MHz, CDCl₃) δ 141.1, 135.6, 133.4, 131.1, 128.4 (2C), 128.4 (2C), 127.4, 126.0, 123.0, 120.2, 118.0, 117.9, 117.8, 110.7, 101.1, 35.4, 34.7, 12.7; HRMS (CI): *m/z* calcd. For C₂₀H₁₈N₂ (M⁺) 286.1470; found 286.1472; IR (neat) 3361, 3026, 2924, 2248 cm⁻¹.

Triisopropyl-(3-methylene-7-phenyl-hept-4-enyloxy)-silane (2n): This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and (3-bromo-but-3-enyloxy)-triisopropylsilane¹¹ to afford the title compound in 85% yield. ¹H NMR (500 MHz, CDCl₃) δ 7.21-7.16 (m, 2H), 7.12-7.09 (m, 3H), 6.00 (d, *J* = 15.8 Hz, 1H), 5.69 (dt, *J* = 15.8, 6.8 Hz, 1H), 4.86 (s, 1H), 4.83 (s, 1H), 3.70 (t, *J* = 7.4 Hz, 2H), 2.63 (t, *J* = 7.5 Hz, 2H), 2.39 (t, *J* = 7.3 Hz, 2H), 2.34 (q, *J* = 7.3 Hz, 2H), 1.05-0.90 (m, 21H); ¹³C NMR (125.8 MHz, CDCl₃) δ 143.0, 141.8, 132.6,

129.3, 128.4 (2C), 128.3 (2C), 125.8, 115.1, 62.8, 36.0, 35.9, 34.7, 18.0 (6C), 12.3 (3C); HRMS (CI): m/z calcd. For $C_{23}H_{37}OSi$ ($M-H^+$) 357.2613, found, 357.2608; IR (neat): 3027, 2942, 2865, 1103 cm^{-1}

***tert*-Butyl-dimethyl-(3-methylene-7-phenyl-hept-4-enyloxy)-silane (2o):** This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and (3-bromo-but-3-enyloxy)-*tert*-butyl-dimethylsilane¹¹ to afford the title compound in 80% yield. 1H NMR (500 MHz, $CDCl_3$) δ 7.26 (t, $J = 7.2$ Hz, 2H), 7.17-7.16 (m, 3H), 6.07 (d, $J = 15.7$ Hz, 1H), 5.75 (dt, $J = 15.7$ and 7.7 Hz, 1H), 4.93 (s, 1H), 4.88 (s, 1H), 3.70 (t, $J = 7.2$ Hz, 2H), 2.71 (t, $J = 7.7$ Hz, 2H), 2.43-2.39 (m, 4H), 0.89 (s, 9H), 0.04 (s, 6H); ^{13}C NMR (125.8 MHz, $CDCl_3$) δ 142.9, 141.8, 132.6, 129.3, 128.4 (2C), 128.3 (2C), 125.8, 115.1, 62.6, 35.9, 35.9, 34.7, 26.0 (3C), 18.4, -5.2 (2C); HRMS (CI): m/z calcd. For $C_{19}H_{29}OSi$ ($M-Me^+$) 301.1985, found, 301.1984; IR (neat) 3027, 2929, 2857, 1095 cm^{-1}

Triethyl-(3-methylene-7-phenyl-hept-4-enyloxy)-silane (2p). This product was obtained by the general method from the reaction of potassium *trans*-4-phenyl-1-but-1-enyl trifluoroborate (**1b**) and (3-bromo-but-3-enyloxy)-triethylsilane¹¹ to afford the title compound in 80% yield. 1H NMR (500 MHz, $CDCl_3$) δ 7.27 (t, $J = 7.5$ Hz, 2H), 7.18-7.16 (m, 3H), 6.07 (d, $J = 15.8$ Hz, 1H), 5.76 (dt, $J = 15.8$ and 6.9 Hz, 1H), 4.93 (s, 1H), 4.88 (s, 1H), 3.70 (t, $J = 7.3$ Hz, 2H), 2.70 (q, $J = 8.2$ Hz, 2H), 2.46-2.38 (m, 4H), 0.95 (t, $J = 8$ Hz, 9H), 0.59 (q, $J = 8$ Hz, 6H); ^{13}C NMR (125.8 MHz, $CDCl_3$) δ 142.9, 141.8, 132.6, 129.3, 128.4 (2C), 128.3 (2C), 125.8, 115.1, 62.2, 35.9, 35.9, 34.6, 6.8 (3C), 4.5 (3C); HRMS (CI): m/z calcd. For $C_{20}H_{32}OSi$ (M^+) 316.2222, found 316.2207; IR (neat) 3026, 2953, 2930, 2876, 1094 cm^{-1} .

2-Benzylidene-but-3-enal (3a). This product was obtained by the general method from the reaction of potassium vinyl trifluoroborate (**1d**) and *trans*- α -bromocinnamaldehyde to afford the title compound in 73% yield. 1H NMR (500 MHz, $CDCl_3$) δ 9.71 (s, 1H), 7.54 (d, $J = 8.8$ Hz, 2H), 7.46-7.41 (m, 3H),

7.22 (s, 1H), 6.58 (dd, $J = 17$ and 11.5 Hz, 1H), 6.16 (d, $J = 17$ Hz, 1H), 5.59 (d, $J = 12$ Hz, 1H); ^{13}C NMR (125.8 MHz, CDCl_3) δ 193.9, 149.8, 136.5, 134.6, 130.3 (2C), 129.9, 128.6 (2C), 127.6, 122.6. HRMS (CI): m/z calcd. For $\text{C}_{11}\text{H}_{11}\text{O}$ ($\text{M}+\text{H}^+$) 159.0810, found 159.0817. IR (neat) 3060, 3029, 2974, 2929, 1682 cm^{-1} .

4-Ethyl-3-methylene-hept-4-en-1-ol (3b). This product was obtained by the general method from the reaction of potassium *trans*-1-ethylbut-1-enyl trifluoroborate (**1e**) and 3-bromo-but-3-en-1-ol to afford the title compound in 66% yield. ^1H NMR (500 MHz, CDCl_3): δ 5.51 (t, $J = 7.1$ Hz, 1H), 5.10 (s, 1H), 4.93 (s, 1H), 3.68 (t, $J = 6.0$ Hz, 2H), 2.52 (t, $J = 6.4$ Hz, 2H), 2.26 (q, $J = 7.5$ Hz, 2H), 2.12 (dt, $J = 14.8$, 7.4 Hz, 2H), 1.01 (t, $J = 7.5$ Hz, 3H), 0.97 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 144.7, 140.0, 129.5, 112.4, 61.4, 37.5, 21.4, 21.0, 14.3, 13.7; HRMS (CI): m/z calcd. For $\text{C}_{10}\text{H}_{18}\text{O}$ (M^+) 154.1357, found 154.1362; IR (neat): 2970, 2938, 2879 cm^{-1} .

3-Methylene-4-phenyl-pent-4-en-1-ol (3c). This product was obtained by the general method from the reaction of potassium α -styryl trifluoroborate (**1f**) and 3-bromo-but-3-en-1-ol to afford the title compound in 60% yield. ^1H NMR (500 MHz, CDCl_3): δ 7.33-7.24 (m, 5H), 5.31 (s, 1H), 5.22 (s, 1H), 5.17 (s, 1H), 5.10 (s, 1H), 3.71 (t, $J = 12.6$ Hz, 2H), 2.53 (t, $J = 12.7$ Hz, 2H), 1.52 (s, 1H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 150.1, 145.4, 140.6, 128.1 (2C), 128.0 (2C), 127.5, 117.7, 114.1, 61.0, 37.7; HRMS (CI): m/z calcd. For $\text{C}_{12}\text{H}_{14}\text{O}$ (M^+) 174.1044, found 174.1038; IR (neat): 3400, 3057, 3029, 2938, 2879 cm^{-1} .

3-Methylene-5-phenyl-pent-4-en-1-ol (3d). This product was obtained by the general method from the reaction of potassium *trans*-styryl trifluoroborate (**1g**) and 3-bromo-but-3-en-1-ol to afford the title compound in 70% yield. ^1H NMR (500 MHz, CDCl_3): δ 7.42 (d, $J = 7.5$ Hz, 2H), 7.32 (t, $J = 7.9$ Hz, 2H), 7.25-7.23 (m, 1H), 6.81 (d, $J = 16.3$ Hz, 1H), 6.61 (d, $J = 16.3$ Hz, 1H), 5.26 (s, 1H), 5.14 (s, 1H),

3.82 (q, $J = 6.4$ Hz, 2H), 2.64 (t, $J = 6.4$ Hz, 2H), 1.47 (t, $J = 6.3$ Hz, 1H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 142.4, 137.1, 130.4, 128.7 (2C), 128.6 (2C), 127.7, 126.5, 118.1, 61.2, 35.4; HRMS (CI): m/z calcd. For $\text{C}_{12}\text{H}_{14}\text{O}$ (M^+) 174.1044, found 174.1040; IR (neat) 3369, 3080, 3058, 3025, 2945, 2884 cm^{-1} .

Cis-3-Methyl-2-(4-phenyl-but-1-enyl)-cyclopent-2-enone (3e). This product was obtained by the general method from the reaction of potassium *cis*-4-phenyl-1-but-1-enyl trifluoroborate (**1c**) and 2-bromo-3-methyl-cyclopent-2-enone to afford the title compound in 94% yield. ^1H NMR (500 MHz, CDCl_3): δ 7.25 (t, $J = 7.3$ Hz, 2H), 7.17-7.14 (m, 3H), 5.85-5.78 (m, 2H), 2.68 (t, $J = 8.1$ Hz, 2H), 2.53-2.52 (m, 2H), 2.42-2.40 (m, 2H), 2.24 (q, $J = 8.2$ Hz, 2H), 1.96 (s, 3H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 208.0, 171.8, 141.7, 137.8, 135.4, 128.3 (2C), 128.2 (2C), 125.7, 118.9, 35.4, 34.6, 31.7, 31.6, 18.3; HRMS (CI): m/z calcd. For $\text{C}_{16}\text{H}_{18}\text{O}$ (M^+) 226.1357, found 226.1364; IR (neat) 3084, 3060, 3025, 2923, 2856, 1697 cm^{-1} .

9-Chloro-3-methylene-non-4-en-1-ol (3f). This product was obtained by the general method from the reaction of potassium *trans*-6-chlorohex-1-enyl trifluoroborate (**1h**) and 3-bromo-but-3-en-1-ol to afford the title compound in 72% yield. ^1H NMR (500 MHz, CDCl_3): δ 6.08 (d, $J = 15.8$ Hz, 1H), 5.71 (dt, $J = 15.8$ and 6.9 Hz, 1H), 5.02 (s, 1H), 4.95 (s, 1H), 3.72 (t, $J = 6.5$ Hz, 2H), 3.54 (t, $J = 6.5$ Hz, 2H), 2.48 (t, $J = 6.5$ Hz, 2H), 2.14 (q, $J = 7.3$ Hz, 2H), 1.78 (dt, $J = 15.0$, 6.7 Hz, 3H), 1.56 (dt, $J = 15.0$, 7.3 Hz, 2H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 142.2, 132.0, 129.9, 115.5, 61.0, 44.8, 35.7, 31.9, 31.9, 26.4; HRMS (CI): m/z calcd. For $\text{C}_9\text{H}_{14}\text{OCl}$ (M-Me^+) 173.0732, found 173.0724. IR (neat) 3384, 3084, 3061, 3025, 2930, 2862 cm^{-1} .

9-Hydroxy-7-methylene-non-5-enoic Acid, Methyl Ester (3g). This product was obtained by the general method from the reaction of potassium *trans*-1-hex-5-enoic acid trifluoroborate, methyl ester (**1i**) and 3-bromo-but-3-en-1-ol to afford the title compound in 70% yield. ^1H NMR (500 MHz, CDCl_3):

δ 6.08 (d, J = 15.8 Hz, 1H), 5.70 (dt, J = 15.8 and 6.94 Hz, 1H), 5.02 (s, 1H), 4.95 (s, 1H), 3.73 (t, J = 6.4 Hz, 2H), 3.67 (s, 3H), 2.49 (t, J = 6.4 Hz, 2H), 2.32 (t, J = 7.5 Hz, 2H), 2.15 (q, J = 7.2 Hz, 2H), 1.73 (dt, J = 14.8, 7.3 Hz, 2H), 1.64 (s, 1H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 173.9, 142.3, 132.4, 129.5, 115.7, 61.1, 51.5, 35.5, 33.4, 32.1, 24.4; HRMS (CI): m/z calcd. For $\text{C}_{11}\text{H}_{17}\text{O}_3$ ($\text{M}-\text{H}^+$) 197.1171, found 197.1182. IR (neat) 3430, 2952, 2883, 1736 cm^{-1} .

2-Benzylidene-4-trimethylsilanyl-but-3-enal (3h). This product was obtained by the general method from the reaction of potassium 2-trimethylsilanyl-vinyl trifluoroborate (**1j**) and *trans*- α -bromocinnamaldehyde to afford the title compound in 82% yield. ^1H NMR (500 MHz, CDCl_3): δ 9.70 (s, 1H), 7.53 (d, J = 6.6 Hz, 2H), 7.44-7.39 (m, 3H), 7.17 (s, 1H), 6.82-6.81 (m, 2H), 0.14 (s, 9H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 193.9, 148.3, 139.1, 137.3, 134.8, 134.7, 130.4 (2C), 129.8, 128.5 (2C), -1.5 (3H); HRMS (CI): m/z calcd. For $\text{C}_{14}\text{H}_{18}\text{OSi}$ (M^+) 230.1126, found 230.1117. IR (neat) 3063, 3027, 2956, 2887, 1687, 1248 cm^{-1} .

8-Hydroxy-6-methylene-oct-4-enenitrile (3i). This product was obtained by the general method from the reaction of potassium 4-cyano-but-1-enyl trifluoroborate (**1l**) and 3-bromo-but-3-en-1-ol to afford the title compound in 80% yield. ^1H NMR (500 MHz, CDCl_3): δ 6.20 (d, J = 15.7 Hz, 1H), 7.71 (dt, J = 15.7 and 6.6 Hz, 1H), 5.11 (s, 1H), 5.05 (s, 1H), 3.75 (t, J = 6.5 Hz, 2H), 2.51-2.42 (m, 6H), 1.25 (s, 1H); ^{13}C NMR (125.8 MHz, CDCl_3): δ 141.6, 134.7, 125.4, 119.0, 117.4, 60.9, 35.3, 28.5, 17.5; HRMS (CI): m/z calcd. For $\text{C}_9\text{H}_{14}\text{NO}$ ($\text{M} + \text{H}^+$) 152.1075, found 152.1074; IR (neat) 3435, 2929, 2835, 2248 cm^{-1} .

References

- [1] Molander, G. A.; Bernardi, C. R. *J. Org. Chem.* **2002**, 67, 8424-8429.

- [2] The corresponding boronic acid was obtained by the published procedure: Brown, H. C.; Bhat, N. G.; Somayaji V. *Organometallics* **1983**, 2, 1311-1316.
- [3] The corresponding boronate ester was obtained by the published procedure: Ohmura, T.; Yamamoto, Y.; Miyaura, N. *J. Am. Chem. Soc.* **2000**, 122, 4990-4991.
- [4] Molander, G. A.; Rodriguez Rivero, M. *Org. Lett.* **2002**, 4, 107-109.
- [5] Commercial sample
- [6] The corresponding boronate ester was obtained by the published procedure: Pereira, S.; Srebnick, M. *Organometallics* **1995**, 14, 3127-3128.
- [7] (Z)-1-Bromo-5-chloro-pent-1-ene was obtained by the published procedure: Brown, H. C.; Subrahmanyam, C.; Hamoaka, T.; Ravindran, N.; Bowman, D. H.; Misumi, S.; Unni, M. K.; Somayaji, V.; Bhat, N. G. *J. Org. Chem.* **1989**, 54, 6068-6075.
- [8] (E)-1-Bromo-5-chloro-pent-1-ene was obtained by the published procedure: Lipshutz, B. H.; Kell, R.; Ellsworth, E. L. *Tetrahedron Lett.* **1990**, 31, 7257-7260.
- [9] 2,3-Dibromo-but-2-ene-dioic acid, dimethyl ester was obtained from dibromomaleic acid according to the published procedure: Hashimoto, N.; Aoyama, T.; Shioiri, T. *Chem. Pharm. Bull.* **1981**, 29, 1475-1478.
- [10] (Z)-5-Bromo-pent-4-enitrile was obtained from (Z)-1-Bromo-5-chloro-pent-1-ene according to the published procedure: Friedman, L.; Shechter, H. *J. Org. Chem.* **1960**, 25, 877-879.
- [11] (3-Bromo-but-3-enyloxy)-triisopropylsilane, (3-bromo-but-enyloxy)-tert-butyl-dimethylsilane and (3-bromo-but-3-enyloxy)-triethylsilane were prepared from 3-bromo-3-en-1-ol by treating respectively with TIPSCl, TBDMSCl and TESCl according to the published procedures: (a) Cunico, R. F.; Bedell, L. *J. Org. Chem.* **1980**, 45, 4797-4798. (b) Corey, E. J.; Venkateswarlu, A. *J. Am. Chem. Soc.* **1972**, 94, 6190. (c) Hart, T. W.; Metcalfe, D. A.; Scheinmann, F. *J. Chem. Soc. Chem. Commun.* **1979**, 156.

Current Data Parameters
 NAME laf94
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20030530
 Time 19:49
 INSTRUM spect
 PROBHD 5 mm QNP 1H
 PULPROG zg30
 TD 32768
 SOLVENT CDCl3
 NS 2
 DS 2
 SWH 3591.954 Hz
 FIDRES 0.109618 Hz
 AQ 4.561356 sec
 RG 655.18
 DB 139.200 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.0000000 sec

CHANNEL f1
 NUC1 1H
 P1 10.00 usec
 PL1 2.00 dB
 SFO1 300.1315007 MHz
 F2 - Processing Parameters
 SI 32768
 SF 300.1300042 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 P1 2.00 usec
 PL 86.57 Hz
 F1 2.108 ppm
 F2 632.63 Hz
 FFCM 0.03231 ppm/cm
 HZCM 9.69714 Hz/cm

Current Data Parameters
 DU /y
 USER gam
 NAME 1
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20050104
 Time 11.11
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 TD 131072
 SOLVENT C6D6
 NS 52
 DS 4
 SWH 100000.000 Hz
 FIDRES 0.762939 Hz
 AQ 0.6554100 sec
 RG 2298.8
 DW 5.000 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.00000000 sec
 D11 0.03000000 sec
 D12 0.0002000 sec

===== CHANNEL f1 =====
 NUC1 19F
 P1 10.60 usec
 PL1 -3.00 dB
 SFO1 470.7901483 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 26.00 dB
 SFO2 500.3920016 MHz

F2 - Processing parameters
 SI 65536
 SF 470.8372320 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 F1P 0.000 ppm
 F2P -200.000 ppm

1c

Current Data Parameters
 DU /y
 USER gam
 NAME cisbf3k
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20030731
 Time 8.14
 INSTRUM spect
 PROBHD 5 mm QNP 1H
 SOLVENT CDCl3
 TE 300.0 K
 PULPROG zg30
 NUC1 1H
 SF01 300.1315007 MHz
 O1P 5.000 ppm
 SW 11.9679 ppm
 AQ 4.5613556 sec
 TD 32768
 FIDRES 0.109618 Hz
 DW 139.200 usec
 RG 4096
 DS 2
 NS 20
 DE 6.00 usec
 D1 1.00000000 sec

===== CHANNEL f1 =====
 P1 10.00 usec
 PL1 2.00 dB

F2 - Processing parameters
 SI 32768
 SF 300.1300041 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 20.00 cm
 F1P 9.000 ppm
 F2P -0.500 ppm

Current Data Parameters
 DU USER /y
 NAME 1F-1
 EXNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20041222
 Time_ 14:05:16
 INSTRUM 400500
 PROBD 5 mm Dual 13
 P1PRPG 6598
 TD 32768
 SOLVENT CCl3
 NS 292
 DS 2
 SNH 3333.332 Hz
 FIDRES 0.30826 Hz
 AQ 0.316284 sec
 RG 16384
 DW 15.000 usec
 DE 19.000 usec
 TE 300.0 K
 H1 40 dB
 H11 2.000000 sec
 CDEPRG WALT16
 F31 107.00 usec
 SFO1 125.832350 MHz
 D11 0.0300000 sec
 S2 18 dB
 F2 15.000 usec
 SFO2 125.882148 MHz
 NUCLEUS 13C
 F2 - Processing Parameters
 SI 32768
 SF 125.832350 MHz
 WD 0
 SSB 1.50 Hz
 GB 0
 PC 1.40
 1D NMR Plot Parameters
 CX 21.00 cm
 FIP 136.956 ppm
 F2P 132.110 ppm

Current Data Parameters
 USER gam
 NAME 1,2-bis(4-fluorophenyl)ethane
 PROCNO 1
 P2 - Acquisition Parameters
 Date_ 20050104
 Time 11:11
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 TOUPOG zgpg30
 TD 65536
 SOLVENT C5D6
 DS 4
 SFO 400.1463 MHz
 AQ 19000.000 Hz
 RG 65536
 DE 2.000 usec
 DE 5.00 usec
 DI 1.0000000 sec
 D1 0.0300000 sec
 D12 0.0002000 sec
 CHANNEL f1
 P1 10.00 usec
 PL1 -1.00 dB
 SFO1 478.7931463 MHz
 CHANNEL f2
 P2 10.00 usec
 PL2 -1.00 dB
 SFO2 478.7931463 MHz
 CPDPRG2 waltz16
 NUC1 1H
 NUC2 13C
 P1 10.00 usec
 PL1 -1.00 dB
 P2 10.00 usec
 PL2 -1.00 dB
 SFO2 500.392016 MHz
 P2 - Processing parameters
 SI 32768
 SF 478.873320 MHz
 MD 32
 EM 1
 GB 0.30 Hz
 ZB 1.00
 FC 1.00
 ID NMR plot parameters
 FID 115.494 ppm
 F1P -125.494 ppm
 F2P -137.731 ppm

Current Data Parameters

DU /y
USER gam
NAME djc008-prod
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20030808
Time 13.23
INSTRUM spect
PROBHD 5 mm GNP 1H
SOLVENT CDCl3
TE 300.0 K
PULPROG zg30
NUC1 1H
SF01 300.1315007 MHz
Q1P 5.000 ppm
SW 11.9679 ppm
AQ 4.5613556 sec
TD 32768
FIDRES 0.109616 Hz
DM 139.200 usec
RG 2048
DS 2
NS 20
DE 6.00 usec
D1 1.00000000 sec

===== CHANNEL f1 =====

P1 10.00 usec
PL1 2.00 dB

F2 - Processing parameters

SF 32768
SF 300.1300041 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters

CX 20.00 cm
F1P 10.000 ppm
F2P -0.500 ppm

ppm

[Integral]

ppm

ppm

```

Current Data Parameters
=====
INSTR  spect
PROBHD 5 mm QNP 1H/1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 92
DS 4
SWH 31446.514 Hz
FIDRES 10.475816 Hz
AQ 1.0420724 sec
RG 4096
DM 15.900 usec
DE 6.00 usec
TE 300.2 K
D1 2.00000000 sec
D11 0.03000000 sec
D12 0.00020000 sec
===== CHANNEL f1 =====
NUC1 13C
P1 7.50 usec
PL 1.00 dB
SFO1 125.8162556 MHz
===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 0.00 dB
PL12 24.00 dB
PL13 24.00 dB
SFO2 500.3820016 MHz
F2 - Processing parameters
SI 17768
SF 125.832283 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40
ID MR plot parameters
CX 22.00 cm
FID 20.000 ppm
F2F -10.000 ppm
  
```

ppm

141.43
141.64
141.82

Current Data Parameters
 DU USER /y
 NAME dm
 EXNO laf15
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20030320
 Time_ 15.50
 INSTRUM spect
 PROBRD 5 mm QNP 1H
 PULPROG zgpg30
 SOLVENT CDCl3
 TEMPER 300.0 K
 NS 2048
 DS 4
 SC 1
 FIDRES 0.109618 Hz
 TD 65536
 RG 32768
 DE 6.00 usec
 D1 1.00000000 sec
 F1 10.00 usec
 F11 2.00 dB
 F2 - Processing Parameters
 SI 32768
 SF 300.130039 MHz
 WDW EM
 SSB 0
 GB 0
 PC 1.00
 ID NMR Plot Parameters
 CX 20.00 cm
 FID 1.00000000 sec
 F2P 1.320 ppm

Current Data Parameters
 USER jaf-gm
 NAME 1af-154f
 PROCNO 1
 P2 - Acquisition Parameters
 Date_ 20031003
 Time_ 15.44
 INSTRUM spect
 PROBHD 5 mm QNP 1H
 PULPROG zgpg30
 TD 65536
 TE 300.0 K
 SFO1 282.3762406 MHz
 O1P -100.000 ppm
 AQ -1.091244 sec
 FIDRES 0.143272 Hz
 DW 8.325 usec
 DE 4094
 NS 5
 DS 9
 SC 1.0000000 usec
 DI 0.03000000 sec
 D11 0.0002000 sec
 D12 0.0002000 sec
 ===== CHANNEL f1 =====
 NUCL1 13C
 P1 12.00 usec
 PL1 0.00 dB
 ===== CHANNEL f2 =====
 CDPRG22 waltz16
 NUCL2 1H
 P2 108.00 usec
 PL2 0.00 dB
 P3 108.00 usec
 PL3 0.00 dB
 SFO2 300.1312009 MHz
 P2 - Processing parameters
 SI 65536
 SF 282.404033 MHz
 GB 0
 SB 0
 LB 0.30 Hz
 PC 1.00
 ID NMR plot parameters
 CX 20.00 cm
 FI 1.00
 FIDP -152.898 ppm

```

Current Data Parameters
Date_ 20050109
Time_ 14:22
NMR 1H
EXPNO 1
PROCNO 1
F2 - Acquisition Parameters
Date_ 20050109
Time_ 14:22
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zgpg30
TD 65536
SOLVENT DMS
NS 1638
DS 4
SWH 6009.615 Hz
AQ 0.025888 sec
RG 406.4
AQ 2.726377 sec
RG 406.4
DM 832.25 usec
DE 3600.0 usec
TE 300.0 K
D1 1.0000000 sec
===== CHANNEL f1 =====
NUC1 1H
P1 7.00 usec
PL1 0.00 dB
SFO1 500.3925019 MHz
F2 - Processing parameters
SI 32768
SF 500.3900189 MHz
WDW EM
SSB 0
GB 0
PC 1.00
1D NMR plot: parameters
CX 22.00 cm
FID 6.869 ppm
F1 5.416 ppm
  
```

Current Data Parameters

DU 1
 NM 1
 NMR 1
 EXFNO 1
 PROCNO 1

Date_ 20041215
 Time_ 12:50:00
 INSTRUM 5 mm Dual 13
 PROBD 5 mm Dual 13
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 16
 DS 2
 SNH 3333.332 Hz
 FIDRES 0.15856 Hz
 AQ 0.051000 sec
 RG 16384
 DW 15.000 usec
 DE 16.000 usec
 TE 300.0 K
 D11 2.00000000 sec
 D12 0.40000000 sec
 CPDPRG waltz16
 P31 107.00 usec
 D11 0.03000000 sec
 S2 18 dB
 SFO1 125.832148 MHz
 NUCLEUS 13C

F2 - Processing parameters
 SI 32768
 SF 125.832148 MHz
 WDW EM
 SSB 0
 GB 1.50 Hz
 GC 0
 PC 1.40

1D NMR Plot Parameters
 CX 21.00 cm
 F1P 137.922 ppm
 F2P 130.554 ppm

Current Data Parameters
 USER jg
 NAME 1f-tm-tbfuo
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20050104
 Time 11:00
 INSTRUM spect
 PROBD 5 mm QNP 1H/1
 PULPROG zgpg30
 TD 65536
 SOLVENT Aceton
 DS 14
 SFO1 470.1493 MHz
 SF2 500.132016 MHz
 F2 - Processing Parameters
 SI 32768
 SF 470.837330 MHz
 EQ 1
 AS 0
 GB 0
 PC 1.00
 1D NMR plot parameters
 FID 1
 F1 139.639 ppm
 F2 147.749 ppm

Current Data Parameters
 DU / /
 USER gain
 NAME 1f-1CN
 EXNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20041222
 Time 18.15
 INSTRUM amx500
 PULPROG zgpg30
 PROCNO 5
 FIDRES 0.246110
 TD 32768
 SOLVENT CDCl3
 NS 97
 DS 1
 SWH 8064.51 Hz
 FIDRES 0.246110 Hz
 AQ 2.0316660 sec
 RG 512
 DW 62.000 use
 DE 88.50 use
 TE 300.2 K
 H1 1.00000000 sec
 D1 12.00 use
 SFO1 500.1362518 MHz
 NUCLEUS 1H

F2 - Processing parameters
 SI 32768
 SF 500.1362518 MHz
 WDW EM
 SSB 0
 LB 1.50 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CX 22.00 cm
 F1 2.891 ppm
 F2 2.170 ppm

```

Current Data Parameters
DU      /y
NAME    1f-tfcdm
EXPNO   1
PROCNO  1

F2 - Acquisition Parameters
Date_   20041215
Time    12.00
INSTRUM spect
PROBHD  5 mm QNP 1H/1
PULPROG zgpg30
TD       65536
SOLVENT CDCl3
NS       402
DS       4
SWH      31446.541 Hz
FIDRES   0.44954 Hz
AQ       1.012024 sec
RG       4096
IN       15.50 usec
DE       6.00 usec
TE       300.0 K
D1       2.00000000 sec
d11      0.00000000 sec
d12      0.00000000 sec

===== CHANNEL f1 =====
NUC1     13C
P1       7.50 usec
PL1      0.00 dB
SFO1     125.8363556 MHz

===== CHANNEL f2 =====
CPDPRG2 MALT16
NUC2     1H
P2       100.00 usec
PL2      0.00 dB
PL12     -26.00 dB
PL13     -26.00 dB
SFO2     500.3820000 MHz

F2 - Processing parameters
SF       125.832292 MHz
WDW      EM
SSB       0
LB       1.00 Hz
GB       0
PC       1.40

1D NMR Plot Parameters
CPDPRG2  MALT16
FID      131.460 cm
F1F      137.861 ppm
F2F

```

Current Data Parameters
 USER gam
 NAME 1f-tfcm
 PROCNO 1
 P2 - Acquisition Parameters
 Date_ 20040818
 Time 14:40
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 TD 65536
 SOLVENT C5D6
 DS 4
 SFO 400.1463 MHz
 SWH 19000.000 Hz
 FIDRES 0.000000 Hz
 AQ 0.655100 sec
 RG 327.68
 DE 5.00 usec
 TE 300.2 K
 D1 1.0000000 sec
 D11 0.0300000 sec
 D12 0.0002000 sec
 CHANNEL f1
 NUC1 13C
 P1 10.00 usec
 PL1 -3.00 dB
 SFO1 477.750145 MHz
 CHANNEL f2
 NUC2 1H
 P2 10.00 usec
 PL2 -2.00 dB
 SFO2 500.392016 MHz
 P2 - Processing parameters
 SI 32768
 SF 477.837320 MHz
 MD 32
 EM 1
 GB 0.30 Hz
 PC 1.00
 LD NMR plot parameters
 F1P -128.704 ppm
 F2P -150.000 ppm

2a

Current Data Parameters
 Date_ 20030717
 Time_ 18.36
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 SOLVENT CDCl3
 PULPROG zgpg30
 NUC1 1H
 SFO1 500.3925019 MHz
 Q1P 5.000 Ppm
 AQC 12.000 Ppm
 AQ 2.7263279 sec
 TD 32768
 FIDRES 0.183399 Hz
 DW 83.200 usec
 RG 35.9
 NS 16
 DE 6.00 usec
 D1 1.00000000 sec
 CHANNEL f1
 P1 7.50 usec
 PL1 0.00 dB
 F2 - Processing parameters
 SI 32768
 SF 500.390118 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00
 1D NMR plot parameters
 CX 22.00 cm
 F1P 3.864 Ppm
 F2P 0.265 Ppm

Current Data Parameters
 DU /y
 USER gam
 Machine laf-couesc02
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20030717
 Time 12:00:00
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 SOLVENT CDC13
 TEMP 300.0 K
 PULPROG zgpg30
 NUC1 125.8369556 MHz
 SF01 109.992 ppm
 OL1 249.8991 ppm
 SW 1.042024 sec
 AQ 0.479836 Hz
 FIDRES 15.900 usec
 RG 4096
 DS 14
 NS 130
 DE 6.00 usec
 D1 2.00000000 sec
 D11 0.03000000 sec
 D12 0.0002000 sec

===== CHANNEL f1 =====
 P1 7.50 usec
 PL1 3.00 dB

===== CHANNEL f2 =====
 CHPRG2 waitz16
 P2 100.00 usec
 PCPD2 0.00 dB
 PL2 26.00 dB
 PL12 26.00 dB
 SFO2 500.3920016 MHz

F2 - Processing parameters
 SI 32768
 SF 125.8231675 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CX 52.00 cm
 F1p 60.086 ppm
 F2p -9.067 ppm

Current Data Parameters

DO	/y
USER	gan
NAME	1af-L1E
EXPNO	1
PROCNO	1

F2 - Acquisition Parameters

Date_	20040302
Time	15.42
INSTRUM	spect
PROBHD	5 mm QNP 1H/1
PULPROG	zgpg30
TD	32768
SOLVENT	CDCl3
NS	16
DS	2
SWH	6009.615 Hz
FIDRES	0.183399 Hz
AQ	2.7265128 sec
RG	128
DW	83.200 usec
DE	6.00 usec
TE	300.0 K
D1	1.0000000 sec

CHANNEL f1

NUC1	1H
P1	7.50 usec
PL1	0.00 dB
SFO1	500.3923019 MHz

F2 - Processing Parameters

SI	32768
SF	500.3900248 MHz
WDW	EM
SSB	0
LB	0.30 Hz
GB	0
PC	1.00

1D NMR plot Parameters

CX	22.00 cm
F1P	9.045 ppm
F2P	-0.386 ppm

Current Data Parameters
 Date_ /y
 USER g/y
 NAME 1a-8-E
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040306
 Time_ 15.53

INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg
 TD 65536
 SOLVENT CDCl3

NS 190
 DS 4

SWH 31446.541 Hz
 FIDRES 0.443866 Hz
 AQ 1.6430726 sec

RG 14596.5
 DW 15.900 usec
 DE 6.00 usec

TE 300.0 K
 D1 2.00000000 sec
 D11 0.03000000 sec

D12 0.00020000 sec

===== CHANNEL f1 =====
 NUC1 13C
 P1 7.50 usec
 PL1 3.00 dB
 SFO1 125.8369556 MHz

===== CHANNEL f2 =====
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 26.00 dB
 PL13 26.00 dB
 SFO2 500.3920010 MHz

F2 - Processing parameters
 SI 32768
 SF 125.8231694 MHz

WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0

FC 1.40

1D NMR plot parameters
 CX 22.00 cm

F1P 46.285 ppm
 F2P 28.129 ppm

Current Data Parameters
 Date_ /y
 User g
 Name laf-24E
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040226
 Time_ 18:05
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 6009.615 Hz
 FIDRES 0.183159 Hz
 AQ 2.726377 sec
 RG 57
 DW 83.200 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.0000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.50 usec
 PL1 0.00 dB
 SFO1 500.3925019 MHz

F2 - Processing parameters
 SI 32768
 SF 500.3900159 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 22.00 cm
 F1P 10.973 ppm
 F2P -1.037 ppm

Current Data Parameters

DU /y
 USER gam
 INSTR 1af-
 EXFNO 2
 PROCNO 1

F2 - Acquisition Parameters

Date_ 20091220
 Time_ 18:11
 INSTRUM spect
 PROBD 5 mm QNP 1H/1
 FULPROG zgpg
 TD 65536
 SOLVENT CDCl3
 NS 119
 DS 4
 SWH 31446.541 Hz
 FIDRES 0.479836 Hz
 AQ 0.021724 sec
 EQ 102174
 RZ 15.900 use
 DE 6.00 use
 TE 300.0 K
 D1 2.0000000 sec
 D11 0.0000000 sec
 D12 0.0002000 sec

CHANNEL f1

NUC1 13C
 P1 7.50 use
 PL 0
 SFO1 125.8369556 MHz

CHANNEL f2

waitz16
 NUC2 1H
 PCPD2 100.00 use
 PL2 0.00 dB
 PL12 26.00 dB
 PL13 26.00 dB
 SFO2 500.3920016 MHz

F2 - Processing Parameters

SI 32768
 SF 125.8231732 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot Parameters

CH 42.00 cm
 F1P 47.932 ppm
 F2P 20.702 ppm

Current Data Parameters
 DU /Y
 USER gan
 NAME lf-ezrew
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20050201
 Time 16.12
 INSTRUM spect
 PULPROG zgpg30
 FIDRES 0.183399 Hz
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 4
 SWH 6009.612 Hz
 FIDRES 0.183399 Hz
 AQ 2.7263477 sec
 RG 64
 DW 83.200 use
 DE 300.00 use
 TE 300.2 K
 D1 1.00000000 sec
 CHANNEL f1
 NUC1 1H
 P1 7.50 use
 PL1 0.00 dB
 SF01 500.13925019 MHz
 F2 - Processing Parameters
 SI 32768
 SF 500.13906519 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00
 1D NMR plot parameters
 CX 22.00 cm
 FIP 3.591 ppm
 F2P 1.736 ppm

```

Current Data Parameters
Date_ 20050201
Time_ 16.19
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 116
DS 4
SWH 31446.541 Hz
FIDRES 0.063000 Hz
AQ 1.042024 sec
RG 1024
DW 15.900 use
DE 6.00 use
TE 300.2 K
D1 2.0000000 sec
D11 0.0300000 sec
D12 0.0002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 7.50 use
PL1 3.00 dB
SFO1 125.8369556 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 100.00 use
PL2 0.00 dB
PL12 26.00 dB
PC12 26.00 dB
SFO2 500.3920016 MHz

F2 - Processing parameters
SI 32768
SF 125.8231752 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
FC 1.40

1D NMR plot parameters
CX 22.00 cm
F1P 45.491 ppm
F2P 28.711 ppm

```

```

Current Data Parameters
=====
DU      /
USER    gam
NAME    laf-174
EXPNO   1
PROCNO  1

F2 - Acquisition Parameters
=====
Date_   20031030
Time    17.22
INSTRUM spect
PROBHD  5 mm QNP
PULPROG zgpg30
TE      300.0 K
NUC1     1H
SFO1    500.3925019 MHz
CL1     5 mm QNP
SM      12.0098
AQ      2.7263477 sec
TD      32768
FIDRES  0.183399 Hz
RG      81.200 usec
DS      40.2
NS      16
DE      6.00 usec
D1      1.00000000 sec

===== CHANNEL f1 =====
P1      7.50 usec
PL1     0.00 dB

F2 - Processing Parameters
=====
SI      32768
SF      500.3900759 MHz
WDW     EM
SSB     0
LB      0.30 Hz
GB      0
PC      1.00

1D NMR plot parameters
=====
CX      22.00 cm
F1P     7.678 ppm
F2P     4.716 ppm
  
```

Current Data Parameters

DU	/y
USBR	gam
NAME	lar-17
EXNO	2
PROCNO	1

F2 - Acquisition Parameters

Date_	20031030
Time	11:30
INSTRUM	5 mm QNP 1H/1
PROBHD	CDC13
SOLVENT	300.0 K
TE	300.2
TEMPROG	zgpg30
NUC1	13C
SFO1	125.8369556 MHz
Q1P	109.992 ppm
SW	249.8991 ppm
AQ	-0.0420724 sec
RG	79836
FIRES	0.479836 Hz
DW	15.900 use
RG	9195.2
DS	4
NS	64
DE	6.00 use
D1	2.00000000 sec
D11	0.03000000 sec
D12	0.00020000 sec

CHANNEL f1

P1	7.50 use
PL1	3.00 dB

CHANNEL f2

NUC2	13C
PROG2	waltz16
PCPD2	100.00 use
PL2	0.00 dB
PL12	26.00 dB
PL13	26.00 dB
SFO2	500.3920016 MHz

F2 - Processing Parameters

SI	32768
SF	125.8231819 MHz
WDW	EM
SSB	0
LB	1.00 Hz
GB	0
PC	1.40

1D NMR Plot Parameters

CX	92.00 cm
F1P	149.690 ppm
F2P	107.784 ppm

Current Data Parameters
NAME lsf-133
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030818
Time 13:27
INSTRUM spect
PROBHD 5 mm QNP 1H/13
SOLVENT CDCl3
TE 300.0 K
PULPROG zg30
NUC1 13C
NUC2 1H
F1 500.1325018 MHz
F2 500.1325018 MHz
O1P 5.000 ppm
SW 12.0098 ppm
AQ 2.7263477 sec
TD 32768
FIDRES 0.13279 Hz
AQ 2.7263477 sec
RG 640.3
DS 2
NS 16
DE 6.00 usec
DI 1.00000000 sec

===== CHANNEL f1 =====
F1 500.1325018 MHz
F2 500.1325018 MHz
SF 500.1325018 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

ID NMR plot parameters
CX 22.00 cm
CY 22.00 cm
F1 4503.51 Hz
F2 -0.500 ppm
F2 -250.20 Hz
PPMCM 0.43182 ppm/cm
HZCM 216.07751 Hz/cm

Current Data Parameters
 DU /y
 USR 9am
 NMR 1a-152
 EXFNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20010816
 Time_ 13:33
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 SOLVENT CDCl3
 TEMP 300.0 K
 PULPROG zgpg30
 NUC1 13C
 SF01 125.8369556 MHz
 O1P 109.992 Ppm
 SW 249.8991 Ppm
 AQ 1.042574 sec
 TQ 0.042574 sec
 FIDRES 0.479836 Hz
 DWD 15.900 usec
 RG 4096
 DS 4
 Ns 64
 DE 6.00 usec
 D1 2.00000000 sec
 D11 0.03000000 sec
 D12 0.00020000 sec

CHANNEL f1
 P1 7.50 usec
 PL1 3.00 dB

CHANNEL f2
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 26.00 dB
 PL13 26.00 dB
 SFO2 500.3920016 MHz

F2 - Processing parameters
 SI 32768
 SF 125.8231761 MHz
 NMR EN
 SS 1.00 Hz
 LB 0
 GB 0
 FC 1.40

1D NMR plot parameters
 CX 22.00 cm
 F1P 48.340 Ppm
 F2P 7.819 Ppm

Current Data Parameters
 DU 1
 INSTR 5
 NAME 1af-125
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20010915
 Time_ 15:53
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 SOLVENT CDCl3
 PULPROG zgpg30
 NUC1 1H
 SF01 500.3925019 MHz
 O1P 5.000 ppm
 SW 12.0098 ppm
 FIDRES 0.183399 Hz
 TD 32768
 DE 1.00000000 sec
 D1 1.00000000 sec
 CHANNEL f1
 F1 500.136125 MHz
 F11 0.00 dB
 F2 - Processing parameters
 SI 32768
 SF 500.3900330 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 EC 1.00
 ID NMR plot parameters
 CX 22.00 cm
 F1P 3.217 ppm
 F2P 0.741 ppm

Current Data Parameters
 DU /Y
 NAME 1af-125
 EXPNO 2
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20030802
 Time 16.00
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 SOLVENT CDCl3
 PULPROG zgpg30
 NUC1 13C
 SF01 125.8369556 MHz
 O1P 109.992
 SW 249.8921 ppm
 FIDRES 0.479836 Hz
 TD 65536
 D1 2.0000000 sec
 DE 6.00 use
 D11 0.0300000 sec
 D12 0.0002000 sec
 CHANNEL f1
 P1 7.50 use
 PL1 3.00 dB
 CHANNEL f2
 CPDPRG2 Wait
 NUC2 1H
 FCFD2 100.00 use
 PL2 0.00 dB
 FID2 26.00 dB
 SFO2 500.3920016 MHz
 F2 - Processing parameters
 SI 32768
 SF 125.8231752 MHz
 WPM 0
 SSB 0
 LB 1.00 Hz
 GB 0
 FC 1.40
 ID NMR plot parameters
 CX 22.00 cm
 FLP 51.224 ppm
 FZP -4.817 ppm

Current Data Parameters
 NAME lf-alcohol
 EXPNO 1
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040802
 Time_ 18:24
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 6009.615 Hz
 FIDRES 0.183379 Hz
 AQ 2.726577 sec
 RG 181
 DW 83.200 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.0000000 sec

CHANNEL f1
 NUC1 1H
 P1 7.50 usec
 PL1 0.00 dB
 SFO1 500.392019 MHz
 F2 - Processing parameters
 SI 32768
 SF 500.3900541 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 EC 1.00
 ID NMR Plot Parameters
 CX 22.00 cm
 FIP 9.000 ppm
 F1 4503.51 Hz
 F2P -0.500 ppm
 F3 233.10 Hz
 F3P 0.000 ppm
 F4 216.07751 Hz/cm
 F4P 0.000 ppm

```

Current Data Parameters
DU / /
USER gam
NAME 1f-alcohol
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040802
Time 18.28
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 108
DS 4
SWH 31446.54 Hz
AQ 0.479836 Hz
FIDRES 1.0420724 sec
RG 4096
DM 15.900 usec
DE 3.00 usec
TE 300.2 K
D1 2.00000000 sec
D11 0.03000000 sec
D12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 7.50 usec
PL1 3.00 dB
SFO1 125.8369556 MHz

===== CHANNEL f2 =====
CDEPRG2 waltz16
NUC2 1H
PCPD2 100.00 usec
PL2 0.00 dB
PL12 26.00 dB
PL13 2.00 dB
SFO2 500.3320016 MHz

F2 - Processing parameters
SI 32768
SF 500.136199 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
FC 1.40

1D NMR plot parameters
CX 22.00 cm
F1P 66.551 ppm
F2P 27.509 ppm

```

```

Current Data Parameters
DU          /Y
USER       gam
NAME       laf-cetecoup
EXPNO      1
PROCNO     1

F2 - Acquisition Parameters
Date_      20030905
Time       18.48
INSTRUM    spect
PROBHD     5 mm QNP
SOLVENT    CDCl3
TE         300.0 K
PULPROG    zg30
NUC1        1H
SFO1       500.39500 MHz
O1P         5.000 ppm
SW          12.0098 ppm
AQ          2.7263477 sec
TD          32768
FIDRES      0.183399 Hz
RG          819.5
DS          2
NS          16
DE          6.00 usec
D1          1.00000000 sec

===== CHANNEL f1 =====
F1          7.50 usec
PL1         0.00 dB

F2 - Processing Parameters
SI          32768
SF          500.3950198 MHz
WDW         EM
SSB         0
RB          0.30 Hz
GB          0
PC          1.00

1D NMR plot parameters
CX          22.00 cm
FIDP        3.266 ppm
F2P         1.661 ppm

```

```

Current Data Parameters
NAME      1f-ai300ai
EXPNO     1
PROCNO    1

F2 - Acquisition Parameters
File_     2001223
Time      17.02
INSTRUM   ax500
PROBHD    5 mm Dual 13
PULPROG   zg
TD         32768
SOLVENT   CDCl3
NS         38
DS         1
SWH        8064.316 Hz
FIDRES     0.246110 Hz
AQ         2.031660 sec
RG         62.000 usec
DE         88.57 usec
TE         300.0 K
HL1        1.000000000
NUC1       13
P1         12.00 usec
SFO1       500.392518 MHz
NUCLEUS    13

F2 - Processing parameters
SI         32768
SF         500.390240 MHz
WDW        EM
SSB        0
LB         1.50 Hz
GB         0
PC         1.40

1D NMR plot parameters
CX         22.00 cm
F1P        62.000 ppm
F1         60.500 ppm
F2P        -250.19 Hz
F2         0.56818 ppm/cm
PPMCM      284.31250 Hz/cm
  
```

Current Data Parameters
 DU USER gam /y
 NAME lf-al
 EXPNO 2
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20041209
 Time 17.08
 INSTRUM ax500
 PULPROG zgpg30
 FIDRES 5 mm Dual 3
 TD 65536
 SOLVENT CDCl3
 NS 47
 DS 2
 SWH 33333.322 Hz
 FIDRES 0.598626 Hz
 AQ 0.9830900 sec
 RG 16384
 DW 15.000 usec
 DE 18.70 usec
 TE 300.2 K
 HL1 40 dB
 D1 2.00000000 sec
 CPDPRG waltz16
 F31 197.00 usec
 SFO1 125.832148 MHz
 D11 0.0300000 sec
 S2 18 dB
 F1 5.00 usec
 SFO1 125.832148 MHz
 NUCLEUS 13C
 F2 - Processing parameters
 SI 32768
 SF 125.8231716 MHz
 WDW EM
 GB 0
 PC 1.40
 ID NMR plot Parameters
 CW 45.761 ppm
 F1P 25.166 ppm
 F2P

```

Current Data Parameters
DU          /y
USER        gam
NAME        _festdppf
EXPNO       1
PROCNO      1

F2 - Acquisition Parameters
Date_       20041001
Time        8.26
PULPROG     zgpg30
PROBHD      5 mm Durr 13
PULPROG     zg
TD           32768
SOLVENT     CDCl3
NS           39
DS           4
SWH          8064.516 Hz
FIDRES       0.246110 Hz
AQ           2.031660 sec
RG           256
RGW          62.000 use
DE           68.57 use
TE           300.0 K
TE          300.0 K
HL1          1 dB
D1           1.0000000 sec
FL           12.00 use
SFO1         500.3922518 MHz
NUCLEUS      1H

F2 - Processing Parameters
SI           32768
SF           500.3900696 MHz
WDW          EM
SSB          0
LB           1.50 Hz
GB           0
PC           1.40

1D NMR plot parameters
CX           22.00 cm
F1P          4.165 ppm
F2P          1.944 ppm

```

Current Data Parameters
 DI /
 USER gam
 NAME -festdppf
 EXPNO 2
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20041001
 Time 8:33
 INSTRUM amx500
 PROCDP Dual 13
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 94
 DS 2
 SWH 3333.322 Hz
 FWHM 0.506626 Hz
 AQ 0.9830900 sec
 RG 32768
 DW 15.000 use
 DE 18.75 use
 TE 300.2 K
 H1 40 dB
 D1 2.00000000 sec
 CPDPRG waitz16
 F31 107.00 use
 SFO1 125.76148 MHz
 D11 0.0300000 sec
 S2 18 dB
 F1 5.00 use
 SFO1 125.8382148 MHz
 NUCLEUS 13C
 F2 - Processing parameters
 SI 32768
 SF 125.8231726 MHz
 WDW EM
 SSB 0
 GB 1.50 Hz
 PC 1.40
 1D NMR plot parameters
 CK 0
 F1P 61.238 ppm
 F2P 14.605 ppm

2l

Current Data Parameters
 DU /Y
 USER gam
 NAME laf-ncis
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20040105
 Time 19.18
 INSTRUM spect
 PULPROG zgpg30
 SOLVENT CDCl3
 TE 300.0 K
 NUC1 1H
 NUC2 13C
 SW 12.0098 EPM
 AQ 2.7263477 sec
 TD 32768
 FIDRES 0.183399 Hz
 SFO 500.136050 MHz
 RG 327.5
 DS 2
 NS 16
 DE 6.00 usec
 D1 1.00000000 sec
 CHANNEL f1
 PL 7.50 usec
 PL1 0.00 dB
 F2 - Processing Parameters
 SI 32768
 SF 500.1360240 MHz
 WDW EM
 DB 0
 SSB 0.30 Hz
 GC 0
 PC 1.00
 1D NMR plot parameters
 CX 22.00 cm
 FID 1.012 EPM
 F2P 1.552 EPM

```

Current Data Parameters
DU / /
USER gam
NAME 1a-172
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20031024
Time 18.45
INSTRUM spect
PROBHD 5 mm QNP
PULPROG zgpg30
TE 300.0 K
NUC1 1H
SFO1 500.3925019 MHz
C1P1 5.0000000 mm
SW 12.0098 kHz
AQ 2.7263477 sec
TD 32768
FIDRES 0.183399 Hz
RG 83.200 usec
DS 40.2
DE 16
D1 1.00000000 sec

===== CHANNEL f1 =====
P1 7.50 usec
PL1 0.00 dB

F2 - Processing parameters
SI 32768
SF 500.3961000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 22.00 cm
FIP 1.060 ppm
FZF 1.965 ppm

```

```

Current Data Parameters
=====
Date_   20031026
Time_   18.48
INSTRUM spect
PROBHD  5 mm QNP 1H/1
SOLVENT CDCl3
PULPROG zgpg30
NUC1     13C
SF01     125.836956 MHz
O1P      100.992 Ppm
RG        4096
AQ        1.0420721 sec
TD        65536
FIDRES    0.479836 Hz
DM        15.900 usec
DE        6.00 usec
D1        2.00000000 sec
D11       0.03000000 sec
D12       0.00020000 sec
=====
CHANNEL f1
P1        7.50 usec
PL1       3.00 dB
=====
CHANNEL f2
=====
CPDPRG2 waltz16
NUC2      1H
PCPD2     100.00 usec
PL2       0.00 dB
PL12      26.00 dB
PL14      2.00 dB
SF02     500.3920016 MHz
=====
F2 - Processing parameters
SI        32768
SF         500.392 MHz
WDW        EM
SSB        0
LB         1.00 Hz
GB         0
FC         1.40
=====
1D NMR plot parameters
CX         22.00 cm
F1P        52.132 Ppm
F2P        -2.768 Ppm
  
```

Current Data Parameters
 Date_ 20041209
 Time_ 16.27
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 6009.615 Hz
 FIDRES 0.18277 Hz
 AQ 2.7263477 sec
 RG 40.3
 DW 83.200 use
 DE 6.00 use
 TE 300.0 K
 D1 1.00000000 sec

CHANNEL f1
 NUC1 1H
 P1 7.50 use
 PL1 0.00 dB
 SFO1 500.3925019 MHz

F2 - Processing parameters
 SI 32768
 SF 500.3900635 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR plot parameters
 CX 22.00 cm
 F1P 3.059 ppm
 F2P 0.425 ppm

Current Data Parameters

DU	/y
NAME	gam
EXPNO	1f--t1p3
PROCNO	1

F2 - Acquisition Parameters

Date_	20050220
Time	16:30
INSTRUM	spect
PROBHD	5 mm QNP 1H/1
PULPROG	zgpg
TD	65536
SOLVENT	CDCl3
NS	83
DS	4
SWH	31446.541 Hz
FIDRES	0.479836 Hz
AQ	1.0422000 sec
RG	4096
DW	15.900 use
DE	6.00 use
TE	300.0 K
D1	2.0000000 sec
D11	0.0000000 sec
D12	0.0002000 sec

CHANNEL f1

NUC1	¹³ C
PL1	7.00 use
PL11	3.00 dB
SFO1	125.8369556 MHz

CHANNEL f2

CPDPRG2	waltz16
NUC2	¹³ C
PCPD2	100.00 use
PL2	0.00 dB
PL12	26.00 dB
PL13	26.00 dB
SFO2	500.3920016 MHz

F2 - Processing Parameters

SI	32768
SF	125.8231704 MHz
WDW	EM
SSB	0
LB	1.00 Hz
GB	0
PC	1.40

1D NMR plot Parameters

CH	5.00 cm
F1P	65.218 ppm
F2P	5.725 ppm

wdd

```

Current Data Parameters
ID      /f
NAME    1af-177
EXPNO   2
PROCNO   1

F2 - Acquisition Parameters
Date_    20041022
Time     9.14
INSTRUM  spect
PROBHD   5 mm QNP 1H/1
PULPROG  zgpg
SOLVENT  CDCl3
NS        42
DS         4
SWH       31446.541 Hz
FIDRES    0.0001000 sec
AQ         1.6420734 sec
RG         4096
DM         15.900 usec
DE         8.00 usec
TE        300.2 K
D1         2.0000000 sec
D11        0.0300000 sec
D12        0.0000000 sec

===== CHANNEL f1 =====
NUC1      13C
P1         7.50 usec
PL1        3.00 dB
SFO1      125.8316556 MHz

===== CHANNEL f2 =====
CPDPRG2  waltz16
NUC2       1H
PCPD2     100.00 usec
PL2        0.00 dB
PL12       2.00 dB
PL13       2.00 dB
PL14       2.00 dB
SFO2      500.3920016 MHz

F2 - Processing parameters
SI         32768
SF         125.831656 MHz
WDW         EM
SSB         0
LB          1.00 Hz
GB          0
PC          1.40

1D NMR plot parameters
CX         22.00 cm
F1F        180.000 ppm
F2F        -18.000 ppm
  
```

Current Data Parameters
 DO /y
 NAME 1a-221
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20040222
 Time_ 14.02
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zg30
 FIDRES 0.163
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 600.615 Hz
 FIDRES 0.163399 Hz
 AQ 2.725600 sec
 RG 90.5
 DW 89.200 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.00000000 sec
 CHANNEL f1
 NUC1 1H
 P1 7.50 usec
 PL1 0.00 dB
 SFO1 500.392019 MHz
 F2 - Processing parameters
 SI 32768
 SF 500.3900251 MHz
 WDM EM
 LB 0.30 Hz
 GB 0
 PC 1.00
 ID NMR plot parameters
 CX 22.00 cm
 FIP 4.088 ppm
 FZP 0.251 ppm

```

Current Data Parameters
INSTRUM  /f/
NAME     14f-221
PROCNO   1
=====
F2 - Acquisition Parameters
Date_    20040122
Time     14:15
INSTRUM  5 mm QNP 1H/1
PROBHD   5 mm QNP 1H/1
PULPROG  zgpg30
TD        65536
SOLVENT  CDCl3
AQ        0.14000000
DS        4
SS        31445.44 Hz
FIDRES    0.472834 Hz
AQ        1.0420724 sec
RG        327.68
DM        15.8000 us
DE        6.000 us
TE        300.2 K
D1        2.00000000 sec
d11       0.04000000 sec
d12       0.00002000 sec
===== CHANNEL f1 =====
NUC1      13C
P1        1.30 us
PL1       -1.00 dB
SFO1      125.830556 MHz
===== CHANNEL f2 =====
CHSPRO2   wait 18
NUC2      1H
P2        100.00 us
PL2       -1.00 dB
SFO2      500.332015 MHz
===== Processing parameters =====
SI        32768
SF         125.831094 MHz
WDW        EM
SSB        0
GB         0
PC         1.40
=====
1D 1H NMR plot parameters
CHRG1     64.810 PF
F2P       1.139 PF

```

```

Current Data Parameters
NAME      lufrvyk1
EXPNO     1
PROCNO    1

F2 - Acquisition Parameters
Date_     20031226
Time      9.26
INSTRUM   spect
PROBHD    5 mm QNP 1H/1
SOLVENT   CDCl3
F1F2      500.0 K
PULPROG   zgpg30
NUC1       1H
SF01      500.3925019 MHz
O1P        5.000 ppm
SWH        12.0098 ppm
AQ         2.7232768 sec
RG          0.183399 Hz
FIRES      83.200 use
D1         181
US         12
DS         6.00 use
DE         1.00000000 sec

===== CHANNEL f1 =====
F1         500.136 MHz
P1         12.00 use
PL1        0.00 dB

F2 - Processing parameters
SI         32768
SF         500.3900156 MHz
WDW        EM
SSB        0
LB         0.30 Hz
GB         0
FC         1.00

1D NMR Plot Parameters
CX         20.00 cm
F1P        10.000 ppm
F1         5003.90 Hz
F2P        -5.000 ppm
F2         0.55000 ppm
PFGMCH     275.21451 Hz/
HZCM

```

```

Current Data Parameters
=====
USER      X
NAME      raf
EXPNO     2
PROCNO    1

F2 - Acquisition Parameters
=====
Date_     20031209
Time      9.31
INSTRUM   spect
PROBHD    5 mm QNP 1H/1
PULPROG   zgpg
TD         65536
SFO1      125.8369556 MHz
NUC1       13C
SFO2      100.6281500 MHz
NUC2       1H
AQ         0.0400000 sec
RG         32768
DE         6.00 usec
D1         2.00000000 sec
D11        0.03000000 sec
D12        0.00020000 sec

=====
CHANNEL f1
P1          7.50 usec
PL1         3.00 dB

=====
CHANNEL f2
=====
=====
PULPROG2   waltz16
NUC2        1H
PCPD2       100.00 usec
PL2         0.00 dB
PL12        26.00 dB
PL13        26.00 dB
SFO2        500.3920016 MHz

F2 - Processing parameters
=====
SI          32768
SF          125.8231170 MHz
WDW         EM
SSB         0
LB          1.00 Hz
GB          0
PC          1.40

ID NMR plot parameters
=====
CX          22.00 cm
F1P         133.657 ppm
F2P         119.018 ppm
  
```

Current Data Parameters
 DU /
 USER gam
 NAME lafi45
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Param
 Date_ 20030923
 Time 10.45
 INSTRUM spect
 PULPROG zgpg30
 SOLVENT CDCl3
 TE 300.0
 FULPROG zg30
 NUC1 1H
 SCAL 500.3925019
 CH1 500.1360888
 SW 12.0098
 AQ 2.7263477
 TD 32768
 FIDRES 0.183399
 RM 83200
 DS 500.2
 NS 16
 DE 6.00
 D1 1.00000000
 CHANNEL f1
 P1 7.50
 PL1 0.00
 F2 - Processing Paramet
 SI 32768
 SF 500.3900143
 WDW EM
 SSB 0
 LB 0.30
 GB 0
 PC 1.00
 ID NMR plot Parameters
 CX 22.00
 FIP 4.177
 FZF 0.622

Current Data Parameters
NAME jmf142
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20080224
Time 10.54
INSTRUM spect
PROBHD 5 mm QNP 1H/1
SOLVENT CDCl3
FL 300.0
F2PRG2 zgpg30
NUC1 13C
SFO1 125.8369556
O1P 109.992
SW 249.8991
AQ 0.02000000
TD 65536
FIDRES 0.479836
DW 15.900
RG 4096
DS 4
SS 14
DE 6.00
D1 2.00000000
D11 0.03000000
D12 0.00020000

===== CHANNEL f1 =====
P1 7.50
PL1 3.00

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 13C
PCPD2 100.00
PL2 0.00
PL12 26.00
PL13 26.00
SFO2 500.3920016

F2 - Processing parameters
SI 32768
SF 125.8211675
WDW EM
SSB 0
LB 1.00
GB 0
FC 1.40

```

Current Data Parameters
DU /
USER gam
NAME lf-gsty-
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20040901
Time 10.33
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 16
DS 2
SWH 6009.61 Hz
FIDRES 0.161399 Hz
AQ 2.7263477 sec
RG 90.5
DM 83.200 usec
DE 6.00 usec
TE 300.2 K
D1 1.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.00 usec
PL1 0.00 dB
SFO1 500.3925019 MHz

F2 - Processing parameters
SI 32768
SF 500.3900244 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 22.00 cm
F1P 7.766 ppm
F2P 6.875 ppm

```

Current Data Parameters
 Date_ /y
 INSTR 1f-9styf
 NAME 1f-9styf
 EXFNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040901
 Time_ 10.38
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg
 TD 65536
 SOLVENT CDCl3
 NS 57
 DS 4
 SWH 31446.541 Hz
 FIDRES 0.448936 Hz
 AQ 1.042026 sec
 RG 2048
 DW 15.900 usec
 DE 6.00 usec
 TE 300.0 K
 D1 2.0000000 sec
 D11 0.0300000 sec
 D12 0.0002000 sec

CHANNEL f1
 NU1 13C
 PCPD1 7.00 usec
 PL1 3.00 dB
 SF01 125.8369556 MHz

CHANNEL f2
 CPDPRG2 waltz16
 NU2 1H
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 26.00 dB
 PL13 26.00 dB
 SF02 500.3920016 MHz

F2 - Processing parameters
 SI 32768
 SF 125.821742 MHz
 WDM 8
 SS 0
 LB 1.00 Hz
 GB 0
 FC 1.40

1D NMR Plot Parameters
 CX 22.00 cm
 F1P 153.486 ppm
 F2P 110.991 ppm

```

Current Data Parameters
Date_ 20040727
Time_ 12.47
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zgpg30
TD 32768
SOLVENT CDCl3
NS 16
DS 2
SWH 600.612 Hz
FIDRES 0.18399 Hz
AQ 2.7263477 sec
RG 181
DE 83.200 usec
TE 300.2 K
D1 1.00000000 sec

CHANNEL f1
NUC1 1H
P1 7.50 usec
PL1 0.00 dB
SFO1 500.3925019 MHz

F2 - Processing parameters
SI 32768
SF 500.3902200 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
FC 1.00

1D NMR plot parameters
CX 22.00 cm
F1P 4.126 ppm
F2P 2.138 ppm

```

1D NMR plot parameters	
CX	21.00 cm
F1P	68.894 ppm
F2P	28.580 ppm

ppm

Current Data Parameters
 IN 1
 NAME 1af-170
 EXNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20031028
 Time 14.32
 INSTRUM spect
 PROCNO 5
 PULPROG zgpg30
 NS 16
 DS 2
 DE 6.00 usec
 DI 1.0000000 sec
 ===== CHANNEL f1 =====
 P1 7.50 usec
 PL 0.00 dB
 F2 - Processing parameters
 SI 32768
 SF 500.360382 MHz
 WDW EM
 SSB 0
 GB 0.30 Hz
 PC 1.00
 ID NO plot parameters
 X 2.00 cm
 Y 5.00 cm
 Z 0.00 cm
 P1 0.00 sec
 P2 -0.500 ppm

wdc

```

Current: Data Parameters
=====
NAME      1af-170
EXPNO     2
PROCNO    1

F2 - Acquisition Parameters
=====
Date_     20031021
Time      14.38
INSTRUM   spect
PROBHD    5 mm QNP1
PULPROG   zgpg
PC        300.0 K
NUC1       13C
SFO1      125.816558 MHz
AQ        1.042074 sec
SI        246.8991 Dm
TD         65536
FIDRES    0.495838 Hz
AQ        15.0000 usec
RG         6094
DS         4
NS         103
DS         6.00 usec
DE         0.00000000 sec
D11        0.03000000 sec
D12        0.00020000 sec

===== CHANNEL f1 =====
F1         125.760 MHz
P1         1.00 dB

===== CHANNEL f2 =====
CPDPRG2   waltz16
NUC2       1H
SFO2       500.136054 MHz
AQ        1.0000 usec
SI         100.00 Dm
P1         0.00 dB
P2         24.00 dB
P3         24.00 dB
SFO2      500.392015 MHz

F2 - Processing parameters
=====
SI         32768
SF         125.831815 MHz
WDW        EM
SSB         0
LB          1.00 Hz
GB          0
PC         1.40

1D NMR plot parameters
=====
CX         22.00 cm
F1P        220.000 ppm
F2P        -10.000 ppm
  
```

ppm

Current Data Parameters
 F1 32768
 F2 32768
 NAME 1a2-171
 EXPR 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20031022
 Time 10:33
 INSTRUM spect
 PULPROG zgpg30
 TE 300.0 K
 NUC1 1H
 SFO1 500.3390045 MHz
 C1P1 5.0000000 mm
 SFO2 500.3390045 MHz
 C2P1 5.0000000 mm
 AQ 2.7261477 sec
 TD 32768
 FIDRES 0.183395 Hz
 AQ 2.7261477 sec
 RG 40.0
 OS 2
 NS 16
 DS 6.00 usec
 DE 1.00000000 sec
 DI 1.00000000 sec
 ===== CHANNEL f1 =====
 P1 7.50 usec
 P2 0.00 dB
 F2 - Processing parameters
 SI 32768
 SF 500.3390045 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00
 ID NP plot parameters
 C1P1 5.0000000 mm
 F1P 9.0000000 MHz
 F2P -0.5000000 MHz

Integral

ppm

Current Data Parameters
 Date_ 20040524
 Time_ 15.14
 INSTRUM spect
 PROBD 5 mm QNP 1H/1
 PULPROG zgpg30
 TD 32768
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 600.635 Hz
 FIDRES 0.11339 Hz
 AQ 2.7263477 sec
 RG 90.5
 DW 83.200 usec
 DE 6.00 usec
 TE 300.2 K
 D1 1.0000000 sec

===== CHANNEL f1 =====
 NUC1 1H
 P1 7.50 usec
 PL1 0.00 dB
 SFO1 500.3925019 MHz

F2 - Processing parameters
 SI 32768
 SF 500.390081 MHz
 WDW EM
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00

1D NMR Plot Parameters
 CX 22.00 cm
 FLIP 3.993 ppm
 FZP 1.324 ppm

Current Data Parameters
 DU /y
 USR gam
 MPR 1f-02
 EXNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040513
 Time_ 17:43
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 ID 65536
 SOLVENT CDCl3
 NS 115
 DS 4
 SWH 31446.541 Hz
 FIDRES 0.479836 Hz
 AQ 1.0420364 sec
 RG 655
 DW 15.900 usec
 DE 6.00 usec
 TE 300.0 K
 D1 2.0000000 sec
 d11 0.0500000 sec
 D12 0.0002000 sec

CHANNEL f1
 NUC1 13C
 P1 7.50 usec
 PL1 0.00 dB
 SFO1 125.8369556 MHz

CHANNEL f2
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 26.00 dB
 PL13 26.00 dB
 SFO2 500.3920016 MHz

F2 - Processing parameters
 SI 32768
 SF 125.8231694 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CK 72.00 cm
 F1P 76.340 ppm
 F2P 12.914 ppm

Current Data Parameters
 NS 1
 USER Y
 NAME ga
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20040525
 Time 14.05
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 TOUFRQG 32768
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 6009.615 Hz
 FIDRES 0.125397 Hz
 AQ 2.7263477 sec
 RG 90.5
 DW 83.200 usec
 DE 6.00 usec
 TE 300.0 K
 D1 1.00000000 sec
 CHANNEL f1
 NUC1 1H
 P1 7.50 usec
 PL 0.00 dB
 SFO1 500.392019 MHz
 F2 - Processing parameters
 SI 32768
 SF 500.3900180 MHz
 SE 32768
 ESF0 0
 SSB 0
 LB 0.30 Hz
 GB 0
 PC 1.00
 1D NMR Plot Parameters
 CX 22.00 cm
 F1P 7.689 ppm
 F2P 6.613 ppm

Current Data Parameters
 DU / y
 NAME 1f-08
 EXPNO 2
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20080310
 Time 14:11
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg30
 FIDRES 0.5516
 SOLVENT CDCl3
 NS 76
 DS 4
 SWH 31446.541 Hz
 FIDRES 0.479836 Hz
 AQ 1.0472544 sec
 RG 15.900 usec
 DE 6.00 usec
 TE 300.0 K
 D1 2.0000000 sec
 D11 0.0000000 sec
 D12 0.0002000 sec

CHANNEL f1
 NUC1 13C
 P1 7.00 usec
 PL1 3.00 dB
 SFO1 125.8369556 MHz

CHANNEL f2
 CPDPRG2 waltz16
 NUC2 1H
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 26.00 dB
 PL13 26.00 dB
 SFO2 500.3920016 MHz

F2 - Processing parameters
 SI 32768
 SF 125.8231713 MHz
 WDW EM
 SSF 0
 LB 1.00 Hz
 GB 0
 FC 1.40

1D NMR plot parameters
 CX 52.00 cm
 F1p 153.010 ppm
 F2p 126.462 ppm

```

Current Data Parameters
Date_ 20040904
Time_ 17.10
INSTRUM spect
PROBHD 5 mm QNP 1H/1
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 600.613 Hz
FIDRES 0.13199 Hz
AQ 2.7263477 sec
RG 128
DE 83.200 usec
TE 300.2 K
D1 1.00000000 sec

===== CHANNEL f1 =====
NUC1 1H
P1 7.50 usec
PL1 0.00 dB
SFO1 500.3925019 MHz

F2 - Processing parameters
SI 32768
SF 500.390016 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 22.00 cm
F1P 2.858 ppm
F2P 2.185 ppm

```

Current Data Parameters
 DU / y
 USER sm
 NAME lf-cmew
 EXPNO 4
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20040907
 Time 17.15
 INSTRUM spect
 PROBHD 5 mm QNP 1H/1
 PULPROG zgpg
 TD 65536
 SOLVENT CDCl3
 NS 90
 DS 4
 SWH 31446.541 Hz
 FIDRES 0.479536 Hz
 AQ 1.042520 sec
 RG 2048
 DW 15.900 usec
 DE 6.00 usec
 TE 300.0 K
 D1 2.000000 sec
 D11 0.030000 sec
 D12 0.0002000 sec

CHANNEL f1
 NUC1 13C
 P1 7.00 usec
 PL1 3.00 dB
 SFO1 125.8369556 MHz

CHANNEL f2
 CPDPRG2 waltz16
 NUC2 13C
 PCPD2 100.00 usec
 PL2 0.00 dB
 PL12 26.00 dB
 PL13 26.00 dB
 SFO2 500.3920016 MHz

F2 - Processing parameters
 SI 32768
 SF 125.8231732 MHz
 WDW EM
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.40

1D NMR plot parameters
 CX 52.00 cm
 F1p 76.096 ppm
 F2p -0.552 ppm

Graphic