SUPPORTING INFORMATION FOR

Stereospecific Rearrangements During the Synthesis of Pyrrolidines and Related

Heterocycles from Cyclizations of Amino Alcohols with Vinyl Sulfones

Thomas G. Back,* Masood Parvez and Huimin Zhai

Department of Chemistry, University of Calgary, Calgary, AB, Canada, T2N 1N4

E-mail: tgback@ucalgary.ca

Contents

Experimental Procedures	SI 3-8
NMR Spectra of 3a	SI 9-10
NMR Spectra of 3b	SI 11-12
NMR Spectra of 3c	SI 13-14
NMR Spectra of 4a	SI 15-16
NMR Spectra of 4b	SI 17-18
NMR Spectra of 4c	SI 19-20
NMR Spectra of 5a	SI 21-22
NMR Spectra of 5c	SI 23-24
NMR Spectra of 6a	SI 25-26
NMR Spectra of 6b	SI 27-28
NMR Spectra of 6c	SI 29-30
NMR Spectra of 8	SI 31-32
NMR Spectra of 9	SI 33-34
NMR Spectra of 10	SI 35-36
NMR Spectra of 12a	SI 37-38
NMR Spectra of 12b	SI 39-40
NMR Spectra of 13a	SI 41-42
NMR Spectra of 14a	SI 43-46
NMR Spectra of 19	SI 47-48
NMR Spectra of 20	SI 49-50
NMR Spectra of 21	SI 51-52
NMR Spectra of 22a	SI 53-54
NMR Spectra of 22b	SI 55-56
NMR Spectra of 24a	SI 57-60
NMR Spectra of 24b	SI 61-62
X-Ray Data for 9	SI 63-78
X-Ray Data for 20	SI 79-88
X-Ray Data for 21	SI 89-98
X-Ray Data for 24b	SI 99-108

Experimental Procedures

(3*R*,4*S*)-3-(Benzenesulfonyl)-*N*,4-dibenzylpyrrolidine (6b). The same procedure was followed as for compound **3a**, starting with 814 mg (5.39 mmol) of amino alcohol **2b**, to give 1.705 g (99%) of **3b** as a yellow oil: IR (film) 3514, 3324, 1445, 1301, 1139, 1082 cm^{-1} ; ^1H NMR (200 MHz) δ 7.95-7.73 (m, 2 H), 7.71-7.46 (m, 3 H), 7.39-7.10 (m, 5 H), 3.57 (dd, J = 10.8, 3.6 Hz, 1 H), 3.38-3.13 (m, 3 H), 3.10-2.88 (m, 2 H), 2.88-2.76 (m, 1 H), 2.70 (crude d, J = 7.4 Hz, 2 H), 2.29 (br s, 2 H); ^{13}C NMR (50 MHz) δ 138.8 (C), 138.1 (C), 133.5 (CH), 129.0 (CH), 128.8 (CH), 128.2 (CH), 127.5 (CH), 126.1 (CH), 62.4 (CH_2), 60.2 (CH), 56.0 (CH_2), 40.5 (CH_2), 37.7 (CH_2); MS (m/z , %) 288 (6), 228 (17), 146 (94), 118 (92) 91 (100); HRMS calcd for $\text{C}_{16}\text{H}_{18}\text{NO}_2\text{S}$ ($\text{M}^+ - \text{CH}_2\text{OH}$): 288.1058. Found: 288.1039.

The same procedure was followed as for compound **4a**, starting with 2.234 g (7.00 mmol) of **3b** to give 2.620 g (92%) of **4b** as a light yellow oil: IR (film) 3495, 1494, 1301, 1145, 1084 cm^{-1} ; ^1H NMR (200 MHz) δ 7.90-7.74 (m, 2 H), 7.71-7.46 (m, 3 H), 7.38-7.19 (m, 8 H), 7.14-7.04 (m, 2 H), 3.88 (d, J = 13.5 Hz, 1 H), 3.53 (d, J = 13.3 Hz, 1 H), 3.47-3.39 (m, 2 H), 3.25-2.84 (m, 7 H), 2.44 (dd, J = 13.5, 9.1 Hz, 1 H); ^{13}C NMR (50 MHz) δ 139.0 (C), 138.7 (C), 138.4 (C), 133.5 (CH), 129.1 (CH), 128.7 (CH), 128.6 (CH), 128.5 (CH), 128.3 (CH), 127.5 (CH), 127.2 (CH), 126.0 (CH), 64.2 (CH), 60.8 (CH_2), 54.7 (CH_2), 54.5 (CH_2), 43.7 (CH_2), 32.4 (CH_2); MS (m/z , %) 409 (M^+ , 0.2), 378 (4), 287 (65), 186 (50), 133 (90), 91 (100); HRMS calcd for $\text{C}_{23}\text{H}_{24}\text{NO}_2\text{S}$ ($\text{M}^+ - \text{CH}_2\text{OH}$): 378.1528. Found: 378.1545.

The same procedure was followed as for compound **5a**, starting with 249 mg (0.608 mmol) of **4b**. The resulting crude chloride **5b** was treated directly with LDA as in the preparation of **6a** to afford 188 mg (79%) of **6b** as a colorless oil: IR (film) 1446, 1298, 1142, 1082 cm^{-1} ; ^1H NMR (200 MHz) δ 7.97-7.78 (m, 2 H), 7.73-7.50 (m, 3 H), 7.33-7.17 (m, 8 H), 7.14-6.94 (m, 2 H), 3.64 (d, J = 13.3 Hz, 1 H), 3.49 (d, J = 13.3 Hz, 1 H), 3.50-3.36 (m, 1 H), 3.10-2.52 (m, 6 H), 2.43 (dd, J = 9.2, 4.7 Hz, 1 H); ^{13}C NMR (50 MHz) δ 139.2 (C), 138.4 (C), 138.3 (C), 133.6 (CH), 129.2 (CH), 128.8 (CH), 128.7 (CH), 128.4 (CH), 128.3 (CH), 128.2 (CH), 127.1 (CH), 126.3 (CH), 67.9 (CH), 59.2 (CH_2), 58.5 (CH_2), 53.7 (CH_2), 41.5 (CH), 40.6 (CH_2); MS (m/z , %) 391 (M^+ , 0.3), 158 (80), 91 (100); HRMS calcd for $\text{C}_{24}\text{H}_{25}\text{NO}_2\text{S}$: 391.1606. Found: 391.1638.

(3*R*,4*S*)-3-(Benzenesulfonyl)-*N*-benzyl-4-isopropylpyrrolidine (6c). The same procedure was followed as for compound **3a**, starting with 966 mg (9.37 mmol) of amino alcohol **2c**, to afford 1.999 g (79%) of **3c** as a yellow oil: IR (film) 3535, 1483, 1312, 1152, 1088 cm^{-1} ; ^1H NMR (200 MHz) δ 8.02-7.83 (m, 2 H), 7.74-7.48 (m, 3 H), 3.56 (dd, J = 10.8, 4.1 Hz, 1 H), 3.35-3.10 (m, 4 H), 3.03-2.87 (m, 1 H), 2.33 (dt, J = 6.8, 4.1 Hz, 1 H), 2.47-1.85 (br s, 2 H), 1.75 (sextet, J = 6.8 Hz, 1 H), 0.94 (d, J = 6.8 Hz, 3 H), 0.88 (d, J = 6.8 Hz, 3 H); ^{13}C NMR (50 MHz) δ 138.5 (C), 133.1 (CH), 128.6 (CH), 127.1 (CH), 63.7 (CH), 60.3 (CH_2), 55.6 (CH_2), 40.5 (CH_2), 28.1 (CH), 18.4 (CH_3), 17.8 (CH_3); MS (m/z , %) 271 (M^+ , 0.3), 240 (17), 141 (28), 98 (100), 86 (97); HRMS calcd for $\text{C}_{12}\text{H}_{18}\text{NO}_2\text{S}$ ($\text{M}^+ - \text{CH}_2\text{OH}$): 240.1058. Found: 240.1057.

The same procedure was followed as for compound **4a**, starting with 1.800 g (6.64 mmol) of **3c**, to afford 2.220 g (93%) of **4c** as a colorless oil: IR (film) 3534, 1445, 1306, 1148, 1086 cm^{-1} ; ^1H NMR (200 MHz) δ 7.81-7.69 (m, 2 H), 7.69-7.44 (m, 3 H), 7.32-7.17 (m, 5 H), 3.85 (d, J = 13.3 Hz, 1 H), 3.63 (d, J = 13.3 Hz, 1 H) superimposed

on 3.75-3.56 (m, 1 H), 3.36 (dd, $J = 10.9, 9.9$ Hz, 1 H), 3.26-3.06 (m, 2 H), 3.05-2.77 (m, 3 H), 2.47 (crude sextet, $J = 4.6$ Hz, 1 H), 1.94-1.75 (m, 1 H), 1.03 (d, $J = 6.7$ Hz, 3 H), 0.88 (d, $J = 6.7$ Hz, 3 H); ^{13}C NMR (50 MHz) δ 139.2 (C), 138.7 (C), 133.2 (CH), 128.8 (CH), 128.5 (CH), 128.1 (CH), 127.3 (CH), 126.9 (CH), 69.4 (CH), 60.0 (CH₂), 55.1 (CH₂), 55.0 (CH₂), 45.8 (CH₂), 27.9 (CH), 21.6 (CH₃), 19.8 (CH₃); MS (m/z , %) 361 (M^+ , 0.7), 330 (19), 239 (17), 132 (31), 91 (100); HRMS calcd for C₁₉H₂₄NO₂S (M^+ -CH₂OH): 330.1528. Found: 330.1543.

The same procedure was followed as for compound **5a**, starting with 2.219 g (6.15 mmol) of **4c** to afford 2.220 g (95%) of **5c** as a colorless solid, mp 53 °C (from chloroform-hexanes); IR (film) 1449, 1305, 1152 cm⁻¹; ^1H NMR (200 MHz) δ 7.98-7.77 (m, 2 H), 7.75-7.43 (m 3 H), 7.29-7.15 (m, 5 H), 3.80 (ddd, $J = 7.3, 6.3, 3.3$ Hz, 1 H), 3.62 (d, $J = 13.7$ Hz, 1 H), 3.54 (d, $J = 13.7$ Hz, 1 H), 3.34-3.18 (m, 2 H), 3.00-2.92 (m, 2 H), 2.75 (dd, $J = 13.8, 6.3$ Hz, 1 H), 2.66 (dd, $J = 13.8, 7.2$ Hz, 1 H), 2.14-1.89 (m, 1 H), 0.96 (d, $J = 6.8$ Hz, 3 H), 0.78 (d, $J = 6.5$ Hz, 3 H); ^{13}C NMR (50 MHz) δ 139.2 (C), 137.9 (C), 133.6 (CH), 129.2 (CH), 128.7 (CH), 128.3 (CH), 127.8 (CH), 127.3 (CH), 67.1 (CH), 59.0 (2xCH₂), 53.3 (CH₂), 47.6 (CH₂), 31.2 (CH), 20.3 (CH₃), 15.9 (CH₃); MS (m/z , %) 379 (M^+ , 0.2), 144 (81), 91 (100). Anal. calcd for C₂₀H₂₆ClNO₂S: C, 63.22; H, 6.90; N, 3.69. Found: C, 63.11; H, 7.05; N, 3.82.

The same procedure was followed as for compound **6a**, starting with 224 mg (0.589 mmol) of **5c**, but using 1.5 equiv of LiHMDS instead of LDA to afford 181 mg (90%) of **6c** as a yellow oil: IR (film) 1448, 1306, 1151, 1090 cm⁻¹; ^1H NMR (200 MHz) δ 7.98-7.82 (m, 2 H), 7.73-7.46 (m 3 H), 7.37-7.09 (m, 5 H), 3.58 (d, $J = 13.2$ Hz, 1 H), 3.45 (d, $J = 13.2$ Hz, 1 H) superimposed on 3.49-3.38 (m, 1 H), 2.98 (dd, $J = 10.4, 5.0$ Hz, 1 H), 2.79-2.60 (m, 2 H), 2.59-2.44 (m, 1 H), 2.35 (dd, $J = 8.7, 5.2$ Hz, 1 H), 1.63 (m, 1 H), 0.88 (d, $J = 6.7$ Hz, 3 H), 0.85 (d, $J = 6.7$ Hz, 3 H); ^{13}C NMR (50 MHz) δ 138.5 (C), 138.4 (C), 133.4 (CH), 128.9 (CH), 128.8 (CH), 128.2 (CH), 128.1 (CH), 126.9 (CH), 66.8 (CH), 59.3 (CH₂), 56.5 (CH₂), 54.6 (CH₂), 45.6 (CH), 31.1 (CH), 21.0 (CH₃), 19.0 (CH₃); MS (m/z , %) 343 (M^+ , 0.6), 201 (34), 158 (91), 132 (25), 91 (100); HRMS calcd for C₂₀H₂₅NO₂S: 343.1606. Found: 343.1613. Anal. calcd for C₂₀H₂₅NO₂S: C, 69.93; H, 7.34; N, 4.08. Found: C, 69.54; H, 7.09; N, 4.15.

(2S,3S,4R)-4-(Benzenesulfonyl)-1,2-dimethyl-3-phenylpyrrolidine (9) and its (2S,3R,4S) isomer 10. (-)-Ephedrine (**7**) (1.652 g, 10.0 mmol) and 1.680 g (10.0 mmol) of vinyl sulfone **1a** were refluxed in 35 mL of xylenes for 2 days. The mixture was evaporated in vacuo and chromatography (50% ethyl acetate-hexanes) afforded 3.2093 g (96%) of **8** as a yellow oil: IR (film) 3513, 1449, 1304, 1144 cm⁻¹; ^1H NMR (200 MHz) δ 7.96-7.83 (m, 2 H), 7.71-7.50 (m 3 H), 7.37-7.15 (m, 5 H), 4.70 (d, $J = 4.3$ Hz, 1 H), 3.25-3.15 (m, 3 H), 3.00-2.89 (m, 2 H), 2.71 (dq, $J = 6.8, 4.4$ Hz, 1 H), 2.18 (s, 3 H), 0.85 (d, $J = 6.8$ Hz, 3 H); ^{13}C NMR (50 MHz) δ 142.1 (C), 139.4 (C), 133.6 (CH), 129.2 (CH), 127.9 (CH), 127.7 (CH), 126.9 (CH), 125.9 (CH), 73.6 (CH), 63.7 (CH), 53.9 (CH₂), 47.9 (CH₂), 38.7 (CH₃), 9.6 (CH₃); MS (m/z , %) 333 (M^+ , 0.2), 316 (1), 226 (30), 84, (100), 42 (99); HRMS calcd for C₁₈H₂₂NO₂S (M^+ -OH): 316.1371. Found: 316.1362.

A solution of **8** (1.182 g, 3.55 mmol) and thionyl chloride (0.65 mL, 8.9 mmol) in 45 mL of chloroform was refluxed for 3 h and then concentrated in vacuo. The residue was dissolved in 10 mL of THF and added to a solution of excess LDA (10.1 mmol) in 10 mL of THF at -78 °C. The mixture was stirred at -78 °C for 2 h and at room temperature for 5 h and was then quenched by filtration through neutral alumina. The

filtrate was concentrated in vacuo, and the residue was chromatographed (25% ethyl acetate-hexanes, then 35% ethyl acetate-hexanes) to afford 326 mg (29%) of diastereomer **10** as a light yellow oil: IR (film) 1444, 1306, 1148 cm^{-1} ; ^1H NMR (200 MHz) δ 7.93-7.74 (m, 2 H), 7.64-7.37 (m, 3 H), 7.21-7.08 (m, 3 H), 7.08-6.96 (m, 2 H), 3.84 (crude sextet, $J = 4.2$ Hz, 1 H), 3.58 (dd, $J = 7.4, 4.3$ Hz, 1 H), 3.48-3.30 (m, 1 H), 2.88-2.62 (m, 2 H), 2.31 (s, 3 H), 0.66 (d, $J = 6.5$ Hz, 3 H); ^{13}C NMR (50 MHz) δ 140.7 (C), 138.7 (C), 133.5 (CH), 129.1 (CH), 128.8 (CH), 128.3 (CH), 128.0 (CH), 126.7 (CH), 69.1 (CH), 65.0 (CH), 55.9 (CH_2), 50.4 (CH), 39.7 (CH_3), 14.8 (CH_3); MS (m/z , %) 174 (3), 156 (37), 128 (19), 115 (100); HRMS calcd for $\text{C}_{12}\text{H}_{16}\text{N}$ ($\text{M}^+ - \text{SO}_2\text{Ph}$): 174.1283. Found: 174.1291. All ^{13}C and ^1H NMR signals were assigned by means of COSY, DEPT and HMQC spectra. The following NOE's were observed: irradiation of MeCH enhanced PhCH (7%); irradiation of PhCH enhanced MeCH (7%) and only slightly enhanced PhSO_2CH (1%); irradiation of PhSO_2CH enhanced one CH_2 signal, but gave no enhancement of PhCH.

Continued elution afforded 446 mg (40%) of **9** as light yellow solid, which crystallized from ethyl acetate-hexanes to afford white crystals: mp 137.5-138 $^\circ\text{C}$ (from ethyl acetate-hexanes); IR (film) 1441, 1304, 1145 cm^{-1} ; ^1H NMR (200 MHz) δ 7.84-7.70 (m, 2 H), 7.59-7.28 (m, 3 H), 7.24-7.04 (m, 3 H), 7.03-6.86 (m, 2 H), 3.82-3.70 (m, 2 H), 3.09 (dd, $J = 9.4, 7.4$ Hz, 1 H), 2.75 (dd, $J = 11.6, 9.9$ Hz, 1 H), 2.31 (s, 3 H), 2.29-2.15 (m, 1 H), 1.00 (d, $J = 5.8$ Hz, 3 H); ^{13}C NMR (50 MHz) δ 139.8 (C), 138.2 (C), 133.4 (CH), 128.9 (CH), 128.5 (CH), 128.4 (CH), 127.7 (CH), 126.9 (CH), 70.2 (CH), 68.5 (CH), 56.2 (CH_2), 54.6 (CH), 39.6 (CH_3), 15.9 (CH_3); MS (m/z , %) 174 (7), 172 (37), 156 (61), 115 (100); HRMS calcd for $\text{C}_{12}\text{H}_{16}\text{N}$ ($\text{M}^+ - \text{SO}_2\text{Ph}$): 174.1283. Found: 174.1279. Anal. calcd for $\text{C}_{18}\text{H}_{21}\text{NO}_2\text{S}$: C, 68.54; H, 6.71; N, 4.44. Found: C, 68.63; H, 6.79; N, 4.48. The structure was confirmed by X-ray crystallography.

2-(Benzenesulfonyl)indolizidine (14a). A solution of (2-piperidine)methanol (**11a**) (1.540 g, 13.39 mmol) and vinyl sulfone **1a** (2.249 g, 13.39 mmol) in 35 mL of xylenes was refluxed for 1 day and concentrated in vacuo to give crude **12a**. Chromatography of the crude product (hexane:ethyl acetate:methanol = 4:1:0.5) gave 3.650 g (96%) of **12a** as a light yellow oil: IR (film) 3510, 1442, 1302, 1145, 1082 cm^{-1} ; ^1H NMR (200 MHz) δ 7.96-7.78 (m, 2 H), 7.69-7.42 (m, 3 H), 3.62 (dd, $J = 11.5, 3.9$ Hz, 1 H), 3.42 (dd, $J = 11.5, 4.4$ Hz, 1 H), 3.35-3.07 (m, 3 H), 3.00-2.77 (m, 1 H), 2.76-2.62 (m, 2 H), 2.33-2.01 (m, 2 H), 1.66-1.10 (m, 6 H); ^{13}C NMR (50 MHz) δ 139.4 (C), 133.3 (CH), 128.9 (CH), 127.5 (CH), 62.6 (CH_2), 60.5 (CH), 52.2 (CH_2), 51.2 (CH_2), 46.5 (CH_2), 27.3 (CH_2), 24.1 (CH_2), 22.9 (CH_2); MS (m/z , %) 283 (M^+ , 2), 252 (17), 141 (13), 110 (92), 82 (100); HRMS calcd for $\text{C}_{13}\text{H}_{18}\text{NO}_2\text{S}$ ($\text{M}^+ - \text{CH}_2\text{OH}$): 252.1058. Found: 252.1065.

A solution of **12a** (3.650 g, 12.85 mmol) and 2.34 mL of thionyl chloride (3.823 g, 32.13 mmol) in 30 mL of chloroform was refluxed for 2 h. The solution was washed with 1 M aqueous KOH solution, water and brine. The organic layer was dried, concentrated and chromatographed (20% ethyl acetate-hexanes) to afford 3.370 g (87%) of **13a**, which crystallized from dichloromethane-hexane to give a light yellow solid: mp 44 $^\circ\text{C}$; IR (film) 1445, 1298, 1144, 1088 cm^{-1} ; ^1H NMR (200 MHz) δ 7.97-7.83 (m, 2 H), 7.69-7.47 (m, 3 H), 3.54-3.37 (m, 2 H), 3.34-3.21 (m, 2 H), 3.13-2.86 (m, 2 H), 2.70-2.57 (m, 1 H), 2.55-2.40 (m, 1 H), 2.28-2.14 (m, 1 H), 1.66-1.15 (m, 6 H); ^{13}C NMR (50 MHz) δ 139.5 (C), 133.7 (CH), 129.2 (CH), 127.8 (CH), 60.2 (CH), 52.5 (CH_2), 51.3

(CH₂), 47.0 (CH₂), 45.0 (CH₂), 28.8 (CH₂), 25.1 (CH₂), 22.4 (CH₂); MS (*m/z*, %) 301 (M⁺, 2), 252 (16), 110 (100), 96 (47), 82 (74). Anal. calcd for C₁₄H₂₀ClNO₂S: C, 55.71; H, 6.68; N, 4.64. Found: C, 55.46; H, 6.52; N, 4.57.

The chloroamine **13a** (281 mg, 0.934 mmol) was dissolved in 5 mL of THF and added to 1.60 mmol of LDA in 5 mL of THF at -78 °C. The mixture was stirred at -78 °C for 2 h and at room temperature for another 2 h and was then quenched by filtration through neutral alumina. The filtrate was concentrated in vacuo and the residue was chromatographed (25% ethyl acetate-hexanes, then 35% ethyl acetate-hexanes) to afford 96 mg (39%) of the less polar diastereomer of **14a** as a colorless oil: IR (film) 1446, 1304, 1283, 1149, 1086 cm⁻¹; ¹H NMR (200 MHz) δ 7.98-7.81 (m, 2 H), 7.72-7.47 (m, 3 H), 3.67-3.52 (m, 1 H), 3.47 (dd, *J* = 10.8, 2.5 Hz, 1 H), 2.98 (dt, *J* = 10.8, 3.1 Hz, 1 H), 2.49-2.30 (m, 1 H), 2.16-1.98 (m, 1 H), 1.99-1.79 (m, 3 H), 1.78-1.61 (m, 2 H), 1.60-1.39 (m, 2 H), 1.24-1.03 (m, 2 H); ¹³C NMR (50 MHz) δ 138.5 (C), 133.4 (CH), 129.0 (CH), 128.6 (CH), 63.5 (CH), 61.0 (CH), 54.1 (CH₂), 52.0 (CH₂), 32.9 (CH₂), 30.2 (CH₂), 24.7 (CH₂), 23.8 (CH₂); MS (*m/z*, %) 265 (M⁺, 0.5), 155 (28), 110 (100), 82 (96); HRMS calcd for C₁₄H₁₉NSO₂ 265.1137. Found: 265.1129.

Continued elution afforded 112 mg (45%) of the more polar diastereomer of **14a** as a colorless oil: IR (film) 1442, 1303, 1148, 1083 cm⁻¹; ¹H NMR (200 MHz) δ 7.96-7.80 (m, 2 H), 7.70-7.47 (m, 3 H), 3.78-3.59 (m, 1 H), 3.14 (dd, *J* = 9.4, 8.3 Hz, 1 H), 3.05-2.92 (m, 1 H), 2.59 (crude t, *J* = 9.1 Hz, 1 H), 2.33 (ddd, *J* = 13.3, 6.0, 3.1 Hz, 1 H), 2.14-1.96 (m, 2 H), 1.85-1.57 (m, 4 H), 1.53-0.98 (m, 3 H); ¹³C NMR (50 MHz) δ 138.8 (C), 133.6 (CH), 129.2 (CH), 128.3 (CH), 63.0 (CH), 60.2 (CH), 54.5 (CH₂), 52.4 (CH₂), 32.5 (CH₂), 30.5 (CH₂), 25.2 (CH₂), 23.8 (CH₂); MS (*m/z*, %) 265 (M⁺, 3), 122 (100), 94 (34); HRMS calcd for C₁₄H₁₉NO₂S: 265.1137. Found: 265.1139.

3-(Benzenesulfonyl)quinolizidine (14b). The same procedure was followed as for compound **12a**, starting with 712 mg (5.52 mmol) of 2-piperidineethanol and 0.929 g (5.53 mmol) of vinyl sulfone **1a**, except that the reaction was refluxed for 2 days. Chromatography afforded 1.559 g (95%) of **12b** as a light yellow oil: IR (film) 3532, 1445, 1301, 1138, 1082 cm⁻¹; ¹H NMR (200 MHz) 7.99-7.85 (m, 2 H), 7.74-7.50 (m, 3 H), 3.84-3.55 (m, 2 H), 3.34-2.93 (m, 4 H), 2.91-2.73 (m, 1 H), 2.67-2.46 (m, 1 H), 2.40-2.15 (m, 1 H), 1.89-1.20 (m, 8 H); ¹³C NMR (50 MHz) δ 139.2 (C), 133.4 (CH), 129.0 (CH), 127.6 (CH), 60.3 (CH₂), 57.6 (CH), 52.7 (CH₂), 50.2 (CH₂), 46.1 (CH₂), 32.0 (CH₂), 28.3 (CH₂), 23.5 (CH₂), 22.2 (CH₂); MS (*m/z*, %) 296 (M⁺, 0.6), 252 (31), 110 (100), 82 (93); HRMS calcd for C₁₃H₁₈NO₂S (M⁺-C₂H₄OH): 252.1058. Found: 252.1064.

The same procedure was followed as in the preparation of **14a** from **12a**. Thus, 813 mg (2.74 mmol) of **12b** afforded 569 mg (74%) of **14b** as a light yellow oil, with spectra in agreement with those reported in the literature.¹²

(2*S*,3*R*,4*S*)-3-(Benzenesulfonyl)-*N*-benzyl-2,4-dimethylpyrrolidine (20) and its (2*R*,3*S*,4*S*) isomer 21. A mixture of 289 mg (3.85 mmol) of **2a** and 702 mg (3.85 mmol) of vinyl sulfone **1b** was refluxed for 2 days in 20 mL of xylenes. Chromatography (hexane:ethyl acetate:methanol = 4:1:0.5) afforded 900 mg (91%) of the corresponding adduct as a mixture of two diastereomers formed in the ratio of ca. 1:1, colorless oil: IR (film) 3408, 1448, 1301, 1144, 1082 cm⁻¹; ¹H NMR (200 MHz, mixture) δ 7.97-7.83 (m, 2 H), 7.72-7.47 (m, 3 H), 3.60-2.99 (m, 5 H), 2.87-2.64 (m, 1 H), 2.18 (br s, 2 H), 1.23 (d, *J* = 6.3 Hz), 1.11 (d, *J* = 6.2 Hz), 1.02 (d, *J* = 6.3 Hz), 0.98 (d, *J* = 6.5 Hz), total of signals from δ 1.23-0.98: 6 H; ¹³C NMR (50 MHz, both diastereomers) δ 139.9 (C), 139.5 (C),

133.7 (CH), 133.5 (CH), 129.2 (CH), 129.1 (CH), 127.6 (CH), 127.5 (CH), 66.0 (CH₂), 65.5 (CH₂), 62.4 (CH₂), 62.2 (CH₂), 51.8 (CH), 51.1 (CH), 46.4 (CH), 45.3 (CH), 22.1 (CH₃), 21.2 (CH₃), 17.9 (CH₃), 16.9 (CH₃); MS (*m/z*, %) 226 (41), 141 (83), 125 (23), 77 (82), 44 (100); HRMS calcd for C₁₁H₁₆NO₂S (M⁺-CH₂OH): 226.0902. Found: 226.0905.

The above product (273 mg, 1.06 mmol) was treated with benzyl bromide as in the preparation of **4a** to afford 259 mg (70%) of **19** (mixture of diastereomers) as a colorless oil: IR (film) 3456, 1451, 1303, 1141, 1079 cm⁻¹; ¹H NMR (200 MHz, mixture) δ 7.92-7.13 (m, 10 H), 3.87-3.23 (m, 6 H), 3.08-2.88 (m, 2 H), 1.93 (br s, 1 H), 1.31 (d, *J* = 7.0 Hz), 1.20 (d, *J* = 6.5 Hz), 1.01 (d, *J* = 6.7 Hz), 0.97 (d, *J* = 6.5 Hz), total of signals from δ 1.31-0.97: 6 H; ¹³C NMR (50 MHz, both diastereomers) δ 139.7 (C), 139.6 (C), 139.4 (C), 139.2 (C), 133.7 (CH), 133.5 (CH), 129.4 (CH), 129.2 (2 x CH), 128.6 (2 x CH), 128.5 (CH), 128.4 (CH), 127.7 (CH), 127.6 (CH), 127.1 (CH), 63.9 (CH₂), 63.6 (CH₂), 60.9 (CH₂), 59.7 (CH₂), 55.1 (CH), 53.8 (CH), 49.6 (CH), 49.5 (CH₂), 48.3 (CH), 48.1 (CH₂), 20.4 (CH₃), 17.2 (CH₃), 13.5 (CH₃), 13.2 (CH₃); MS (*m/z*, %) 347 (M⁺, 0.5), 316 (19), 91 (100); HRMS calcd for C₁₈H₂₂NO₂S (M⁺-CH₂OH): 316.1371. Found: 316.1388.

A solution of **19** (644 mg, 1.86 mmol) and thionyl chloride (0.34 mL, 4.7 mmol) in 25 mL of chloroform was refluxed for 3 h and then concentrated in vacuo. The residue was dissolved in 5 mL of THF and added to a solution of excess LDA (5.6 mmol) in 6 mL of THF at -78 °C. The mixture was stirred at -78 °C for 2 h and at room temperature for 2 h, and was then quenched by filtration through neutral alumina. The filtrate was concentrated in vacuo, and the residue was chromatographed (14% ethyl acetate-hexanes), to afford 320 mg (52%) of a mixture of **20** and an unidentified diastereomer as an 83:17 mixture (NMR integration). Recrystallization of this mixture from chloroform-methanol provided pure **20** as light yellow crystals, mp 134-135 °C. Further chromatography with 20% ethyl acetate-hexanes afforded 62 mg (10%) of **21**, which gave white crystals from ethyl acetate-hexanes, mp 139-142 °C.

Compound **20**: IR (film) 1442, 1300, 1146, 1082 cm⁻¹; ¹H NMR (200 MHz) δ 8.01-7.84 (m, 2 H), 7.75-7.50 (m, 3 H), 7.39-7.16 (m, 5 H), 3.98 (d, *J* = 13.3 Hz, 1 H), 3.18 (d, *J* = 13.3 Hz, 1 H), 3.02 (quintet, *J* = 6.4 Hz, 1 H), 2.87 (dd, *J* = 7.2, 3.8 Hz, 1 H), 2.66-2.34 (m, 3 H), 1.18 (d, *J* = 6.0 Hz, 3 H), 0.96 (d, *J* = 6.8 Hz, 3 H); ¹³C NMR (50 MHz) δ 138.9 (C), 138.6 (C), 133.6 (CH), 129.2 (CH), 128.6 (CH), 128.4 (CH), 128.1 (CH), 126.9 (CH), 77.4 (CH), 60.2 (CH), 59.4 (CH₂), 56.9 (CH₂), 33.3 (CH), 21.5 (CH₃), 19.5 (CH₃); MS (*m/z*, %) 329 (M⁺, 2), 187 (38), 172 (90), 146 (70), 91 (100); HRMS calcd for C₁₉H₂₃NO₂S: 329.1450. Found: 329.1437. Anal. calcd for C₁₉H₂₃NO₂S: C, 69.27; H, 7.04; N, 4.25. Found: C, 69.34; H, 7.10; N, 4.28. The structure was confirmed by X-ray crystallography.

Compound **21**: IR (film) 1445, 1301, 1145, 1082 cm⁻¹; ¹H NMR (200 MHz) δ 7.97-7.82 (m, 2 H), 7.72-7.51 (m, 3 H), 7.39-7.19 (m, 5 H), 3.93 (d, *J* = 12.8 Hz, 1 H), 3.29 (d, *J* = 12.3 Hz, 1 H) superimposed on 3.36-3.26 (m, 1 H), 3.01 (quintet, *J* = 5.8 Hz, 1 H), 2.90 (dd, *J* = 8.4, 5.8 Hz, 1 H), 2.78-2.53 (m, 1 H), 2.40 (dd, *J* = 10.9, 8.6 Hz, 1 H), 1.39 (d, *J* = 7.2 Hz, 3 H), 0.77 (d, *J* = 6.2 Hz, 3 H); ¹³C NMR (50 MHz) δ 140.3 (C), 139.0 (C), 133.5 (CH), 129.1 (CH), 128.8 (CH), 128.5 (CH), 128.3 (CH), 127.0 (CH), 72.1 (CH), 61.8 (CH), 60.8 (CH₂), 58.2 (CH₂), 35.6 (CH), 20.1 (CH₃), 13.2 (CH₃); MS (*m/z*, %) 329 (M⁺, 7), 173 (78), 158 (63), 146 (55), 91 (100). Anal. calcd for

C₁₉H₂₃NO₂S: C, 69.27; H, 7.04; N, 4.25. Found: C, 68.88; H, 6.82; N, 4.16. The structure was confirmed by X-ray crystallography.

(+/-)-2-cis-(Benzenesulfonyl)-2-trans-methylindolizidine (24a) and its 2-epi isomer 25a. A mixture of 433.6 mg (3.77 mmol) of (2-piperidine)methanol (**11a**) and 686 mg (3.77 mmol) of vinyl sulfone **1b** in 25 mL of xylenes was refluxed for 3 days. After removal of solvent in vacuo, chromatography (hexane:ethyl acetate:methanol = 4:1:0.5) afforded 324 mg (29%) of **22a** as a mixture of two diastereomers formed in the ratio of 60:40 as a light yellow oil; IR (film) 3501, 1446, 1301, 1142, 1081 cm⁻¹; ¹H NMR (200 MHz, mixture) δ 8.00-7.81 (m, 2 H), 7.71-7.46 (m, 3 H), 3.71 (minor isomer, dd, *J* = 11.6, 4.1 Hz, 1 H), 3.56-3.20 (m, 3 H), 2.94 (major isomer, dd, *J* = 13.3, 4.9 Hz, 1 H), 2.87-2.63 (m, 2 H), 2.60 (br s, 1 H), 2.50-2.00 (m, 3 H), 1.70-1.35 (m, 3 H), 1.35-1.04 (m, 2 H), 1.30 (minor isomer, d, *J* = 6.8 Hz, 3 H), 1.28 (major isomer, d, *J* = 6.8 Hz, 3 H); ¹³C NMR (50 MHz) major diastereomer: δ 137.9 (C), 133.5 (CH), 129.0 (CH), 128.6 (CH), 62.0 (CH₂), 61.4 (CH), 58.7 (CH), 52.9 (CH₂), 50.4 (CH₂), 26.1 (CH₂), 23.2 (CH₂), 22.5 (CH₂), 12.7 (CH₃); minor diastereomer: δ 138.7 (C), 133.4 (CH), 128.9 (CH), 128.5 (CH), 62.0 (CH₂), 61.8 (CH), 58.4 (CH), 54.0 (CH₂), 52.1 (CH₂), 26.3 (CH₂), 23.6 (CH₂), 22.7 (CH₂), 13.0 (CH₃); MS (*m/z*, %) 296 (M⁺-1, 1), 266 (18), 124 (100), 96 (58), 82 (80); HRMS calcd for C₁₄H₂₀NO₂S (M⁺-CH₂OH): 266.1215. Found: 266.1221.

A solution of **22a** (332 mg, 1.11 mmol) and thionyl chloride (0.26 mL, 3.56 mmol) in 25 mL of chloroform was refluxed for 3 h and then concentrated in vacuo. The residue was dissolved in 5 mL of THF and added to a solution of excess LDA (3.4 mmol) in 6 mL of THF at -78 °C. The mixture was stirred at -78 °C for 2 h and at room temperature for 6 h, and was then quenched by filtration through neutral alumina. The filtrate was concentrated in vacuo, and the residue was chromatographed (25% ethyl acetate-hexanes) to afford 134 mg (43%) of the less polar isomer **24a** as a yellow oil: IR (film) 1443, 1300, 1140 1075 cm⁻¹; ¹H NMR (200 MHz) 7.99-7.85 (m, 2 H), 7.71-7.46 (m, 3 H), 3.69 (d, *J* = 11.1 Hz, 1 H), 2.85 (dt, *J* = 10.8, 2.9 Hz, 1 H), 2.21 (dd, *J* = 12.3, 10.4 Hz, 1 H), 2.03 (d, *J* = 11.0 Hz, 1 H), 1.98-1.74 (m, 2 H), 1.72-1.58 (m, 3 H), 1.48-1.40 (m, 1 H), 1.43 (s, 3 H), 1.33-1.22 (m, 1 H), 1.17-0.96 (m, 2 H); ¹³C NMR (50 MHz) δ 136.8 (C), 133.3 (CH), 130.1 (CH), 128.5 (CH), 66.3 (C), 62.5 (CH), 62.4 (CH₂), 52.0 (CH₂), 41.1 (CH₂), 30.2 (CH₂), 24.8 (CH₂), 23.9 (CH₂), 22.8 (CH₃); MS (*m/z*, %) 279 (M⁺, 2), 136 (100), 108 (49); HRMS calcd for C₁₅H₂₁NO₂S: 279.1293. Found: 279.1313.

Further elution with 30% ethyl acetate-hexanes afforded 88 mg (28%) of the more polar isomer **25a** as a yellow oil: IR (film) 1444, 1300, 1152; 1076 cm⁻¹; ¹H NMR (200 MHz) 7.97-7.87 (m, 2 H), 7.72-7.51 (m, 3 H), 2.97 (d, *J* = 9.2 Hz, 1 H) superimposed on 3.05-2.91 (m, 1 H), 2.74 (d, *J* = 9.4 Hz, 1 H), 2.67 (dd, *J* = 13.3, 6.0 Hz, 1 H), 2.23-2.10 (m, 1 H), 2.04 (td, *J* = 11.1, 3.4 Hz, 1 H), 1.82-1.70 (m, 2 H), 1.65-1.57 (m, 1 H), 1.48 (s, 3 H), 1.32 (dd, *J* = 13.5, 10.6 Hz, 2 H), 1.22-1.06 (m, 2 H); ¹³C NMR (50 MHz) δ 136.7 (C), 133.6 (CH), 130.1 (CH), 128.9 (CH), 66.2 (C), 64.6 (CH), 61.8 (CH₂), 52.6 (CH₂), 41.2 (CH₂), 30.8 (CH₂), 25.2 (CH₃), 25.1 (CH₂), 23.9 (CH₂); MS (*m/z*, %) 279 (M⁺, 1.4), 136 (100), 108 (76); HRMS calcd for C₁₅H₂₁NO₂S: 279.1293. Found: 279.1279.

Desulfonylation of 6a. Product **6a** was desulfonylated with sodium amalgam by the procedure of Trost et al.¹⁰ to afford (*R*)-*N*-benzyl-3-methylpyrrolidine in 45% yield, with NMR spectra as reported in the literature;^{11a} [α]_D -8.6 ° (c 1.75, EtOH), lit.^{11b} [α]_D -10.5 ° (c 3.25, EtOH).

^1H NMR Spectrum of **3a**

^{13}C NMR Spectrum of **3a**

^1H NMR Spectrum of **3b**

^{13}C NMR Spectrum of **3b**

^1H NMR Spectrum of **3c**

^{13}C NMR Spectrum of **3c**

^1H NMR Spectrum of **4a**

^{13}C NMR Spectrum of **4a**

^1H NMR Spectrum of **4b**

^{13}C NMR Spectrum of **4b**

^1H NMR Spectrum of **4c**

^{13}C NMR Spectrum of **4c**

^1H NMR Spectrum of **5a**

^{13}C NMR Spectrum of **5a**

¹H NMR Spectrum of **5c**

^{13}C NMR Spectrum of **5c**

^1H NMR Spectrum of **6a**

^{13}C NMR Spectrum of **6a**

^1H NMR Spectrum of **6b**

^{13}C NMR Spectrum of **6b**

^1H NMR Spectrum of **6c**

^{13}C NMR Spectrum of **6c**

^1H NMR Spectrum of **8**

^{13}C NMR Spectrum of **8**

^1H NMR Spectrum of **9**

^{13}C NMR Spectrum of **9**

¹H NMR Spectrum of **10**

^{13}C NMR Spectrum of **10**

^1H NMR Spectrum of **12a**

^{13}C NMR Spectrum of **12a**

^1H NMR Spectrum of **12b**

^{13}C NMR Spectrum of **12b**

¹H NMR Spectrum of **13a**

^{13}C NMR Spectrum of **13a**

¹H NMR Spectrum of Less Polar Isomer of **14a**

^{13}C NMR Spectrum of Less Polar Isomer of **14a**

¹H NMR Spectrum of More Polar Isomer of **14a**

^{13}C NMR Spectrum of More Polar Isomer of **14a**

^1H NMR Spectrum of **19**

^{13}C NMR Spectrum of **19**

^1H NMR Spectrum of **20**

^{13}C NMR Spectrum of **20**

^1H NMR Spectrum of **21**

^{13}C NMR Spectrum of **21**

^1H NMR Spectrum of **22a**

^{13}C NMR Spectrum of **22a**

^1H NMR Spectrum of **22b**

^{13}C NMR Spectrum of **22b**

^1H NMR Spectrum of **24a** (major epimer)

^{13}C NMR Spectrum of **24a** (major epimer)

^1H NMR Spectrum of **24a** (minor epimer)

^{13}C NMR Spectrum of **24a** (minor epimer)

^1H NMR Spectrum of **24b**

^{13}C NMR Spectrum of **24b**

ORTEP Diagram of **9**

Experimental:

A colorless prismatic crystal of $C_{18}H_{21}NO_2S$ was coated with Paratone 8277 oil (Exxon) and mounted on a glass fiber. All measurements were made on a Nonius KappaCCD diffractometer with graphite monochromated Mo- $K\alpha$ radiation. Cell constants obtained from the refinement¹ of 19100 reflections in the range $1.0 < \theta < 27.5^\circ$ corresponded to a primitive monoclinic cell; details of crystal data and structure refinement have been provided in Table 1. The data were collected² at a temperature of 173(2) K using the ω and ϕ scans to a maximum θ value of 27.5° . The data were corrected for Lorentz and polarization effects and for absorption using multi-scan method¹. Since the crystal did not show any sign of decay during data collection a decay correction was deemed unnecessary.

The structure was solved by the direct methods³ and expanded using Fourier techniques⁴.

There are two independent molecules in the asymmetric unit. The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included at geometrically idealized positions and were not refined. The final cycle of full-matrix least-squares refinement using SHELXL97⁵ converged (largest parameter shift was 0.00 times its esd) with unweighted and weighted agreement factors, $R = 0.0425$ and $wR = 0.0812$ (all data), respectively, and goodness of fit, $S = 1.030$. The absolute structure was established by the Flack method⁶ with the absolute configuration at the chiral centers: C2, C2' (R), C3, C3' (S) and C4, C4' (S). The Flack parameter for the inverted structure was 1.00(5). Therefore, the inverted structure was rejected as the one present in the crystal. The weighting scheme was based on counting statistics and the final difference Fourier map was essentially featureless. The figures were plotted with the aid of ORTEPII⁷.

References:

1. Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307.
2. Hooft, R. (1998). *COLLECT*. Nonius BV, Delft. The Netherlands.
3. Altomare, A., Cascarano, M., Giacovazzo, C. & Guagliardi, A. (1993). *SIR92*. J. Appl. Cryst., **26**, 343
4. Beurskens, P.T., Admiraal, G., Beurskens, G., Bosman, W.P., de Gelder, R., Israel, R. & Smits, J.M.M. (1994). The *DIRDIF-94* program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
5. Sheldrick, G.M. (1997). *SHELXL97*. University of Göttingen, Germany.
6. Flack, H. D. (1983). *Acta Cryst.* A39, 876.

7. Johnson, C.K. (1976). *ORTEP II*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.

Table 1. Crystal data and structure refinement for $C_{18}H_{21}NO_2S$.

Identification code	T. Back – 5 (RZ – 145)		
Empirical formula	$C_{18}H_{21}NO_2S$		
Formula weight	315.42		
Temperature	173(2) K		
Wavelength	0.71073 Å		
Crystal system	Monoclinic		
Space group	P21		
Unit cell dimensions	$a = 8.682(2)$ Å	$\alpha = 90^\circ$.	
	$b = 10.123(2)$ Å	$\beta = 93.99(2)^\circ$.	
	$c = 18.874(5)$ Å	$\gamma = 90^\circ$.	
Volume	$1654.8(7)$ Å ³		
Z	4		
Density (calculated)	1.266 Mg/m ³		
Absorption coefficient	0.202 mm ⁻¹		
F(000)	672		
Crystal size	0.20 x 0.18 x 0.10 mm ³		
Theta range for data collection	5.9 to 27.5°.		
Index ranges	$-11 \leq h \leq 10$, $-13 \leq k \leq 13$, $-24 \leq l \leq 24$		
Reflections collected	19100		
Independent reflections	7234 [R(int) = 0.050]		
Completeness to $\theta = 27.5^\circ$	97.8 %		
Absorption correction	Multi-scan method		
Max. and min. transmission	0.9801 and 0.9607		
Refinement method	Full-matrix least-squares on F ²		
Data / restraints / parameters	7234 / 1 / 397		
Goodness-of-fit on F ²	1.030		
Final R indices [I > 2σ(I)]	R1 = 0.0425, wR2 = 0.0725		
R indices (all data)	R1 = 0.0716, wR2 = 0.0812		
Absolute structure parameter	-0.01(5)		
Largest diff. peak and hole	0.235 and -0.302 e.Å ⁻³		

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{18}\text{H}_{21}\text{NO}_2\text{S}$. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

Atom	x	y	z	$U(\text{eq})$
S(1)	4423(1)	-753(1)	-108(1)	23(1)
O(1)	4316(2)	578(2)	-379(1)	29(1)
O(2)	3653(2)	-1783(2)	-528(1)	31(1)
N(1)	8265(2)	-80(2)	-606(1)	28(1)
C(1)	7182(3)	-1187(2)	-676(1)	28(1)
C(2)	6407(3)	-1215(2)	32(1)	22(1)
C(3)	7392(2)	-276(2)	534(1)	22(1)
C(4)	8874(2)	-132(2)	137(1)	25(1)
C(5)	9422(3)	-161(3)	-1130(2)	43(1)
C(6)	9812(3)	1084(2)	356(1)	37(1)
C(7)	7677(2)	-708(2)	1298(1)	25(1)
C(8)	7709(3)	223(3)	1841(2)	41(1)
C(9)	8058(3)	-143(3)	2545(2)	52(1)
C(10)	8400(3)	-1436(3)	2715(2)	48(1)
C(11)	8370(3)	-2363(3)	2182(1)	44(1)
C(12)	8016(3)	-2003(2)	1484(1)	35(1)
C(13)	3728(2)	-760(2)	750(1)	24(1)
C(14)	3134(3)	-1915(3)	1027(1)	37(1)
C(15)	2576(3)	-1875(3)	1692(1)	47(1)
C(16)	2576(3)	-710(3)	2076(1)	44(1)
C(17)	3190(3)	427(3)	1807(1)	37(1)
C(18)	3772(3)	404(2)	1139(1)	29(1)
S(1')	2336(1)	2684(1)	4884(1)	28(1)
O(1')	2595(2)	1284(2)	4851(1)	40(1)
O(2')	1557(2)	3209(2)	5472(1)	41(1)
N(1')	6346(2)	2540(2)	5446(1)	26(1)
C(1')	5090(3)	3453(2)	5575(1)	30(1)
C(2')	4136(2)	3544(2)	4853(1)	24(1)
C(3')	5184(2)	2947(2)	4302(1)	22(1)
C(4')	6768(2)	2854(2)	4724(1)	24(1)

C(5')	7624(3)	2692(3)	5992(1)	36(1)
C(6')	7799(3)	1813(2)	4426(1)	28(1)
C(7')	5236(2)	3666(2)	3603(1)	23(1)
C(8')	5217(3)	5032(2)	3549(1)	28(1)
C(9')	5329(3)	5659(3)	2901(1)	33(1)
C(10')	5496(3)	4930(3)	2293(1)	34(1)
C(11')	5530(3)	3564(3)	2329(1)	32(1)
C(12')	5396(2)	2941(2)	2978(1)	28(1)
C(13')	1286(2)	3178(2)	4089(1)	27(1)
C(14')	1364(3)	2439(3)	3478(1)	37(1)
C(15')	494(3)	2832(3)	2866(1)	47(1)
C(16')	-438(3)	3932(3)	2876(2)	47(1)
C(17')	-512(3)	4665(3)	3489(2)	43(1)
C(18')	353(2)	4291(3)	4102(1)	35(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for $\text{C}_{18}\text{H}_{21}\text{NO}_2\text{S}$.

S(1)-O(1)	1.4423(16)
S(1)-O(2)	1.4463(16)
S(1)-C(13)	1.767(2)
S(1)-C(2)	1.786(2)
N(1)-C(5)	1.460(3)
N(1)-C(1)	1.463(3)
N(1)-C(4)	1.465(3)
C(1)-C(2)	1.537(3)
C(2)-C(3)	1.556(3)
C(3)-C(7)	1.509(3)
C(3)-C(4)	1.540(3)
C(4)-C(6)	1.518(3)
C(7)-C(12)	1.384(3)
C(7)-C(8)	1.391(3)
C(8)-C(9)	1.393(4)
C(9)-C(10)	1.375(4)
C(10)-C(11)	1.375(4)
C(11)-C(12)	1.382(3)
C(13)-C(18)	1.388(3)
C(13)-C(14)	1.393(3)
C(14)-C(15)	1.378(4)
C(15)-C(16)	1.384(4)
C(16)-C(17)	1.380(4)
C(17)-C(18)	1.390(4)
S(1')-O(1')	1.4375(19)
S(1')-O(2')	1.4409(17)
S(1')-C(13')	1.771(2)
S(1')-C(2')	1.794(2)
N(1')-C(1')	1.462(3)
N(1')-C(5')	1.469(3)
N(1')-C(4')	1.469(3)
C(1')-C(2')	1.546(3)
C(2')-C(3')	1.552(3)
C(3')-C(7')	1.509(3)

C(3')-C(4')	1.544(3)
C(4')-C(6')	1.517(3)
C(7')-C(8')	1.386(3)
C(7')-C(12')	1.405(3)
C(8')-C(9')	1.387(3)
C(9')-C(10')	1.380(4)
C(10')-C(11')	1.385(3)
C(11')-C(12')	1.389(3)
C(13')-C(14')	1.380(3)
C(13')-C(18')	1.389(3)
C(14')-C(15')	1.394(4)
C(15')-C(16')	1.377(4)
C(16')-C(17')	1.380(4)
C(17')-C(18')	1.388(3)

O(1)-S(1)-O(2)	117.56(10)
O(1)-S(1)-C(13)	108.18(11)
O(2)-S(1)-C(13)	109.01(10)
O(1)-S(1)-C(2)	109.58(10)
O(2)-S(1)-C(2)	107.22(10)
C(13)-S(1)-C(2)	104.52(10)
C(5)-N(1)-C(1)	111.37(19)
C(5)-N(1)-C(4)	115.24(19)
C(1)-N(1)-C(4)	104.15(17)
N(1)-C(1)-C(2)	104.72(17)
C(1)-C(2)-C(3)	105.22(18)
C(1)-C(2)-S(1)	110.05(15)
C(3)-C(2)-S(1)	114.44(14)
C(7)-C(3)-C(4)	113.94(17)
C(7)-C(3)-C(2)	116.82(18)
C(4)-C(3)-C(2)	101.75(17)
N(1)-C(4)-C(6)	112.53(19)
N(1)-C(4)-C(3)	102.24(17)
C(6)-C(4)-C(3)	113.22(19)
C(12)-C(7)-C(8)	117.5(2)
C(12)-C(7)-C(3)	122.3(2)

C(8)-C(7)-C(3)	120.1(2)
C(7)-C(8)-C(9)	121.0(3)
C(10)-C(9)-C(8)	120.3(3)
C(9)-C(10)-C(11)	119.1(3)
C(10)-C(11)-C(12)	120.6(3)
C(11)-C(12)-C(7)	121.4(2)
C(18)-C(13)-C(14)	120.8(2)
C(18)-C(13)-S(1)	118.85(18)
C(14)-C(13)-S(1)	120.37(19)
C(15)-C(14)-C(13)	118.6(2)
C(14)-C(15)-C(16)	121.0(2)
C(17)-C(16)-C(15)	120.3(2)
C(16)-C(17)-C(18)	119.6(2)
C(13)-C(18)-C(17)	119.7(2)
O(1')-S(1')-O(2')	118.63(11)
O(1')-S(1')-C(13')	108.37(11)
O(2')-S(1')-C(13')	107.87(11)
O(1')-S(1')-C(2')	109.75(11)
O(2')-S(1')-C(2')	107.64(10)
C(13')-S(1')-C(2')	103.55(10)
C(1')-N(1')-C(5')	110.65(19)
C(1')-N(1')-C(4')	104.73(17)
C(5')-N(1')-C(4')	113.74(17)
N(1')-C(1')-C(2')	104.53(18)
C(1')-C(2')-C(3')	105.35(17)
C(1')-C(2')-S(1')	110.94(15)
C(3')-C(2')-S(1')	112.68(15)
C(7')-C(3')-C(4')	113.64(17)
C(7')-C(3')-C(2')	116.93(18)
C(4')-C(3')-C(2')	102.69(16)
N(1')-C(4')-C(6')	112.67(18)
N(1')-C(4')-C(3')	102.79(16)
C(6')-C(4')-C(3')	112.23(18)
C(8')-C(7')-C(12')	117.4(2)
C(8')-C(7')-C(3')	123.0(2)
C(12')-C(7')-C(3')	119.5(2)

C(7')-C(8')-C(9')	121.3(2)
C(10')-C(9')-C(8')	120.4(2)
C(9')-C(10')-C(11')	119.7(2)
C(10')-C(11')-C(12')	119.6(2)
C(11')-C(12')-C(7')	121.5(2)
C(14')-C(13')-C(18')	121.1(2)
C(14')-C(13')-S(1')	120.13(18)
C(18')-C(13')-S(1')	118.72(17)
C(13')-C(14')-C(15')	118.9(3)
C(16')-C(15')-C(14')	120.3(3)
C(15')-C(16')-C(17')	120.6(2)
C(16')-C(17')-C(18')	119.9(3)
C(17')-C(18')-C(13')	119.3(2)

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{18}\text{H}_{21}\text{NO}_2\text{S}$.

The anisotropic displacement factor exponent takes the form:

$$-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$$

Atom	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
S(1)	24(1)	20(1)	25(1)	1(1)	-2(1)	0(1)
O(1)	35(1)	21(1)	32(1)	7(1)	0(1)	4(1)
O(2)	33(1)	32(1)	28(1)	-5(1)	-6(1)	-6(1)
N(1)	33(1)	24(1)	28(1)	2(1)	4(1)	-1(1)
C(1)	31(1)	25(1)	30(1)	-6(1)	2(1)	3(1)
C(2)	26(1)	13(1)	27(1)	-1(1)	-1(1)	1(1)
C(3)	24(1)	16(1)	26(1)	-2(1)	-2(1)	2(1)
C(4)	25(1)	21(1)	29(1)	2(1)	0(1)	0(1)
C(5)	49(2)	43(2)	40(2)	5(1)	15(1)	-1(1)
C(6)	33(1)	34(2)	43(2)	3(1)	0(1)	-9(1)
C(7)	21(1)	26(1)	27(1)	-3(1)	1(1)	-2(1)
C(8)	44(2)	38(2)	39(2)	-14(1)	-4(1)	7(1)
C(9)	53(2)	71(2)	31(2)	-25(2)	-2(1)	3(2)
C(10)	43(2)	75(2)	24(2)	7(2)	0(1)	-7(2)
C(11)	57(2)	42(2)	34(2)	9(1)	0(1)	-6(1)
C(12)	47(2)	29(2)	27(1)	1(1)	0(1)	0(1)
C(13)	23(1)	22(1)	27(1)	0(1)	-3(1)	-1(1)
C(14)	43(2)	30(2)	39(2)	2(1)	3(1)	-9(1)
C(15)	59(2)	42(2)	40(2)	7(1)	12(1)	-16(1)
C(16)	49(2)	55(2)	31(1)	1(2)	11(1)	-3(2)
C(17)	38(1)	37(2)	37(2)	-9(1)	4(1)	2(1)
C(18)	27(1)	26(1)	33(1)	2(1)	2(1)	2(1)
S(1')	24(1)	34(1)	27(1)	6(1)	4(1)	-3(1)
O(1')	38(1)	29(1)	51(1)	14(1)	-2(1)	-6(1)
O(2')	31(1)	65(1)	28(1)	4(1)	10(1)	-3(1)
N(1')	26(1)	29(1)	22(1)	4(1)	2(1)	0(1)
C(1')	29(1)	35(2)	26(1)	-4(1)	2(1)	-1(1)
C(2')	24(1)	23(1)	26(1)	1(1)	4(1)	-4(1)
C(3')	21(1)	21(1)	24(1)	-2(1)	3(1)	-2(1)
C(4')	23(1)	21(1)	28(1)	1(1)	3(1)	-2(1)

C(5')	32(1)	45(2)	30(1)	-1(1)	-2(1)	0(1)
C(6')	25(1)	26(1)	32(1)	4(1)	4(1)	1(1)
C(7')	20(1)	28(1)	23(1)	0(1)	1(1)	1(1)
C(8')	34(1)	28(2)	24(1)	-2(1)	4(1)	1(1)
C(9')	40(1)	27(1)	31(1)	4(1)	2(1)	2(1)
C(10')	35(1)	44(2)	24(1)	7(1)	0(1)	-1(1)
C(11')	35(1)	39(2)	23(1)	-5(1)	2(1)	1(1)
C(12')	29(1)	27(1)	29(1)	-3(1)	1(1)	2(1)
C(13')	19(1)	34(1)	27(1)	4(1)	4(1)	-4(1)
C(14')	28(1)	50(2)	34(1)	-2(1)	5(1)	-1(1)
C(15')	35(1)	77(2)	29(1)	-3(2)	3(1)	-5(2)
C(16')	27(1)	74(2)	39(2)	20(2)	-2(1)	-8(1)
C(17')	30(1)	44(2)	54(2)	18(1)	1(1)	-3(1)
C(18')	29(1)	34(1)	40(1)	6(1)	2(1)	-8(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{18}\text{H}_{21}\text{NO}_2\text{S}$.

Atom	x	y	z	U(eq)
H(1A)	7734	-2025	-751	34
H(1B)	6408	-1045	-1079	34
H(2)	6470	-2131	230	27
H(3)	6867	603	531	27
H(4)	9527	-939	215	30
H(5A)	10130	590	-1070	52
H(5B)	8909	-143	-1609	52
H(5C)	10003	-987	-1063	52
H(6A)	10197	1002	855	44
H(6B)	9158	1872	298	44
H(6C)	10687	1165	58	44
H(8)	7489	1121	1729	49
H(9)	8059	504	2910	62
H(10)	8654	-1684	3195	57
H(11)	8594	-3261	2295	53
H(12)	8006	-2659	1123	41
H(14)	3116	-2713	762	44
H(15)	2184	-2658	1890	56
H(16)	2152	-693	2526	53
H(17)	3215	1220	2076	45
H(18)	4197	1182	950	34
H(1'A)	5501	4330	5723	36
H(1'B)	4452	3108	5948	36
H(2')	3929	4494	4735	29
H(3')	4819	2026	4196	26
H(4')	7294	3733	4724	28
H(5'A)	8457	2080	5894	43
H(5'B)	7256	2498	6460	43
H(5'C)	8011	3601	5986	43
H(6'A)	8040	2064	3944	33
H(6'B)	7266	959	4411	33

H(6'C)	8759	1743	4729	33
H(8')	5126	5549	3964	34
H(9')	5289	6595	2876	39
H(10')	5588	5364	1852	41
H(11')	5645	3056	1913	39
H(12')	5414	2004	2999	34
H(14')	2001	1676	3475	44
H(15')	544	2338	2440	56
H(16')	-1035	4187	2457	57
H(17')	-1154	5425	3491	51
H(18')	308	4791	4526	41

Table 6. Torsion angles [°] for C₁₈H₂₁NO₂S.

C(5)-N(1)-C(1)-C(2)	162.32(19)
C(4)-N(1)-C(1)-C(2)	37.5(2)
N(1)-C(1)-C(2)-C(3)	-12.3(2)
N(1)-C(1)-C(2)-S(1)	111.44(17)
O(1)-S(1)-C(2)-C(1)	-61.72(17)
O(2)-S(1)-C(2)-C(1)	66.91(16)
C(13)-S(1)-C(2)-C(1)	-177.45(15)
O(1)-S(1)-C(2)-C(3)	56.50(18)
O(2)-S(1)-C(2)-C(3)	-174.86(15)
C(13)-S(1)-C(2)-C(3)	-59.23(18)
C(1)-C(2)-C(3)-C(7)	-140.42(18)
S(1)-C(2)-C(3)-C(7)	98.65(19)
C(1)-C(2)-C(3)-C(4)	-15.7(2)
S(1)-C(2)-C(3)-C(4)	-136.63(16)
C(5)-N(1)-C(4)-C(6)	67.9(3)
C(1)-N(1)-C(4)-C(6)	-169.79(19)
C(5)-N(1)-C(4)-C(3)	-170.31(18)
C(1)-N(1)-C(4)-C(3)	-48.0(2)
C(7)-C(3)-C(4)-N(1)	164.92(18)
C(2)-C(3)-C(4)-N(1)	38.30(19)
C(7)-C(3)-C(4)-C(6)	-73.8(2)
C(2)-C(3)-C(4)-C(6)	159.61(19)
C(4)-C(3)-C(7)-C(12)	-77.0(3)
C(2)-C(3)-C(7)-C(12)	41.3(3)
C(4)-C(3)-C(7)-C(8)	98.5(2)
C(2)-C(3)-C(7)-C(8)	-143.2(2)
C(12)-C(7)-C(8)-C(9)	-0.5(4)
C(3)-C(7)-C(8)-C(9)	-176.2(2)
C(7)-C(8)-C(9)-C(10)	0.9(4)
C(8)-C(9)-C(10)-C(11)	-1.0(4)
C(9)-C(10)-C(11)-C(12)	0.7(4)
C(10)-C(11)-C(12)-C(7)	-0.3(4)
C(8)-C(7)-C(12)-C(11)	0.2(3)
C(3)-C(7)-C(12)-C(11)	175.8(2)

O(1)-S(1)-C(13)-C(18)	-24.49(19)
O(2)-S(1)-C(13)-C(18)	-153.41(17)
C(2)-S(1)-C(13)-C(18)	92.21(19)
O(1)-S(1)-C(13)-C(14)	154.94(18)
O(2)-S(1)-C(13)-C(14)	26.0(2)
C(2)-S(1)-C(13)-C(14)	-88.36(19)
C(18)-C(13)-C(14)-C(15)	0.7(4)
S(1)-C(13)-C(14)-C(15)	-178.7(2)
C(13)-C(14)-C(15)-C(16)	1.1(4)
C(14)-C(15)-C(16)-C(17)	-2.3(4)
C(15)-C(16)-C(17)-C(18)	1.7(4)
C(14)-C(13)-C(18)-C(17)	-1.3(3)
S(1)-C(13)-C(18)-C(17)	178.18(18)
C(16)-C(17)-C(18)-C(13)	0.1(4)
C(5')-N(1')-C(1')-C(2')	160.74(18)
C(4')-N(1')-C(1')-C(2')	37.8(2)
N(1')-C(1')-C(2')-C(3')	-14.8(2)
N(1')-C(1')-C(2')-S(1')	107.41(17)
O(1')-S(1')-C(2')-C(1')	-75.65(18)
O(2')-S(1')-C(2')-C(1')	54.76(18)
C(13')-S(1')-C(2')-C(1')	168.83(16)
O(1')-S(1')-C(2')-C(3')	42.19(18)
O(2')-S(1')-C(2')-C(3')	172.61(15)
C(13')-S(1')-C(2')-C(3')	-73.33(17)
C(1')-C(2')-C(3')-C(7')	-137.23(19)
S(1')-C(2')-C(3')-C(7')	101.7(2)
C(1')-C(2')-C(3')-C(4')	-12.1(2)
S(1')-C(2')-C(3')-C(4')	-133.20(15)
C(1')-N(1')-C(4')-C(6')	-166.91(18)
C(5')-N(1')-C(4')-C(6')	72.2(2)
C(1')-N(1')-C(4')-C(3')	-45.9(2)
C(5')-N(1')-C(4')-C(3')	-166.82(19)
C(7')-C(3')-C(4')-N(1')	161.99(18)
C(2')-C(3')-C(4')-N(1')	34.7(2)
C(7')-C(3')-C(4')-C(6')	-76.7(2)
C(2')-C(3')-C(4')-C(6')	156.05(18)

C(4')-C(3')-C(7')-C(8')	-81.5(3)
C(2')-C(3')-C(7')-C(8')	37.9(3)
C(4')-C(3')-C(7')-C(12')	94.9(2)
C(2')-C(3')-C(7')-C(12')	-145.7(2)
C(12')-C(7')-C(8')-C(9')	1.0(3)
C(3')-C(7')-C(8')-C(9')	177.4(2)
C(7')-C(8')-C(9')-C(10')	-1.4(4)
C(8')-C(9')-C(10')-C(11')	0.9(4)
C(9')-C(10')-C(11')-C(12')	0.0(4)
C(10')-C(11')-C(12')-C(7')	-0.4(4)
C(8')-C(7')-C(12')-C(11')	0.0(3)
C(3')-C(7')-C(12')-C(11')	-176.6(2)
O(1')-S(1')-C(13')-C(14')	-24.8(2)
O(2')-S(1')-C(13')-C(14')	-154.42(18)
C(2')-S(1')-C(13')-C(14')	91.7(2)
O(1')-S(1')-C(13')-C(18')	153.16(17)
O(2')-S(1')-C(13')-C(18')	23.6(2)
C(2')-S(1')-C(13')-C(18')	-90.34(19)
C(18')-C(13')-C(14')-C(15')	0.4(3)
S(1')-C(13')-C(14')-C(15')	178.32(19)
C(13')-C(14')-C(15')-C(16')	-0.6(4)
C(14')-C(15')-C(16')-C(17')	0.6(4)
C(15')-C(16')-C(17')-C(18')	-0.3(4)
C(16')-C(17')-C(18')-C(13')	0.0(3)
C(14')-C(13')-C(18')-C(17')	-0.1(3)
S(1')-C(13')-C(18')-C(17')	-178.05(17)

ORTEP Diagram of **20**

Experimental:

A colorless prismatic crystal of $C_{19}H_{23}NO_2S$ was coated with Paratone 8277 oil (Exxon) and mounted on a glass fiber. All measurements were made on a Nonius KappaCCD diffractometer with graphite monochromated Mo- $K\alpha$ radiation. Cell constants obtained from the refinement¹ of 9074 reflections in the range $6.0 < \theta < 27.6^\circ$ corresponded to a primitive orthorhombic cell; details of crystal data and structure refinement have been provided in Table 1. The data were collected² at a temperature of 173(2) K using the ω and ϕ scans to a maximum θ value of 27.6° . The data were corrected for Lorentz and polarization effects and for absorption using multi-scan method¹. Since the crystal did not show any sign of decay during data collection a decay correction was deemed unnecessary.

The structure was solved by the direct methods³ and expanded using Fourier techniques⁴. The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included at geometrically idealized positions and were not refined. The final cycle of full-matrix least-squares refinement using SHELXL97⁵ converged (largest parameter shift was 0.00 times its esd) with unweighted and weighted agreement factors, $R = 0.0505$ and $wR = 0.1086$ (all data), respectively, and goodness of fit, $S = 0.955$. The absolute structure was established by the Flack method⁶ with the absolute configuration at the chiral centers: C1 (S), C2 (R) and C3 (S). The Flack parameter for the inverted structure was 1.01(9). Therefore, the inverted structure was rejected as the one present in the crystal. The weighting scheme was based on counting statistics and the final difference Fourier map was essentially featureless. The figures were plotted with the aid of ORTEPII⁷.

References:

8. Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307.
9. Hooft, R. (1998). *COLLECT*. Nonius BV, Delft. The Netherlands.
10. Altomare, A., Cascarano, M., Giacovazzo, C. & Guagliardi, A. (1993). *SIR92. J. Appl. Cryst.*, **26**, 343
11. Beurskens, P.T., Admiraal, G., Beurskens, G., Bosman, W.P., de Gelder, R., Israel, R. & Smits, J.M.M. (1994). The *DIRDIF-94* program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
12. Sheldrick, G.M. (1997). *SHELXL97*. University of Göttingen, Germany.
13. Flack, H. D. (1983). *Acta Cryst.* A39, 876.
14. Johnson, C.K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.

Table 1. Crystal data and structure refinement for C₁₉H₂₃NO₂S.

Identification code	T. Back – 6 (RZ – 147a)
Empirical formula	C ₁₉ H ₂₃ NO ₂ S
Formula weight	329.44
Temperature	173(2) K
Wavelength	0.71073 Å
Crystal system	Orthorhombic
Space group	P2 ₁ 2 ₁ 2 ₁
Unit cell dimensions	a = 5.7550(12) Å b = 39.7980(13) Å c = 7.712(4) Å
Volume	1766.3(9) Å ³
Z	4
Density (calculated)	1.239 Mg/m ³
Absorption coefficient	0.192 mm ⁻¹
F(000)	704
Crystal size	0.18 x 0.18 x 0.17 mm ³
Theta range for data collection	6.0 to 27.6°.
Index ranges	-6 ≤ h ≤ 7, -51 ≤ k ≤ 51, -10 ≤ l ≤ 9
Reflections collected	9074
Independent reflections	4004 [R(int) = 0.083]
Completeness to theta = 27.6°	98.1 %
Absorption correction	Multi-scan method
Max. and min. transmission	0.9680 and 0.9662
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	4004 / 0 / 208
Goodness-of-fit on F ²	0.955
Final R indices [I > 2σ(I)]	R1 = 0.0505, wR2 = 0.0889
R indices (all data)	R1 = 0.1177, wR2 = 0.1086
Absolute structure parameter	-0.01(9)
Largest diff. peak and hole	0.227 and -0.306 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

Atom	x	y	z	$U(\text{eq})$
S(1)	6387(1)	8333(1)	902(1)	27(1)
O(1)	8792(3)	8402(1)	1329(2)	36(1)
O(2)	5707(3)	8335(1)	-908(2)	39(1)
N(1)	5044(4)	8973(1)	4492(3)	25(1)
C(1)	4868(5)	8614(1)	4025(3)	25(1)
C(2)	4556(5)	8623(1)	2022(3)	24(1)
C(3)	5131(5)	8986(1)	1424(3)	26(1)
C(4)	6300(6)	9132(1)	3048(3)	31(1)
C(5)	2867(6)	8445(1)	4971(4)	38(1)
C(6)	2944(5)	9174(1)	874(4)	36(1)
C(7)	6162(6)	9026(1)	6179(3)	31(1)
C(8)	5859(5)	9380(1)	6850(3)	25(1)
C(9)	7564(5)	9529(1)	7878(4)	31(1)
C(10)	7233(6)	9846(1)	8597(4)	37(1)
C(11)	5219(6)	10022(1)	8258(4)	36(1)
C(12)	3507(6)	9880(1)	7208(4)	33(1)
C(13)	3830(5)	9562(1)	6515(3)	29(1)
C(14)	5682(5)	7932(1)	1759(3)	24(1)
C(15)	3562(5)	7784(1)	1328(3)	28(1)
C(16)	3116(5)	7458(1)	1883(4)	33(1)
C(17)	4781(6)	7286(1)	2841(4)	35(1)
C(18)	6841(5)	7436(1)	3297(4)	36(1)
C(19)	7314(5)	7764(1)	2770(4)	30(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$.

S(1)-O(1)	1.449(2)
S(1)-O(2)	1.450(2)
S(1)-C(14)	1.773(3)
S(1)-C(2)	1.786(3)
N(1)-C(7)	1.468(3)
N(1)-C(4)	1.472(3)
N(1)-C(1)	1.476(3)
C(1)-C(5)	1.520(4)
C(1)-C(2)	1.556(4)
C(2)-C(3)	1.550(3)
C(3)-C(6)	1.525(4)
C(3)-C(4)	1.536(4)
C(7)-C(8)	1.511(3)
C(8)-C(9)	1.393(4)
C(8)-C(13)	1.398(4)
C(9)-C(10)	1.392(4)
C(10)-C(11)	1.379(4)
C(11)-C(12)	1.394(4)
C(12)-C(13)	1.385(4)
C(14)-C(19)	1.392(4)
C(14)-C(15)	1.395(4)
C(15)-C(16)	1.390(4)
C(16)-C(17)	1.391(4)
C(17)-C(18)	1.374(4)
C(18)-C(19)	1.395(4)
O(1)-S(1)-O(2)	118.40(12)
O(1)-S(1)-C(14)	107.74(12)
O(2)-S(1)-C(14)	107.63(12)
O(1)-S(1)-C(2)	109.32(13)
O(2)-S(1)-C(2)	107.59(13)
C(14)-S(1)-C(2)	105.43(13)
C(7)-N(1)-C(4)	113.1(2)
C(7)-N(1)-C(1)	112.82(19)

C(4)-N(1)-C(1)	105.5(2)
N(1)-C(1)-C(5)	111.3(2)
N(1)-C(1)-C(2)	103.09(19)
C(5)-C(1)-C(2)	113.6(2)
C(3)-C(2)-C(1)	107.0(2)
C(3)-C(2)-S(1)	109.40(19)
C(1)-C(2)-S(1)	113.35(19)
C(6)-C(3)-C(4)	113.7(2)
C(6)-C(3)-C(2)	111.3(2)
C(4)-C(3)-C(2)	101.8(2)
N(1)-C(4)-C(3)	103.7(2)
N(1)-C(7)-C(8)	112.9(2)
C(9)-C(8)-C(13)	118.3(2)
C(9)-C(8)-C(7)	120.5(3)
C(13)-C(8)-C(7)	121.1(3)
C(8)-C(9)-C(10)	120.9(3)
C(11)-C(10)-C(9)	120.0(3)
C(10)-C(11)-C(12)	119.9(3)
C(13)-C(12)-C(11)	119.9(3)
C(12)-C(13)-C(8)	120.9(3)
C(19)-C(14)-C(15)	121.4(2)
C(19)-C(14)-S(1)	119.1(2)
C(15)-C(14)-S(1)	119.5(2)
C(16)-C(15)-C(14)	118.9(3)
C(17)-C(16)-C(15)	119.8(3)
C(18)-C(17)-C(16)	121.0(3)
C(17)-C(18)-C(19)	120.1(3)
C(14)-C(19)-C(18)	118.8(3)

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$.

The anisotropic displacement factor exponent takes the form:

$$-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$$

Atom	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
S(1)	33(1)	20(1)	28(1)	0(1)	2(1)	2(1)
O(1)	23(1)	32(1)	53(1)	1(1)	5(1)	-2(1)
O(2)	63(2)	32(1)	21(1)	0(1)	1(1)	6(1)
N(1)	36(2)	17(1)	21(1)	-1(1)	0(1)	-5(1)
C(1)	32(2)	17(1)	26(1)	2(1)	-1(2)	1(1)
C(2)	30(2)	19(1)	26(1)	0(1)	-1(1)	0(1)
C(3)	35(2)	18(1)	25(1)	3(1)	8(1)	0(1)
C(4)	37(2)	23(1)	33(2)	4(1)	0(2)	-11(1)
C(5)	52(2)	29(1)	34(2)	0(1)	6(2)	-14(2)
C(6)	46(2)	26(1)	38(2)	6(1)	3(2)	9(1)
C(7)	40(2)	25(1)	29(2)	-2(1)	-7(2)	-1(1)
C(8)	31(2)	24(1)	21(1)	2(1)	1(1)	-5(1)
C(9)	27(2)	35(2)	31(2)	-1(1)	-1(2)	-2(1)
C(10)	42(2)	36(2)	33(2)	-7(1)	0(2)	-12(2)
C(11)	49(2)	23(1)	36(2)	-7(1)	14(2)	-7(2)
C(12)	35(2)	27(1)	36(2)	2(1)	4(2)	0(1)
C(13)	34(2)	26(1)	26(1)	3(1)	-4(2)	-4(1)
C(14)	29(2)	22(1)	23(1)	-5(1)	1(1)	5(1)
C(15)	30(2)	25(1)	29(2)	-4(1)	-4(2)	1(1)
C(16)	35(2)	30(2)	33(2)	-10(1)	2(2)	-4(1)
C(17)	49(2)	22(1)	36(2)	-2(1)	3(2)	-2(2)
C(18)	42(2)	28(1)	39(2)	6(1)	-2(2)	6(1)
C(19)	32(2)	27(1)	31(2)	-1(1)	-4(1)	0(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$.

Atom	x	y	z	U(eq)
H(1)	6356	8497	4319	30
H(2)	2898	8572	1734	29
H(3)	6265	8979	442	31
H(4A)	7973	9074	3080	37
H(4B)	6131	9380	3086	37
H(5A)	3182	8443	6220	57
H(5B)	2702	8213	4557	57
H(5C)	1426	8569	4745	57
H(6A)	2295	9069	-169	55
H(6B)	3335	9409	625	55
H(6C)	1796	9165	1811	55
H(7A)	7842	8977	6073	37
H(7B)	5497	8867	7031	37
H(9)	8974	9411	8092	37
H(10)	8394	9941	9321	44
H(11)	4998	10239	8740	43
H(12)	2122	10001	6968	39
H(13)	2657	9467	5802	34
H(15)	2443	7904	667	34
H(16)	1678	7353	1609	39
H(17)	4490	7060	3185	43
H(18)	7946	7316	3972	43
H(19)	8724	7871	3095	36

Table 6. Torsion angles [°] for C₁₉H₂₃NO₂S.

C(7)-N(1)-C(1)-C(5)	78.6(3)
C(4)-N(1)-C(1)-C(5)	-157.4(2)
C(7)-N(1)-C(1)-C(2)	-159.3(2)
C(4)-N(1)-C(1)-C(2)	-35.4(3)
N(1)-C(1)-C(2)-C(3)	12.6(3)
C(5)-C(1)-C(2)-C(3)	133.1(2)
N(1)-C(1)-C(2)-S(1)	133.3(2)
C(5)-C(1)-C(2)-S(1)	-106.2(2)
O(1)-S(1)-C(2)-C(3)	61.6(2)
O(2)-S(1)-C(2)-C(3)	-68.1(2)
C(14)-S(1)-C(2)-C(3)	177.22(19)
O(1)-S(1)-C(2)-C(1)	-57.7(2)
O(2)-S(1)-C(2)-C(1)	172.49(19)
C(14)-S(1)-C(2)-C(1)	57.8(2)
C(1)-C(2)-C(3)-C(6)	-108.1(3)
S(1)-C(2)-C(3)-C(6)	128.7(2)
C(1)-C(2)-C(3)-C(4)	13.4(3)
S(1)-C(2)-C(3)-C(4)	-109.8(2)
C(7)-N(1)-C(4)-C(3)	169.0(2)
C(1)-N(1)-C(4)-C(3)	45.2(3)
C(6)-C(3)-C(4)-N(1)	84.9(3)
C(2)-C(3)-C(4)-N(1)	-34.9(3)
C(4)-N(1)-C(7)-C(8)	73.6(3)
C(1)-N(1)-C(7)-C(8)	-166.8(2)
N(1)-C(7)-C(8)-C(9)	-148.7(3)
N(1)-C(7)-C(8)-C(13)	33.8(4)
C(13)-C(8)-C(9)-C(10)	1.8(4)
C(7)-C(8)-C(9)-C(10)	-175.7(3)
C(8)-C(9)-C(10)-C(11)	-1.7(4)
C(9)-C(10)-C(11)-C(12)	0.6(4)
C(10)-C(11)-C(12)-C(13)	0.3(4)
C(11)-C(12)-C(13)-C(8)	-0.1(4)
C(9)-C(8)-C(13)-C(12)	-0.9(4)
C(7)-C(8)-C(13)-C(12)	176.6(2)

O(1)-S(1)-C(14)-C(19)	5.2(2)
O(2)-S(1)-C(14)-C(19)	133.9(2)
C(2)-S(1)-C(14)-C(19)	-111.5(2)
O(1)-S(1)-C(14)-C(15)	-171.3(2)
O(2)-S(1)-C(14)-C(15)	-42.5(2)
C(2)-S(1)-C(14)-C(15)	72.1(2)
C(19)-C(14)-C(15)-C(16)	-1.9(4)
S(1)-C(14)-C(15)-C(16)	174.4(2)
C(14)-C(15)-C(16)-C(17)	-0.4(4)
C(15)-C(16)-C(17)-C(18)	2.0(4)
C(16)-C(17)-C(18)-C(19)	-1.3(5)
C(15)-C(14)-C(19)-C(18)	2.6(4)
S(1)-C(14)-C(19)-C(18)	-173.7(2)
C(17)-C(18)-C(19)-C(14)	-1.0(4)

ORTEP Diagram of **21**

Experimental:

A colorless prismatic crystal of $C_{19}H_{23}NO_2S$ was coated with Paratone 8277 oil (Exxon) and mounted on a glass fiber. All measurements were made on a Nonius KappaCCD diffractometer with graphite monochromated Mo- $K\alpha$ radiation. Cell constants obtained from the refinement¹ of 10194 reflections in the range $1.0 < \theta < 27.6^\circ$ corresponded to a primitive orthorhombic cell; details of crystal data and structure refinement have been provided in Table 1. The data were collected² at a temperature of 173(2) K using the ω and ϕ scans to a maximum θ value of 27.6° . The data were corrected for Lorentz and polarization effects and for absorption using multi-scan method¹. Since the crystal did not show any sign of decay during data collection a decay correction was deemed unnecessary.

The structure was solved by the direct methods³ and expanded using Fourier techniques⁴. The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included at geometrically idealized positions and were not refined. The final cycle of full-matrix least-squares refinement using SHELXL97⁵ converged (largest parameter shift was 0.00 times its esd) with unweighted and weighted agreement factors, $R = 0.0524$ and $wR = 0.0954$ (all data), respectively, and goodness of fit, $S = 1.009$. The absolute structure was established by the Flack method⁶ with the absolute configuration at the chiral centers: C1 (R), C2 (S) and C3 (S). The Flack parameter for the inverted structure was 1.07(10). Therefore, the inverted structure was rejected as the one present in the crystal. The weighting scheme was based on counting statistics and the final difference Fourier map was essentially featureless. The figures were plotted with the aid of ORTEPII⁷.

References:

15. Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307.
16. Hooft, R. (1998). *COLLECT*. Nonius BV, Delft. The Netherlands.
17. Altomare, A., Cascarano, M., Giacovazzo, C. & Guagliardi, A. (1993). *SIR92. J. Appl. Cryst.*, **26**, 343
18. Beurskens, P.T., Admiraal, G., Beurskens, G., Bosman, W.P., de Gelder, R., Israel, R. & Smits, J.M.M. (1994). The *DIRDIF-94* program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
19. Sheldrick, G.M. (1997). *SHELXL97*. University of Göttingen, Germany.
20. Flack, H. D. (1983). *Acta Cryst.* A39, 876.
21. Johnson, C.K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.

Table 1. Crystal data and structure refinement for C₁₉H₂₃NO₂S.

Identification code	T. Back – 4 (RZ – 147b)
Empirical formula	C ₁₉ H ₂₃ NO ₂ S
Formula weight	329.44
Temperature	173(2) K
Wavelength	0.71073 Å
Crystal system	Orthorhombic
Space group	P2 ₁ 2 ₁ 2 ₁
Unit cell dimensions	a = 5.6376(4) Å b = 15.1734(10) Å c = 20.0417(8) Å
Volume	1714.50(18) Å ³
Z	4
Density (calculated)	1.276 Mg/m ³
Absorption coefficient	0.198 mm ⁻¹
F(000)	704
Crystal size	0.20 x 0.18 x 0.18 mm ³
Theta range for data collection	6.0 to 27.6°.
Index ranges	-7 ≤ h ≤ 6, -17 ≤ k ≤ 19, -26 ≤ l ≤ 22
Reflections collected	10194
Independent reflections	3871 [R(int) = 0.088]
Completeness to theta = 27.6°	97.8 %
Absorption correction	Multi-scan method
Max. and min. transmission	0.9652 and 0.9614
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	3871 / 0 / 208
Goodness-of-fit on F ²	1.009
Final R indices [I > 2σ(I)]	R1 = 0.0524, wR2 = 0.0796
R indices (all data)	R1 = 0.1219, wR2 = 0.0954
Absolute structure parameter	-0.08(10)
Largest diff. peak and hole	0.226 and -0.292 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

Atom	x	y	z	$U(\text{eq})$
S(1)	38(1)	-3643(1)	-1380(1)	23(1)
O(1)	-2259(3)	-3881(1)	-1644(1)	31(1)
O(2)	1110(4)	-4203(1)	-886(1)	32(1)
N(1)	965(4)	-3260(2)	-3202(1)	22(1)
C(1)	1204(5)	-2800(2)	-2555(1)	22(1)
C(2)	2082(5)	-3514(2)	-2062(1)	21(1)
C(3)	2366(5)	-4360(2)	-2488(1)	27(1)
C(4)	617(5)	-4195(2)	-3054(1)	28(1)
C(5)	2878(5)	-2010(2)	-2604(1)	31(1)
C(6)	2168(7)	-5257(2)	-2151(2)	42(1)
C(7)	-948(5)	-2877(2)	-3603(1)	29(1)
C(8)	-1292(5)	-3329(2)	-4265(1)	22(1)
C(9)	-3350(5)	-3798(2)	-4398(1)	28(1)
C(10)	-3686(5)	-4207(2)	-5009(1)	29(1)
C(11)	-1980(5)	-4152(2)	-5497(1)	27(1)
C(12)	68(6)	-3677(2)	-5371(1)	28(1)
C(13)	400(5)	-3272(2)	-4761(1)	25(1)
C(14)	-194(6)	-2565(2)	-1036(1)	20(1)
C(15)	-2160(5)	-2051(2)	-1171(1)	24(1)
C(16)	-2332(5)	-1219(2)	-893(1)	30(1)
C(17)	-535(5)	-900(2)	-488(1)	30(1)
C(18)	1426(5)	-1414(2)	-354(1)	28(1)
C(19)	1599(5)	-2257(2)	-625(1)	26(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$.

S(1)-O(2)	1.438(2)
S(1)-O(1)	1.444(2)
S(1)-C(14)	1.779(3)
S(1)-C(2)	1.797(3)
N(1)-C(4)	1.463(3)
N(1)-C(7)	1.465(3)
N(1)-C(1)	1.478(3)
C(1)-C(5)	1.529(4)
C(1)-C(2)	1.548(4)
C(2)-C(3)	1.550(4)
C(3)-C(4)	1.523(4)
C(3)-C(6)	1.524(4)
C(7)-C(8)	1.507(4)
C(8)-C(13)	1.379(4)
C(8)-C(9)	1.386(4)
C(9)-C(10)	1.387(4)
C(10)-C(11)	1.375(4)
C(11)-C(12)	1.385(4)
C(12)-C(13)	1.382(4)
C(14)-C(15)	1.382(4)
C(14)-C(19)	1.386(4)
C(15)-C(16)	1.384(4)
C(16)-C(17)	1.385(4)
C(17)-C(18)	1.379(4)
C(18)-C(19)	1.393(4)
O(2)-S(1)-O(1)	118.72(12)
O(2)-S(1)-C(14)	107.86(12)
O(1)-S(1)-C(14)	107.79(13)
O(2)-S(1)-C(2)	108.53(12)
O(1)-S(1)-C(2)	108.94(12)
C(14)-S(1)-C(2)	104.01(12)
C(4)-N(1)-C(7)	113.4(2)
C(4)-N(1)-C(1)	107.0(2)

C(7)-N(1)-C(1)	111.2(2)
N(1)-C(1)-C(5)	111.7(2)
N(1)-C(1)-C(2)	105.0(2)
C(5)-C(1)-C(2)	113.1(2)
C(1)-C(2)-C(3)	105.08(19)
C(1)-C(2)-S(1)	110.86(18)
C(3)-C(2)-S(1)	113.31(18)
C(4)-C(3)-C(6)	115.4(2)
C(4)-C(3)-C(2)	102.0(2)
C(6)-C(3)-C(2)	119.2(2)
N(1)-C(4)-C(3)	102.9(2)
N(1)-C(7)-C(8)	113.5(2)
C(13)-C(8)-C(9)	118.3(3)
C(13)-C(8)-C(7)	121.1(3)
C(9)-C(8)-C(7)	120.6(3)
C(8)-C(9)-C(10)	120.9(3)
C(11)-C(10)-C(9)	120.4(3)
C(10)-C(11)-C(12)	119.0(3)
C(13)-C(12)-C(11)	120.4(3)
C(8)-C(13)-C(12)	121.0(3)
C(15)-C(14)-C(19)	120.7(2)
C(15)-C(14)-S(1)	120.1(2)
C(19)-C(14)-S(1)	119.2(2)
C(14)-C(15)-C(16)	119.5(3)
C(15)-C(16)-C(17)	120.2(3)
C(18)-C(17)-C(16)	120.2(3)
C(17)-C(18)-C(19)	119.9(3)
C(14)-C(19)-C(18)	119.4(3)

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$.

The anisotropic displacement factor exponent takes the form:

$$-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$$

Atom	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
S(1)	26(1)	24(1)	20(1)	1(1)	1(1)	-4(1)
O(1)	23(1)	33(1)	36(1)	-6(1)	3(1)	-9(1)
O(2)	47(1)	27(1)	22(1)	7(1)	-2(1)	2(1)
N(1)	25(1)	24(1)	17(1)	-3(1)	-2(1)	4(1)
C(1)	23(2)	23(2)	19(1)	1(1)	1(1)	-1(1)
C(2)	20(2)	24(2)	20(1)	2(1)	-1(1)	0(1)
C(3)	32(2)	25(2)	26(2)	-4(1)	-2(1)	10(2)
C(4)	34(2)	25(2)	25(2)	-9(1)	-4(1)	5(1)
C(5)	38(2)	31(2)	25(2)	1(1)	-4(1)	-1(2)
C(6)	67(2)	26(2)	32(2)	-1(2)	-4(2)	13(2)
C(7)	32(2)	33(2)	24(1)	-1(2)	-5(1)	10(1)
C(8)	26(2)	22(2)	18(1)	2(1)	-5(1)	7(1)
C(9)	26(2)	34(2)	24(2)	4(1)	2(1)	3(2)
C(10)	25(2)	26(2)	35(2)	1(2)	-9(1)	-3(1)
C(11)	36(2)	26(2)	20(2)	-2(1)	-6(1)	1(2)
C(12)	35(2)	29(2)	21(1)	5(1)	5(2)	3(2)
C(13)	27(2)	24(2)	25(2)	3(1)	-5(1)	-2(1)
C(14)	23(2)	24(1)	14(1)	2(1)	3(1)	-4(2)
C(15)	23(2)	29(2)	21(1)	0(1)	2(1)	-4(1)
C(16)	26(2)	31(2)	31(2)	4(2)	5(1)	5(2)
C(17)	38(2)	25(2)	26(2)	-2(1)	4(1)	-5(2)
C(18)	33(2)	29(2)	23(2)	1(2)	-5(1)	-9(2)
C(19)	27(2)	33(2)	18(1)	4(1)	-2(1)	1(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{19}\text{H}_{23}\text{NO}_2\text{S}$.

Atom	x	y	z	U(eq)
H(1)	-395	-2590	-2409	26
H(2)	3666	-3338	-1882	25
H(3)	3986	-4332	-2690	33
H(4A)	989	-4566	-3446	33
H(4B)	-1032	-4313	-2910	33
H(5A)	2989	-1721	-2167	38
H(5B)	2263	-1591	-2933	38
H(5C)	4455	-2212	-2742	38
H(6A)	2386	-5723	-2484	50
H(6B)	598	-5315	-1946	50
H(6C)	3393	-5309	-1806	50
H(7A)	-596	-2247	-3682	35
H(7B)	-2447	-2911	-3347	35
H(9)	-4545	-3839	-4065	34
H(10)	-5104	-4528	-5091	34
H(11)	-2203	-4436	-5915	33
H(12)	1253	-3630	-5706	34
H(13)	1814	-2948	-4681	30
H(15)	-3384	-2267	-1453	29
H(16)	-3686	-864	-980	36
H(17)	-654	-326	-302	36
H(18)	2658	-1193	-77	34
H(19)	2933	-2618	-528	31

Table 6. Torsion angles [°] for C₁₉H₂₃NO₂S.

C(4)-N(1)-C(1)-C(5)	147.3(2)
C(7)-N(1)-C(1)-C(5)	-88.4(3)
C(4)-N(1)-C(1)-C(2)	24.4(3)
C(7)-N(1)-C(1)-C(2)	148.7(2)
N(1)-C(1)-C(2)-C(3)	1.8(3)
C(5)-C(1)-C(2)-C(3)	-120.3(3)
N(1)-C(1)-C(2)-S(1)	-121.0(2)
C(5)-C(1)-C(2)-S(1)	116.9(2)
O(2)-S(1)-C(2)-C(1)	-170.79(17)
O(1)-S(1)-C(2)-C(1)	58.6(2)
C(14)-S(1)-C(2)-C(1)	-56.1(2)
O(2)-S(1)-C(2)-C(3)	71.3(2)
O(1)-S(1)-C(2)-C(3)	-59.3(2)
C(14)-S(1)-C(2)-C(3)	-174.00(19)
C(1)-C(2)-C(3)-C(4)	-25.7(3)
S(1)-C(2)-C(3)-C(4)	95.5(2)
C(1)-C(2)-C(3)-C(6)	-154.1(3)
S(1)-C(2)-C(3)-C(6)	-32.9(3)
C(7)-N(1)-C(4)-C(3)	-164.3(2)
C(1)-N(1)-C(4)-C(3)	-41.4(3)
C(6)-C(3)-C(4)-N(1)	171.4(2)
C(2)-C(3)-C(4)-N(1)	40.6(3)
C(4)-N(1)-C(7)-C(8)	-58.7(3)
C(1)-N(1)-C(7)-C(8)	-179.3(2)
N(1)-C(7)-C(8)-C(13)	-67.9(3)
N(1)-C(7)-C(8)-C(9)	113.5(3)
C(13)-C(8)-C(9)-C(10)	0.8(4)
C(7)-C(8)-C(9)-C(10)	179.4(3)
C(8)-C(9)-C(10)-C(11)	-0.3(4)
C(9)-C(10)-C(11)-C(12)	-0.4(4)
C(10)-C(11)-C(12)-C(13)	0.6(4)
C(9)-C(8)-C(13)-C(12)	-0.6(4)
C(7)-C(8)-C(13)-C(12)	-179.2(3)
C(11)-C(12)-C(13)-C(8)	0.0(4)

O(2)-S(1)-C(14)-C(15)	-142.3(2)
O(1)-S(1)-C(14)-C(15)	-13.0(2)
C(2)-S(1)-C(14)-C(15)	102.6(2)
O(2)-S(1)-C(14)-C(19)	36.7(2)
O(1)-S(1)-C(14)-C(19)	166.0(2)
C(2)-S(1)-C(14)-C(19)	-78.4(2)
C(19)-C(14)-C(15)-C(16)	0.2(4)
S(1)-C(14)-C(15)-C(16)	179.14(19)
C(14)-C(15)-C(16)-C(17)	0.7(4)
C(15)-C(16)-C(17)-C(18)	-0.6(4)
C(16)-C(17)-C(18)-C(19)	-0.2(4)
C(15)-C(14)-C(19)-C(18)	-1.0(4)
S(1)-C(14)-C(19)-C(18)	179.99(19)
C(17)-C(18)-C(19)-C(14)	1.1(4)

ORTEP Diagram of **24b**

Experimental:

A colorless prismatic crystal of $C_{16}H_{23}NO_2S$ was coated with Paratone 8277 oil (Exxon) and mounted on a glass fiber. All measurements were made on a Nonius KappaCCD diffractometer with graphite monochromated $Mo-K\alpha$ radiation. Cell constants obtained from the refinement¹ of 6614 reflections in the range $3.4 < \theta < 27.5^\circ$ corresponded to a primitive monoclinic cell; details of crystal data and structure refinement have been provided in Table 1. The data were collected² at a temperature of 173(2) K using the ω and ϕ scans to a maximum θ value of 27.5° . The data were corrected for Lorentz and polarization effects and for absorption using multi-scan method¹. Since the crystal did not show any sign of decay during data collection a decay correction was deemed unnecessary.

The structure was solved by the direct methods³ and expanded using Fourier techniques⁴.

There are two independent molecules in the asymmetric unit. The non-hydrogen atoms were refined anisotropically. Hydrogen atoms were included at geometrically idealized positions and were not refined. The final cycle of full-matrix least-squares refinement using SHELXL97⁵ converged (largest parameter shift was 0.00 times its esd) with unweighted and weighted agreement factors, $R = 0.049$ and $wR = 0.114$ (all data), respectively, and goodness of fit, $S = 0.95$. The weighting scheme was based on counting statistics and the final difference Fourier map was essentially featureless. The figures were plotted with the aid of ORTEPII⁶.

References:

22. Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307.
23. Hooft, R. (1998). *COLLECT*. Nonius BV, Delft. The Netherlands.
24. Altomare, A., Cascarano, M., Giacovazzo, C. & Guagliardi, A. (1993). *SIR92*. J. Appl. Cryst., **26**, 343
25. Beurskens, P.T., Admiraal, G., Beurskens, G., Bosman, W.P., de Gelder, R., Israel, R. & Smits, J.M.M. (1994). The *DIRDIF-94* program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, The Netherlands.
26. Sheldrick, G.M. (1997). *SHELXL97*. University of Göttingen, Germany.
27. Johnson, C.K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.

Table 1. Crystal data and structure refinement for C₁₆H₂₃NO₂S.

Identification code	T. Back – 10 (RZ – 5)	
Empirical formula	C ₁₆ H ₂₃ NO ₂ S	
Formula weight	293.41	
Temperature	173(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2 ₁ /n	
Unit cell dimensions	a = 11.310(4) Å	
	b = 12.234(5) Å	β = 106.815(17)°.
	c = 11.597(6) Å	
Volume	1536.0(11) Å ³	
Z	4	
Density (calculated)	1.269 Mg/m ³	
Absorption coefficient	0.21 mm ⁻¹	
F(000)	632	
Crystal size	0.10 x 0.08 x 0.08 mm ³	
Theta range for data collection	3.4 to 27.5°.	
Index ranges	-14 ≤ h ≤ 14, -15 ≤ k ≤ 15, -15 ≤ l ≤ 15	
Reflections collected	6614	
Independent reflections	3503 [R(int) = 0.0953]	
Completeness to theta = 27.5°	99.5 %	
Absorption correction	Multi-scan method	
Max. and min. transmission	0.983 and 0.979	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	3503 / 0 / 181	
Goodness-of-fit on F ²	0.95	
Final R indices [I > 2σ(I)]	R1 = 0.049, wR2 = 0.088	
R indices (all data)	R1 = 0.151, wR2 = 0.114	
Largest diff. peak and hole	0.25 and -0.33 e.Å ⁻³	

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{16}\text{H}_{23}\text{NO}_2\text{S}$. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

Atom	x	y	z	$U(\text{eq})$
S(1)	2333(1)	421(1)	4752(1)	30(1)
O(1)	3204(2)	-475(1)	4997(2)	35(1)
O(2)	1061(1)	185(1)	4111(2)	38(1)
N(1)	2062(2)	586(2)	8139(2)	26(1)
C(1)	1516(2)	-227(2)	8761(2)	36(1)
C(2)	1618(2)	105(2)	10051(2)	40(1)
C(3)	2953(2)	318(2)	10753(2)	39(1)
C(4)	3478(2)	1173(2)	10087(2)	35(1)
C(5)	3351(2)	834(2)	8794(2)	28(1)
C(6)	3820(2)	1728(2)	8129(2)	32(1)
C(7)	3692(2)	1432(2)	6819(2)	30(1)
C(8)	2371(2)	1059(2)	6172(2)	25(1)
C(9)	1984(2)	172(2)	6923(2)	28(1)
C(10)	1432(2)	1996(2)	5913(2)	32(1)
C(11)	2873(2)	1413(2)	3919(2)	27(1)
C(12)	2033(3)	2059(2)	3088(2)	38(1)
C(13)	2462(3)	2828(2)	2429(3)	46(1)
C(14)	3716(3)	2949(2)	2607(3)	48(1)
C(15)	4552(3)	2297(2)	3425(3)	41(1)
C(16)	4126(2)	1522(2)	4079(2)	33(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for $\text{C}_{16}\text{H}_{23}\text{NO}_2\text{S}$.

S(1)-O(2)	1.4455(17)
S(1)-O(1)	1.4460(17)
S(1)-C(11)	1.766(2)
S(1)-C(8)	1.812(3)
N(1)-C(1)	1.466(3)
N(1)-C(5)	1.467(3)
N(1)-C(9)	1.477(3)
C(1)-C(2)	1.522(4)
C(2)-C(3)	1.516(3)
C(3)-C(4)	1.519(3)
C(4)-C(5)	1.522(3)
C(5)-C(6)	1.521(3)
C(6)-C(7)	1.527(3)
C(7)-C(8)	1.535(3)
C(8)-C(10)	1.532(3)
C(8)-C(9)	1.533(3)
C(11)-C(16)	1.381(3)
C(11)-C(12)	1.388(3)
C(12)-C(13)	1.386(4)
C(13)-C(14)	1.381(4)
C(14)-C(15)	1.383(4)
C(15)-C(16)	1.385(3)
O(2)-S(1)-O(1)	118.00(10)
O(2)-S(1)-C(11)	107.85(12)
O(1)-S(1)-C(11)	107.73(11)
O(2)-S(1)-C(8)	108.18(11)
O(1)-S(1)-C(8)	108.22(11)
C(11)-S(1)-C(8)	106.29(11)
C(1)-N(1)-C(5)	112.49(19)
C(1)-N(1)-C(9)	108.66(18)
C(5)-N(1)-C(9)	109.97(19)
N(1)-C(1)-C(2)	112.3(2)
C(3)-C(2)-C(1)	110.7(2)

C(2)-C(3)-C(4)	108.7(2)
C(3)-C(4)-C(5)	111.9(2)
N(1)-C(5)-C(6)	109.08(19)
N(1)-C(5)-C(4)	111.3(2)
C(6)-C(5)-C(4)	110.90(19)
C(5)-C(6)-C(7)	112.83(19)
C(6)-C(7)-C(8)	110.8(2)
C(10)-C(8)-C(9)	110.2(2)
C(10)-C(8)-C(7)	113.29(19)
C(9)-C(8)-C(7)	109.3(2)
C(10)-C(8)-S(1)	108.13(17)
C(9)-C(8)-S(1)	106.02(16)
C(7)-C(8)-S(1)	109.61(16)
N(1)-C(9)-C(8)	110.60(18)
C(16)-C(11)-C(12)	120.5(2)
C(16)-C(11)-S(1)	119.8(2)
C(12)-C(11)-S(1)	119.7(2)
C(13)-C(12)-C(11)	119.4(3)
C(14)-C(13)-C(12)	119.9(3)
C(13)-C(14)-C(15)	120.7(3)
C(14)-C(15)-C(16)	119.6(3)
C(11)-C(16)-C(15)	119.9(3)

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{16}\text{H}_{23}\text{NO}_2\text{S}$.

The anisotropic displacement factor exponent takes the form:

$$-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$$

Atom	U ¹¹	U ²²	U ³³	U ²³	U ¹³	U ¹²
S(1)	25(1)	35(1)	28(1)	-3(1)	5(1)	0(1)
O(1)	33(1)	30(1)	41(1)	-2(1)	10(1)	10(1)
O(2)	21(1)	55(1)	34(1)	-10(1)	1(1)	-7(1)
N(1)	25(1)	31(1)	24(1)	4(1)	8(1)	-2(1)
C(1)	30(2)	40(2)	36(2)	8(1)	7(1)	-7(1)
C(2)	39(2)	47(2)	37(2)	9(1)	16(2)	-3(1)
C(3)	40(2)	46(2)	28(2)	5(1)	8(1)	-3(2)
C(4)	34(2)	40(2)	30(2)	-3(1)	7(1)	-5(1)
C(5)	24(2)	30(1)	27(2)	-1(1)	4(1)	0(1)
C(6)	30(2)	37(2)	27(2)	-2(1)	5(1)	-10(1)
C(7)	26(2)	34(1)	28(2)	0(1)	7(1)	-5(1)
C(8)	24(2)	28(1)	22(2)	-1(1)	7(1)	-2(1)
C(9)	23(1)	30(1)	28(2)	1(1)	4(1)	-2(1)
C(10)	31(2)	34(2)	30(2)	1(1)	9(1)	5(1)
C(11)	28(2)	33(1)	22(2)	-2(1)	9(1)	2(1)
C(12)	39(2)	47(2)	31(2)	3(1)	13(2)	13(1)
C(13)	62(2)	44(2)	35(2)	7(1)	20(2)	17(2)
C(14)	77(2)	35(2)	41(2)	1(1)	32(2)	0(2)
C(15)	45(2)	45(2)	38(2)	-7(1)	21(2)	-7(2)
C(16)	32(2)	44(2)	25(2)	0(1)	10(1)	3(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for $\text{C}_{16}\text{H}_{23}\text{NO}_2\text{S}$.

Atom	x	y	z	U(eq)
H(1A)	1939	-937	8768	43
H(1B)	635	-327	8312	43
H(2A)	1123	774	10047	48
H(2B)	1280	-484	10449	48
H(3A)	3436	-367	10830	46
H(3B)	3001	583	11572	46
H(4A)	4362	1289	10517	42
H(4B)	3041	1874	10085	42
H(5)	3858	162	8809	34
H(6A)	4701	1872	8550	38
H(6B)	3354	2409	8150	38
H(7A)	4279	839	6792	35
H(7B)	3901	2076	6399	35
H(9A)	1126	-61	6516	33
H(9B)	2529	-472	6990	33
H(10A)	1437	2353	6670	38
H(10B)	605	1705	5523	38
H(10C)	1652	2529	5378	38
H(12)	1171	1975	2973	46
H(13)	1894	3272	1855	55
H(14)	4007	3486	2163	57
H(15)	5413	2380	3537	49
H(16)	4694	1066	4636	40

Table 6. Torsion angles [°] for C₁₆H₂₃NO₂S.

C(5)-N(1)-C(1)-C(2)	55.0(3)
C(9)-N(1)-C(1)-C(2)	177.0(2)
N(1)-C(1)-C(2)-C(3)	-56.1(3)
C(1)-C(2)-C(3)-C(4)	55.7(3)
C(2)-C(3)-C(4)-C(5)	-56.0(3)
C(1)-N(1)-C(5)-C(6)	-176.70(19)
C(9)-N(1)-C(5)-C(6)	62.0(2)
C(1)-N(1)-C(5)-C(4)	-54.0(3)
C(9)-N(1)-C(5)-C(4)	-175.29(19)
C(3)-C(4)-C(5)-N(1)	55.4(3)
C(3)-C(4)-C(5)-C(6)	177.0(2)
N(1)-C(5)-C(6)-C(7)	-56.2(3)
C(4)-C(5)-C(6)-C(7)	-179.1(2)
C(5)-C(6)-C(7)-C(8)	51.4(3)
C(6)-C(7)-C(8)-C(10)	72.3(3)
C(6)-C(7)-C(8)-C(9)	-51.1(3)
C(6)-C(7)-C(8)-S(1)	-166.87(16)
O(2)-S(1)-C(8)-C(10)	-50.12(18)
O(1)-S(1)-C(8)-C(10)	-179.05(15)
C(11)-S(1)-C(8)-C(10)	65.47(18)
O(2)-S(1)-C(8)-C(9)	68.06(17)
O(1)-S(1)-C(8)-C(9)	-60.86(18)
C(11)-S(1)-C(8)-C(9)	-176.34(15)
O(2)-S(1)-C(8)-C(7)	-174.06(16)
O(1)-S(1)-C(8)-C(7)	57.01(18)
C(11)-S(1)-C(8)-C(7)	-58.47(19)
C(1)-N(1)-C(9)-C(8)	171.84(18)
C(5)-N(1)-C(9)-C(8)	-64.6(2)
C(10)-C(8)-C(9)-N(1)	-67.0(2)
C(7)-C(8)-C(9)-N(1)	58.2(3)
S(1)-C(8)-C(9)-N(1)	176.26(15)
O(2)-S(1)-C(11)-C(16)	-158.05(19)
O(1)-S(1)-C(11)-C(16)	-29.7(2)
C(8)-S(1)-C(11)-C(16)	86.1(2)

O(2)-S(1)-C(11)-C(12)	20.5(2)
O(1)-S(1)-C(11)-C(12)	148.8(2)
C(8)-S(1)-C(11)-C(12)	-95.4(2)
C(16)-C(11)-C(12)-C(13)	-0.9(4)
S(1)-C(11)-C(12)-C(13)	-179.36(19)
C(11)-C(12)-C(13)-C(14)	-0.4(4)
C(12)-C(13)-C(14)-C(15)	1.1(4)
C(13)-C(14)-C(15)-C(16)	-0.6(4)
C(12)-C(11)-C(16)-C(15)	1.4(4)
S(1)-C(11)-C(16)-C(15)	179.86(19)
C(14)-C(15)-C(16)-C(11)	-0.6(4)