Supporting Information

for

Structurally simple pyridine-*N*-oxides as efficient organocatalysts for the enantioselective allylation of aromatic aldehydes

Luca Pignataro, Maurizio Benaglia,* Rita Annunziata, Mauro Cinquini, and Franco Cozzi

*Dipartimento di Chimica Organica e Industriale - Universita' degli Studi di Milano
via Golgi 19, I-20133 Milano, Italy*

Table of Contents:

Synthesis of catalysts 1b , 2b , 4a , 4b , 6a , 6b , 7a , 7b	pag. S2
Synthesis of <i>N</i> -oxides 10 , 11	pag. S5
HPCL analysis details	pag. S6 ^{S1}

General Methods. TLC was performed on Merck silica gel 60 TLC plates F254 and visualized using UV or phosphomolibdic acid. Flash chromatography was carried out on silica gel (230-400 mesh). ^1H - and ^{13}C -NMR were recorded at 300 and 75 MHz, respectively, with the indicated solvent and at the indicated temperature. Chemical shifts were determined relative to tetramethylsilane (for hydrogen atoms) and residual solvent peaks (for carbon atoms). Optical rotations were obtained on a Perkin-Elmer 241 polarimeter at 589 nm. HPLC for ee determination was performed on Agilent 1100 instrument under the indicated conditions.

Synthesis of **1b** and **2b**.

Suzuki coupling. Following the procedure described for **1a**, product **2-(2,6-dimethoxyphenyl)-pyridine 1b** was obtained in 81% yield as a white solid, m.p. 81-82°C. IR (DCM): 1605, 1456, 1298, 1130 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ 8.71 (d, J = 5.9 Hz, 1H), 7.70 (7, J = 5.6 Hz, 1H), 7.30 (m, 2H), 7.20 (dd, J = 7.5, 4.8 Hz, 1H), 6.63 (d, J =8.0 Hz, 2H), 3.70 (s, 6H). ^{13}C NMR (75 MHz, CDCl_3): δ 158.0, 154.5, 149.6, 135.6, 129.6, 126.1, 121.6, 119.0, 104.1, 55.9. Elem. anal. calcd. for $\text{C}_{13}\text{H}_{13}\text{NO}_2$: C, 72.54; H, 6.09; N, 6.51; Found: C, 72.38; H, 6.18; N, 6.53.

Demethylation. Following the same procedure described for **2a**, product **2-(2,6-hydroxyphenyl)-pyridine 2b** was obtained in 87% yield, as a white solid, m.p. 182°C. IR (DCM): 3350, 1655, 1465, 1030 cm^{-1} . ^1H NMR (300 MHz, DMSO): δ 12.35 (bs, 2H), 8.61 (m, 2H), 7.96 (7, J =7.5 Hz, 1H), 7.36 (dd, J = 6.5, 4.9 Hz, 1H), 7.06 (t, J =8.5 Hz, 1H), 6.40 (d, J =8.5 Hz, 2H). ^{13}C NMR (75 MHz, DMSO): δ 159.1, 156.2, 145.6, 138.1, 130.8, 125.6, 121.3, 108.3 107.4. Elem. anal. calcd. for $\text{C}_{11}\text{H}_9\text{NO}_2$: C, 70.58; $\text{H}^{\text{S}2}$, 4.85; N, 7.48. Found: C, 70.48; H, 4.91; N, 7.43.

Synthesis of **4a** and **4b**.

Esterification. Following the procedure described for **3a**, (*S,S*)-**3-methyl-2-[2',6'-bis(*N*-Cbz-phenylalaninoxy)phenyl]-pyridine 4a** was obtained (50% yield) as a white solid after flash chromatography with a 60:40 hexanes:EtOAc mixture as eluant; m.p. 61°-63 °C; $[\alpha]^{23}_{\text{D}}$ -62.4 (c 0.2

in DCM). IR: 1769, 1725 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ 8.45 (d, J = 4.4 Hz, 1H), 7.45 (m, 2H), 7.15-7.30 (m, 20H), 7.10 (t, J = 5.4 Hz, 1H), 7.05 (m, 2H), 6.90 (m, 2H), 5.10 (m, 2H), 4.90-5.00 (m, 4H), 4.50 (m, 2H), 2.75-2.85 (m, 2H), 2.50-2.55 (m, 2H), 2.10 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): δ 169.0, 156.0, 149.0, 143.0, 140.1, 138.6, 138.0, 135.6, 135.0, 130.9, 130.7, 129.8, 129.7, 129.5, 129.1, 128.6, 127.7, 125.0, 120.1, 119.0, 118.7, 67.8, 67.6, 55.8, 55.5, 37.8, 37.3, 29.5. Elem. anal. calcd. for $\text{C}_{46}\text{H}_{41}\text{N}_3\text{O}_8$: C, 72.33; H, 5.41; N, 5.50. Found: C, 72.45; H, 5.48; N, 5.28.

N-Oxydation. Following the procedure described for **3b**, (*S,S*)-**3-methyl-2-[2',6'-bis(*N*-Cbz-phenylalaninoxy)phenyl]-pyridine-*N*-oxide 4b** was obtained (75% yield) as a white solid after flash chromatography with a 98:2 DCM:MeOH mixture as eluant; m.p. 71°-73 °C; $[\alpha]^{23}_{\text{D}}$ -2.4 (c 0.2 in DCM). IR: 1780, 1711 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ 8.24 (d, J = 5.4 Hz, 1H), 7.54 (t, J = 5.4 Hz, 1H), 7.15-7.30 (m, 20H), 7.00-7.10 (m, 4H), 5.50 (m, 1H), 5.40 (m, 1H), 5.00 (m, 4H), 4.50 (m, 2H), 2.75-3.10 (m, 4H), 2.05 (s, 3H). ^{13}C NMR (75 MHz, CDCl_3): δ 168.0, 155.0, 149.0, 142.0, 139.1, 137.6, 137.0, 135.5, 135.4, 130.8, 130.5, 129.7, 129.6, 129.4, 129.1, 128.6, 128.5, 127.1, 124.9, 120.1, 119.0, 66.8, 66.7, 55.1, 54.8, 37.8, 37.3, 29.5. Elem. anal. calcd. for $\text{C}_{46}\text{H}_{41}\text{N}_3\text{O}_9$: C, 70.85; H, 5.30; N, 5.39. Found: C, 70.45; H, 5.33; N, 5.29.

Synthesis of **6a** and **6b**.

Esterification. To a stirred solution of 2-(2,6-hydroxyphenyl)-3-methylpyridine (100 mg, 0.497 mmol) and a catalytic amount of DMAP in dry THF (6 mL), DCC (218 mg, 1.05 mmol) and *N*-BOC-proline (225 mg, 1.05 mmol) were added. The mixture was stirred for 24 h under nitrogen, the precipitate was filtered, the solvent was evaporated, and the residue was purified by flash chromatography with a 97:3 DCM:MeOH mixture as eluant. (*S,S*)-**3-Methyl-2-[2',6'-bis(*N*-Boc-prolinoxy)phenyl]-pyridine 6a** (0.37 mmol, 75%) was obtained as a white solid: m^{S3} .p. 78-80°C; $[\alpha]^{23}_{\text{D}}$ -112.0 (c 0.2 in DCM). IR: 1771, 1695, cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ 8.50 (bd, J = 4.5 Hz, 1H), 7.40-7.65 (m, 2H), 7.10-7.30 (m, 3H), 4.25-4.35 (m, 2H), 3.30-3.48 (m, 4H), 2.12 (s, 3H),

1.20-1.95 (m, 8H), 1.45 (s, 18H). ^{13}C NMR (75 MHz, CDCl_3): δ 170.2, 154.5, 154.1, 151.0, 148.9, 148.6, 146.6, 146.3, 137.6, 137.4, 134.6, 134.2, 129.3, 126.6, 122.9, 122.8, 120.5, 120.4, 119.6, 119.4, 80.0, 79.8, 58.7, 46.2, 46.1, 30.2, 29.1, 28.3, 23.9, 23.0, 18.3, 18.2. Elem. anal. calcd. for $\text{C}_{32}\text{H}_{41}\text{N}_3\text{O}_8$: C, 64.52; H, 6.94; N, 7.05. Found: C, 64.95; H, 6.88; N, 7.08.

N-Oxydation. Following the procedure described for **3b**, (*S,S*)-**3-methyl-2-[2',6'-bis(*N*-Cbz-prolinoxy)phenyl]-pyridine-*N*-oxide 6b** was obtained (85% yield) as a white solid after flash chromatography with a 96:4 DCM:MeOH mixture as eluant; m.p. 112-113 °C; $[\alpha]^{23}_{\text{D}}$ -88.4 (c 0.25 in DCM). IR: 1772, 1696, cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ 8.21 (d, J = 5.5 Hz, 1H), 7.55-7.60 (m, 1H), 7.10-7.30 (m, 2H), 7.20 (m, 1H), 7.15 (d, J = 6.5 Hz, 1H), 4.25-4.45 (m, 2H), 3.30-3.50 (m, 4H), 2.10 (s, 3H), 1.50-1.95 (m, 8H), 1.49 (s, 18H). ^{13}C NMR (75 MHz, CDCl_3): δ 170.5, 170.3, 154.3, 154.1, 149.2, 149.0, 142.4, 142.2, 138.9, 138.4, 137.2, 136.8, 130.8, 130.6, 126.3, 125.5, 124.9, 124.6, 120.7, 120.4, 119.8, 119.4, 80.2, 79.8, 58.9, 58.6, 46.4, 46.1, 30.2, 29.5, 28.4, 24.1, 23.9, 18.7, 18.2. Elem. anal. calcd. for $\text{C}_{32}\text{H}_{41}\text{N}_3\text{O}_9$: C, 62.83; H, 6.76; N, 6.87. Found: C, 62.78; H, 6.62; N, 6.89.

Synthesis of **7a** and **7b**.

Esterification. Following the procedure described for **5a**, (*S,S*)-**2-[2',6'-bis(*N*-Cbz-prolinoxy)phenyl]-pyridine 7a** was obtained from **2b** in 85% yield as a white solid after flash chromatography with a 97:3 DCM:MeOH mixture as eluant; m.p. 51°-53 °C; $[\alpha]^{23}_{\text{D}}$ -118.4 (c 0.2 in DCM). IR: 1771, 1704 cm^{-1} . ^1H NMR (300 MHz, CDCl_3): δ 8.60 (m, 1H), 7.50 (m, 1H), 7.20-7.40 (m, 14H), 6.50 (t, J =5.3 Hz, 1H), 5.15 (A part of an AB system, J = 7.6 Hz, 1H), 5.05 (B part of an AB system, J = 7.6 Hz, 1H), 5.05 (bs, 2H), 4.40 (m, 2H), 3.40-3.50 (m, 4H), 2.05 (m, 2H), 1.50-1.95 (m, 6H). ^{13}C NMR (75 MHz, CDCl_3): δ 170.3, 154.7, s4 152.1, 149.3, 148.7, 136.6, 135.9, 129.3,

128.5, 128.0, 127.7, 127.5, 125.5, 122.5, 120.5, 120.2, 67.1, 59.1, 46.7, 30.2, 23.9. Elem. anal. calcd. for C₃₇H₃₅N₃O₈: C, 68.40; H, 5.43; N, 6.47. Found: C, 68.45; H, ^{S4} 5.48; N, 6.55.

N-Oxydation. Following the procedure described for **3b**, (*S,S*)-2-[2',6'-bis(*N*-Cbz-prolinoxy)-phenyl]-pyridine-*N*-oxide **7b** was obtained (83% yield) as a white solid after flash chromatography with a 96:4 DCM:MeOH mixture as eluant; m.p. 61°-63 °C; [α]_D²³ -120.6 (c 0.1 in DCM). IR: 1773, 1704 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 8.30 (m, 1H), 7.50-7.40 (m, 1H), 7.20-7.40 (m, 14H), 6.70 (m, 1H), 5.05-5.20 (m, 4H), 4.30-4.40 (m, 2H), 3.40-3.60 (m, 4H), 2.05 (m, 2H), 1.50-1.90 (m, 6H). ¹³C NMR (75 MHz, CDCl₃): δ 170.3, 154.8, 152.1, 149.2, 142.5, 139.6, 136.5, 128.8, 128.6, 128.5, 125.8, 125.5, 124.5, 120.5, 120.2, 119.6, 67.3, 59.2, 46.5, 30.7, 23.5. Elem. anal. calcd. for C₃₇H₃₅N₃O₉: C, 66.76; H, 5.30; N, 6.31. Found: C, 66.78; H, 5.32; N, 6.29.

2-(2,6-dimethoxyphenyl)-3-methylpyridine N-oxide 10. Following the procedure described for **3b**, this product was obtained by oxydation of **1a** in 66% yield as a white solid after flash chromatography with a 97:3 DCM:MeOH mixture as eluant; m.p. 200-205°C (dec); IR: 1610, 1455, 1295, 1130 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 8.26 (bt, 1H), 7.40 (t, *J* = 8.4, 1H), 7.13-7.21 (m, 2H), 6.68 (d, *J* = 8.4, 2H), 3.77 (s, 6H), 2.07 (s, 3H). ¹³C NMR (75 MHz, CDCl₃): δ 157.9, 148.0, 137.4, 137.0, 131.1, 126.7, 123.6, 104.3, 56.0, 18.9. Elem. anal. calcd. for C₁₄H₁₅NO₃: C, 68.55; H, 6.16; N, 5.71. Found: C, 68.80; H, 6.32; N, 5.69.

2-(2,6-dimethylphenyl)-3-methylpyridine N-oxide 11. Following the procedure described for **3b**, this product was obtained by oxydation of 2-(2,6-dimethylphenyl)-6-methylpyridine¹³ in 67% yield as a white solid after flash chromatography with a 97:3 DCM:MeOH mixture as eluant; m.p. 133-135 °C; IR: 1605, 1455, 1300cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 8.26 (t, *J*=4.6, 1H), 7.12-7.24 (m, 5H), 2.01 (s, 6H), 1.95 (s, 3H). ¹³C NMR (75 MHz, CDCl₃): δ 149.5, 137.6, 136.0, 135.0,

129.0, 127.7, 127.0, 123.8, 19.0, 18.5. Elem. anal. calcd. for $C_{14}H_{15}NO$: C, 78.84; H, 7.09; N, 6.57. Found: C, 78.99; H, 7.22; N, 6.40.

1-Phenyl-3-buten-1-ol 8a. This product was purified with a hexanes: EtOAc 90:10 mixture as eluant. It had 1H NMR data in agreement with those reported in the literature.^{5c} The enantiomeric excess was determined by HPLC on a Chiracel OD column (hexane:isopropanol 95:5; flow rate 0.8 ml/min; λ 220 nm): t_R = 10.4 min, t_S = 11.3 min.

1-(4-Methylphenyl)-3-buten-1-ol 8b. This product was purified with a hexanes: EtOAc 90:10 mixture as eluant. It had 1H NMR data in agreement with those reported in the literature.^{5c} The enantiomeric excess was determined by HPLC on a Chiracel OD column (hexane:isopropanol 99:1; flow rate 0.8 ml/min; λ 220 nm): t_R = 26.3 min, t_S = 25.0 min.

1-(4-Bromophenyl)-3-buten-1-ol 8c. This product was purified with a hexanes: EtOAc 90:10 mixture as eluant. It had 1H NMR data in agreement with those reported in the literature.¹⁴ The enantiomeric excess was determined by HPLC on a Chiracel OJ column (hexane:isopropanol 99:1; flow rate 0.8 ml/min; λ 220 nm): t_R = 29.0 min, t_S = 26.0 min.

1-(4-Fluorophenyl)-3-buten-1-ol 8d. This product was purified with a hexanes: EtOAc 90:10 mixture as eluant. It had 1H NMR data in agreement with those reported in the literature.^{5f} For determining its enantiomeric excess it was necessary to convert the product into its acetate (acetic anhydride in pyridine, 16h, room temperature, quantitative yield). The ee was determined by HPLC on a Chiralpak AD column (hexane:isopropanol 99:1; flow rate 0.8 ml/min; λ 220 nm): t_R = 6.4 min, t_S = 7.4 min.

1-(4-Nitrophenyl)-3-buten-1-ol 8e. This product was purified with a hexanes: EtOAc 80:20 mixture as eluant. It had 1H NMR data in agreement with those reported in the literature.^{5k} The

enantiomeric excess was determined by HPLC on a Chiralpak AD column (hexane:isopropanol 97:3; flow rate 0.8 ml/min; λ 220 nm): $t_R = 31.9$ min^{S6}, $t_S = 34.7$ min.

1-(3-Nitrophenyl)-3-buten-1-ol 8f. This product was purified with a hexanes: EtOAc 90:10 mixture as eluant. It had ¹H NMR data in agreement with those reported in the literature.^{5k,15} The enantiomeric excess was determined by HPLC on a Chiralpak AD column (hexane:isopropanol 97:3; flow rate 0.8 ml/min; λ 220 nm): $t_R = 26.3$ min, $t_S = 24.7$ min.

1-(3-Methylphenyl)-3-buten-1-ol 8g. This product was purified with a hexanes: EtOAc 90:10 mixture as eluant. It had ¹H NMR data in agreement with those reported in the literature.¹⁶ The enantiomeric excess was determined by HPLC on a Chiracel OD column (hexane:isopropanol 99:1; flow rate 0.8 ml/min; λ 220 nm): $t_R = 20.1$ min, $t_S = 29.0$ min.

1-Pentafluorophenyl-3-buten-1-ol 8h. This product was purified with a hexanes:EtOAc 95:5 mixture as eluant. It had ¹H NMR data in agreement with those reported in the literature.¹⁷ For determining its enantiomeric excess it was necessary to convert the product into its acetate (acetic anhydride in pyridine, 16h, room temperature, quantitative yield). The ee was determined by HPLC on a Chiracel OD column (hexane:isopropanol 99:1; flow rate 0.8 ml/min; λ 220 nm): $t_S = 5.2$ min, $t_R = 5.9$ min.

(E)-1-Phenyl-1,5-hexadien-3-ol 8i. This product was purified with a hexanes:EtOAc 90:10 mixture as eluant. It had ¹H NMR data in agreement with those reported in the literature.^{5h} The ee was determined by HPLC on a Chiracel OD column (hexane:isopropanol 99:1; flow rate 0.8 ml/min; λ 220 nm): $t_R = 10.3$ min, $t_S = 16.1$ min.

Anti- and syn-1-phenyl-2-methyl-3-buten-1-ol 9. The diastereoisomeric ratio was determined on the crude products exploiting the methyl doublets at 0.87 (*anti*) and 1.01 (*syn*) ppm.^{5h} After purification by flash chromatography with a hexanes:EtOAc 95:5 mixture as eluant, chiral HPLC

analysis (Chiracel AD column; hexane:isopropanol 95:5; flow rate 0.8 ml/min; λ 220 nm)
confirmed the isomer ratio and allowed to establish the ee of the *anti* isomer as 69%: $t_{1R,2R}$ = 8.2
min, $t_{1S,2S}$ = 8.8 min.