Suzuki-Miyaura Cross-Coupling of 1,1-Dichloro-1-alkenes with 9-Alkyl-9-BBN.

Frédéric Liron, Céline Fosse, Alban Pernolet and Emmanuel Roulland*

*Institut de Chimie des Substances Naturelles, CNRS,
Avenue de la Terrasse, 91198 Gif-sur-Yvette, France*

emmanuel.roulland@icsn.cnrs-gif.fr

Supporting Information

Table of Contents:

General and Material.....S3	Typical procedure for non-selective cross-coupling of 1,1-dichloro alkenes:..... S10
Experimental procedures and characterization data:	20, 28 S11
	12S11
Typical procedure for the preparation of 1,1-dichloro alkenes (method A):.....S3	Typical procedure for second coupling, leading to tri-substituted olefins:..... S12
2S3	32 S12
4 S3	31 S12
	33-Z S12
Typical procedure for the preparation of 1,1-dichloro alkenes (method B):.....S4	33-E S13
1 S4	Typical procedure for Suzuki cross-coupling with phenylboronic acid:..... S13
3 S4	35S13
5 S4	36S14
6 S5	
Typical procedure for stereoselective mono coupling:..... S5	¹ H and ¹³ C NMR:
11S5	1 (¹ H NMR).....S15
18S5	1 (¹³ C NMR)..... S16
15S6	2 (¹ H NMR).....S17
19 S6	2 (¹³ C NMR)..... S18
29, 30 S6	3 (¹ H NMR)..... S19
20S7	3 (¹³ C NMR)..... S20
16 S7	4 (¹ H NMR).....S21
17 S7	4 (¹³ C NMR)..... S22
21 S8	5 (¹ H NMR).....S23
22 S8	5 (¹³ C NMR)..... S24
27 S8	6 (¹ H NMR).....S25
26S9	6 (¹³ C NMR)..... S26
25-cis, 25-trans S9	11 (¹ H NMR).....S27
23-cis, 23-trans S10	11 (¹³ C NMR).....S28
24-cis+transS10	12 (¹ H NMR).....S29
	12 (¹³ C NMR)..... S30
	18 (¹ H NMR).....S31
	18 (¹³ C NMR)..... S32

19 (¹ H NMR).....	S33
19 (¹³ C NMR).....	S34
15 (¹ H NMR).....	S35
15 (¹³ C NMR).....	S36
20 (¹ H NMR).....	S37
20 (¹³ C NMR).....	S38
28 (¹ H NMR).....	S39
28 (¹³ C NMR).....	S40
23-cis (¹ H NMR).....	S41
23-cis (¹³ C NMR).....	S42
23-trans (¹ H NMR).....	S43
23-trans (¹³ C NMR).....	S44
24-cis+trans (¹ H NMR).....	S45
24-cis+trans (¹³ C NMR).....	S46
25-cis (¹ H NMR).....	S47
25-cis (¹³ C NMR).....	S48
25-trans (¹ H NMR).....	S49
25-trans (¹³ C NMR).....	S50
29 (¹ H NMR).....	S51
29 (¹³ C NMR).....	S52
30 (¹ H NMR).....	S53
30 (¹³ C NMR).....	S54
16 (¹ H NMR).....	S55

16 (¹³ C NMR).....	S56
22 (¹ H NMR).....	S57
22 (¹³ C NMR).....	S58
17 (¹ H NMR).....	S59
17 (¹³ C NMR).....	S60
21 (¹ H NMR).....	S61
21 (¹³ C NMR).....	S62
26 (¹ H NMR).....	S63
26 (¹³ C NMR).....	S64
27 (¹ H NMR).....	S65
27 (¹³ C NMR).....	S66
33-E (¹ H NMR).....	S67
33-E (¹³ C NMR).....	S68
33-Z (¹ H NMR).....	S69
33-Z (¹³ C NMR).....	S70
35 (¹ H NMR).....	S71
35 (¹³ C NMR).....	S72
36 (¹ H NMR).....	S73
36 (¹³ C NMR).....	S74
31 (¹ H NMR).....	S75
31 (¹³ C NMR).....	S76
32 (¹ H NMR).....	S77
32 (¹ H NMR).....	S78

Experimental section

General experimental procedures: All reactions were carried out under an argon atmosphere. Air or water-sensitive reagents were transferred using Schlenk techniques. NMR spectra were recorded (BRUKER 300 MHz, ^1H , and 75 MHz, ^{13}C) in CDCl_3 (which also provided the lock signal at $\delta = 7.26$ and $\delta = 77.16$ ppm). IR spectra were recorded on a Perkin Elmer FT-IR system using diamond window Dura SamplIR. Mass spectra were determined with Micromass LCT. Silica gel (36-70 μm) was used for flash chromatography. TLC plates (Merck 60 F₂₅₄ aluminum sheets) were visualized by ultraviolet and spraying with phosphomolybdic acid (5% in ethanol) or KMnO_4 (5% in water). THF and ether were distilled under argon from sodium/benzophenone prior to use. Dichloromethane was distilled from CaH_2 under argon prior to use.

Typical procedure for the preparation of 1,1-dichloro alkenes (method A): DBU (15.0 mL, 99.8 mmol) was added dropwise to a solution of the aldehyde (45.0 mmol) in chloroform (9.0 mL, 111.9 mmol). The reaction mixture was stirred at rt for 5 h then Ac_2O (13.0 mL, 138.5 mmol) was added. The solution was stirred for 16 h. The mixture was diluted with chloroform and washed with 2N HCl. The organic phases were dried over Na_2SO_4 and concentrated *in vacuum*. The residue was dissolved in AcOH (45.0 mL, 789.1 mmol) and Zn powder (6.0 g, 91.8 mmol) was added and the mixture was warmed to 50 $^\circ\text{C}$ for 1 h. After cooling to rt, the reaction mixture was neutralized by 2N NaOH and extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 , concentrated *in vacuum* and distilled off with a Büchi oven, yielding the desired 1,1-dichloro-1-alkene.

1,1-Dichloro-2-(4-methoxyphenyl)ethane (2):

Prepared according to method A from *p*-anisaldehyde. Distillation afforded 8.4 g (93% overall) of 1,1-dichloro-1-alkene **2** as a colorless oil.

IR (film) 2956, 2836, 1604, 1508, 1250, 1176, 1031 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 3.81 (s, 3H), 6.78 (s, 1H), 6.89 (dd, $J = 2.1, 6.6$ Hz, 2H), 7.49 (dd, $J = 2.1, 6.6$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 55.3, 113.9, 118.8, 126.0, 128.6, 130.4, 159.6; MS (EI) m/z 20 (M^+ , ^{37}Cl), 202 (M^+ , ^{35}Cl), 189 (M-Me), 187 (M-Me), 161, 159, 137, 131, 123, 89; Anal. Calcd. for $\text{C}_9\text{H}_8\text{Cl}_2\text{O}$: C 53.23; H 3.97, found C 53.47; H 3.92.

1,1-Dichloro-2-(4-fluorophenyl)ethane (4):

Prepared according to method A from 4-fluorobenzaldehyde. Distillation afforded 7.8 g (79% overall) of 1,1-dichloro-1-alkene **4** as a colorless oil.

IR (film) 3037, 2359, 1602, 1506, 1231, 1160, 910 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 6.82 (s, 1H), 7.03-7.10 (m, 2H), 7.49-7.55 (m, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 115.5 (d, $J = 21.3$ Hz), 120.9, 127.4, 129.5 (d, $J = 3.8$ Hz), 130.5 (d, $J = 8.3$ Hz), 162.3 (d, $J = 247.6$ Hz); MS (EI) m/z 192 (M^+ , ^{37}Cl), 190 (M^+ , ^{35}Cl), 167, 155 (M-Cl), 125, 123, 120, 109, 95; Anal. calcd. for $\text{C}_8\text{H}_5\text{Cl}_2\text{F}$: C 50.30; H 2.64; found C: 50.23; H 2.48.

Typical procedure for the preparation of 1,1-dichloro alkenes (method B): The aldehyde (50.0 mmol) and CCl_4 (10 mL, 103.3 mmol) were added to a suspension of Al foil (1.835 g, 70.4 mmol) and PbBr_2 (208 mg, 0.6 mol%) in DMF (200 mL). A strong exothermic reaction occurred and Al was consumed within 1 h. The reaction mixture was quenched with water and extracted with Et_2O . After extraction, the combined organic layers were dried over Na_2SO_4 and concentrated *in vacuum*. The residue was dissolved in pyridine (12 mL, 149.2 mmol) and Ac_2O (8.4 mL, 88.8 mmol) was added. The reaction mixture was stirred for 4 h, then hydrolyzed by 2N HCl. The solution was extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuum*. The solid residue was dissolved in AcOH (40 mL, 700.1 mmol) and Zn powder (6.51 g, 99.6 mmol) was added. The reaction mixture was warmed at 50 °C for 1 h. The solution was neutralized by 2N NaOH and extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuum*. The residue was distilled off with a Büchi oven, yielding the expected 1,1-dichloro-1-alkene.

(2,2-dichlorovinyl)benzene (1):

Prepared according to method B from benzaldehyde. Distillation afforded 7.33 g (87 % overall) of 1,1-dichloro-1-alkene **1** as a colorless oil.

IR (film) 3058, 3027, 1608, 1446, 900 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 6.58 (s, 1H), 7.34 (m, 3H), 7.52 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 121.1, 128.60, 128.69, 128.77; MS (EI) m/z 173 (M^+ , ^{37}Cl), 171 (M^+ , ^{35}Cl); Anal. calcd. for $\text{C}_{10}\text{H}_8\text{Cl}_2$: C 55.53; H 3.49; found C 54.84; H 3.52.

1,1-Dichloro-2-(4-methoxycarbonylphenyl)ethane (3):

Prepared according to method B from 4-methoxycarbonylbenzaldehyde. Distillation afforded 2.20 g (80% overall) of 1,1-dichloro-1-alkene **3** as a slightly yellow solid

m.p.: 57-58 °C; IR (solid) 2952, 1715, 1597, 1438, 1272, 1115 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 3.93 (s, 3H), 6.90 (s, 1H), 7.60 (dd, $J = 1.8, 6.3$ Hz, 2H), 8.03 (dd, $J = 1.8, 6.6$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 52.3, 123.3, 127.8, 128.6, 129.1, 129.7, 137.7, 160.3; MS (EI) m/z 232 (M^+ , ^{37}Cl), 230 (M^+ , ^{35}Cl), 212, 201, 199, 165, 163, 136, 121, 101; HRMS (DCI) m/z calcd. for $\text{C}_{10}\text{H}_9\text{Cl}_2\text{O}_2$ $[\text{M}+\text{H}]^+$ 234.9920, found 234.9915.

1,1-Dichloro-2-(2,4-dimethoxyphenyl)ethane (5):

Prepared according to method B from 2,4-dimethoxybenzaldehyde. Distillation afforded 7.70 g (65% overall) of 1,1-dichloro-1-alkene **5** as a colorless oil.

IR (film) 2940, 2837, 1605, 1575, 1498, 1455, 1293, 1264, 1207, 1159, 1121, 1032 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 3.81 (s, 3H), 3.83 (s, 3H), 6.43 (d, $J = 2.4$ Hz, 1H), 6.50 (dd, $J = 2.4, 8.7$ Hz, 1H), 7.02 (s, 1H), 7.70 (d, $J = 8.6$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 55.4, 55.5, 98.1, 104.2, 115.3, 118.9, 123.3, 129.7, 158.0, 161.1; MS (EI) m/z 234 (M^+ , ^{37}Cl), 232

(M^+ , ^{35}Cl), 198 ($M-\text{Cl}$), 182, 167, 155, 121; HRMS (DCI) m/z calcd. for $\text{C}_{10}\text{H}_{11}\text{Cl}_2\text{O}_2$ [$M+\text{H}$] $^+$ 233.0137, found 233.0139.

1,1-dichlorohept-1-ene (6):

Prepared according to method B from hexanal. Distillation afforded 2.70 g (49.5% overall) of 1,1-dichloro-1-alkene **6** as a colorless oil.

IR (film) 2957, 2926, 2858, 1621, 1458, 882 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.90 (m, 3H), 1.31 (m, 4H), 1.41 (m, $J = 7.1$ Hz, 2H), 2.16 (q, $J = 7.3$ Hz, 2H), 5.85 (t, $J = 7.5$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 22.6, 28.0, 29.7, 31.4, 119.8, 130.3; MS (EI) m/z 168 (M^+ , ^{37}Cl), 166 (M^+ , ^{35}Cl); HRMS (DCI) m/z calcd. for $\text{C}_7\text{H}_{12}\text{Cl}_2$ [M] $^+$ 166.0316, found 166.0315.

Typical procedure for stereoselective mono coupling: A terminal alkene (1.2 mmol) was added to a solution of 9-BBN (0.5 M in THF, 2.4 mL, 1.2 mmol). The solution was stirred at rt for 1 h. A solution of the 1,1-dichloro-1-alkene (1.0 mmol) in THF (2 mL), was added followed by $\text{Pd}_2(\text{dba})_3$ (23.1 mg, 2.5 mol%), XantPhos (28.5 mg, 5 mol%), KF (180 mg, 3.0 mmol) and K_3PO_4 (635 mg, 3.0 mmol) (method A). Alternatively, CsF (454.4 mg, 3.0 mmol) and Cs_2CO_3 (979.6 mg, 3.0 mmol) were used instead of KF and K_3PO_4 (method B). The solution was refluxed until completion (see Table 3 for reaction times). After cooling to rt, the reaction mixture was quenched with water and extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuum*. The residue was purified by column chromatography, affording the mono coupled product.

(Z)-2-Chloro-1-(4-methoxyphenyl)-5-phenylpent-1-ene (11):

Prepared according to typical procedure from allylbenzene (495 μL , 4.17 mmol) and 1,1-dichloro-1-alkene **2** (609 mg, 3.00 mmol). Purified by column chromatography (heptane/ CH_2Cl_2 , 85/15) afforded **1** as a colorless solid (741 mg, 86%, method A; 758 mg, 88%, method B).

m.p.: 95-96 $^{\circ}\text{C}$; IR (solid) 2931, 2360, 1605, 1508, 1247, 1176, 1032 cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz) δ_{H} 2.00 (m, 2H), 2.49 (t, $J = 7.2$ Hz, 2H), 2.67 (t, $J = 7.5$ Hz, 2H), 3.81 (s, 3H), 6.39 (s, 1H), 6.88 (dd, $J = 2.1, 6.6$ Hz, 2H), 7.20 (d, $J = 6.9$ Hz, 2H), 7.25-7.32 (m, 3H), 7.57 (dd, $J = 2.1, 6.9$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 29.1, 34.6, 40.5, 55.2, 113.5, 124.1, 125.9, 127.7, 128.3, 128.5, 130.3, 132.5, 141.8, 158.8; MS (EI) m/z 288 (M^+ , ^{37}Cl), 286 (M^+ , ^{35}Cl), 181, 159, 147, 145, 131, 121, 115; Anal. calcd. for $\text{C}_{18}\text{H}_{19}\text{ClO}$: C 75.38; H 6.68, found C 75.32; H 6.86.

(Z)-2-Chloro-1-(4-methoxycarbonylphenyl)-5-phenylpent-1-ene (18):

Prepared according to typical procedure from allylbenzene (165 μL , 1.39 mmol) and 1,1-dichloro-1-alkene **3** (230 mg, 1.00 mmol). Purification by column chromatography

(heptane/CH₂Cl₂, 7/3) yielded **18** as a colorless oil (210 mg, 65%, method A; 235 mg, 73%, method B).

IR (film) 2925, 1718, 1606, 1433, 1274, 1181, 1106 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ_H 2.01 (m, 2H), 2.54 (t, *J* = 7.5 Hz, 2H), 2.69 (t, *J* = 7.5 Hz, 2H), 3.91 (s, 3H), 6.49 (s, 1H), 7.19-7.23 (m, 3H), 7.28-7.33 (m, 2H), 7.65 (dd, *J* = 1.8, 6.6 Hz, 2H), 8.01 (dd, *J* = 1.8, 6.6 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃) δ_C 29.0, 34.6, 40.6, 52.1, 124.0, 125.9, 128.40, 128.42, 128.8, 129.4, 136.9, 139.6, 141.5, 166.8; MS (EI) *m/z* 316 (M⁺, ³⁷Cl), 314 (M⁺, ³⁵Cl), 285, 283, 279 (M-Cl), 249, 247; HRMS (DCI) *m/z* calcd. for C₁₉H₁₉ClO₂ [M]⁺ 314.1073, found 314.1066.

(Z)-2-Chloro-1,5-diphenylpent-1-ene (15)

Prepared according to typical procedure from allylbenzene (165 μL, 1.39 mmol) and 1,1-dichloro-1-alkene **1** (173 mg, 1.0 mmol). Purification by column chromatography (heptane) afforded **15** as a colorless oil (170 mg, 69%; method A; 197 mg, 80%, method B).

IR (film) 2929, 1494, 1446, 1092, 906, 729 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ_H 2.01 (m, 2H), 2.52 (t, *J* = 7.5 Hz, 2H), 2.68 (t, *J* = 7.5 Hz, 2H), 6.46 (s, 1H), 7.09-7.38 (m, 8H), 7.56-7.61 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ_C 29.1, 34.6, 40.5, 124.8, 125.9, 127.5, 128.1, 128.4, 128.5, 129.0, 134.5, 135.1, 141.8; MS (EI) *m/z* 258 (M⁺, ³⁷Cl), 256 (M⁺, ³⁵Cl), 221 (M-Cl), 129, 115, 104; HRMS (EI) *m/z* calcd. for C₁₇H₁₇Cl [M]⁺ 256.1019, found 256.1010.

(Z)-2-Chloro-1-(4-fluorophenyl)-5-phenylpent-1-ene (19):

Prepared according to typical procedure from allylbenzene (165 μL, 1.39 mmol) and 1,1-dichloro-1-alkene **4** (190 mg, 1.00 mmol). Purification by column chromatography (heptane) afforded **19** as a colorless oil (225 mg, 82%, method A; 186 mg, 68%, method B).

IR (film) 2930, 1602, 1506, 1453, 1226, 1159, 1104 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ_H 1.95-2.05 (m, 2H), 2.50 (t, *J* = 7.2 Hz, 2H), 2.67 (t, *J* = 6.9 Hz, 2H), 6.41 (s, 1H), 6.97-7.06 (m, 2H), 7.17-7.32 (m, 5H), 7.53-7.59 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ_C 29.1, 34.6, 40.4, 115.1 (d, *J* = 21.3 Hz), 123.6, 125.9, 128.4, 128.45, 130.6, 130.7, 131.1, 131.15, 131.2, 134.3, 141.7, 161.9 (d, *J* = 245.9 Hz); MS (EI) *m/z* 276 (M⁺, ³⁷Cl), 274 (M⁺, ³⁵Cl), 239 (M-Cl), 104; HRMS (EI) *m/z* calcd. for C₁₇H₁₆ClF [M]⁺ 274.0925, found 274.0932.

(Z)-1,2-Dichloro-5-phenylpent-1-ene (29) and (Z) 5-chloro-1,8-diphenyloct-4-ene (30):

Prepared according to typical procedure from allylbenzene (660 μL, 5.96 mmol) and trichloroethylene **9** (130 mg, 0.99 mmol). Purification by column chromatography (heptane) afforded the mono coupled product **29** (16 mg, 29%) and the bis coupled product **30** (61 mg, 21%) as colorless oils.

Data for **29**:

IR (film) 2939, 1614, 1496, 1453, 1098, 869, 747 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ_H 1.87-1.97 (m, 2H), 2.40 (t, *J* = 7.3 Hz, 2H), 2.62 (t, *J* = 7.5 Hz, 2H), 6.13 (s, 1H), 7.14-7.22 (m,

3H), 7.25-7.32 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 28.6, 34.4, 36.9, 113.8, 126.0, 128.4, 130.6, 137.1, 138.1, 141.2; MS (EI) m/z 216 (M^+ , ^{37}Cl), 214 (M^+ , ^{35}Cl), 205, 179 (M-Cl), 143, 128, 115, 105; Anal. calcd. for $\text{C}_{11}\text{H}_{12}\text{Cl}_2$: C 61.42; H 5.62; found C 61.66; H 5.80.

Data for **30**:

IR (film) 2927, 1602, 1495, 1452, 908, 732 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 1.66-1.74 (m, 2H), 1.83-1.93 (m, 2H), 2.19-2.26 (m, 2H), 2.32-2.37 (m, 2H), 2.58-2.66 (m, 4H), 5.47 (t, $J = 6.9$ Hz, 1H), 7.17-7.20 (m, 5H), 7.27-7.31 (m, 5H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 28.2, 29.0, 30.5, 34.6, 35.5, 38.9, 125.5, 125.7, 125.8, 128.31, 128.35, 128.4, 128.5, 134.7, 141.9, 142.3; MS (EI) m/z 300 (M^+ , ^{37}Cl), 298 (M^+ , ^{35}Cl), 263 (M-Cl), 194, 185, 171, 157, 144, 128, 116, 106; Anal. calcd. for $\text{C}_{20}\text{H}_{23}\text{Cl}$: C 80.38; H 7.76; found C 79.66; H 7.78

(Z)-2-Chloro-1-(2,4-dimethoxyphenyl)-5-phenylpent-1-ene (20):

Prepared according to typical procedure from allylbenzene (165 μL , 1.39 mmol) and 1,1-dichloro-1-alkene **5** (233 mg, 0.99 mmol). Reaction time: 44 h. Purification by column chromatography (heptane/ CH_2Cl_2 , 7/3) afforded **20** as a colorless oil (260 mg, 82%, method A; 181 mg, 57%, method B).

IR (film) 2944, 1607, 1577, 1499, 1453, 1288, 1262, 1157, 1128, 1118, 1032 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 1.94-2.04 (m, 2H), 2.52 (t, $J = 6.9$ Hz, 2H), 2.68 (t, $J = 7.5$ Hz, 2H), 3.80 (s, 3H), 3.82 (s, 3H), 6.44 (d, $J = 2.4$ Hz, 1H), 6.50 (dd, $J = 2.4, 8.4$ Hz, 1H), 6.60 (s, 1H), 7.16-7.32 (m, 5H), 7.78 (d, $J = 8.4$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 29.2, 34.6, 40.4, 55.4, 55.5, 98.2, 103.9, 117.0, 119.4, 125.8, 128.25, 128.3, 128.6, 130.5, 133.2, 158.0, 160.3; MS (EI) m/z 318 (M^+ , ^{37}Cl), 316 (M^+ , ^{35}Cl), 211, 177, 151, 115, 103, 91; HRMS (DCI) m/z calcd. for $\text{C}_{19}\text{H}_{21}\text{ClO}_2$ $[\text{M}]^+$ 316.1230, found 316.1244.

(Z)-2-Chloro-5-benzyloxy-1-(4-methoxyphenyl)pent-1-ene (16):

Prepared according to typical procedure from allylbenzylether (165 μL , 1.08 mmol) and 1,1-dichloro-1-alkene **2** (203 mg, 1.00 mmol). Purification by column chromatography (heptane/ CH_2Cl_2 , 7/3) afforded **16** as a light yellow oil (263 mg, 83%, method A; 250mg, 79%, method B).

IR (film) 2926, 2853, 1606, 1509, 1248, 1177, 1101, 1030 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 1.92-2.01 (m, 2H), 2.59 (t, $J = 7.5$ Hz, 2H), 3.53 (t, $J = 6.3$ Hz, 2H), 3.81 (s, 3H), 4.51 (s, 2H), 6.40 (s, 1H), 6.87 (dd, $J = 1.8, 6.9$ Hz, 2H), 7.25-7.35 (m, 5H), 7.53 (dd, $J = 2.1, 6.9$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 27.7, 37.8, 55.2, 68.8, 73.0, 113.5, 124.2, 127.6, 127.7, 128.4, 130.2, 132.2, 138.5, 158.8; MS (EI) m/z 316 (M^+ , ^{35}Cl), 227, 225, 207, 181, 172, 159, 145, 115; Anal. calcd. for $\text{C}_{19}\text{H}_{21}\text{ClO}_2$: C 72.03; H 6.68; found C 71.71; H 6.99.

(Z)-2-Chloro-1-(4-methoxyphenyl)non-1-ene (17):

Prepared according to typical procedure from hept-1-ene (170 μ L, 1.28 mmol) and 1,1-dichloro-1-alkene **2** (203 mg, 1.00 mmol). Purification by column chromatography (heptane/ CH_2Cl_2 , 85/15) afforded **17** as a colorless oil (223 mg, 83%, method A; 164 mg, 61%, method B).

IR (film) 2925, 2854, 1606, 1508, 1248, 1177, 1034 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.86-0.91 (m, 3H), 1.27-1.34 (m, 8H), 1.59-1.69 (m, 2H), 2.45 (t, $J = 7.2$ Hz, 2H), 3.81 (s, 3H), 6.39 (s, 1H), 6.85-6.89 (m, 2H), 7.53-7.58 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 22.6, 27.7, 28.6, 29.1, 31.8, 55.2, 113.5, 123.5, 127.9, 130.2, 133.3, 158.7; MS (EI) m/z 268 (M^+ , ^{37}Cl), 266 (M^+ , ^{35}Cl), 183, 181, 145, 121, 115, 103; HRMS (DCI) m/z calcd. for $\text{C}_{16}\text{H}_{23}\text{ClO}$ [M] $^+$ 266.1437, found 266.1437.

(Z)-2-Chloro-1-(2,4-dimethoxyphenyl)non-1-ene (21):

Prepared according to typical procedure from hept-1-ene (170 μ L, 1.28 mmol) and 1,1-dichloro-1-alkene **5** (233 mg, 1.00 mmol). Purification by column chromatography (heptane/ CH_2Cl_2 , 7/3) afforded **21** as a slightly yellow oil (248 mg, 80%, method A; 245 mg, 79%, method B).

IR (film) 2925, 2854, 1608, 1578, 1458, 1289, 1263, 1206, 1157, 1128, 1035 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.87-0.91 (m, 3H), 1.24-1.36 (m, 8H), 1.60-1.69 (m, 2H), 2.48 (t, $J = 7.5$ Hz, 2H), 3.80 (s, 3H), 3.82 (s, 3H), 6.43 (d, $J = 2.4$ Hz, 1H), 6.50 (dd, $J = 2.1, 8.7$ Hz, 1H), 6.57 (s, 1H), 7.77 (d, $J = 8.7$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 22.6, 27.7, 28.6, 29.0, 31.8, 55.3, 55.5, 98.1, 100.0, 103.8, 117.1, 118.6, 130.4, 133.9, 158.0, 160.2; MS (EI) m/z 298 (M^+ , ^{37}Cl), 296 (M^+ , ^{35}Cl), 261 (M-Cl), 211, 175, 151, 138, 121, 115; HRMS (DCI) m/z calcd. for $\text{C}_{17}\text{H}_{25}\text{ClO}_2$ [M] $^+$ 296.1543, found 296.1554.

(Z)-2-Chloro-1-(4-fluorophenyl)non-1-ene (22):

Prepared according to typical procedure from hept-1-ene (170 μ L, 1.28 mmol) and 1,1-dichloro-1-alkene **4** (190 mg, 1.00 mmol). Purification by column chromatography (heptane) afforded **22** as a slightly yellow oil (200 mg, 74%, method A; 235 mg, 87%, method B).

IR (film) 2925, 2855, 1602, 1506, 1229 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.87-0.91 (m, 3H), 1.26-1.36 (m, 8H), 1.60-1.69 (m, 2H), 2.47 (t, $J = 7.2$ Hz, 2H), 6.42 (s, 1H), 6.99-7.06 (m, 2H), 7.53-7.60 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 22.6, 27.6, 28.6, 29.0, 31.8, 41.1, 115.0 (d, $J = 21.3$ Hz), 123.0, 130.6 (d, $J = 8.2$ Hz), 131.3, 135.0, 161.8 (d, $J = 245.4$ Hz); MS (EI) m/z 256 (M^+ , ^{37}Cl), 254 (M^+ , ^{35}Cl), 147, 135, 133, 109; HRMS (DCI) m/z calcd. for $\text{C}_{15}\text{H}_{20}\text{ClF}$ [M] $^+$ 254.1237, found 254.1228.

(Z)-7-Chloro-tetradec-6-ene (27):

Prepared according to typical procedure from hept-1-ene (170 μ L, 1.28 mmol) and 1,1-dichloro-1-alkene **6** (167 mg, 1.00 mmol). Purification by column chromatography (heptane) afforded **27** as a slightly yellow oil (170 mg, 74%, method A; 149 mg, 65%, method B).

IR (film) 2955, 2923, 2855, 2359, 2340, 1660, 1465 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.88 (t, $J = 6.6$ Hz, 3H), 0.89 (t, $J = 6.9$ Hz, 3H), 1.24-1.40 (m, 14H), 1.48-1.58 (m, 2H), 2.15 (q, $J = 7.2$ Hz, 2H), 2.29 (dt, $J = 0.6, 7.2$ Hz, 2H), 5.43 (t, $J = 6.9$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.0, 14.1, 22.5, 22.6, 27.4, 28.4, 28.5, 29.0, 31.4, 31.8, 39.5, 125.4, 134.7; MS (EI) m/z 189, 172, 153, 135, 107; HRMS (DCI) m/z calcd. for $\text{C}_{14}\text{H}_{27}\text{Cl}$ $[\text{M}]^+$ 230.1801, found 230.1801.

(Z)-4-Chloro-1-benzyloxydec-4-ene (26):

Prepared according to typical procedure from allylbenzylether (180 μL , 1.17 mmol) and 1,1-dichloro-1-alkene **6** (167 mg, 1.00 mmol). Purification by column chromatography (heptane/ CH_2Cl_2 , 85/15) afforded **26** as a slightly yellow oil (170 mg, 65%, method A; 120 mg, 46%, method B).

IR (film) 2924, 2854, 1453, 1102 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.89 (t, $J = 6.6$ Hz, 3H), 1.21-1.40 (m, 6H), 1.81-1.90 (m, 2H), 2.10-2.17 (m, 2H), 2.42 (t, $J = 7.2$ Hz, 2H), 3.47 (t, $J = 6.0$ Hz, 2H), 4.49 (s, 2H), 5.44 (t, $J = 6.9$ Hz, 1H), 7.26-7.36 (m, 5H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.0, 22.5, 27.5, 28.3, 28.4, 31.4, 36.1, 68.2, 72.9, 126.2, 127.5, 127.6, 128.3, 133.7, 138.5; MS (EI) m/z 280 (M^+ , ^{35}Cl), 268, 254; HRMS (DCI, NH_3) m/z calcd. for $\text{C}_{17}\text{H}_{29}\text{ClNO}$ $[\text{M}+\text{NH}_4]^+$ 298.1938, found 298.1935.

***Cis* (Z)-Methyl-3-(2-chloronon-1-enyl)-2,2-dimethyl-(1-cyclopropane)carboxylate (25-cis) and *trans* (Z)-Methyl-3-(2-chloronon-1-enyl)-2,2-dimethyl-(1-cyclopropane)carboxylate (25-trans):**

Prepared according to typical procedure from hept-1-ene (170 μL , 1.28 mmol) and 1,1-dichloro-1-alkene **7** (223 mg, 1.00 mmol). Reaction time: 144 h. Purification by column chromatography (heptane/ CH_2Cl_2 , 7/3) afforded **25-cis** and **25-trans** (179 mg, 63%, method A; 199 mg, 70%, method B) as colorless oils.

Data for 25-cis:

IR (film) 2925, 2856, 1727, 1435, 1192, 1142 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.88 (t, $J = 6.6$ Hz, 3H), 1.24 (s, 6H), 1.26-1.30 (m, 8H), 1.56 (m, 2H), 1.75 (d, $J = 8.7$ Hz, 1H), 2.16 (t, $J = 8.7$ Hz, 1H), 2.35 (t, $J = 7.2$ Hz, 2H), 3.65 (s, 3H), 5.82 (d, $J = 8.7$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 14.9, 22.6, 27.0, 27.5, 28.5, 29.0, 31.2, 31.8, 32.2, 39.7, 51.2, 119.6, 136.9, 171.5; MS (EI) m/z 288 (M^+ , ^{37}Cl), 286 (M^+ , ^{35}Cl), 251 (M-Cl), 227, 129, 107; HRMS (DCI) m/z calcd. for $\text{C}_{16}\text{H}_{27}\text{ClO}_2$ $[\text{M}]^+$ 286.1700, found 286.1709.

Data for 25-trans:

IR (film) 2925, 2856, 1728, 1438, 1225, 1164, 1114 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.88 (t, $J = 6.6$ Hz, 3H), 0.80 (s, 3H), 1.15-1.30 (m, 11H), 1.50 (d, $J = 5.4$ Hz, 1H), 1.53-1.58 (m, 2H), 2.28-2.34 (m, 3H), 3.69 (s, 3H), 5.19 (d, $J = 7.8$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 20.2, 22.5, 22.7, 27.4, 28.5, 28.7, 29.0, 31.8, 32.5, 34.6, 39.5, 122.2, 137.8, 172.3; MS (EI) m/z 288 (M^+ , ^{37}Cl), 286 (M^+ , ^{35}Cl), 251, 227, 129, 107; HRMS (DCI) m/z calcd. for $\text{C}_{16}\text{H}_{28}\text{ClO}_2$ $[\text{M}+\text{H}]^+$ 287.1778, found 287.1778.

cis (Z)-Methyl-3-(2-chloro-5-phenylpent-1-enyl)-2,2-dimethyl-(1-cyclopropane)carboxylate (**23-cis**) and *trans* (Z)-Methyl-3-(2-chloro-5-phenylpent-1-enyl)-2,2-dimethyl-(1-cyclopropane)carboxylate (**23-trans**):

Prepared according to typical procedure from allylbenzene (165 μ L, 1.39 mmol) and 1,1-dichloro-1-alkene **7** (223 mg, 1.00 mmol). Purification by column chromatography (heptane/ CH_2Cl_2 , 7/3) afforded **23-cis** and **23-trans** (176 mg, 58%, method A; 240 mg, 79%, method B) as colorless oils.

Data for **23-cis**:

IR (film) 2946, 1725, 1437, 1168, 1113 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 1.16 (s, 3H), 1.29 (s, 3H), 1.50 (d, $J = 3$ Hz, 1H), 1.87-1.90 (m, 2H), 2.32-2.37 (m, 3H), 2.61 (t, $J = 7.2$ Hz, 2H), 3.69 (s, 3H), 5.20 (d, $J = 4.5$ Hz, 1H), 7.16-7.20 (m, 3H), 7.26-7.29 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 20.2, 22.5, 28.6, 28.9, 32.5, 34.6, 38.8, 51.6, 122.8, 125.9, 128.3, 128.4, 137.1, 141.8, 172.2; MS (EI) m/z 306 (M^+ , ^{35}Cl), 271 (M-Cl), 247, 211, 155, 105; HRMS (DCI) m/z calcd. for $\text{C}_{18}\text{H}_{23}\text{ClO}_2$ [M] $^+$ 306.1386, found 306.1384.

Data for **23-trans**:

IR (film) 2950, 1724, 1436, 1191, 1138 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 1.24 (d, $J = 1.5$ Hz, 3H), 1.25 (d, $J = 1.5$ Hz, 3H), 1.77 (dd, $J = 1.5, 8.4$ Hz, 1H), 1.83-1.94 (m, 2H), 2.17 (dt, $J = 1.2, 8.6$ Hz, 1H), 2.39 (t, $J = 7.2$ Hz, 2H), 2.60 (t, $J = 7.8$ Hz, 2H), 3.64 (s, 3H), 5.86 (d, $J = 8.6$ Hz, 1H), 7.15-7.20 (m, 3H), 7.25-7.30 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.9, 27.0, 28.9, 29.1, 31.2, 32.2, 34.6, 39.1, 51.2, 120.3, 125.8, 128.3, 128.5, 136.3, 141.9, 171.4; MS (EI) m/z 308 (M^+ , ^{37}Cl), 306 (M^+ , ^{35}Cl), 271 (M-Cl), 247, 211, 155, 129, 117; HRMS (DCI) m/z calcd. for $\text{C}_{18}\text{H}_{23}\text{ClO}_2$ [M] $^+$ 306.1386, found 306.1389.

cis and *trans* (Z)-Methyl-3-(2-chloro-5-benzyloxypent-1-enyl)-2,2-dimethyl-(1-cyclopropane)carboxylate (**24**):

Prepared according to typical procedure from allylbenzylether (180 μ L, 1.17 mmol) and 1,1-dichloro-1-alkene **7** (223 mg, 1.00 mmol). Reaction time: 161 h. Purification by column chromatography (heptane/ CH_2Cl_2 , 1/2) afforded **24** as a colorless oil (195 mg, 59%, method A; 215 mg, 65%, method B as a mixture of stereoisomers at cyclopropyl ring).

IR (film) 2949, 2924, 2856, 1725, 1437, 1166 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 1.12 (s, 2H), 1.23 (d, $J = 1.5$ Hz, 1H), 1.26 (s, 2H), 1.48 (d, $J = 5.4$ Hz, 0.7H), 1.59 (s, 1H), 1.76 (d, $J = 8.4$ Hz, 0.3H), 1.81-1.92 (m, 2H), 2.12-2.18 (m, 0.3H), 2.28-2.33 (m, 0.7H), 2.40-2.50 (m, 2H), 3.45-3.50 (m, 2H), 3.64 (s, 1H), 3.69 (s, 2H), 4.49 (s, 1.3H), 4.50 (s, 0.7H), 5.20 (d, $J = 8.1$ Hz, 0.7H), 5.87 (d, $J = 8.4$ Hz, 0.3H), 7.30-7.35 (m, 5H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 20.2, 22.5, 27.6, 27.7, 28.5, 28.6, 32.1, 32.5, 34.5, 36.1, 36.4, 51.2, 51.6, 68.7, 69.0, 120.2, 122.9, 127.5, 127.6, 127.65, 128.4, 136.9, 138.4, 172.2; MS (EI) m/z 337 (M^+ , ^{35}Cl), 319, 301, 269, 241, 211, 179; HRMS (DCI) m/z calcd. for $\text{C}_{19}\text{H}_{25}\text{ClO}_3$ [$\text{M}+\text{H}$] $^+$ 337.1571, found 337.1576.

Typical procedure for non-selective cross-coupling of 1,1-dichloro alkenes: A terminal alkene (1.2 mmol) was added to a solution of 9-BBN (0.5 M in THF, 2.4 mL, 1.2 mmol). The solution was stirred at rt for 1 h. To a solution of 1,1-dichloro-1-alkene (1.0 mmol) in THF (2

mL) was added Pd₂Cl₂(dppp) (16.6 mg, 2.5 mol%), KF (180 mg, 3.0 mmol) and K₃PO₄ (635 mg, 3.0 mmol). The solution was refluxed until tlc shown completion. After cooling to rt, the reaction mixture was quenched with water and extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄ and concentrated *in vacuum*. The residue was purified by column chromatography, affording both mono and bis coupled products.

(Z)-2-Chloro-1-(2,4-dimethoxyphenyl)-5-phenylpent-1-ene (20) and 1-(2,4-dimethoxyphenyl)-5-phenyl-2-(3-phenylpropyl)pent-1-ene (28):

Prepared according to typical procedure from allylbenzene (165 μ L, 1.39 mmol) and 1,1-dichloro-1-alkene **5** (233 mg, 1.00 mmol). Reaction time: 24 h. Purification by column chromatography (heptane/CH₂Cl₂, 7/3) afforded both the mono coupled product **20** (75 mg, 24%) and the bis coupled product **28** (73 mg, 18%) as colorless oils.

Data for **20**:

IR (film) 2944, 1607, 1577, 1499, 1453, 1288, 1262, 1157, 1128, 1118, 1032 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ _H 1.94-2.04 (m, 2H), 2.52 (t, *J* = 6.9 Hz, 2H), 2.68 (t, *J* = 7.5 Hz, 2H), 3.80 (s, 3H), 3.82 (s, 3H), 6.44 (d, *J* = 2.4 Hz, 1H), 6.50 (dd, *J* = 2.4, 8.4 Hz, 1H), 6.60 (s, 1H), 7.16-7.32 (m, 5H), 7.78 (d, *J* = 8.4 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ _C 29.2, 34.6, 40.4, 55.4, 55.5, 98.2, 103.9, 117.0, 119.4, 125.8, 128.25, 128.3, 128.6, 130.5, 133.2, 158.0, 160.3; MS (EI) *m/z* 318 (M⁺, ³⁷Cl), 316 (M⁺, ³⁵Cl), 211, 177, 151, 115, 103, 91; HRMS (DCI) *m/z* calcd. for C₁₉H₂₂ClO₂ [M+H]⁺ 317.1308, found 317.1313.

Data for **28**:

IR (film) 2930, 1606, 1577, 1497, 1452, 1284, 1259, 1206, 1156, 1120, 1034 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ _H 1.68-1.87 (m, 4H), 2.19-2.25 (m, 4H), 2.55 (t, *J* = 7.8 Hz, 2H), 2.65 (t, *J* = 7.8 Hz, 2H), 3.78 (s, 3H), 3.81 (s, 3H), 6.24 (s, 1H), 6.38 (dd, *J* = 1.8, 8.1 Hz, 1H), 6.44 (m, 1H), 6.96 (d, *J* = 8.1 Hz, 1H), 7.10-7.31 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ _C 29.8, 29.9, 30.4, 35.7, 35.9, 36.3, 55.3, 55.4, 98.3, 103.7, 120.1, 120.7, 125.6, 128.2, 128.3, 128.4, 128.5, 130.2, 141.3, 142.4, 142.7, 158.0, 159.4; MS (EI) *m/z* 400 (M⁺), 295, 192, 177, 151, 91; HRMS (DCI) *m/z* calcd. for C₂₈H₃₂O₂ [M]⁺ 400.2402, found 400.2398.

2-(3-Phenylpropyl)-1-(4-methoxyphenyl)-5-phenylpent-1-ene (12):

Prepared according to typical procedure from allylbenzene (135 μ L, 1.13 mmol) and alkenyl chloride **2** (203 mg, 1.00 mmol). Reaction time: 22 h. Purification by column chromatography (heptane/CH₂Cl₂, 85/15) afforded **12** as a colorless oil (170 mg, 58%).

IR (film) 2928, 2360, 1605, 1508, 1452, 1246, 1175, 1032 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ _H 1.78 (m, 4H), 2.23 (m, 4H), 2.61 (m, 4H), 3.80 (s, 3H), 6.22 (s, 1H), 6.79 (dd, *J* = 2.1, 6.3 Hz, 2H), 7.03-7.06 (m, 2H), 7.12-7.30 (m, 10H); ¹³C NMR (75 MHz, CDCl₃) δ _C 29.9, 30.1, 35.6, 35.9, 36.8, 55.2, 113.5, 125.0, 125.7, 128.2, 128.4, 129.6, 130.9, 141.3, 142.2, 142.5,

157.7; MS (EI) m/z 370 (M^+), 249, 173, 162, 147, 121; HRMS (DCI) m/z calcd. for $C_{27}H_{30}O$ [M] $^+$ 370.2297, found 370.2300.

Typical procedure for second coupling, leading to tri-substituted olefins: A terminal alkene (1.2 mmol) was added to a solution of 9-BBN (0.5 M in THF, 2.4 mL, 1.2 mmol). The solution was stirred at rt for 1 h. Chloro-olefin (1.0 mmol) was added dissolved in THF (2 mL), and $PdCl_2(dppp)$ (5 mol%) or $Pd_2(dba)_3$ (5 mol%) with 2-(dicyclohexylphosphino)biphenyl (10 mol%), KF (180 mg, 3.0 mmol) and K_3PO_4 (636 mg, 3.0 mmol) were successively added. The reaction mixture was refluxed until completion. After cooling to rt, the solution was quenched with water and extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuum*. The residue was purified by column chromatography, yielding the stereoselectively tri-substituted olefin.

(E)-2-Heptyl-1-(4-methoxycarbonylphenyl)-5-phenylpent-1-ene (32):

Prepared according to typical procedure from hept-1-ene (170 μ L, 1.28 mmol) and alkenyl chloride **18** (202 mg, 0.59 mmol). Reaction time: 26 h. Purification by column chromatography (heptane/ CH_2Cl_2 , 7/3) afforded **32** as a colorless oil (160 mg, 69%).

IR (film) 2923, 2853, 1720, 1604, 1433, 1273, 1176, 1101 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ_H 0.85-0.89 (m, 3H), 1.23-1.28 (m, 8H), 1.43-1.46 (m, 2H), 1.80-1.90 (m, 2H), 2.19-2.25 (m, 4H), 2.67 (t, $J = 7.5$ Hz, 2H), 3.90 (s, 3H), 6.28 (s, 1H), 7.14-7.32 (m, 7H), 7.95-7.99 (m, 2H); ^{13}C NMR (75 MHz, $CDCl_3$) δ_C 14.0, 22.6, 28.2, 29.0, 29.7, 29.8, 30.8, 31.8, 35.6, 37.0, 51.9, 124.4, 125.7, 127.4, 128.3, 128.4, 128.5, 129.4, 142.3, 143.4, 145.8, 167.1; MS (EI) m/z 378 (M^+), 190, 175, 131, 129, 104, 91; HRMS (EI) m/z calcd. for $C_{26}H_{34}O_2$ [M] $^+$ 378.2558, found 378.2545.

(E)-2-(3-Phenylpropyl)-5-benzyloxy-1-(4-methoxyphenyl)pent-1-ene (31):

Prepared according to typical procedure from allylbenzene (165 μ L, 1.39 mmol) and alkenyl chloride **16** (172 mg, 0.54 mmol). Reaction time: 46 h. Purification by column chromatography (heptane/ CH_2Cl_2 , 1/1) afforded **31** as a colorless oil (136 mg, 69%).

IR (film) 2929, 2850, 1605, 1507, 1452, 1245, 1174, 1100, 1029 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ_H 1.74-1.85 (m, 4H), 2.21-2.29 (m, 4H), 2.59 (d, $J = 7.5$ Hz, 2H), 3.49-3.54 (m, 2H), 3.80 (s, 3H), 4.51 (s, 2H), 6.21 (s, 1H), 6.76-6.80 (m, 2H), 7.13-7.35 (m, 12H); ^{13}C NMR (75 MHz, $CDCl_3$) δ_C 28.2, 29.9, 30.2, 33.7, 35.9, 55.2, 69.9, 72.9, 113.5, 125.0, 125.7, 127.5, 127.6, 128.2, 128.3, 128.5, 129.6, 130.9, 138.6, 141.0, 142.2, 157.7; MS (EI) m/z 400 (M^+), 294, 282, 176, 147, 121, 91; HRMS (DCI) m/z calcd. for $C_{28}H_{32}O_2$ [M] $^+$ 400.2402, found 400.2397.

(Z)-2-(3-Phenylpropyl)-1-(4-fluorophenyl)non-1-ene (33-Z):

Prepared according to typical procedure from allylbenzene (135 μ L, 1.13 mmol) and alkenyl chloride **22** (192 mg, 0.75 mmol). Reaction time: 95 h. Purification by column chromatography (heptane) afforded **33-Z** as a colorless oil (190 mg, 74%).

IR (film) 2924, 2854, 1505, 1224 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.89 (t, $J = 6.6$ Hz, 3H), 1.24-1.31 (m, 8H), 1.41-1.51 (m, 2H), 1.71-1.82 (m, 2H), 2.11-2.24 (m, 4H), 2.58 (t, $J = 7.5$ Hz, 2H), 6.20 (s, 1H), 6.87-6.95 (m, 2H), 7.01-7.07 (m, 2H), 7.12-7.29 (m, 5H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 22.7, 28.2, 29.2, 29.4, 29.9, 30.0, 31.9, 35.9, 37.2, 114.9 (d, $J = 21.2$ Hz), 124.0, 125.8, 128.3 (d, $J = 13.0$ Hz), 130.0, 130.1, 134.5 (d, $J = 3.3$ Hz), 142.1, 143.3, 161.0 (d, $J = 244.4$ Hz); MS (EI) m/z 338 (M^+), 234, 150, 135, 103; Anal. calcd. for $\text{C}_{24}\text{H}_{31}\text{F}$: C 85.16; H 9.23; found C 85.08; H 9.45.

(E)-2-(3-phenylpropyl)-1-(4-fluorophenyl)non-1-ene (33-E):

Prepared according to typical procedure from hept-1-ene (60 μL , 0.44 mmol) and alkenyl chloride **19** (87 mg, 0.32 mmol). Reaction time: 76 h. Purification by column chromatography (heptane) afforded **33-E** as a colorless oil (76 mg, 74%).

IR (film) 2924, 2854, 1505, 1224 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.85-0.89 (m, 3H), 1.24-1.26 (m, 8H), 1.36-1.46 (m, 2H), 1.78-1.88 (m, 2H), 2.14-2.22 (m, 4H), 2.66 (d, $J = 7.8$ Hz, 2H), 6.21 (s, 1H), 6.94-7.02 (m, 2H), 7.10-7.32 (m, 7H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 22.6, 28.2, 29.1, 29.7, 29.9, 30.5, 31.8, 35.6, 36.7, 114.8 (d, $J = 21.3$ Hz), 124.0, 125.8, 128.4 (d, $J = 9.8$ Hz), 130.0, 130.1, 134.5 (d, $J = 3.2$ Hz), 142.5, 143.4; MS (EI) m/z 338 (M^+), 150, 135, 109, 104; HRMS (DCI) m/z calcd. for $\text{C}_{24}\text{H}_{31}\text{F}$ [M] $^+$ 338.2410, found 338.2424.

Typical procedure for Suzuki cross-coupling with phenylboronic acid 34: Alkenyl chloride (1.0 mmol.), phenylboronic acid **34** (1.2 equiv.), $\text{Pd}_2(\text{dba})_3$ (5 mol%), 2-(dicyclohexylphosphinyl)biphenyl (10 mol%) KF (176 mg, 3.0 mmol) and K_3PO_4 (636 mg, 3.0 mmol) were refluxed in THF for several hours. After cooling to rt, the reaction mixture was quenched with water and extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 and concentrated *in vacuo*. The residue was purified by column chromatography.

(Z)-2-Phenyl-1-(4-methoxyphenyl)non-1-ene (35):

Prepared according to typical procedure from alkenyl chloride **17** (204 mg, 0.76 mmol) and phenylboronic acid (121 mg, 1.00 mmol). Reaction time: 89 h. Purification by column chromatography (heptane/ CH_2Cl_2 , 85/15) afforded **35** as a colorless oil (134 mg, 58%).

IR (film) 2953, 2924, 2853, 1606, 1508, 1247, 1175, 1035 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ_{H} 0.84-0.89 (m, 3H), 1.24-1.41 (m, 10H), 2.45 (dt, $J = 1.2, 7.5$ Hz, 2H), 3.71 (s, 3H), 6.36 (s, 1H), 6.59-6.64 (m, 2H), 6.80-6.85 (m, 2H), 7.13-7.17 (m, 2H), 7.23-7.33 (m, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ_{C} 14.1, 22.6, 28.0, 29.1, 29.2, 31.8, 40.7, 55.1, 113.2, 125.4, 126.6, 128.5, 128.6, 130.0, 130.2, 141.6, 141.7, 157.8; MS (EI) m/z 308 (M^+), 224, 178, 165, 145, 121, 114; Anal. calcd. for $\text{C}_{22}\text{H}_{28}\text{O}$: C 85.66; H 9.15; found C 85.74; H 9.39.

(Z)-2,5-diphenyl-1-(4-methoxyphenyl)pent-1-ene (36):

Prepared according to typical procedure from allkenyl chloride **11** (287 mg, 1.00 mmol) and phenylboronic acid (145 mg, 1.19 mmol). Reaction time: 97 h. Purification by column chromatography (heptane/CH₂Cl₂, 85/15) afforded **36** as a colorless oil (220 mg, 66%) and some starting material **11** was recovered (75 mg, 26%).

IR (film) 2927, 2854, 1605, 1508, 1440, 1247, 1175, 1033 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ_H 1.69-1.75 (m, 2H), 2.52 (d, *J* = 4.5 Hz, 2H), 2.64 (d, *J* = 4.5 Hz, 2H), 3.71 (s, 3H), 6.37 (s, 1H), 6.61-6.63 (m, 2H), 6.83-6.85 (m, 2H), 7.14-7.17 (m, 4H), 7.24-7.31 (m, 6H); ¹³C NMR (75 MHz, CDCl₃) δ_C 29.6, 35.4, 40.2, 55.1, 113.2, 125.7, 125.9, 126.8, 128.2, 128.4, 128.5, 128.6, 130.0, 140.9, 141.4, 142.4, 157.9; MS (EI) *m/z* 328 (M⁺), 224, 121, 115; HRMS (DCI) *m/z* calcd. for C₂₄H₂₄O [M]⁺ 328.1828, found 328.1833.