Supporting Information for:

Syntheses of Stereochemically Diverse Nine-Membered Ring-Containing Biaryls

Shyam Krishnan and Stuart L. Schreiber*

*Howard Hughes Medical Institute, Department of Chemistry and Chemical Biology,
Harvard University, and the Broad Institute of Harvard and MIT, Cambridge,
Massachusetts 02138*

Table of Contents

1. Experimental techniques and apparatus.....	S1
2. General procedure for compound cleavage from solid support.....	S4
3. Synthesis of building blocks.....	S4
4. Loading of 2-Bromo-5-hydroxybenzaldehyde.....	S6
5. Reductive amination with primary amines.....	S7
6. Reductive amination with secondary amines.....	S12
7. Reductive alkylation of secondary amines.....	S17
8. Mitsunobu Reaction.....	S23
9. Biaryl Coupling.....	S31
10. Synthesis of Dimer (15).....	S39
11. Spectra.....	S45

1. Experimental Techniques and Apparatus

Except as otherwise indicated, reactions were carried out under argon with dry, freshly distilled solvents. Anhydrous 2-methyltetrahydrofuran (Aldrich, racemic) was stored over activated molecular sieves. Dry solvents were dispensed from a delivery system that passes the solvents through packed columns (tetrahydrofuran, diethyl ether,

acetonitrile, and methylene chloride: dry neutral alumina; hexane, benzene, and toluene: dry neutral alumina and Q5 reactant; dimethylformamide: activated molecular sieves). *n*-BuLi in hexane (Aldrich), *t*-BuLi in pentane (Aldrich) and *i*-PrMgCl in THF (Aldrich) were titrated with Salicylaldehyde Phenylhydrazone¹ or 4-biphenylmethanol.² (±)-Calycotomine was purchased from Acros Organics. All other amino alcohols were purchased from Aldrich. 4-pentenal was purchased from Lancaster. 2-Bromo-5-hydroxybenzaldehyde was synthesized using a previously reported procedure³. 5-(*t*-Butyldimethylsilanoxy)-pentanal was synthesized as described below. All other aliphatic aldehydes and bromophenols were purchased from Aldrich. All other reagents were purified in accordance with the instructions in “*Purification of Laboratory Chemicals*”⁴ or used as obtained from commercial sources.

Yields of solution phase reactions refer to chromatographically and spectroscopically pure compounds. Yields of solid (gel) phase reactions were determined by ¹H NMR, using 2,5-dimethylfuran⁵ as an internal standard with a relaxation delay (*d*₁) of 10s. All solution-phase reactions were monitored by thin layer chromatography (TLC) using glass plates precoated with Merck silica gel 60 F₂₅₄ or aluminum oxide 60 F₂₅₄. Visualization was by the quenching of UV fluorescence (*I*_{max} = 254 nm) or by staining with ceric ammonium molybdate or potassium permanganate or Dragendorff’s reagent (0.08% w/v bismuth subnitrate and 2% w/v KI in 3M aq. AcOH). Retention factors (*R*_f) are quoted to

(1) Love, B.E.; Jones, E.G.; *J. Org. Chem.* **1999**, *64*, 3755-3756.

(2) Vogel’s Textbook of Practical Organic Chemistry, Fifth Edition, Longman Scientific & Technical, 1989.

(3) Léo, P.-M.; Morin, C.; Philouze, C. *Org. Lett.* **2002**, *4*, 2711-2714.

(4) Armarego, W. L. F.; Perrin, D. D. *Purification of Laboratory Chemicals, Fourth Edition*; Butterworth-Heinemann: Oxford, 1997.

(5) Gerritz, S.W.; Seffler, A.M. *J. Comb. Chem.* **2000**, *2*, 39-41.

0.01. Melting points were obtained using a Mel-Temp II melting point apparatus and are uncorrected. Infrared spectra were recorded either as a thin film between NaCl plates or as a KBr disc on a Nicolet 5PC FT-IR spectrometer with internal referencing. Absorption maxima (ν_{max}) are reported in wavenumbers (cm^{-1}) and the following abbreviations are used: w, weak; m, medium; s, strong; br, broad. Proton magnetic resonance spectra were recorded on Varian Mercury400 (400MHz), Bruker Avance DMX500 (500MHz), and Varian Unity/Inova500 (500MHz) spectrometers. Proton assignments are supported by ^1H - ^1H COSY, HMQC and HMBC spectra where necessary. Stereochemical assignments are supported by NOESY (1D and 2D) and ROESY (2D) spectra. Chemical shifts (δ_{H}) are quoted in ppm and are referenced to tetramethylsilane (internal). Coupling constants (J) are reported in Hertz to the nearest 0.5Hz. Data are reported as follows: chemical shift, integration, multiplicity [br, broad; s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; or as a combination of these (*e.g.* dd, dt, *etc.*)], coupling constant(s) and assignment. Diastereotopic protons are assigned as X and X', where the ' indicates the lower field proton. Carbon magnetic resonance spectra were recorded on Varian Mercury400 (100MHz) and Varian Unity/Inova500 (125MHz) spectrometers. Carbon spectra assignments are supported where necessary by ^{13}C - ^1H (HMQC) correlations. Chemical shifts (δ_{C}) are quoted in ppm to the nearest 0.01 ppm, and are referenced to tetramethylsilane (internal). Low resolution mass spectra were obtained with JEOL AX-505H, SX-102A (CI/EI), Micromass Platform II and LCT (APCI/ES/LCMS) spectrometers. Only molecular ions, fragments from molecular ions and other major peaks are reported. High resolution mass spectra was obtained with a Micromass LCT (ES) spectrometer, and reported mass values are within the error limits of ± 5 ppm mass units.

2. General procedure for Cleavage of Compound from Resin:

5-40 mg of beads were transferred to a 1.7mL microcentrifuge eppendorf or 20mL polypropylene vial and freshly prepared HF.Py/Py/THF (90:5:5 by volume, 100 μ L per 10mg beads) was added. The beads were gently agitated for 1.5h and the excess HF quenched by addition of methoxytrimethylsilane (100 μ L per 10mg beads) and the beads further agitated for 1h. The beads were filtered and further washed with freshly distilled THF (x 2) and MeCN (x 2). The washings and filtrate were combined, filtered through a pad of SiO₂ and concentrated under reduced pressure to yield the library compound.

Nomenclature of Library Compounds:

Library compounds are named with the building blocks that comprise them. For example, **A1B4C5** refers to a product resulting from the combination of the starting 2-bromo-5-hydroxybenzaldehyde (**A1**) and building blocks **B4** and **C5** (Figure 2 in the text). The prefix 'c' is used for the cyclic biaryl-containing medium rings.

3. Synthesis of Building Blocks:

5-(*tert*-Butyldimethylsilyloxy)-pentanal (Building Block C5):

(Z)-5-penten-1-ol (6.17g, 48.12 mmol, 1.0 equiv.), Imidazole (9.83g, 144.4 mmol, 3 equiv.) and 4-DMAP (293.9mg, 2.40 mmol, 0.05 equiv.) were dissolved in 20mL anhydrous DMF in a 100mL round bottom flask. The solution was cooled in an ice bath

and a solution of *tert*-Butyldimethylsilylchloride (8.70g, 57.74 mmol, 1.2 equiv.) in 20mL anhydrous DMF was gradually added with constant stirring. The ice bath was then removed and the solution allowed to warm to room temperature and stirred for 6h. The reaction mixture was poured into 200mL Et₂O and washed with saturated aqueous NaHCO₃ (4 x 50mL) and brine (100mL). The aqueous washes were extracted with Et₂O (50mL). The Et₂O extracts were combined and dried over MgSO₄ and concentrated under reduced pressure. The residue obtained was chromatographed over SiO₂ (100:1 Hexanes/EtOAc) to yield (Z)- *tert*-Butyldimethyl-oct-5-enyloxysilane as a colorless oil (11.60g, 99%). This was dissolved in CH₂Cl₂/MeOH (10:1, 110mL) along with NaHCO₃ (406.0mg, 4.83 mmol, 0.1 equiv.) and the solution cooled to -78⁰C. Ozone was bubbled into the solution with constant stirring. After 3.5h, the solution turned blue indicated consumption of the olefin. Dimethylsulfide (35.4 mL, 0.483 mol, 10.0 equiv.) was then added and the solution allowed to gradually warm to room temperature, and then concentrated under reduced pressure. The residue obtained was taken up in 200mL Et₂O and washed with saturated aqueous NaHCO₃ (100mL) and brine (100mL), then dried over MgSO₄ and concentrated under reduced pressure and the residue obtained chromatographed over SiO₂ (Hexanes/EtOAc, 100:1, then 10:1) to yield 5-(*tert*-Butyldimethylsilanyloxy)-pentanal as a colorless liquid (9.40g, 91% over two steps).

R_f = 0.37 (SiO₂; Hexanes: EtOAc; 10:1).

IR (film)/cm⁻¹: 2954 (s), 2930(s), 2858(s), 1728(s), 1472(m), 1256(s), 1102(s), 836(s).

δ_H (400MHz, CDCl₃): 0.04 (s, 6H, SiCH₃), 0.89 (s, 9H, C(CH₃)₃), 1.54 (m, 2H, TBSOCH₂CH₂-), 1.69 (m, 2H, -CH₂CH₂CHO), 2.44 (app. dt, 2H, J7.5Hz, 1.6Hz, CH₂CHO), 3.62 (app. t, J6Hz, 2H, TBSOCH₂-), 9.75 (app. t, 1H, J1.5Hz, -CHO).

δ_c (100MHz, CDCl_3): 18.3, 18.6, 25.9, 32.1, 43.6, 62.6, 202.6.

m/z (ESI): 217 (MH^+ ; 100%).

HRMS found 217.1624. $\text{C}_{11}\text{H}_{25}\text{O}_2\text{Si}^+$ requires 217.1627.

4. Loading of 2-Bromo-5-hydroxybenzaldehyde:

500-600 μm polystyrene (1% divinylbenzene) silyl linker functionalized macrobeads (500mg, theoretical loading 1.4mmol/g, 0.7mmol) in a polypropylene tube capped with a septum were swollen in 15mL $\text{TMSCl}/\text{CH}_2\text{Cl}_2$ (2.5% v/v) for 20 min. The solution was then drained under a positive pressure of argon and the beads washed with anhydrous CH_2Cl_2 (3 x 5mL x 5 min). A solution of TfOH in CH_2Cl_2 (3% v/v, 12.6mL, 4.26 mmol, 6.0 equiv.) was then added and the beads allowed to stand for 30 min under argon with occasional agitation whereupon the beads turned dark-orange. The solution was then drained under a positive pressure of argon and the beads washed with anhydrous CH_2Cl_2 (3 x 5mL x 5 min) to remove excess TfOH . Freshly distilled 2,6-Lutidine (0.66mL, 5.68 mmol, 8.0 equiv.) was then added whereupon the beads turned colorless. A solution of 2-Bromo-5-Hydroxybenzaldehyde in anhydrous $\text{CH}_2\text{Cl}_2/\text{DMF}$ (428.8mg, 2.133 mmol, 3.0 equiv. dissolved in 5mL $\text{CH}_2\text{Cl}_2/\text{DMF}$ (4:1)) was then added and the beads allowed to stand under Ar for 1h with occasional agitation. The septum of the tube was then quickly replaced with a polypropylene cap and the beads tumbled at ambient temperature for 15h. The solution was then drained and the beads washed with DMF (3 x

1h), 1:1 THF/DMF (1 x 12h, 2 x 2h), THF (3 x 1h) and CH₂Cl₂ (4 x 1h). The beads were then dried *in vacuo* (house-vacuum for 2h followed by high vacuum for 12h) to yield free-flowing pale reddish-brown beads (548mg) containing the immobilized aldehyde.

MAS-NMR δ_H (600MHz, CDCl₃): 10.27 (s, -CHO).

51.5 mg of beads were transferred to a 20mL polypropylene vial and 0.8mL of HF.Py/Py/THF (90:5:5 by volume) was added. The beads were gently agitated for 1.5h and the excess HF quenched by addition of methoxytrimethylsilane (0.8mL) and the beads further agitated for 1h. The beads were filtered and further washed with freshly distilled THF (x 2) and MeCN (x 2). The washings and filtrate were combined, filtered through a pad of SiO₂ and concentrated under reduced pressure to yield 12.6mg of the starting aromatic aldehyde (> 95% pure by ¹H NMR). The on bead loading was calculated to be 1.22 mmol/g.

5. Reductive Amination with Primary amines:

A glass vial, screwcapped with a teflon-lined septum containing the polymer supported aldehyde (500mg, 1.22 mmol/g, 0.61mmol) was charged with a solution of amino alcohol in appropriate solvent (5mL). The suspension was then agitated at ambient temperature for 16-24h. The solution was then drained and the beads washed with THF/DMF/TEA (50:50:1, 2 x 20 min) and THF/TEA (100:1, 2 x 20 min). The vial containing the beads was then purged with Argon; then the beads resuspended in 3mL

anhydrous THF. A solution of Sodium Cyanoborohydride (**CAUTION:** Highly Toxic!), (383 mg, 6.0 mmol, 10 equiv.) in 3mL methanol was then added and the suspension agitated gently. 300 μ L glacial Acetic Acid (5% v/v w.r.t to 1: 1 THF/MeOH, 5.3 mmol, 8.8 equiv.) was then added and the suspension gently agitated at ambient temperature for 16-20h. The solution was then drained and the beads washed with 50/50/1 THF/DMF/TEA (3 x 30 min), 100/1 THF/TEA (3 x 30 min), 1/1 THF/MeCN (2 x 20 min), MeCN (2 x 20 min) and CH₂Cl₂ (4 x 30 min). The beads were then dried *in vacuo* (House vacuum for 2h followed by High vacuum for 12h).

Fig.1. Amino alcohol building blocks tested in reductive aminations. **B1-B13** were selected for library synthesis. **B14-B21** produced the corresponding reductive amination products in low purity and were not selected for library synthesis.

(R)-4-Bromo-3-[2-hydroxy-1-phenylethylamino)-methyl]-phenol, A1B1.

1:1 THF/Trimethyl Orthoformate was used as the solvent for Schiff Base formation.

Colorless oil. Yield: 71%; $R_f = 0.24$ (SiO_2 ; Chloroform: Isopropanol; 20:1).

IR (film)/ cm^{-1} : 3284 sb, 2925s, 1576s, 1453s, 1294s, 1025s, 701s.

δ_H (500MHz, CDCl_3): 3.61-3.65 (m, 2H), 3.73-3.78 (m, 2H), 3.86 (dd, $J_{8.5\text{Hz}}$, 4Hz, 1H), 6.61 (dd, $J_{8.5\text{Hz}}$, 3Hz, 1H), 6.86 (d, $J_{3\text{Hz}}$, 1H), 7.29-7.39 (m, 6H, Ar-H).

δ_c (100MHz, CDCl_3): 51.2, 64.0, 66.7, 114.1, 116.3, 117.6, 127.5, 128.0, 128.8, 133.7, 139.6, 139.7, 155.4.

m/z (ESI): 322 (MH^+ ; 28%), 324 (MH^+ , ^{81}Br ; 28%).

HRMS found 322.0448. $\text{C}_{15}\text{H}_{17}\text{BrNO}_2^+$ (MH^+ ; ^{79}Br) requires 322.0442.

(S,S)-4-Bromo-3-[(2S,3S)-2-methoxy-3-phenyl-2-hydroxypropylamino)-methyl]-phenol, A1B3. 1:1 Benzene/Trimethylorthoformate used as solvent.

Colorless oil. Yield: 57%; $R_f = 0.41$ (SiO_2 ; Benzene : Isopropanol; 5:1).

δ_H (400MHz, C_6D_6): 2.75 (app. quintet, $J_{4\text{Hz}}$, 1H, NCHCH_2OMe), 2.88 (s, 3H, OCH_3), 2.91 (dd, $J_{9.7\text{Hz}}$, 4.5Hz, 1H, $\text{CHH}'\text{OMe}$), 3.07 (dd, $J_{9.5\text{Hz}}$, 3.5Hz, 1H, $\text{CHH}'\text{OMe}$), 3.70 (d, $J_{14\text{Hz}}$, 1H, $\text{NCHH}'\text{Ar}$), 3.83 (d, $J_{14\text{Hz}}$, 1H, $\text{NCHH}'\text{Ar}$), 4.71 (d, $J_{8\text{Hz}}$, 1H, OCHPh),

6.58 (dd, $J_{8.5\text{Hz}}$, 3Hz, 1H, Ar-H), 7.04 (d, $J_{3\text{Hz}}$, 1H, Ar-H), 7.08 (m, 1H, Ar-H), 7.14-7.20 (m, 3H, Ar-H), 7.35 (d, $J_{7.5\text{Hz}}$, 1H, Ar-H).

δ_c (100MHz, C_6D_6): 51.4, 58.6, 63.9, 69.6, 73.2, 113.2, 117.1, 117.8, 127.3, 128.5, 128.6, 133.8, 139.0, 142.1, 157.2.

m/z (ESI): 366 (MH^+ ; 35%), 368 (MH^+ , ^{81}Br ; 30%).

HRMS found 366.0713. $\text{C}_{17}\text{H}_{21}\text{BrNO}_3^+$ (MH^+ ; ^{79}Br) requires 366.0705.

4-Bromo-3-[(2S-hydroxy-1R-methyl-2-phenyl-ethylamino)-methyl]-phenol,

A1B4, 4. Trimethylorthoformate was used as solvent.

White solid. Yield: 76%; R_f = 0.37 (SiO_2 , Benzene: Isopropanol; 5:1).

δ_H (400MHz, CD_3OD): 1.12 (d, $J_{6.5\text{Hz}}$, 3H, CHCH_3), 2.83 (m, 1H, NCHCH_3), 3.72 (ABq, $J_{AB}13.5\text{Hz}$, 2H, NCH_2Ar), 4.53 (d, $J_{6.5\text{Hz}}$, 1H, OCHPh), 6.61 (dd, $J_{9\text{Hz}}$, 3Hz, 1H, Ar-H), 6.80 (d, $J_{3\text{Hz}}$, 1H, Ar-H), 7.23–7.32 (m, 6H, Ar-H).

δ_c (100MHz, CD_3OD): 15.2, 51.8, 58.8, 77.1, 113.6, 117.3, 119.0, 127.8, 128.7, 129.5, 134.5, 140.1, 143.5, 158.4.

m/z (ESI): 336 (MH^+ , ^{79}Br ; 100%), 338 (MH^+ , ^{81}Br ; 99%).

HRMS found 336.0594. $\text{C}_{16}\text{H}_{19}\text{BrNO}_2$ (MH^+ , ^{79}Br) requires 336.0599.

***Trans*-4-Bromo-3-[(2-hydroxycyclohexylamino)-methyl]-phenol, A1B6.**

White solid. Yield: 69% ; $R_f = 0.51$ (SiO₂, Benzene: Isopropanol; 1:1).

δ_H (500MHz, CD₃OD): 1.25-1.33 (m, 4H, C(3)H, C(4)H, C(5)H, C(6)H), 1.74 (m, 2H, C(4)H', C(5)H'), 1.97 (m, 1H, C(6)H'), 2.17 (m, 1H, C(3)H'), 2.56 (m, 1H, C(2)H), 3.40 (m, 1H, C(1)H), 4.00 (ABq, J_{AB} 13Hz, 2H, C(7)H₂), 6.70 (dd, J 8.5Hz, 3Hz, 1H, C(9)H), 6.94 (d, J 3Hz, 1H, C(10)H), 7.38 (d, J 8.5Hz, C(8)H).

δ_c (125MHz, CD₃OD): 25.3, 25.4, 29.4, 35.3, 50.4, 63.6, 73.0, 113.8, 118.3, 119.4, 134.8, 137.4, 158.7.

m/z (ESI): 300 (MH⁺; 100%), 302 (MH⁺, ⁸¹Br; 95%).

HRMS found 300.0599. C₁₃H₁₉BrNO₂ (MH⁺, ⁷⁹Br) requires 300.0602.

6. Reductive Amination with Secondary Amines:

A glass vial, screwcapped with a teflon-lined septum containing the polymer supported aldehyde (75mg, 1.22 mmol/g, 92μmol) was purged with Argon and then charged with a solution of amino alcohol in appropriate solvent (2mL). The suspension

was then warmed at 50-60°C for 16-24h. The solution was then drained and the beads washed with THF/DMF/TEA (50:50:1, 2 x 20 min) and THF/TEA (100:1, 2 x 20 min). The vial containing the beads was then purged with Argon; then the beads resuspended in 1.5 mL anhydrous THF. A solution of Sodium Cyanoborohydride (**CAUTION: Highly Toxic!**), (1M in MeOH, 1.5mL, 1.5mmol, 16.5 equiv.) was then added and the suspension agitated gently. 150µL glacial Acetic Acid (5% v/v w.r.t to 1: 1 THF/MeOH, 2.65 mmol, 29.0 equiv.) was then added and the suspension gently agitated at ambient temperature for 16-20h. The solution was then drained and the beads washed with 50/50/1 THF/DMF/TEA (3 x 30 min), 100/1 THF/TEA (3 x 30 min), 1/1 THF/MeCN (2 x 20 min), MeCN (2 x 20 min) and CH₂Cl₂ (4 x 30 min). The beads were then dried *in vacuo* (House vacuum for 2h followed by High vacuum for 12h). Cleavage of compound from resin was followed by purification and spectroscopic characterization.

4-Bromo-3-[(2S-hydroxy-1R-methyl-2-phenylethyl)-methylamino]-methylphenol, A1B7. Toluene was used as solvent.

White solid. Yield: 73% ; R_f = 0.26 (SiO₂, Benzene: Isopropanol; 10:1).

δ_H (500MHz, CD₃OD): 1.12 (d, J = 6.5Hz, 3H, CHCH₃), 2.27 (s, 3H, NCH₃), 2.95 (m, 1H, NCHCH₃), 3.64 (ABq, J_{AB} = 14Hz, 2H, NCH₂Ar), 4.79 (d, J = 6Hz, 1H, OCHPh), 6.55 (dd, J = 8.5Hz, 3Hz, 1H, Ar-H), 6.72 (d, J = 3Hz, 1H, Ar-H), 7.19–7.29 (m, 6H, Ar-H).

δ_c (125MHz, CD₃OD): 9.5, 38.5, 59.0, 65.1, 76.2, 114.1, 116.7, 118.9, 127.5, 128.0, 129.0, 134.0, 140.6, 145.5, 158.1.

m/z (ESI): 350 (MH⁺; 99%), 352 (MH⁺, ⁸¹Br; 100%).

HRMS found 350.0764. C₁₇H₂₁BrNO₂ (⁷⁹Br) requires 350.0755.

***Rac*-4-Bromo-3-(1-hydroxymethyl-6,7-dimethoxy-3,4-dihydro-1H-isoquinolin-2-ylmethyl)-phenol, A1B9.** 1:1 Toluene/THF was used as solvent.

Colorless oil. Yield: 55%; R_f = 0.41 (SiO₂; Benzene : Isopropanol; 5:1).

δ_H (400MHz, C₆D₆): 2.10 (m, 1H, ArCHH'CH₂N), 2.69 (m, 2H, ArCHH'CH₂N, ArCH₂CHH'N), 2.98 (m, 1H, ArCH₂CHH'N), 3.39 (s, 3H, OCH₃). 3.42 (s, 3H, OCH₃). 3.57 (m, 1H), 3.66 (m, 2H), 3.76 (ABq, J_{AB}14Hz, 2H, ArCH₂N), 6.25 (s, 1H, Ar-H), 6.35 (s, 1H, Ar-H), 6.66 (dd, J_{8.5}Hz, 3Hz, 1H, Ar-H), 7.20 (d, J_{8.5}Hz, 1H, Ar-H), 7.30 (d, J = 8.5Hz, 1H, Ar-H).

δ_c (100MHz, C₆D₆): 23.1, 42.9, 55.6, 55.7, 57.1, 61.8, 64.0, 111.3, 112.6, 114.2, 117.1, 118.5, 125.4, 126.4, 134.2, 138.7, 148.7, 149.0, 157.2.

m/z (ESI): 408 (MH⁺; 28.5%), 410 (MH⁺, ⁸¹Br; 25.8%).

HRMS found 408.0802. C₁₉H₂₃BrNO₄ (MH⁺:⁷⁹Br) requires 408.0810.

(S)-4-Bromo-3-(3-hydroxymethyl-3,4-dihydro-1H-isoquinolin-2-ylmethyl)-

phenol, A1B10. 1:1 Toluene/THF was used as solvent.

White solid. Yield: 86%; $R_f = 0.38$ (SiO₂, Benzene: Isopropanol; 10:1).

IR (film)/cm⁻¹: 3274 (b,s), 2922 (s), 1575 (s), 1438 (s), 1294 (s), 1025 (s), 747 (s).

δ_H (500MHz, CD₃OD): 2.77 (dd, $J_{16.5\text{Hz}}$, 5Hz, 1H, C(3)HH'), 3.02 (dd, $J_{16.5\text{Hz}}$, 5Hz, 1H, C(3)HH'), 3.14 (app. quintet, $J_{6\text{Hz}}$, 1H, C(2)H), 3.53 (dd, $J_{11.0\text{Hz}}$, 7.1Hz, 1H, C(1)HH'), 3.68 (d, $J_{16\text{Hz}}$, 1H), 3.77-3.88 (m, 4H), 6.60 (dd, $J_{8.5\text{Hz}}$, 3Hz, Ar-H), 6.98 (d, $J_{7\text{Hz}}$, 1H, Ar-H), 7.02 (d, $J_{3\text{Hz}}$, 1H, Ar-H), 7.08-7.14 (m, 3H, Ar-H), 7.32 (d, $J_{8.5\text{Hz}}$, 1H, Ar-H).

δ_c (125MHz, CD₃OD): 29.9, 51.9, 58.3, 60.4, 62.8, 114.1, 117.0, 118.8, 126.9, 127.4, 127.6, 130.0, 134.4, 134.9, 135.4, 140.4, 158.2.

m/z (ESI): 348 (MH⁺, ⁷⁹Br; 100%), 350 (MH⁺, ⁸¹Br; 100%).

HRMS found 348.0598. C₁₇H₁₉BrNO₂ (MH⁺, ⁷⁹Br) requires 348.0599.

(S)-4-Bromo-3-(3-hydroxymethyl-3,4-dihydro-1H-isoquinolin-2-ylmethyl)-

phenol, A1B11. 1:1 Toluene/THF was used as solvent.

Colorless oil. Yield: 62%; $R_f = 0.37$ (SiO₂, Benzene: Isopropanol; 1:1).

δ_{H} (500MHz, CD₃OD): 1.44-1.53 (m, 1H), 1.58-1.71 (m, 3H), 1.75-1.84 (m, 2H), 2.58 (m, 1H, C(6)HH'), 2.88 (m, 1H, C(2)H), 2.99 (m, 1H, C(6)HH'), 3.74 (dd, *J*12Hz, 4.5Hz, 1H, C(1)HH'), 3.81 (m, 2H, C(1)HH' and C(7)HH'), 4.35 (d, *J*14Hz, 1H, C(1)HH'), 6.68 (dd, *J*9Hz, 3Hz, 1H, C(9)H), 7.06 (d, *J*3Hz, 1H, C(10)H), 7.37 (d, *J*9Hz, 1H, C(8)H).

δ_{C} (125MHz, CD₃OD): 23.5, 24.4, 28.1, 52.5, 57.3, 62.8, 65.3, 114.6, 118.3, 119.9, 134.7, 136.6, 158.5.

m/z (ESI): 300 (MH⁺, ⁷⁹Br; 100%), 302 (MH⁺, ⁸¹Br; 96%).

HRMS found 300.0594. C₁₃H₁₉BrNO₂ (MH⁺, ⁷⁹Br) requires 300.0599.

(*S,S*)-4-Bromo-3-[(2-hydroxy-1-methyl-2-phenylethyl)-methylamino]-

methyl}-phenol, A1B13. 1:1 Toluene/THF was used as solvent.

White solid. Yield: 63%; *R*_f = 0.38 (SiO₂, Hexanes : Ethyl Acetate; 1:1).

δ_{H} (500MHz, CD₃OD): 0.81 (d, *J*7Hz, 3H, CHCH₃), 2.28 (s, 3H, NCH₃), 2.80 (m, 1H, NCHCH₃), 3.68 (ABq, *J*_{AB}13Hz, 2H, NCH₂Ar), 4.32 (d, *J*10Hz, 1H, OCHPh), 6.67 (dd, *J*8.5Hz, 3Hz, 2H, Ar-H), 7.24–7.32 (m, 5H, Ar-H), 7.38 (d, *J*8.5Hz, 1H, Ar-H).

δ_{C} (125MHz, CD₃OD): 7.9, 35.3, 59.7, 65.0, 76.0, 114.6, 117.4, 119.7, 128.5, 129.0, 129.3, 134.7, 139.7, 143.0, 158.3.

m/z (ESI): 350 (MH⁺; 100%), 352 (MH⁺, ⁸¹Br; 99%).

HRMS found 350.0745. C₁₇H₂₁BrNO₂ (MH⁺, ⁷⁹Br) requires 350.0755.

7. Reductive Alkylation of Secondary Amines:

A glass vial, screwcapped with a teflon-lined septum containing the polymer supported aminoalcohol (70mg, 0.70 mmol/g) was charged with a solution of aliphatic aldehyde in Trimethyl Orthoformate (1mL). The suspension was agitated for 48h at ambient temperature. The solution was removed and the beads washed with 50:50:1 THF/DMF/TEA (2 x 20 min) and 100:1 THF/TEA (2 x 20 min), then resuspended in THF (1mL). A stock solution of Sodium Cyanoborohydride in Methanol (1M, 1mL, 1mmol, equiv.) was then added, followed by 100 μ L of glacial Acetic Acid (1.77mmol, equiv.). The suspension was agitated at ambient temperature for 16 hours and the solution drained. The beads were washed with 50:50:1 THF/DMF/TEA (3 x 30 min), 100:1 THF/TEA (3 x 30 min), 1:1 THF/MeCN (2 x 20 min), MeCN (2 x 20 min), CH₂Cl₂ (4 x 30 min) and dried *in vacuo* (house-vacuum for 2h, then high-vacuum for 12h). Cleavage of compound from resin was followed by purification and spectroscopic characterization.

Fig.2. Aldehyde building blocks tested in reductive alkylation of primary amines. Building blocks **C1-C6** were selected for library synthesis. Reactions with **C7** and **C8** resulted in ca. 10% of a side-product from reduction of the olefin of the α,β -unsaturated aldehyde and were not selected. A major side product from **C9** resulted from aldol reaction of the starting aldehyde prior to reductive amination.

4-Bromo-3-[(cyclopropylmethyl)-(2*S*-hydroxy-1*R*-methyl-2-phenylethyl)-amino]-methyl-phenol, A1B4C1.

Colorless oil. Yield: 54%; R_f = 0.38 (SiO₂, Benzene: Isopropanol; 10:1).

δ_H (500MHz, CDCl₃): 0.05 (m, 2H), 0.41 (m, 1H), 0.50 (m, 1H), 0.78 (m, 1H), 1.12 (d, J 7Hz, 3H, NCHCH₃), 2.24 (dd, J 13Hz, 7.5Hz, 1H, NCH'HCH), 2.43 (dd, J 13Hz, 6Hz,

1H, NCHHCH), 3.32 (m, 1H, NCHCH₃), 3.64 (ABq, J_{AB} 15.5Hz, 2H, NCH₂Ar), 4.71 (d, J 6Hz, 1H, OCHPh), 5.22 (bs, ArOH), 6.57 (dd, J 9Hz, 3Hz, 1H, ArH), 6.74 (d, J 3Hz, 1H, ArH), 7.27–7.38 (m, 6H, ArH).

δ_c (125MHz, CDCl₃): 2.9, 5.3, 9.7, 10.2, 54.5, 56.0, 60.3, 74.9, 113.5, 115.6, 116.9, 126.8, 127.2, 128.1, 133.2, 140.0, 142.6, 155.5.

m/z (ESI): 390 (MH⁺, ⁷⁹Br; 100%), 392 (MH⁺, ⁸¹Br; 99%).

HRMS found 390.1077. C₂₀H₂₅BrNO₂ (MH⁺, ⁷⁹Br) requires 390.1068.

4-Bromo-3-[(2S-hydroxy-1R-methyl-2-phenylethyl)-(3-methylbutyl)-amino]-methyl}-phenol, A1B4C2.

Colorless oil. Yield: 66%; R_f = 0.39 (Triethylamine-washed SiO₂, Ethyl Acetate: Hexanes: Triethylamine; 60:40:1).

δ_H (500MHz, CDCl₃): 0.81 (m, 6H, CH(CH₃)₂), 1.09 (d, J 7Hz, 3H, NCHCH₃), 1.31 (m, 2H, NCH₂CH₂), 1.43 (m, 1H, CH(CH₃)₂), 2.39 (m, 2H, NCH₂CH₂), 3.02 (quintet, J 7Hz, 1H, NCHCH₃), 3.58 (ABq, J_{AB} 15.5Hz, 2H, NCH₂Ar), 4.65 (d, J 6Hz, 1H, OCHPh), 4.68 (bs, ArOH), 6.52 (dd, J 8.5Hz, 3Hz, 1H, ArH), 6.57 (d, J 3Hz, 1H, ArH), 7.25–7.34 (m, 6H, ArH).

δ_c (100MHz, CDCl₃): 10.0, 22.8, 22.8, 26.5, 37.1, 49.3, 55.1, 60.5, 74.5, 113.5, 115.8, 117.2, 126.7, 127.1, 128.1, 133.3, 140.0, 142.7, 155.6.

m/z (ESI): 406 (MH⁺, ⁷⁹Br; 100%), 408 (MH⁺, ⁸¹Br; 99%).

HRMS found 406.1376. C₂₁H₂₉BrNO₂ (MH⁺, ⁷⁹Br) requires 406.1381.

4-Bromo-3-[(3R,7-dimethyl-oct-6-enyl)-(2S-hydroxy-1R-methyl-2-phenylethyl)-amino]-methyl-phenol, A1B4C3.

Colorless oil. Yield: 82%; R_f = 0.38 (SiO₂, Benzene: Isopropanol; 10:1).

δ_H (500MHz, CDCl₃): 0.81 (d, *J*7Hz, 3H, CH₂CHCH₃), 1.08–1.14 (m, 1H), 1.14 (overlaps with multiplet; d, *J*7Hz, 3H, NCHCH₃), 1.21–1.43 (m, 4H), 1.59 (s, 3H, C(Me)CH₃), 1.69 (s, 3H, C(Me)CH₃), 1.88–1.99 (m, 2H, Me₂C:CHCH₂), 2.45 (m, 2H, NCH₂CH₂), 2.99 (m, 1H, NCHCH₃), 3.54 (d, *J*16Hz, 1H, NCH'HA_r), 3.64 (d, *J*16Hz, 1H, NCH'HA_r), 4.67 (d, *J*6.5Hz, 1H, OCHPh), 5.08 (m, 1H, Me₂C:CH), 6.40 (s, 1H, Ar-H), 6.54 (dd, *J*8.5Hz, 3Hz, 1H, Ar-H), 7.27–7.35 (m, 6H, Ar-H).

δ_c (100MHz, CDCl₃): 9.9, 17.7, 19.8, 25.6, 25.8, 31.0, 35.2, 37.2, 49.0, 55.2, 61.1, 75.1, 113.6, 115.4, 117.1, 124.8, 126.9, 127.1, 128.1, 131.3, 133.2, 143.3, 155.2.

m/z (ESI): 474 (MH⁺, ⁷⁹Br; 80%), 476 (MH⁺, ⁸¹Br; 100%).

HRMS found 474.2006. C₂₆H₃₇BrNO₂ (MH⁺, ⁷⁹Br) requires 474.2007.

4-Bromo-3-[(2*S*-hydroxy-1*R*-methyl-2-phenylethyl)-(3-phenylpropyl)-amino]-methyl-phenol, A1B4C4, 5.

Colorless oil. Yield: 89%; $R_f = 0.23$ (Triethylamine-washed SiO_2 , Ethyl Acetate: Hexanes: Triethylamine; 60:40:1).

δ_H (500MHz, CDCl_3): 1.08 (d, J 7Hz, 3H, NCHCH_3), 1.71 (m, 2H, NCH_2CH_2), 2.44 (m, 4H, NCH_2CH_2 and PhCH_2CH_2), 2.98 (quintet, J 7Hz, 1H, NCHCH_3), 3.58 (ABq, J_{AB} 15.5Hz, 2H, NCH_2Ar), 4.60 (d, J 6.5Hz, 1H, OCHPh), 6.44 (d, J 3Hz, 1H, Ar-H), 6.51 (dd, J 9Hz, 3Hz, 1H, Ar-H), 7.22–7.31 (m, 6H, Ar-H).

δ_c (100MHz, CDCl_3): 10.0, 29.8, 33.6, 50.6, 55.2, 60.9, 74.9, 113.6, 115.7, 117.1, 125.9, 126.8, 127.1, 128.1, 128.3, 128.4, 133.3, 140.0, 142.0, 142.9, 155.3.

m/z (ESI): 454 (MH^+ , ^{79}Br ; 100%), 456 (MH^+ , ^{81}Br ; 98%).

HRMS found 454.1386. $\text{C}_{25}\text{H}_{29}\text{BrNO}_2$ (MH^+ , ^{79}Br) requires 454.1381.

4-Bromo-3-[(2*S*-hydroxy-1*R*-methyl-2-phenylethyl)-pent-4-enyl-amino]-methyl-phenol, A1B4C5.

Colorless oil. Yield: 94%; $R_f = 0.28$ (Triethylamine-washed SiO_2 , Ethyl Acetate: Hexanes: Triethylamine; 60:40:1).

δ_{H} (500MHz, CDCl_3): 1.11 (d, J 7Hz, 3H, NCHCH_3), 1.50 (m, 2H, NCH_2CH_2), 1.92 (m, 2H, $\text{NCH}_2\text{CH}_2\text{CH}_2$), 2.40 (m, 2H, NCH_2CH_2), 3.01 (quintet, J 7Hz, 1H, NCHCH_3), 3.58 (ABq, J_{AB} 15.5Hz, 2H, NCH_2Ar), 4.65 (d, J 6Hz, 1H, OCHPh), 4.94 (m, 2H, CH:CH_2), 5.72 (m, 1H, CH:CH_2), 6.49 (m, 1H, Ar-H), 6.53 (dd, J 8.5Hz, 2.9Hz, 1H, Ar-H), 7.25–7.36 (m, 5H, Ar-H).

δ_{C} (100MHz, CDCl_3): 10.0, 27.4, 31.5, 50.6, 55.2, 60.8, 74.9, 113.6, 114.8, 115.6, 117.0, 126.8, 127.2, 128.1, 133.3, 138.4, 140.1, 142.9, 155.3.

m/z (ESI): 404 (MH^+ , ^{79}Br ; 97%), 406 (MH^+ , ^{81}Br ; 100%).

HRMS found 404.1231. $\text{C}_{21}\text{H}_{27}\text{BrNO}_2$ (MH^+ , ^{79}Br) requires 404.1225.

4-Bromo-3-[(2S)-hydroxy-1R-methyl-2-phenylethyl]-(5-hydroxypentyl)-amino]-methyl}-phenol, A1B4C6.

Colorless oil. Yield: 82%; R_f = 0.26 (Triethylamine-washed SiO_2 , Ethyl Acetate: Methanol:Triethylamine; 60:40:1).

δ_{H} (400MHz, C_6D_6): 0.82-1.02 (m, 4H, $\text{NCH}_2\text{CHH}'$, $\text{NCH}_2\text{CH}_2\text{CH}_2$ and $\text{CHH}'\text{CH}_2\text{OH}$), 1.06 (d, J 6.5Hz, 3H, NCHCH_3), 1.18 (m, 1H, $\text{CHH}'\text{CH}_2\text{OH}$), 1.32 (m, 1H, $\text{NCH}_2\text{CHH}'$), 2.18 (m, 2H, NCH_2CH_2), 2.97 (quintet, J 6.5Hz, 1H, NCHCH_3), 3.17 (m, 1H, $\text{CH}_2\text{CHH}'\text{OH}$), 3.34 (m, 1H, $\text{CH}_2\text{CHH}'\text{OH}$), 3.67 (ABq, J_{AB} 16.5Hz, 2H, NCH_2Ar), 4.35 (d, J 6.5Hz, 1H, OCHPh), 6.72 (dd, J 9Hz, 3Hz, 1H, Ar-H), 7.02 (m, 2H, Ar-H), 7.10 (m, 2H, Ar-H), 7.16 (m, 2H, Ar-H), 7.31 (d, J 8.5Hz, 1H, Ar-H).

δ_c (100MHz, C_6D_6): 9.2, 22.0, 26.7, 31.3, 49.6, 54.8, 59.8, 63.0, 75.7, 112.9, 116.3, 117.2, 127.1, 127.4, 127.9, 133.7, 140.5, 144.7, 157.2.

HRMS found 422.1327. $C_{22}H_{29}BrNO_2$ (MH^+ , ^{79}Br) requires 422.1331.

8. Mitsunobu Reaction:

A glass vial, screwcapped with a teflon-lined septum containing the polymer supported aminoalcohol (70mg, 0.70mmol/g) was charged with 2 mL of a stock solution of tributylphosphine (3.1mL, 12.6mmol) and bromophenol (12.6mmol) in THF (60 mL). The suspension was gently agitated for 10 min and then cooled to $-78^\circ C$. 1 mL of a stock solution of TMAD in THF (2.16g, 12.6mmol in 31mL) was then added dropwise with constant agitation. The reaction mixture was then allowed to slowly warm to room temperature and agitated for 24h at ambient temperature and the solution drained. The beads were washed with 100:1 DMF/TEA (3 x 30 min), 50:50:1 THF/DMF/TEA (3 x 30 min), 100:1 THF/TEA (3 x 30 min), and CH_2Cl_2 (4 x 30 min); then dried *in vacuo* (house-vacuum for 2h, then high-vacuum for 12h). Cleavage of compound from resin was followed by purification and spectroscopic characterization.

Fig.3. Phenol building blocks tested in the Mitsunobu reaction. Reactions with building blocks **D1-D5** resulted in complete conversion to products with high purity; **D1** and **D2** were selected for library synthesis. Reaction with **D6** resulted in 87% conversion and **D7-D9** yielded no product. These building blocks were not selected for library synthesis.

4-Bromo-3-((2*S*-(2-bromophenoxy)-1*R*-methyl-2-phenylethyl)-methanimino)-methylphenol, A1B7D1.

Colorless oil. Yield:52%; R_f = 0.23 (Triethylamine-washed SiO_2 , Hexanes: Ethyl Acetate: Triethylamine; 50:50:1).

δ_{H} (400MHz, CDCl_3): 1.35 (d, J 6.5Hz, 3H, NCHCH_3), 2.39 (s, 3H, NCH_3), 3.12 (dq, J 6.5Hz, 5.5Hz, 1H, NCHCH_3), 3.63 (d, J 15.5Hz, 1H, $\text{ArCHH}'\text{N}$), 3.78 (d, J 15.5Hz, 1H, $\text{ArCHH}'\text{N}$), 5.30 (d, J 5.5Hz, 1H, OCHPh), 6.33 (d, J 3.5Hz, 1H, Ar-H), 6.54 (dd, J 8.5Hz, 3.1Hz, 1H, Ar-H), 6.62 (dd, J 8Hz, 1.5Hz, 1H, Ar-H), 6.73 (dt, J 7.5Hz, 1.5Hz, 1H, Ar-H), 7.03 (m, 1H, Ar-H), 7.27-7.35 (m, 6H, Ar-H), 7.50 (dd, J 8Hz, 2Hz, 1H).

δ_{C} (100MHz, CDCl_3): 9.3, 37.8, 58.1, 64.1, 81.7, 112.4, 113.8, 114.3, 115.3, 116.8, 121.6, 126.8, 127.2, 128.1, 128.5, 133.2, 133.3, 139.8, 140.5, 153.9, 154.9.

m/z (ESI): 504 (MH^+ , ^{79}Br , ^{79}Br ; 58%), 506 (MH^+ , ^{79}Br , ^{81}Br ; 100%), 508 (MH^+ , ^{81}Br , ^{81}Br ; 53%).

HRMS found 504.0177. $\text{C}_{23}\text{H}_{24}\text{Br}_2\text{NO}_2$ (MH^+ , ^{79}Br) requires 504.0174.

4-Bromo-3-({[2S-(2-bromopyridin-3-yloxy)-1R-methyl-2-phenylethyl]-methyl-amino}-methyl)-phenol, A1B7D2.

Colorless oil. Yield: 82%; R_f = 0.43 (SiO_2 , Ethyl Actetate).

δ_{H} (500MHz, CDCl_3): 1.35 (d, J 7Hz, 3H, NCHCH_3), 2.39 (s, 3H, NCH_3), 3.15 (dq, J 7Hz, 5Hz, 1H, NCHCH_3), 3.65 (d, J 15Hz, 1H, $\text{ArCHH}'\text{N}$), 3.76 (d, J 15Hz, 1H, $\text{ArCHH}'\text{N}$), 5.31 (d, J 5Hz, 1H, OCHPh), 6.49 (d, J 3Hz, 1H, Ar-H), 6.57 (dd, J 8.5Hz, 3Hz, 1H, Ar-H), 6.85 (dd, J 8.5Hz, 1.5Hz, 1H, Ar-H), 6.98 (dd, J 8Hz, 4.5Hz, 1H, Ar-H), 7.29 (m, 5H, Ar-H), 7.34 (d, J 7Hz, 1H, Ar-H), 7.87 (dd, J 4.5Hz, 1.5Hz, 1H, Ar-H).

δ_{C} (125MHz, CDCl_3): 9.2, 37.9, 58.3, 64.0, 82.5, 113.8, 115.5, 117.0, 121.1, 123.2, 126.6, 127.7, 128.3, 128.7, 133.2, 139.2, 139.5, 141.0, 151.1, 155.2.

m/z (ESI): 505 (MH^+ , ^{79}Br , ^{79}Br ; 50%), 507 (MH^+ , ^{79}Br , ^{81}Br ; 100%).

HRMS found 505.0126. C₂₂H₂₃Br₂N₂O₂ (MH⁺, ⁷⁹Br) requires 505.0122.

Stereochemistry of the Mitsunobu reaction of 1,2-aminoalcohols with phenols:

The reaction of 1,2-aminoalcohols under Mitsunobu conditions to yield products corresponding to overall retention of configuration at the reacting center is precedented^x. This is presumably due to the involvement of an aziridinium ion intermediate. In our hands, Mitsunobu reactions of (1*S*,2*R*)-*N,N*-dimethylephedrine with 2- and 4-nitrophenols yielded the same products as were obtained with S_N_{Ar} reactions with 2- and 4-fluoronitrobenzenes (Fig.1).

Fig.4. Mitsunobu reactions of (1*S*,2*R*)-*N,N*-dimethylephedrine with nitrophenols and S_N_{Ar} reactions with the corresponding fluoronitrobenzenes yield the same products.

Mitsunobu Reaction of amino alcohols with nitrophenols:

(1*S*,2*R*)-*N,N*-dimethylephedrine (1.0 equiv.) was dissolved in THF (0.1M) with the nitrophenol (1.5 equiv.) and triphenylphosphine (1.5 equiv.). Diisopropylazodicarboxylate (1.5 equiv.) was then added by syringe and the reaction

(6) For Mitsunobu reactions proceeding via aziridinium ion intermediates, see (a) Freedman, J.; Vaal, M.J.; Huber, E.W. *J. Org. Chem.* **1991**, 56, 670-672. (b) Okuda, M.; Tomioka, K. *Tetrahedron Lett.* **1994**, 35, 4585-4586. (c) Poelert, M. A.; Hulshof, L.A.; Kellogg, R.M. *Rec. Trav. Chim. Pays-Bas.* **1994**, 113, 355-364.

mixture stirred for 12h, then diluted with EtOAc and washed with 1M NaOH (x 2) and brine (x 1). The EtOAc layer was then dried over K₂CO₃ and concentrated under reduced pressure. The residue thus obtained was chromatographed over SiO₂ to yield the aminoether.

SN_{Ar} reaction of amino alcohols with nitrofluorobenzenes:

(*1S,2R*)-*N,N*-dimethylephedrine (1.0 equiv.) was dissolved in THF (0.1M) and the solution cooled to -78°C. KHMDS (1.5 equiv.) was added via syringe, followed by the nitrofluorobenzene (1.2 equiv.). The reaction mixture was allowed to warm to room temperature over 12h, then poured into saturated aqueous sodium bicarbonate and extracted with EtOAc (x 3). The EtOAc extracts were combined, washed with brine (x 1), then dried over K₂CO₃ and concentrated and the residue chromatographed over SiO₂ to yield the aminoether.

(1*R*,2*S*)-Dimethyl-[1-methyl-2-(2-nitrophenoxy)-2-phenylethyl]-amine.

Yield: 83% (Mitsunobu), 98% (SN_{Ar}). R_f = 0.37 (TEA-washed SiO₂, Hexanes/EtOAc/TEA : 50:50:1).

IR (film/cm⁻¹): 2937m, 2825m, 2781m, 1605s, 1525s, 1485s, 1452s, 1353s, 1277s, 1253s, 991s, 860s, 743s, 701s.

δ_H (500MHz, CDCl₃): 1.18 (d, *J*7Hz, 3H, -CHCH₃), 2.36 (s, 6H, -NCH₃), 2.96 (dq, *J*7Hz, 3Hz, 1H, -NCHCH₃), 5.51 (d, *J*3Hz, 1H, -OCHPh), 6.79 (d, *J*9Hz, 1H, Ar-H), 6.92 (t, *J*8Hz, 1H, Ar-H), 7.25-7.35 (m, 6H, Ar-H), 7.82 (m, 1H, Ar-H).

δ_c (125MHz, $CDCl_3$): 7.8 (-CHCH₃), 41.7 (-NCH₃), 65.6 (-NCHCH₃), 81.5 (-OCHPh), 115.6, 120.1, 125.6, 126.0, 127.8, 128.7, 133.9, 139.2, 140.1, 150.8.

m/z (ESI): 301 (MH^+ :100%).

HRMS found 301.1547. $C_{17}H_{21}N_2O_3$ (MH^+) requires 301.1552.

$[\alpha]_D$ (23°C) = -65.7 (c = 0.9, $CHCl_3$).

(1R,2S)-Dimethyl-[1-methyl-2-(4-nitrophenoxy)-2-phenylethyl]-amine.

Yield: 54% (Mitsunobu), 81% (SN_{Ar}). R_f = 0.44 (TEA-washed SiO_2 , Hexanes/EtOAc/TEA : 50:50:1).

IR (film/ cm^{-1}): 1590s, 1512s, 1340s, 1254s, 1111s.

δ_H (500MHz, $CDCl_3$): 1.15 (d, J 7Hz, 3H, -CHCH₃), 2.38 (s, 6H, -NCH₃), 2.93 (dq, J 7Hz, 3Hz, 1H, -NCHCH₃), 5.44 (d, J 3Hz, 1H, -OCHPh), 6.88 (app. d, J 9.5Hz, 2H, Ar-H), 7.27 (m, 2H, Ar-H), 7.34 (m, 3H, Ar-H), 8.09 (app. d, J 9.5Hz, 2H, Ar-H).

δ_c (125MHz, $CDCl_3$): 8.2 (-CHCH₃), 41.8 (-NCH₃), 65.3 (-NCHCH₃), 81.4 (-OCHPh), 115.8, 125.8, 126.0, 127.8, 128.7, 139.2, 141.5, 162.8.

m/z (ESI): 301 (MH^+ :100%).

HRMS found 301.1554. $C_{17}H_{21}N_2O_3$ (MH^+) requires 301.1552.

$[\alpha]_D$ (23°C) = +15.3 (c = 1.06, $CHCl_3$).

Mitsunobu reactions of Ephedrine and Pseudoephedrine yielded different products. Pseudoephedrine yielded 5% of a product resulting from attack at the carbon bearing the methyl group (Fig.2.).

Fig.5. Mitsunobu reactions of (1*S*,2*R*)-*N,N*-dimethylephedrine and (1*S*,2*S*)-*N,N*-dimethylpseudoephedrine with 2-Bromophenol yield different products.

Mitsunobu Reaction of amino alcohols with 2-bromophenol:

The amino alcohol (1.0 equiv.) was dissolved in Benzene (0.1M) with the 2-bromophenol (1.5 equiv.) and triphenylphosphine (1.5 equiv.). Solid 1,1'-Azodicarbonyldipiperidine (1.5 equiv.) was then added and the reaction mixture stirred for 12h, then diluted with EtOAc and washed with 1M NaOH (x 2) and brine (x 1). The EtOAc layer was then dried over K₂CO₃ and concentrated under reduced pressure. The residue thus obtained was chromatographed over SiO₂ to yield the aminoether.

(1*R*,2*S*)-Dimethyl-[1-methyl-2-(2-bromophenoxy)-2-phenylethyl]-amine. Colorless oil.

Yield: 92%. $R_f = 0.33$ (TEA-washed SiO_2 , Hexanes/EtOAc/TEA : 66:33:1).

IR (film/ cm^{-1}): 1476s, 1276s, 1245s, 1030m, 995m, 744s, 700m.

δ_{H} (500MHz, CDCl_3): 1.24 (d, J 7.5Hz, 3H, $-\text{CHCH}_3$), 2.41 (s, 6H, $-\text{NCH}_3$), 2.96 (m, 1H, $-\text{NCHCH}_3$), 5.44 (d, J 2.5Hz, 1H, $-\text{OCHPh}$), 6.58 (dd, J 8.5Hz, 1.5Hz, 1H, Ar-H), 6.73 (m, 1H, Ar-H), 7.03 (m, 1H, Ar-H), 7.22-7.33 (m, 5H, Ar-H), 7.52 (dd, 1H, J 8Hz, 1.5Hz, 1H, Ar-H).

δ_{C} (125MHz, CDCl_3): 8.1 ($-\text{CHCH}_3$), 41.7 ($-\text{NCH}_3$), 65.9 ($-\text{NCHCH}_3$), 80.1 ($-\text{OCHPh}$), 112.5, 114.2, 121.5, 126.0, 127.4, 128.2, 128.5, 133.3, 140.0, 153.6.

HRMS found 334.0808. $\text{C}_{17}\text{H}_{21}\text{BrNO}$ (MH^+) requires 334.0806.

$[\alpha]_{\text{D}}$ (23°C) = -120.5 ($c = 2.33$, CHCl_3).

(1S,2S)-Dimethyl-[1-methyl-2-(2-bromophenoxy)-2-phenylethyl]-amine. Colorless oil.

Yield: 93%. $R_f = 0.31$ (TEA-washed SiO_2 , Hexanes/EtOAc/TEA : 83:17:1).

IR (film/ cm^{-1}): 1475s, 1244s, 1030m, 746s, 702m.

δ_{H} (500MHz, CDCl_3): 0.94 (d, J 7Hz, 3H, $-\text{CHCH}_3$), 2.44 (s, 6H, $-\text{NCH}_3$), 3.18 (q, J 7Hz, 1H, $-\text{NCHCH}_3$), 5.23 (d, J 7Hz, 1H, $-\text{OCHPh}$), 6.70 (m, 2H, Ar-H), 7.03 (dt, J 8Hz, 1.5Hz, 1H, Ar-H), 7.29 (m, 5H, Ar-H), 7.47 (dd, 1H, J 8Hz, 2Hz, 1H, Ar-H).

δ_{C} (125MHz, CDCl_3): 12.7, 42.1, 63.9, 82.7, 112.8, 114.7, 121.6, 127.3, 127.8, 128.1, 128.4, 133.3, 139.1, 154.2.

HRMS found 334.0811. $C_{17}H_{21}BrNO$ (MH^+) requires 334.0806.

$[\alpha]_D$ (23°C) = -59.0 ($c = 2.37$, $CHCl_3$).

(1*R*, 2*R*)-[2-(2-Bromophenoxy)-1-phenylpropyl]-dimethylamine. Colorless oil.

Yield: 5%. $R_f = 0.38$ (TEA-washed SiO_2 , Hexanes/EtOAc/TEA : 83:17:1).

IR (film/ cm^{-1}): 1475s, 1245s, 1030m, 748s, 704m.

δ_H (500MHz, $CDCl_3$): 1.13 (d, J 6Hz, 3H, -CHCH $\underline{H}_3$), 2.30 (s, 6H, -NCH $\underline{H}_3$), 3.59 (d, J 6.5Hz, 1H, -NCH $\underline{H}$ Ph), 4.90(dq, J 6.5Hz, 6Hz, 1H, -OCH $\underline{H}$ Me), 6.82 (dt, J 8Hz, 1.5Hz 1H, Ar- $\underline{H}$), 6.98 (d, J 8.5Hz, 1H, Ar- $\underline{H}$), 7.22-7.34 (m, 6H, Ar- $\underline{H}$), 7.54 (dd, 1H, J 8Hz, 2Hz, 1H, Ar- $\underline{H}$).

δ_c (125MHz, $CDCl_3$): 16.4, 43.8, 74.1, 75.8, 113.9, 115.5, 122.1, 127.4, 128.0, 128.3, 129.7, 133.7, 137.7, 154.4.

HRMS found 334.0806. $C_{17}H_{21}BrNO$ (MH^+) requires 334.0806.

$[\alpha]_D$ (23°C) = -17.5 ($c = 0.28$, $CHCl_3$).

9. Biaryl Coupling:

The polymer supported cyclization precursor (20mg, 0.60mmol/g, 0.012mmol) in a glass vial, screwcapped with a teflon-lined septum was swollen in anhydrous toluene (2 x 5mL x 20 min) and then dried *in vacuo* at 50°C for 30 min. The vial was placed in a bath maintained at 0°C (cryocool) and a pre-cooled solution of ⁱPrⁿBu₂MgLi¹ (4.0 mL of a solution ca. 0.3M in 2-MeTHF, ca. 100 equiv) was added *via* syringe and the beads incubated in this solution at 0°C for 6-8h. The solution was then removed *via* cannula under a positive pressure of Ar, and a pre-cooled solution of CuCN.2LiBr (1.0M in 2-MeTHF, 1.2mL, 100 equiv.) added. The beads were incubated in this solution at 0°C for 6h and the solution then removed *via* cannula under a positive pressure of Ar. A pre-cooled solution of *m*-dinitrobenzene (1.0M in 2-MeTHF, 1.2mL, 100 equiv.) was then added and the vials transferred to a cold cabinet maintained at 6 °C where they were agitated for 12h. The beads were then filtered and washed with 8-Hydroxyquinoline (5% w/v in 50:50:1 THF (stabilized):DMF:TEA; 2 x 10 min, 2 x 12h) with gentle agitation; then further washed with 50:50:1 THF/DMF/TEA (3 x 30 min), 100:1 THF/TEA (3 x 30 min), and CH₂Cl₂ (4 x 30 min); then dried *in vacuo* (house-vacuum for 2h, then high-vacuum for 12h). Cleavage of compound from resin was followed by purification and spectroscopic characterization.

(7) To 45 mL anhydrous 2-MeTHF was added ⁱPrMgCl (1.59M in THF, 19.0mL, 30.21mmol, 1.0 equiv.) and the solution cooled to -30°C. ⁿBuLi (2.1M in Hexanes, 28.0mL, 58.8mmol, 1.95 equiv.) was then added and the mixture stirred at -30°C till a clear yellow solution resulted. This solution was used immediately for further reactions.

(P, 6S, 7R)-7-Methyl-6-phenyl-8-(3-phenyl-propyl)-6,7,8,9-tetrahydro-5-oxa-1,8-diaza-dibenzo[*a,c*]cyclononen-11-ol, *c*-A1B4C4D2, 7.

Colorless oil. Yield: 80%. R_f = 0.30 (Triethylamine washed SiO_2 , Hexanes/EtOAc/ Et_3N : 25:75:1).

δ_{H} (500MHz, C_6D_6): 0.98 (d, J 7Hz, 3H, CHCH_3), 1.67 (app. quintet, J 7Hz, 2H, C(15) $\underline{\text{H}}_2$), 2.39 (app. quintet, J 7Hz, 1H, C(14) $\underline{\text{H}}\text{H}'$), 2.47 (app. quintet, J 6.5Hz, 1H, C(14) $\text{H}\underline{\text{H}}'$), 2.53 (app. quintet, J 6.5Hz, 1H, C(16) $\underline{\text{H}}\text{H}'$), 2.62 (app. quintet, J 6.5Hz, 1H, C(16) $\text{H}\underline{\text{H}}'$), 2.96 (q, J 7Hz, 1H, C(7) $\underline{\text{H}}$), 3.32 (d, J 14Hz, 1H, C(9) $\underline{\text{H}}\text{H}'$), 3.48 (d, J 14Hz, 1H, C(9) $\text{H}\underline{\text{H}}'$), 4.98 (s, 1H, C(6) $\underline{\text{H}}$), 6.52 (dd, J 8Hz, 4.5Hz, 1H, C(3) $\underline{\text{H}}$), 6.70 (dd, J 8Hz, 1Hz, 1H, C(4) $\underline{\text{H}}$), 6.82 (dd, J 8.5Hz, 2Hz, 1H, C(12) $\underline{\text{H}}$), 7.08 (m, 2H, Ar- $\underline{\text{H}}$), 7.12-7.19 (m, 8H, Ar- $\underline{\text{H}}$), 7.30 (d, J 2Hz, 1H, C(10) $\underline{\text{H}}$), 7.66 (d, J 8.5Hz, 1H, C(13) $\underline{\text{H}}$), 8.31 (dd, J 4.5Hz, 1Hz, 1H, C(2) $\underline{\text{H}}$).

δ_{C} (125MHz, C_6D_6): 14.7 ($-\text{CHCH}_3$), 30.5 (C-15), 33.6 (C-16), 55.3 (C-9), 57.6 (C-14), 67.5 (C-7), 85.2 (C-1), 115.4 (C-12), 117.5 (C-10), 123.3 (C-3), 125.8, 126.1, 126.5 (C-4), 127.2, 127.9, 128.1, 128.3, 128.5, 128.9, 131.3 (C-13), 141.2 (quat.), 142.6 (quat.), 144.1 (C-2), 153.9 (quat.), 154.0 (quat.), 158.0 (quat.).

m/z (ESI): 451 (MH^+ :100%).

HRMS found 451.2390. $\text{C}_{30}\text{H}_{31}\text{N}_2\text{O}_2$ (MH^+) requires 451.2385.

Fig.6. nOe's observed on irradiation of C(6)H for *c*-**A1B4C4D2**, **7** (NOESY1D, 500MHz).

(*M*, 6*R*)-8-Pent-4-enyl-6-phenyl-6,7,8,9-tetrahydro-5-oxa-1,8-diaza-dibenzo[*a,c*]cyclononen-11-ol, *c*-A1B2C5D2**, **8**.**

Pale yellow oil. Yield: 91%. R_f = 0.32 (Triethylamine washed SiO_2 , Hexanes/EtOAc/ Et_3N : 20:80:1).

δ_H (500MHz, CDCl_3): 1.21 (m, 1H, C(15)HH'), 1.39 (m, 1H, C(15)HH'), 1.84 (m, 2H, C(16)H₂), 2.44 (m, 2H, C(14)H₂), 2.69 (d, J 14.5Hz, 1H, C(7)HH'), 3.15 (dd, J 14.5Hz, 10Hz, 1H, C(7)HH'), 3.32 (d, J 13.5Hz, 1H, C(9)HH'), 3.51 (d, J 13.5Hz, 1H, C(9)HH'), 4.91-4.97 (m, 2H, C(18)H₂), 4.94 (d, J 10Hz, 1H, C(6)H; overlaps with the C(18)H₂ multiplet), 5.70 (tdd, J 17Hz, 10.5Hz, 6.5Hz, 1H, C(17)H), 6.60 (m, 2H, C(10)H and C(12)H), 7.09 (d, J 9.5Hz, 1H, Ar-H), 7.24-7.29 (m, 5H, Ar-H), 7.32 (m, 2H, Ar-H), 8.31 (dd, J 4.5Hz, 2Hz, 1H, C(2)H).

δ_c (100MHz, $CDCl_3$): 26.5 (C-15), 31.1 (C-16), 55.8 (C-9), 57.2 (C-14), 61.6 (C-7), 85.0 (C-6), 114.7, 115.0, 117.1, 123.5, 125.7, 127.1, 127.7, 128.3, 128.4, 129.3, 130.8, 138.1, 138.3, 140.2, 142.1, 155.9, 157.0.

m/z (ESI): 387 (MH^+ :29%).

HRMS found 387.2066. $C_{25}H_{27}N_2O_2$ (MH^+) requires 387.2072.

Fig.7. nOe's diagnostic of (*M*, 6*R*) stereochemistry for *c*-**A1B2C5D2**, **8** (ROESY, 500MHz).

(*P*, 6*R*)-8-Pent-4-enyl-6-phenyl-6,7,8,9-tetrahydro-5-oxa-8-aza-dibenzo[*a,c*]-cyclononen-11-ol, *c*-A1B2C5D1**, **9**.**

Colorless oil. Yield: 76%; R_f = 0.38 (Triethylamine-washed SiO_2 , Hexanes: Ethyl Acetate: Triethylamine; 50:50:1).

δ_H (500MHz, $CDCl_3$): 1.48-1.62 (m, 2H, C(15) $\underline{H}_2$), 2.02 (m, 2H, C(16) $\underline{H}_2$), 2.62 (m, 2H, C(14) $\underline{H}_2$), 2.85 (d, J_{15Hz} , 1H, C(7) $\underline{H}H'$), 3.25 (dd, J_{15Hz} , 8Hz, 1H, C(7) $\underline{H}H'$), 3.32 (d, J_{13Hz} , 1H, C(9) $\underline{H}H'$), 3.59 (d, J_{13Hz} , 1H, C(9) $\underline{H}H'$), 4.92-5.01 (m, 2H, C(18) $\underline{H}_2$), 5.02

(d, J 8Hz, 1H, C(6)H), 5.77 (tdd, J 17Hz, 10.5Hz, 6.5Hz, 1H, C(17)H), 6.82 (dd, J 8.5Hz, 2.5Hz, 1H, C(12)H), 6.86 (d, J 8.5Hz, 1H, C(13)H), 7.10 (m, 2H, Ar-H), 7.21-7.28 (m, 3H, Ar-H), 7.29-7.35 (m, 4H, Ar-H).

δ_c (125MHz, $CDCl_3$): 26.7 (C-15), 31.3 (C-16), 55.6 (C-9), 55.9 (C-14), 61.5 (C-7), 83.0 (C-6), 114.7, 114.8, 116.9, 119.2, 123.4, 125.4, 127.3, 128.4, 128.7, 129.6, 130.7, 131.8, 134.6, 138.1, 138.3, 142.0, 155.2, 157.6.

m/z (ESI): 386 (MH^+ :44%).

HRMS found 386.2115. $C_{26}H_{28}NO_2$ (MH^+) requires 386.2120.

Fig.8. nOe's diagnostic of (*P*, 6*R*) stereochemistry for *c*-A1B2C5D1, **9** (NOESY, 500MHz).

(*P*, 6*S*, 7*R*)-7,8-Dimethyl-6-phenyl-6,7,8,9-tetrahydro-5-oxa-1,8-diazadibenzo-
[*a,c*]cyclononen-11-ol, *c*-A1B7D2, **10**.

Yellow oil. Yield:58%; R_f = 0.31 (SiO_2 , Ethyl Actetate).

δ_{H} (500MHz, CDCl_3): 0.81 (d, J 7Hz, 3H, CHCH_3), 2.65 (s, 3H, NCH_3), 3.04 (q, J 7Hz, C(7)H), 3.30 (d, J 14Hz, 1H, C(9)HH'), 3.68 (d, J 14Hz, 1H, C(9)HH'), 5.20 (s, 1H, C(6)H), 6.66 (dd, J 8.5Hz, 2Hz, 1H, Ar-H), 6.99 (d, J 1.5Hz, 1H, Ar-H), 7.12-7.18 (m, 2H, Ar-H), 7.25-7.36 (m, 6H, Ar-H), 8.38 (d, J 4Hz, 1H, C(2)H).

δ_{C} (125MHz, CDCl_3): 13.5 (CHCH_3), 46.6 ($-\text{NCH}_3$), 56.6 (C-7 or C-9), 67.0 (C-7 or C-9), 83.4 (C-6), 115.4, 117.4, 123.4, 125.4, 126.8, 127.2, 127.9, 128.3, 130.5, 139.3, 141.4, 143.7, 152.8, 153.7, 157.4.

m/z (ESI): 347 (MH^+ :100%).

HRMS found 347.1756. $\text{C}_{22}\text{H}_{23}\text{N}_2\text{O}_2$ (MH^+) requires 347.1759.

Fig.9. nOe's diagnostic of (P, 6*S*, 7*R*) stereochemistry for ***c*-A1B7D2, 10** (NOESY, 500MHz).

(*M*, 6*S*, 7*R*)-8-(5-Hydroxypentyl)-7-methyl-6-phenyl-6,7,8,9-tetrahydro-5-oxa-1,8-diaza-dibenzo[*a,c*]cyclononen-11-ol, *c*-A1B4C6D2, 11.

Colorless oil. Yield: 69%. R_f = 0.31 (Triethylamine washed SiO_2 , EtOAc/MeOH/ Et_3N : 95:5:1).

δ_{H} (500MHz, CDCl_3): 0.77 (d, J 7Hz, 3H, CHCH_3), 1.44 (m, 2H, $\text{C}(17)\text{H}_2$), 1.51-1.58 (m, 4H, $\text{C}(15)\text{H}_2$ and $\text{C}(16)\text{H}_2$), 2.73 (m, 2H, $\text{C}(14)\text{H}_2$), 3.09 (q, J 7Hz, 1H, $\text{C}(7)\text{H}$), 3.37 (d, J 14Hz, 1H, $\text{C}(9)\text{HH}'$), 3.59 (m, 2H, $\text{C}(18)\text{H}_2$), 3.65 (d, J 14Hz, 1H, $\text{C}(9)\text{HH}'$), 5.16 (s, 1H, $\text{C}(6)\text{H}$), 6.79 (dd, J 8.5Hz, 2.5Hz, 1H, Ar-H), 7.08 (m, 2H, Ar-H), 7.12 (dd, J 8Hz, 5Hz, 1H, $\text{C}(3)\text{H}$), 7.27 (m, 1H, Ar-H), 7.31-7.36 (m, 5H, Ar-H), 8.35 (dd, J 5Hz, 1.5Hz, 1H, $\text{C}(2)\text{H}$).

δ_{C} (125MHz, CDCl_3): 13.9 ($-\text{CHCH}_3$), 23.4 (C-17), 27.8 (C-15 or C-16), 32.1 (C-15 or C-16), 54.6 (C-9), 57.7 (C-14), 62.4 (C-18), 67.0 (C-7), 85.0 (C-6), 115.2, 117.1, 123.3, 125.4, 126.6, 127.2, 128.2, 128.4, 130.5, 140.3, 141.7, 143.8 (C-3), 152.9, 153.6, 157.0.

m/z (ESI): 419 (MH^+ :100%).

HRMS found 419.2344. $\text{C}_{30}\text{H}_{31}\text{N}_2\text{O}_2$ (MH^+) requires 419.2334.

Stereochemistry assigned in analogy to *c*-A1B4C4D2, 7.

10. Synthesis of Dimer (15)

i. Double Reductive Amination:

3-Cyclopentenemethanol triisopropylsilyl ether (85.2mg, 0.335mmol) was dissolved in $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (10:1, 10mL) along with anhydrous NaHCO_3 (9.0mg, 0.101mmol, 0.3 equiv.) and the solution cooled to -78°C . Ozone was bubbled into the solution with constant stirring. After 5min, the solution turned blue indicating consumption of the olefin. Dimethylsulfide (0.25mL, 3.35mol, 10.0 equiv.) was then added and the solution allowed to gradually warm to room temperature, then concentrated under reduced pressure. The residue was azeotroped with Benzene (1 x 10mL). A solution of (1*S*, 2*R*)-*N*-(2-Bromobenzyl)-norephedrine (429.0mg, 1.339mmol, 4.0 equiv. in 5mL MeOH) was added, followed by glacial AcOH (0.5mL). Sodium Cyanoborohydride (s)(126.0mg, 2.01mmol, 6.0 equiv.) was added in small portions with stirring. After the addition was complete, the reaction mixture for stirred for 24h, then the excess reductant quenched by addition of 1M HCl (1mL). The reaction mixture was then poured into 1M NaOH (mL) and extracted with EtOAc (4 x 20mL). The EtOAc extracts were washed with brine (1 x 50mL), dried over K_2CO_3 and concentrated under reduced pressure. The residue thus obtained was chromatographed over SiO_2 (TEA-washed,

Hexanes/EtOAc/TEA : 75:25:1) to yield the diamine (**13**) as a colorless oil, 162.0mg (54% for two steps) R_p , 0.30 TEA-washed SiO_2 , Hexanes/EtOAc/TEA : 75:25:1).

IR (film)/ cm^{-1} : 3416b, 2940s, 2864s, 1463s, 1025s, 750s.

δ_H (500MHz, CDCl_3): 1.00 (m, 21H, C(9)H and C(10)H₃), 1.05 (m, 6H, C(3)H₃ and C(3')H₃), 1.19-1.34 (m, 3H), 1.41 (m, 1H), 1.50 (m, 1H), 2.31-2.44 (m, 4H, C(5)H₂ and C(5')H₂), 3.00 (m, 2H, C(2)H and C(2')H), 3.36 (m, C(8)H₂), 3.64 (m, 4H, C(4)H₂ and C(4')H₂), 4.65 (d, J 5.5Hz, 1H, C(1)H), 4.68 (d, J 5.5Hz, 1H, C(1')H), 7.04 (m, 2H, Ar-H), 7.14-7.32 (m, 14H, Ar-H), 7.47 (m, 2H, Ar-H).

δ_c (125MHz, CDCl_3): 9.95 (C-3), 10.03 (C-3'), 12.0 (C-9), 18.1 (C-10), 29.70, 29.74, 37.8, 49.4, 55.03, 55.10, 60.64, 60.69, 66.0, 74.73, 74.80, 124.01, 124.07, 126.3, 127.1, 127.3, 128.0, 128.22, 128.26, 130.4, 130.5, 132.6, 139.15, 139.18, 142.87, 142.91.

HRMS found 893.3282. $\text{C}_{47}\text{H}_{67}\text{Br}_2\text{N}_2\text{O}_3\text{Si}$ (MH^+) requires 893.3287.

ii. Double Mitsunobu reaction:

δ_c (125MHz, $CDCl_3$): 10.0, 10.2, 12.0, 18.2, 30.5, 37.3, 48.5, 48.9, 54.82, 54.87, 61.56, 61.62, 65.6, 81.4, 81.6, 112.43, 112.45, 114.2, 121.48, 121.50, 123.79, 123.83, 126.43, 126.48, 127.2, 127.4, 127.86, 127.89, 128.2, 130.25, 130.29, 132.32, 132.34, 133.3, 139.8, 140.1, 140.15, 153.92, 153.96.

m/z (ESI): 603 ($1/2 \times MH_2^{2+}$:100%).

iii. Bis-(biaryl) (15):

Cyclization precursor (**14**) (151.1mg, 0.1254 mmol) in a flame-dried 50mL r.b. flask equipped with a magnetic stirrer bar was azeotroped with benzene (3 x 10mL) and dried in vacuo at 50°C for 3h, then purged with argon. 10mL of anhydrous 2-MeTHF was then added to generate a clear solution which was then cooled to -78°C (substrate concentration 12.5mM). A solution of $t\text{-BuLi}$ in pentane (1.68M, 0.6mL, 1.00 mmol, 8.0 equiv.) was added and the resultant pale yellow solution stirred at -78°C for 30 min. A solution of $\text{CuCN} \cdot 2\text{LiBr}$ (1.0M in CuCN , 2.0M in LiBr , in anhydrous 2-MeTHF; 0.4mL, 4.0 equiv. CuCN) was then added and the solution allowed to warm to -40°C over 2h. A solution of $m\text{-DNB}$ (1.0M in anhydrous 2-MeTHF; 1.1mL, 10.0 equiv.) was then added and the dark brown solution allowed to warm to room temperature. After 12h, the reaction mixture was poured into 10% NH_4OH /saturated aq. NH_4Cl (30 mL) and the phases separated. The aqueous phase was extracted with EtOAc (3 x 30mL) and the

extracts and the organic phase combined and washed brine (50mL); then dried over K_2CO_3 before concentration under reduced pressure. The dark brown oil obtained was chromatographed over SiO_2 (Benzene/Hexanes (1:1 to 7:1) to remove excess oxidant and oxidant by-products, followed by elution with Hexanes/EtOAc/TEA (100:0:1 to 90:10:1)) to yield bis-biaryl (**15**) as a colorless oil, 65.4mg (59% yield, 77% yield per cyclization).

R_f = 0.38 (Triethylamine washed SiO_2 , Hexanes/EtOAc/ Et_3N : 95:5:1).

IR (film)/ cm^{-1} : 2940s, 2864s, 1495s, 1480s, 1440s, 1372s, 1257s, 1225s, 752s, 738s.

δ_H (500MHz, $CDCl_3$): 0.69 (app. t, J 8Hz, 6H, C(14) $\underline{H}_3$ and C(14') $\underline{H}_3$), 1.01-1.13 (m, 21H, C(19) $\underline{H}$ and C(20) $\underline{H}_3$), 1.63 (app. septet, J 7Hz, 2H, C(16) $\underline{H}_2$), 1.77 (app. septet, J 7Hz, 2H, C(16') $\underline{H}_2$), 1.95 (app. quintet, J 6Hz, 1H, C(17) $\underline{H}$), 2.86 (m, 4H, C(15) $\underline{H}_2$ and C(15') $\underline{H}_2$), 3.12 (app. septet, J 7Hz, 2H, C(7) $\underline{H}$ and C(7') $\underline{H}$), 3.44 (d, J 13.5Hz, 1H, C(9) $\underline{H}H'$), 3.49 (d, J 13.5Hz, 1H, C(9') $\underline{H}H'$), 3.74 (m, 4H, C(9) $\underline{H}H'$, C(9') $\underline{H}H'$ and C(18) $\underline{H}_2$), 5.20 (s, 1H, C(6) $\underline{H}$), 5.22 (s, 1H, C(6') $\underline{H}$), 6.77 (m, 2H, Ar- $\underline{H}$), 7.13 (app. q, J 7Hz, 2H, Ar- $\underline{H}$), 7.19-7.37 (m, 20H, Ar- $\underline{H}$), 7.64 (d, J 7.5Hz, 1H, Ar- $\underline{H}$), 7.67 (d, J 7.5Hz, 1H, Ar- $\underline{H}$).

δ_c (125MHz, $CDCl_3$): 12.1 (C-19), 14.6 and 14.7 (C-14 and C-14'), 18.2 (C-20), 30.35 and 30.43 (C-16 and C-16'), 36.9 (C-17), 55.1 and 55.3 (C-9 and C-9'), 56.4 and 56.7 (C-15 and C-15'), 66.0 (C-18), 67.0 and 67.4 (C-7 and C-7'), 84.4 (C-6 and C-6'), 118.8,

123.7, 125.56, 125.60, 126.86, 126.90, 126.95, 126.98, 127.01, 127.04, 128.28, 128.32, 128.39, 129.02, 129.04, 129.23, 129.26, 129.8, 130.45, 130.49, 134.1, 138.0, 139.30, 139.36, 142.8, 156.9.

m/z (ESI): (MH⁺:60%).

HRMS found 885.5391. C₅₉H₇₃N₂O₃Si (MH⁺) requires 885.5390.

Fig. 10. nOe's observed for bis-biaryl **15** (NOESY, 500MHz). Red arrows indicate a transannular nOe characteristic of (M, 6R, 7S) stereochemistry typically observed in analogous monomeric biaryl medium rings.

III-28

exp2 s2pu1

```

SAMPLE
date Jan 23 2004 dfrq 500.211 DEC. & VT
solvent CDC13 dn H1
file exp dpwr 32
ACQUISITION
sfrq 500.211 dm nnn
tn H1 dmm C
at 2.048 dmf 8770
np 32768 dseq
sw 8000.0 dres 1.0
fb 4000 homo n
bs 8 temp 22.0
ss 2
tpwr 58 lb PROCESSING 0.10
pw 5.0 wtfile
d1 0 proc ft
tof 0 fn not used
nt 32 math f
ct 32
alock n werr
gain not used wexp
FLAGS vbs
il n wnt wft
in n
dp y
hs nn
DISPLAY
sp -250.5
wp 5502.0
vs 28
sc 0
wc 250
hzmm 22.01
is 197.88
rfl 1500.0
rfp 0
th 1
ins 1.000
ai cdc ph

```

III-28

exp1 std13c

```

SAMPLE
date Jan 24 2004 dfrq 400.046 DEC. & VT
solvent CDC13 dn H1
file exp dpwr 46
ACQUISITION
sfrq 100.601 dm yyy
tn C13 dmm V
at 1.280 dmf 11000
np 64000 temp 22.0
sw 25000.0 lb PROCESSING 1.00
fb 13800 wtfile
bs 16 pw 61 proc ft
tpwr 2.0 fn not used
d1 1.000
tof 0 werr
nt 15000 wexp
ct 14736 wbs
alock n wnt
gain not used
FLAGS
il n
in n
dp y
DISPLAY
sp -0.7
wp 20117.8
vs 743
sc 0
wc 250
hzmm 80.47
is 500.00
rfl 10746.3
rfp 7752.6
th 8
ins 100.000
nm ph

```

III-179-2

exp4 std1h

SAMPLE DEC. & VT
 date Apr 22 2004 dfrq 400.046
 solvent Benzene dn H1
 file exp dpwr 40
 ACQUISITION dof 0
 sfrq 400.046 dm nnn
 tn H1 dmm c
 at 2.733 dmf 7500
 np 32768 temp 20.0
 sw 5995.2
 fb not used lb 0.10
 bs 8 wtfile
 ss 2 proc ft
 tpwr 60 fn not used
 pw 10.0
 d1 10.000 werr
 tof 0 wexp
 nt 32 wbs
 ct 32 wnt
 alock n
 gain not used
 FLAGS
 il n
 in n
 dp y
 DISPLAY
 sp -200.2
 wp 4400.4
 vs 82
 sc 0
 wc 250
 hzmm 17.60
 is 445.87
 rfl 959.9
 rfp 0
 th 2
 ins
 ai cdc ph 4.000

III-179-2

exp5 std13c

SAMPLE DEC. & VT
 date Apr 22 2004 dfrq 400.046
 solvent Benzene dn H1
 file exp dpwr 46
 ACQUISITION dof 0
 sfrq 100.601 dm yyy
 tn C13 dmm w
 at 1.280 dmf 11000
 np 64000 temp 22.0
 sw 25000.0
 fb 13800 lb 1.00
 bs 16 wtfile
 tpwr 61 proc ft
 pw 2.0 fn not used
 d1 1.000
 tof 0 werr
 nt 3000 wexp
 ct 720 wbs
 alock n
 gain not used
 FLAGS
 il n
 in n
 dp y
 DISPLAY
 sp -0.1
 wp 20117.6
 vs 1571
 sc 0
 wc 250
 hzmm 80.47
 is 500.00
 rfl 15841.4
 rfp 12875.7
 th 8
 ins
 nm ph 100.000

III-162-3

exp1 std1h

SAMPLE DEC. & VT
 date Apr 2 2004 dfrq 400.048
 solvent CD300 dn H1
 file exp dpwr 40
 ACQUISITION dof 0
 sfrq 400.048 dm nnn
 tn H1 dmm c
 at 2.733 dmf 7500
 np 32768 temp 20.0
 sw 5995.2
 fb not used lb PROCESSING
 bs 8 wtfile 0.10
 ss 2 proc ft
 tpwr 60 fn not used
 pw 10.0
 d1 0.120 werr
 tof 0 wexp
 nt 32 wbs
 ct 32 wnt
 alock n
 gain not used
 FLAGS
 il n
 in n
 dp y
 DISPLAY
 sp -200.3
 wp 4400.4
 vs 117
 sc 0
 wc 250
 hzmm 17.60
 ls 1730.15
 rf1 2298.5
 rfp 1320.2
 th 1
 ins 2.000
 al cdc ph

III-162-3

exp2 std13c

SAMPLE DEC. & VT
 date Apr 2 2004 dfrq 400.048
 solvent CD300 dn H1
 file exp dpwr 40
 ACQUISITION dof 0
 sfrq 100.601 dm yyy
 tn C13 dmm w
 at 1.280 dmf 11000
 np 64000 temp 22.0
 sw 25000.0
 fb 13800 lb PROCESSING
 bs 16 wtfile 1.00
 tpwr 61 proc ft
 pw 2.0 fn not used
 d1 0 werr
 tof 0 wexp
 nt 10000 wbs
 ct 688 wnt
 alock n
 gain not used
 FLAGS
 il n
 in n
 dp y
 DISPLAY
 sp -0.5
 wp 20117.8
 vs 166
 sc 0
 wc 250
 hzmm 80.47
 ls 500.00
 rf1 7786.0
 rfp 4929.0
 th 3
 ins 100.000
 nm ph

III-182-2

exp3 s2pu1

SAMPLE DEC. & VT
 date Apr 28 2004 dfrq 499.752
 solvent CD300 dn H1
 file exp dpwr 34
 ACQUISITION dof 0
 sfrq 499.752 dm nnn
 tn H1 dmm c
 at 2.048 dmf 10214
 np 32768 dseq
 sw 8000.0 dres 1.0
 fb 4000 homo n
 bs 8 temp 23.0
 ss 2 PROCESSING
 tpwr 54 lb 0.10
 pw 6.0 wtfile
 d1 10.000 proc ft
 tof 0 fn not used
 nt 8 math f
 ct 8
 alock n werr
 gain not used wexp
 FLAGS n wbs
 il n wnt wft
 in n
 dp y
 hs nn
 DISPLAY
 sp -250.2
 wp 5497.1
 vs 54
 sc 0
 wc 250
 hzmm 21.99
 is 1186.55
 rfl 3164.6
 rfp 1649.2
 th 2
 ins 2.000
 ai ph

III-182-2

exp4 s2pu1

SAMPLE DEC. & VT
 date Apr 28 2004 dfrq 499.752
 solvent CD300 dn H1
 file exp dpwr 34
 ACQUISITION dof 0
 sfrq 125.674 dm yyy
 tn C13 dmm w
 at 1.092 dmf 10000
 np 65536 dseq
 sw 29996.3 dres 1.0
 fb 17000 homo n
 bs 16 temp 23.0
 tpwr 57 PROCESSING
 pw 6.0 lb 1.00
 d1 1.000 wtfile
 tof 0 proc ft
 nt 4000 fn not used
 ct 1248 math f
 alock n werr
 gain 50 wexp
 FLAGS n wbs
 il n wnt
 in n
 dp y
 hs nn
 DISPLAY
 sp -0.5
 wp 25131.7
 vs 69
 sc 0
 wc 250
 hzmm 100.53
 is 500.00
 rfl 9075.4
 rfp 6157.4
 th 5
 ins 100.000
 nm cdc ph

III-168-2

exp1 s2pu1

SAMPLE DEC. & VT
 date Apr 6 2004 dfrq 499.752
 solvent CD300 dn H1
 file exp dpwr 34
 ACQUISITION dof 0
 sfrq 499.752 dm nnn
 tn H1 dmm c
 at 2.048 dmf 10214
 np 32768 dseq 1.0
 sw 8000.0 dres n
 fb 4000 homo n
 bs 8 temp 23.0
 ss 2
 tpwr 54 lb PROCESSING 0.10
 pw 6.0 wtfile
 d1 0.200 proc ft
 tof 0 fn not used f
 nt 16 math
 ct 16
 alock n werr
 gain not used wexp
 FLAGS wbs
 il n wnt wft
 in n
 dp y
 hs nn
 DISPLAY
 sp -250.2
 wp 5497.1
 vs 49
 sc 0
 wc 250
 hzmm 21.99
 is 238.09
 rfl 3165.0
 rfp 1649.2
 th 2
 ins 3.000
 al ph

III-168-2

exp2 s2pu1

SAMPLE DEC. & VT
 date Apr 6 2004 dfrq 499.752
 solvent CD300 dn H1
 file exp dpwr 34
 ACQUISITION dof 0
 sfrq 125.674 dm yyy
 tn C13 dmm w
 at 1.092 dmf 10000
 np 65536 dseq 1.0
 sw 29996.3 dres n
 fb 17000 homo n
 bs 16 temp 23.0
 tpwr 57
 pw 6.0 lb PROCESSING 1.00
 d1 0 wtfile
 tof 0 proc ft
 nt 5000 fn not used f
 ct 1040 math
 alock n werr
 gain 50 wexp
 FLAGS n wbs
 il n wnt
 in n
 dp y
 hs nn
 DISPLAY
 sp -0.5
 wp 25131.7
 vs 112
 sc 0
 wc 250
 hzmm 100.53
 is 500.00
 rfl 9072.7
 rfp 6157.4
 th 4
 ins 100.000
 nm cdc ph

SAMPLE DEC. & VT
 date Apr 28 2004 dfrq 499.752
 solvent CD3OD dn H1
 file exp dpwr 34
 ACQUISITION dof 0
 sfrq 499.752 dm nnn
 tn H1 dmm c
 at 2.048 dmf 10214
 np 32768 dseq
 sw 8000.0 dres 1.0
 fb 4000 homo n
 bs 8 temp 23.0
 ss 2
 tpwr 54 lb PROCESSING 0.10
 pw 6.0 wtfile
 d1 10.000 proc ft
 tof 0 fn not used
 nt 8 math f
 ct 8
 alock n werr
 gain not used wexp
 FLAGS wbs
 il n wnt wft
 in n
 dp y
 hs nn
 DISPLAY
 sp -250.2
 wp 5497.1
 vs 34
 sc 0
 wc 250
 hzmm 21.99
 ls 271.24
 rfl 3164.6
 rfp 1649.2
 th 1
 ins 1.000
 al ph

SAMPLE DEC. & VT
 date Apr 28 2004 dfrq 499.752
 solvent CD3OD dn H1
 file exp dpwr 34
 ACQUISITION dof 0
 sfrq 125.674 dm yyy
 tn C13 dmm w
 at 1.032 dmf 10000
 np 65536 dseq
 sw 29996.3 dres 1.0
 fb 17000 homo n
 bs 16 temp 23.0
 tpwr 57
 pw 6.0 lb PROCESSING 1.00
 d1 1.000 wtfile
 tof 0 proc ft
 nt 4000 fn not used
 ct 272 math f
 alock n werr
 gain 50 wexp
 FLAGS n wbs
 il n wnt
 in n
 dp y
 hs nn
 DISPLAY
 sp -2923.5
 wp 29995.3
 vs 89
 sc 0
 wc 250
 hzmm 119.99
 ls 500.00
 rfl 9081.8
 rfp 6157.4
 th 5
 ins 100.000
 nm cdc ph

11

STANDARD PROTON PARAMETERS

Pulse Sequence: gCOSH
 Solvent: CDCl₃
 Temp. 22.0 C / 295.1 K
 INOVA-500 "Inova500b"

Relax. delay 1.000 sec
 Acq. time 0.128 sec
 Width 8000.0 Hz
 2D Width 8000.0 Hz
 4 repetitions
 128 increments
 OBSERVE H1, 500.2085438 MHz
 DATA PROCESSING
 Sq. sine bell 0.064 sec
 F1 DATA PROCESSING
 Sq. sine bell 0.016 sec
 FT size 2048 x 2048
 Total time 10 min, 6 sec

15

STANDARD PROTON PARAMETERS

Pulse Sequence: gHMOC
 Solvent: CDCl₃
 Temp. 22.0 C / 295.1 K
 User: 1-14-87
 INOVA-500 "Inova500b"

Relax. delay 1.000 sec
 Acq. time 0.128 sec
 Width 8000.0 Hz
 2D Width 21384.7 Hz
 22 repetitions
 2 x 128 increments
 OBSERVE H1, 500.2085433 MHz
 DECOUPLE C13, 125.7869397 MHz
 Power 45 dB
 on during acquisition
 off during delay
 GARP-1 modulated
 DATA PROCESSING
 Gauss apodization 0.059 sec
 F1 DATA PROCESSING
 Gauss apodization 0.011 sec
 FT size 2048 x 2048
 Total time 1 hr, 51 min, 35 sec

STANDARD PROTON PARAMETERS

Data Collected on:
nmrsun2-inova500
Archive directory:
/export/home/ds2/vnmrsys/data
Sample directory:

File: II-184-2-NOESY

Pulse Sequence: NOESY

Solvent: CDCl₃

Temp. 22.0 C / 295.1 K

Relax. delay 1.000 sec

Mixing 0.800 sec

Acq. time 0.128 sec

Width 8000.0 Hz

2D Width 8000.0 Hz

42 repetitions

2 x 200 increments

OBSERVE H1, 500.2085385 MHz

DATA PROCESSING

Gauss apodization 0.059 sec

F1 DATA PROCESSING

Gauss apodization 0.023 sec

FT size 2048 x 2048

Total time 9 hr, 5 min

STANDARD PROTON PARAMETERS

Pulse Sequence: NOESY

Solvent: CDCl₃

Temp. 22.0 C / 295.1 K

INOVA-500 "inova500b"

Relax. delay 1.000 sec

Mixing 0.800 sec

Acq. time 0.128 sec

Width 8000.0 Hz

2D Width 8000.0 Hz

42 repetitions

2 x 200 increments

OBSERVE H1, 500.2085385 MHz

DATA PROCESSING

Gauss apodization 0.059 sec

F1 DATA PROCESSING

Gauss apodization 0.023 sec

FT size 2048 x 2048

Total time 9 hr, 8 min, 12 sec