Synthesis, characterization of 1, 2 – dimethyl imidazolium type of ionic liquids and its catalytic activities

Pandurangan Ganapathi and Kilivelu Ganesan*

PG & Research Department of Chemistry, Presidency College (Autonomous), Chennai-600 005, India

E-mail: kiliveluganesan@yahoo.co.in

Supporting information:

General procedure for alkylation: 1, 2- dimethylimidazole (1.040×10^{-2} mol/L; 1.0 equi.) is mixed with benzyl bromide (1.092×10^{-2} mol/L; 1.05 equi.) / 4-nitrobenzyl bromide (1.040×10^{-2} mol/L; 1.05 equi.) in presence of 30 mL of dry CH_3CN under refluxing condition for about 2-2.5 hours afforded the *N*-alkylated quaternary ammonium bromide **1a** & **1b** in 95-97% yield.

1,2-dimethyl(3-methylene benzene)-imidazolium bromide 1a: 2.75g; 95% Semisolid; ^1H NMR (D_2O); δ : 2.45(s, 3H); 3.66(s, 3H); 5.22(s, 2H); 7.20(d, $J=1.8$ Hz, 2H) 7.24 (s, 2H) 7.33(t, $J=7.2$ Hz, 3H); ^{13}C NMR δ : 9.19, 34.75, 51.42, 121.10, 122.37, 127.76, 129.26, 133.73, 144.58; MS:267; Elemental analysis:- Molecular Formula ($\text{C}_{12}\text{H}_{15}\text{N}_2\text{Br}$); Calculated: C: 53.93; H: 5.61; N: 10.48; Found C: 53.86; H: 5.52; N: 10.40.

1,2-dimethyl(3-methylene 4'-nitro benzene)-imidazolium bromide 1b: 3.25g; 97%; MP : 268-270 °C. ^1H NMR ($\text{DMSO}-d_6$); δ : 2.58(s, 3H); 3.78(s, 3H); 5.61(s, 2H); 7.57(d, $J=1.8$ Hz, 2H); 7.73 (d, $J=2.0$ Hz, 2H); 8.24(d, $J=1.8$ Hz, 2H); ^{13}C NMR δ :10.0, 35.44, 50.28, 121.79, 123.38, 124.44, 129.30, 142.40, 145.56, 147.83; MS:312; Elemental analysis: - Molecular Formula ($\text{C}_{12}\text{H}_{14}\text{N}_3\text{Br}$); Calculated: C: 46.15; H: 4.48; N: 13.46; Found C: 46.06; H: 4.40; N: 13.38.

General procedure for metathesis reaction: *N*-Alkylated quaternary ammonium bromide (1.0 equi.) is treated with different anion containing inorganic salts such as NaBF_4 , KPF_6 , LiCF_3SO_3 (1.05 equi.) in the presence of 10 mL deionized water at room temperature for 1 hour afforded the anion exchanged ionic liquids. After the metathesis reaction followed by Soxhlet extraction with dry THF for about 1 hour to give metallic bromide free ionic liquids (**2a-f**) with 90-93% yield.

1,2-dimethyl(3-methylene benzene)-imidazolium tetrafluoroborate 2a: 0.67g; 93%; Semisolid; ^1H NMR (D_2O); δ : 2.43(s, 3H); 3.68(s, 3H); 5.26(s, 2H); 7.18(d, $J=1.9$ Hz, 2H) 7.21 (s, 2H) 7.4 (t, $J=7.4$ Hz, 3H); ^{13}C NMR δ : 9.17, 34.78, 51.44, 121.08, 122.35, 127.74, 129.27, 133.71, 144.55; MS:274; Elemental analysis:- Molecular Formula ($\text{C}_{12}\text{H}_{15}\text{BF}_4\text{N}_2$); Calculated: C: 52.55; H: 5.47; N: 10.22; Found C: 52.48; H: 5.32; N: 10.14.

1,2-dimethyl(3-methylene benzene)-imidazolium hexafluorophosphate 2b: 0.76g; 90%; Semisolid; ^1H NMR (D_2O); δ : 2.46(s, 3H); 3.65(s, 3H); 5.20(s, 2H); 7.19(d, $J=1.9$ Hz, 2H), 7.23 (s, 2H) 7.35(t, $J=7.6$ Hz, 3H); ^{13}C NMR δ : 9.16, 34.70, 51.35, 121.02, 122.31, 127.72, 129.20, 133.68, 144.53; MS:332; Elemental analysis: Molecular Formula ($\text{C}_{12}\text{H}_{15}\text{PF}_6\text{N}_2$); Calculated: C: 43.38; H: 4.51; N: 8.41; Found C: 43.26; H: 4.40; N: 8.32.

1,2-dimethyl(3-methylene benzene)-imidazolium trifluoromethanesulfonate 2c: 0.72g; 90%; Semisolid. ^1H NMR (D_2O); δ : 2.45(s, 3H); 3.65(s, 3H); 5.24(s, 2H); 7.23(d, $J=1.8$ Hz, 2H) 7.29 (s, 2H) 7.37 (t, $J=7.5$ Hz, 3H); ^{13}C NMR δ : 9.19, 34.77, 51.45, 121.17, 122.30, 127.74, 129.27, 133.75, 144.60; MS:336; Elemental analysis:- Molecular Formula ($\text{C}_{13}\text{H}_{15}\text{F}_3\text{O}_3\text{SN}_2$); Calculated: C: 46.42; H: 4.46; N: 8.33; Found C: 46.37; H: 4.34; N: 8.28.

1,2-dimethyl(3-methylene 4'-nitro benzene)-imidazolium tetrafluoroborate 2d: 0.63g; 91%; MP : 85-87 °C. ^1H NMR ($\text{DMSO}-d_6$); δ : 2.54(s, 3H); 3.74(s, 3H); 5.58(s, 2H); 7.55 (d, $J=1.8$ Hz, 2H); 7.70(d, $J=2.0$ Hz, 2H); 8.21(d, $J=1.8$ Hz, 2H); ^{13}C NMR δ :10.1, 35.40, 50.26, 121.78, 123.35, 124.42, 129.32, 142.38, 145.56, 147.81; MS:319; Elemental analysis:- Molecular Formula ($\text{C}_{12}\text{H}_{14}\text{BF}_4\text{N}_3\text{O}_2$); Calculated: C: 45.14; H: 4.38; N: 13.16; Found C: 45.06; H: 4.26; N: 13.10.

1,2-dimethyl(3-methylene 4'-nitro benzene)-imidazolium hexafluorophosphate 2e: 0.74g; 91%; MP : 175-178 °C. ^1H NMR ($\text{DMSO}-d_6$); δ : 2.56(s, 3H); 3.75(s, 3H); 5.63(s, 2H); 7.59 (d, $J=1.8$ Hz, 2H); 7.74 (d, $J=2.0$ Hz, 2H); 8.25(d, $J=1.8$ Hz, 2H); ^{13}C NMR δ :10.12, 35.45, 50.29, 121.76, 123.33, 124.41, 129.33, 142.38, 145.53, 147.80;

MS:377; Elemental analysis:- Molecular Formula ($C_{12}H_{14}PF_6N_3O_2$); Calculated: C: 38.19; H: 3.71; N: 11.14; Found C: 38.17; H: 3.64; N: 11.06.

1,2-dimethyl(3-methylene 4'-nitro benzene)-imidazolium trifluoromethanesulfonate 2f: 0.70g; 91%; Semisolid. 1H NMR ($DMSO-d_6$); δ : 2.58(s, 3H); 3.70(s, 3H); 5.57(s, 2H); 7.53(d, $J=1.8$ Hz, 2H); 7.74 (d, $J=2.0$ Hz, 2H); 8.23 (d, $J=1.8$ Hz, 2H); ^{13}C NMR δ :10.0, 35.41, 50.23, 121.75, 123.34, 124.47, 129.39, 142.42, 145.59, 147.85; MS:381; Elemental analysis:- Molecular Formula ($C_{13}H_{14}F_3O_5SN_3$); Calculated: C: 40.94; H: 3.67; N: 11.02; Found C: 40.86; H: 3.54; N: 10.94.

General procedure for Mannich reaction: Aniline / 4- nitroaniline (1.0 equi; 1.387×10^{-3} mol/L), α - naphthol (1.0 equi; 1.387×10^{-3} mol/L) and paraformaldehyde (2.05 equi; 2.843×10^{-3} mol/L) are mixed with optimized catalytic amount of ionic liquid at room temperature afforded the Mannich product of oxazine derivatives in 85 - 90 % yield.

3,4-dihydro-3-(3-nitrophenyl)-2H-naphtho[2,1-e][1,3] oxazine 9: 1H NMR ($DMSO-d_6$); δ :4.73(s, 2H); 5.45(s, 2H); 6.98-7.95(m,10H); ^{13}C NMR; δ : 49.1, 79.4, 112.7,115.3, 117.5, 119.8, 120.7, 124.2, 125.2, 125.5, 126.0, 127.6, 129.2, 132.9, 147.9, 148.8; MS:306; Elemental analysis:- Molecular Formula ($C_{18}H_{14}N_2O_3$); Calculated: C: 70.58; H: 4.57; N: 9.15; Found C: 70.40; H: 4.44; N: 9.10.

3,4-dihydro-3-(4-nitrophenyl)-2H-naphtho[2,1-e][1,3] oxazine 10: 1H NMR ($DMSO-d_6$); δ :4.73(s, 2H); 5.46(s, 2H); 6.96-7.98(m,10H); ^{13}C NMR; δ : 49.1, 79.4, 112.7, 115.3, 117.5, 119.8, 120.7, 124.2, 125.2, 125.5, 126.0, 127.6, 129.2, 132.9, 147.9, 148.8; MS:306; Elemental analysis:- Molecular Formula ($C_{18}H_{14}N_2O_3$); Calculated: C: 70.58; H: 4.57; N: 9.15; Found C: 70.36; H: 4.39; N: 9.08.

1H NMR spectrum of 1,2-dimethyl(3-methylene benzene)-imidazolium bromide 1a:

Current Data Parameters
NAME DPG-3
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20140626
Time 17.00
INSTRUM spect
PROBHD 5 mm PABBO BB/
PULPROG zg30
TD 65536
SOLVENT D2O
NS 16
DS 2
SWH 8012.820 Hz
FIDRES 0.122266 Hz
AQ 4.0894465 sec
RG 11.24
DW 62.400 usec
DE 6.50 usec
TE 295.7 K
D1 1.00000000 sec
TD0 1

----- CHANNEL f1 -----
SFO1 400.2324716 MHz
NUC1 1H
P1 14.50 usec
PLW1 12.00000000 W

F2 - Processing parameters
SI 65536
SF 400.2300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

C^{13} NMR spectrum of 1,2-dimethyl(3-methylene benzene)-imidazolium bromide 1a:

Current Data Parameters

NAME DPG-3
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters

Date_ 20140626
Time 17.30
INSTRUM spect
PROBHD 5 mm PABBO BB/
PULPROG zgpg30
TD 65536
SOLVENT D2O
NS 512
DS 4
SWH 24038.461 Hz
FIDRES 0.366798 Hz
AQ 1.3631488 sec
RG 204.46
DW 20.800 usec
DE 6.50 usec
TE 296.4 K
D1 2.00000000 sec
D11 0.03000000 sec
TD0 1

===== CHANNEL f1 =====

SFO1 100.6479769 MHz
NUC1 13C
P1 9.50 usec
PLW1 54.00000000 W

===== CHANNEL f2 =====

SFO2 400.2316009 MHz
NUC2 1H
CPDPRG[2] waltz16
PCPD2 90.00 usec
PLW2 12.00000000 W
PLW12 0.31147999 W
PLW13 0.25229999 W

F2 - Processing parameters

SI 32768
SF 100.6379135 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

¹H NMR spectrum of 1,2-dimethyl(3-methylene 4'-nitro benzene)-imidazolium bromide 1b:

Current Data Parameters
NAME DPG-6
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20140626
Time 17.34
INSTRUM spect
PROBHD 5 mm PABBO BB/
PULPROG zg30
TD 65536
SOLVENT DMSO
NS 16
DS 2
SWH 8012.820 Hz
FIDRES 0.122266 Hz
AQ 4.0894465 sec
RG 31.36
DW 62.400 usec
DE 6.50 usec
TE 295.6 K
D1 1.00000000 sec
TD0 1

----- CHANNEL f1 -----
SFO1 400.2324716 MHz
NUC1 1H
P1 14.50 usec
PLW1 12.00000000 W

F2 - Processing parameters
SI 65536
SF 400.2300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00

¹³C NMR spectrum of 1,2-dimethyl(3-methylene 4'-nitro benzene)-imidazolium bromide 1b:

Current Data Parameters
NAME DPG-6
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20140626
Time 18.04
INSTRUM spect
PROBHD 5 mm PABBO BB/
PULPROG zgpg30
TD 65536
SOLVENT DMSO
NS 512
DS 4
SWH 24038.461 Hz
FIDRES 0.366798 Hz
AQ 1.3631488 sec
RG 204.46
DM 20.800 usec
DE 6.50 usec
TE 296.5 K
D1 2.00000000 sec
D11 0.03000000 sec
TD0 1

===== CHANNEL f1 =====
SFO1 100.6479769 MHz
NUC1 13C
P1 9.50 usec
PLW1 54.00000000 W

===== CHANNEL f2 =====
SFO2 400.2316009 MHz
NUC2 1H
CPDPRG2 waltz16
PCPD2 90.00 usec
PLW2 12.00000000 W
PLW12 0.31147999 W
PLW13 0.25229999 W

F2 - Processing parameters
SI 32768
SF 100.6379135 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40