Note: the version of SI posted on July 16, 2004, contained errors. The corrected version was posted August 10, 2004.

Synthesis and Characterization of a Series of Homo-oligopeptide Peroxyesters

Fernando Formaggio,^{*,†,‡} Marco Crisma,[‡] Laura Scipionato,[†]
Sabrina Antonello,[†] Flavio Maran,[†] and Claudio Toniolo^{†,‡}

Department of Chemistry, University of Padova, 35131 Padova, Italy and Institute of Biomolecular Science, CNR, Padova Unit, 35131 Padova, Italy

fernando.formaggio@unipd.it

SUPPORTING INFORMATION

Contents

Synthesis and characterization of	
Pht-Aib-OO'Bu	S-2
PhAc- <i>cis</i> -1,4-Chx-OO'Bu	S-2
PhAc- <i>trans-rac</i> -1,2-Chx-OO'Bu	S-3
Pht- <i>trans-rac</i> -1,2-Chx-OO'Bu	S-3
Pht-Aib- <i>trans-rac</i> -1,2-Chx-OO'Bu	S-4
Pht-(Aib) ₂ - <i>trans-rac</i> -1,2-Chx-OO'Bu	S-4
Pht-(Aib) ₃ - <i>trans-rac</i> -1,2-Chx-OO'Bu	S-5
Pht-(Aib) ₄ - <i>trans-rac</i> -1,2-Chx-OO'Bu	S-5
Pht-(Aib) ₅ - <i>trans-rac</i> -1,2-Chx-OO'Bu	S-6
Pht-(Aib) ₆ - <i>trans-rac</i> -1,2-Chx-OO'Bu	S-6
Figure S1: FT-IR absorption spectra of the Pht-(Aib) _n - <i>trans-rac</i> -1,2-Chx-OO'Bu (<i>n</i> = 1-6) peptide series in CDCl ₃ solution	S-7
Figure S2: X-Ray diffraction structure of PhAc- <i>trans-rac</i> -1,2-Chx-OO'Bu (molecule B)	S-8
Table S1. Crystal data and structure refinement for PhAc- <i>cis</i> -1,4-Chx-OO'Bu	S-9
Table S2. Crystal data and structure refinement for PhAc- <i>trans-rac</i> -1,2-Chx-OO'Bu	S-10

Pht-Aib-OO'Bu*¹

To a solution of Pht-Aib-Cl (0.30 g, 1.2 mmol) in anhydrous CH₂Cl₂, at 0 °C, a 5.5 M solution of HOO'Bu in decane (0.23 ml, 1.2 mmol) and DMAP (0.29 g, 2.4 mmol) were added. The reaction was stirred at rt for 1 d. Then, the solvent was removed *in vacuo* and the residue dissolved in EtOAc. The solution was washed with 10% KHSO₄, H₂O, 5% NaHCO₃ and H₂O, dried over Na₂SO₄ and evaporated to dryness. The product was purified by flash chromatography.

Yield 68%; oil.

Rf₁ (CHCl₃/EtOH 9:1) 0.95, **Rf₂** (1-ButOH/AcOH/H₂O 3:1:1) 0.95, **Rf₃** (toluene/EtOH 7:1) 0.85, **Rf₄** (EtOAc/petroleum ether 1:3) 0.60.

IR (KBr): 1780, 1719, 1612 cm⁻¹.

¹H NMR (CDCl₃, 250 MHz): δ 7.84 and 7.71 (2m, 4H, Pht CH); 1.95 (s, 6H, 2 βCH₃); 1.35 (s, 9H, OO'Bu).

¹³C NMR (CD₃CN, 50 MHz): δ 170.95, 169.11, 135.56, 132.64, 123.95, 85.02, 61.10, 26.32, 25.12.

MS (ESI-TOF): [M+Na]⁺_{calc.} = 328.13; [M+Na]⁺_{sper.} = 328.12.

PhAc-*cis*-1,4-Chx-OO'Bu

To a solution of PhAc-*cis*-1,4-Chx-OH (0.2 g, 0.76 mmol) in anhydrous CH₂Cl₂, EDC·HCl (0.29 g, 1.52 mmol) was added, followed after a few min by a 5.5 M solution of HOO'Bu in decane (0.69 ml, 3.8 mmol) and DMAP (0.37 g, 3.04 mmol). The reaction was refluxed for 2 h. Then, the solvent was removed *in vacuo* and the residue was applied directly to a flash chromatography column (eluant: EtOAc/PE 1:1).

Yield 65 %; **m. p.** 135-136 °C.

Rf₁ 0.95, **Rf₃** 0.85, **Rf₄** 0.60.

IR (KBr): 3302, 1772, 1636, 1551 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): δ 7.39-7.22 (m, 5H, PhAc CH); 5.47 (d, 1H, NH); 3.95-3.90 (m, 1H, βCH); 3.55 (d, 2H, PhAc CH₂); 2.52-2.50 (m, 1H, αCH); 1.77-1.41 (m, 8H, Chx CH₂); 1.30 (s, 9H, OO'Bu).

¹³C NMR (CDCl₃, 50 MHz): δ 171.98, 170.19, 134.9, 129.22, 129.00, 127.35, 83.35, 45.65, 43.89, 37.89, 28.85, 26.08, 25.01.

MS (ESI-TOF): [M+H]⁺_{calc.} = 334.20 [M+H]⁺_{sper.} = 334.20; [2M+H]⁺_{calc.} = 667.40, [2M+H]⁺_{sper.} = 667.37.

* All derivatives and peptides described in this work are based on a *racemic* β-amino acid.

¹ Rüchardt, C.; Hamprecht, G. *Chem. Ber.* **1968**, *101*, 3957-3962.

PhAc-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from PhAc-*trans-rac*-1,2-Chx-OH (0.16 g, 0.61 mmol), EDC·HCl (0.23 g, 1.22 mmol), HOO'Bu (0.55 ml of a 5.5 M solution in decane, 3.0 mmol) and DMAP (0.30 g, 2.44 mmol). Eluant for flash chromatography purification: EtOAc/PE 2:3.

Yield 94 %; **m. p.** 125-126 °C.

R_f1 0.95, **R_f3** 0.85, **R_f4** 0.60.

IR (KBr): 3269, 1768, 1647, 1557 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): 7.35-7.19 (m, 5H, PhAc CH); 5.35 (d, 1H, NH); 4.55-3.87 (m, 1H, βCH); 3.52 (s, 2H, PhAc CH₂); 2.45-2.27 (m, 1H, αCH); 2.03-1.91, 1.74-1.57 and 1.41-1.15 (3m, 8H, Chx CH₂); 1.30 (s, 9H, OO'Bu).

¹³C NMR (CDCl₃, 50 MHz): δ 171.96, 170.21, 134.76, 128.88, 129.05, 127.45, 83.36, 45.68, 44.06, 37.98, 28.96, 26.18, 25.15.

MS (ESI-TOF): [M+H]⁺_{calc.} = 334.20, [M+H]⁺_{sper.} = 334.22; [2M+H]⁺_{calc.} = 667.40, [2M+H]⁺_{sper.} = 667.42.

Pht-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from Pht-*trans-rac*-1,2-Chx-OH (0.16 g, 0.6 mmol), EDC·HCl (0.23 g, 1.22 mmol), HOO'Bu (0.55 ml of a 5.5 M solution in decane, 3.0 mmol) and DMAP (0.30 g, 2.44 mmol). Eluant for flash chromatography purification: EtOAc/PE 1:9.

Yield 70 %; oil.

R_f1 0.95, **R_f3** 0.85, **R_f4** 0.60.

IR (KBr): 1771, 1710, 1611 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): δ 7.83 and 7.67 (2m, 4H, Pht CH); 4.45-4.31 (m, 1H, βCH); 3.62-3.47 (m, 1H, αCH); 2.17-1.23 (m, 8H, Chx CH₂); 1.13 (s, 9H, OO'Bu).

¹³C NMR (CDCl₃, 50 MHz): δ 170.91, 167.71, 133.81, 131.78, 123.07, 83.29, 50.97, 42.87, 29.84, 29.49, 25.77, 25.03, 24.44.

MS (ESI-TOF): [M+H]⁺_{calc.} = 346.16; [M+H]⁺_{sper.} = 346.21.

Pht-Aib-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from Pht-Aib-*trans-rac*-1,2-Chx-OH (0.24 g, 0.67 mmol), EDC·HCl (0.26 g, 1.34 mmol), HOO'Bu (0.61 ml of a 5.5 M solution in decane, 3.35 mmol) and DMAP (0.33 g, 2.70 mmol). Eluant for flash chromatography purification: EtOAc/PE 1:2.

Yield 88 %; **m. p.** 145-146 °C.

R_f1 0.90, **R_f2** 0.85, **R_f3** 0.50.

IR (KBr): 3377, 1774, 1739, 1716, 1682, 1534 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): 7.79 and 7.65 (2m, 4H, Pht CH); 5.82 (d, 1H, Chx NH); 4.03-3.85 (m, 1H, βCH); 2.62-2.51 (m, 1H, αCH); 1.80 and 1.75 (2s, 6H, 2 βCH₃); 2.17-1.23 (m, 8H, Chx CH₂); 1.28 (s, 9H, OO'Bu).

¹³C NMR (CDCl₃, 50 MHz): δ 172.22, 170.98, 168.46, 133.94, 131.87, 123.02, 83.48, 61.39, 60.32, 50.76, 46.69, 31.64, 28.98, 26.11, 24.86, 24.52, 24.38, 24.17.

MS (ESI-TOF): [M+H]⁺_{calc.} = 431.40; [M+H]⁺_{sper.} = 431.25.

Pht-(Aib)₂-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from Pht-(Aib)₂-*trans-rac*-1,2-Chx-OH (0.30 g, 0.68 mmol), EDC·HCl (0.26 g, 1.34 mmol), HOO'Bu (0.62 ml of a 5.5 M solution in decane, 3.40 mmol) and DMAP (0.33 g, 2.70 mmol). Eluant for flash chromatography purification: EtOAc/PE 1:1.

Yield 85 %; **m. p.** 167-169 °C.

R_f1 0.75, **R_f2** 0.85, **R_f3** 0.40.

IR (KBr): 3379, 1774, 1709, 1659, 1520 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): 7.89 and 7.73 (2m, 4H, Pht CH); 6.88 (d, 1H, Chx NH); 5.97 (s, 1H, Aib NH); 4.07-3.89 (m, 1H, βCH); 2.80-2.66 (m, 1H, αCH); 1.81 and 1.79 (2s, 6H, 2 βCH₃); 1.53 and 1.41 (2s, 6H, 2 βCH₃); 2.25-1.13 (m, 8H, Chx CH₂); 1.27 (s, 9H, OO'Bu).

MS (ESI-TOF): [M+H]⁺_{calc.} = 516.51; [M+H]⁺_{sper.} = 516.36.

Pht-(Aib)₃-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from Pht-(Aib)₃-*trans-rac*-1,2-Chx-OH (0.10 g, 0.19 mmol), EDC·HCl (0.07 g, 0.38 mmol), HOO'Bu (0.17 ml of a 5.5 M solution in decane, 0.95 mmol) and DMAP (0.09 g, 0.76 mmol). Eluant for flash chromatography purification: EtOAc/PE 1:1.

Yield 75 %; **m. p.** 138-140 °C.

R_f1 0.55, **R_f2** 0.85, **R_f3** 0.30.

IR (KBr): 3383, 1774, 1709, 1664, 1532 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): 7.81 and 7.71 (2m, 4H, Pht CH); 7.24 (d, 1H, Chx NH); 6.97, 6.44 (2s, 2H, Aib NH); 4.07-3.90 (m, 1H, βCH); 2.59-2.46 (m, 1H, αCH); 1.88 and 1.77 (2s, 6H, 2 βCH₃); 1.44 and 1.36 (2s, 12H, 4 βCH₃); 2.11-1.15 (m, 8H, Chx CH₂); 1.21 (s, 9H, OO'Bu).

¹³C NMR (CDCl₃, 50 MHz): δ 175.00, 172.95, 172.88, 172.26, 168.92, 134.45, 131.66, 123.32, 83.06, 61.46, 57.42, 57.05, 48.76, 46.63, 31.64, 30.84, 29.62, 26.80, 26.38, 26.14, 25.16, 24.83, 24.45, 24.18, 24.07, 23.91, 22.95.

MS (ESI-TOF): [M+H]⁺_{calc.} = 601.62; [M+H]⁺_{sper.} = 601.35.

Pht-(Aib)₄-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from Pht-(Aib)₄-*trans-rac*-1,2-Chx-OH (0.30 g, 0.50 mmol), EDC·HCl (0.19 g, 1.0 mmol), HOO'Bu (0.45 ml of a 5.5 M solution in decane, 2.5 mmol) and DMAP (0.24 g, 2.0 mmol). Eluant for flash chromatography purification: EtOAc.

Yield 78 %; **m. p.** 165-167 °C.

R_f1 0.60, **R_f2** 0.85, **R_f3** 0.30.

IR (KBr): 3371, 3337, 1774, 1711, 1669, 1524 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): 7.86 and 7.76 (2m, 4H, Pht CH); 7.24 (s, 2H, Aib NH); 7.07 (d, 1H, Chx NH); 5.94 (s, 1H, Aib NH); 4.05-3.96 (m, 1H, βCH); 2.67-2.55 (m, 1H, αCH); 1.87 and 1.77 (2s, 6H, 2 βCH₃); 1.60, 1.53, 1.47 and 1.40 (4s, 18H, 6 βCH₃); 2.04-1.16 (m, 8H, Chx CH₂); 1.27 (s, 9H, OO'Bu).

¹³C NMR (CDCl₃, 50 MHz): δ 174.51, 173.51, 173.26, 172.60, 171.39, 168.99, 134.72, 131.53, 123.46, 83.13, 61.39, 57.48, 57.04, 56.83, 48.87, 47.29, 31.89, 29.78, 27.23, 26.77, 26.66, 26.22, 24.60, 24.41, 24.17, 23.85, 23.67, 23.55.

MS (ESI-TOF): [M+H]⁺_{calc.} = 686.73; [M+H]⁺_{sper.} = 601.39.

Pht-(Aib)₅-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from Pht-(Aib)₅-*trans-rac*-1,2-Chx-OH (0.24 g, 0.35 mmol), EDC·HCl (0.13 g, 0.70 mmol), HOO'Bu (0.32 ml of a 5.5 M solution in decane, 1.75 mmol) and DMAP (0.17 g, 1.4 mmol). Eluant for flash chromatography purification: EtOAc.

Yield 81 %; **m. p.** 158-159 °C.

R_{f1} 0.60, **R_{f2}** 0.85, **R_{f3}** 0.30.

IR (KBr): 3329, 1774, 1760, 1709, 1667, 1530 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): 7.85 and 7.75 (2m, 4H, Pht CH); 7.56 (s, 1H, Aib NH); 7.28 (d, 1H, Chx NH); 7.22, 6.51 and 5.30 (3s, 3H, Aib NH); 4.12-3.96 (m, 1H, βCH); 2.66-2.48 (m, 1H, αCH); 1.88 and 1.79 (2s, 6H, 2 βCH₃); 2.04-1.16 (m, 32H, Chx CH₂ and Aib βCH₃); 1.27 (s, 9H, OO'Bu).

MS (ESI-TOF): [M+H]⁺_{calc.} = 771.84; [M+H]⁺_{sper.} = 771.44.

Pht-(Aib)₆-*trans-rac*-1,2-Chx-OO'Bu

This compound was prepared as described above for PhAc-*cis*-1,4-Chx-OO'Bu, starting from Pht-(Aib)₆-*trans-rac*-1,2-Chx-OH (0.05 g, 0.06 mmol), EDC·HCl (0.03 g, 0.16 mmol), HOO'Bu (0.06 ml of a 5.5 M solution in decane, 0.33 mmol) and DMAP (0.03 g, 0.25 mmol). Eluant for flash chromatography purification: EtOAc/PE 8:2.

Yield 66 %; **m. p.** 147-148 °C.

R_{f1} 0.55, **R_{f2}** 0.85, **R_{f3}** 0.30.

IR (KBr): 3319, 1775, 1711, 1663, 1529 cm⁻¹.

¹H NMR (CDCl₃, 200 MHz): 7.78-7.56 (m, 4H, Pht CH); 7.56, 7.49, 7.31 and 7.22 (4s, 4H, Aib NH); 7.17 (d, 1H, Chx NH); 6.97 (s, 1H, Aib NH); 4.05-3.90 (m, 1H, Chx βCH); 2.67-2.45 (m, 1H, Chx αCH); 1.85 and 1.76 (2s, 6H, 2 βCH₃); 2.04-1.16 (m, 38H, Chx CH₂ and Aib βCH₃); 1.19 (s, 9H, OO'Bu).

MS (ESI-TOF): [M+H]⁺_{calc.} = 856.95; [M+H]⁺_{sper.} = 856.70.

Figure S1: FT-IR absorption spectra (N-H stretching region) of the Pht-(Aib)_n-*trans-rac*-1,2-Chx-OO^tBu ($n = 1-6$) peptide series in CDCl₃ solution. Peptide concentration: 1mM.

Figure S2: X-Ray diffraction structure of PhAc-*trans-rac*-1,2-Chx-OO'Bu (molecule **B**).

Table S1. Crystal data and structure refinement for PhAc-*cis*-1,4-Chx-OO'Bu.

Identification code	mc49	
Empirical formula	C ₁₉ H ₂₇ N O ₄	
Formula weight	333.42	
Temperature	293(2) K	
Wavelength	1.54178 Å	
Crystal system	Monoclinic	
Space group	P2(1)/a	
Unit cell dimensions	a = 9.667(2) Å	α = 90°.
	b = 17.814(3) Å	β = 105.49(2)°.
	c = 11.034(2) Å	γ = 90°.
Volume	1831.1(6) Å ³	
Z	4	
Density (calculated)	1.209 Mg/m ³	
Absorption coefficient	0.680 mm ⁻¹	
F(000)	720	
Crystal size	0.40 x 0.40 x 0.25 mm ³	
Theta range for data collection	4.16 to 60.05°.	
Index ranges	-10 ≤ h ≤ 10, -1 ≤ k ≤ 20, 0 ≤ l ≤ 12	
Reflections collected	3064	
Independent reflections	2714 [R(int) = 0.0227]	
Completeness to theta = 60.05°	99.7 %	
Absorption correction	None	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	2714 / 0 / 205	
Goodness-of-fit on F ²	1.038	
Final R indices [I > 2σ(I)]	R1 = 0.0446, wR2 = 0.1232	
R indices (all data)	R1 = 0.0490, wR2 = 0.1266	
Largest diff. peak and hole	0.221 and -0.178 e.Å ⁻³	

Table S2. Crystal data and structure refinement for PhAc-*trans-rac*-1,2-Chx-OO'Bu.

Identification code	mc47	
Empirical formula	C ₁₉ H ₂₇ N O ₄	
Formula weight	333.42	
Temperature	293(2) K	
Wavelength	1.54178 Å	
Crystal system	Monoclinic	
Space group	P2(1)	
Unit cell dimensions	a = 9.717(2) Å	$\alpha = 90^\circ$.
	b = 10.789(2) Å	$\beta = 91.73(7)^\circ$.
	c = 18.670(3) Å	$\gamma = 90^\circ$.
Volume	1956.4(6) Å ³	
Z	4	
Density (calculated)	1.132 Mg/m ³	
Absorption coefficient	0.637 mm ⁻¹	
F(000)	720	
Crystal size	0.40 x 0.30 x 0.15 mm ³	
Theta range for data collection	2.37 to 60.02°.	
Index ranges	-10 ≤ h ≤ 10, -1 ≤ k ≤ 12, 0 ≤ l ≤ 20	
Reflections collected	3507	
Independent reflections	3412 [R(int) = 0.0403]	
Completeness to theta = 60.02°	99.7 %	
Absorption correction	None	
Refinement method	Full-matrix-block least-squares on F ²	
Data / restraints / parameters	3412 / 1 / 409	
Goodness-of-fit on F ²	1.044	
Final R indices [I > 2sigma(I)]	R1 = 0.0692, wR2 = 0.1912	
R indices (all data)	R1 = 0.0773, wR2 = 0.1995	
Absolute structure parameter	-0.2(4)	
Largest diff. peak and hole	0.242 and -0.236 e.Å ⁻³	