Synthesis and Confirmation of the Absolute Stereochemistry of the (–)-Aflastatin A C₉–C₂₇ Degradation Polyol

David A. Evans,* William C. Trenkle, Jing Zhang and Jason D. Burch

Dept of Chemistry & Chemical Biology, Harvard University, Cambridge, Massachusetts, 02138

Supporting Information

General Information: All reactions were carried out under an atmosphere of nitrogen in flame- or oven-dried glassware with magnetic stirring, unless otherwise noted. THF, Et₂O, CH₂Cl₂ and toluene are purified by filtration through a column of activated alumina under an argon atmosphere.¹ Reagents were purified prior to use following the guidelines of Perrin and Armarego.² Analytical thin layer chromatography (TLC) was performed on EM Reagent 0.25 mm silica gel 60-F plates. Visualization was accomplished with UV light and exposure to aqueous ceric ammonium molybdate solution or anisaldehyde followed by heating. Flash chromatography was performed using EM silica gel 60 (230-240 mesh). Solvents for extraction and chromatography were HPLC grade. Optical rotations were measured on a Jasco DIP-0181 digital polarimeter with a sodium lamp and reported as follows: $[\alpha]_D^{25}$ (c g/100 ml, solvent). Infrared spectra were recorded on a Perkin Elmer model 1600 FT-IR spectrometer. Low or high resolution mass spectra were obtained on a JEOL AX-505 or SX-102 high resolution magnetic sector mass spectrometer by the Harvard University Mass Spectrometry Laboratory. ¹H NMR spectra were recorded on Inova-500 (500 MHz) or Mercury-400 (400 MHz) spectrometers. Chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance as the internal standard (CDCl₃: δ 7.26 ppm, C₆D₆: δ 7.15 ppm, CD₃OD: δ 3.30 ppm). Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, bs = broad singlet, m = multiplet), coupling constants (Hz), integration. ¹³C NMR spectra were recorded on Varian INOVA-500 (125 MHz) or Mercury-400 (100 MHz) spectrometers with complete proton decoupling. Chemical shifts are reported in ppm from tetramethylsilane with the solvent as the internal standard (CDCl₃: δ 77.0 ppm, C₆D₆: δ 128.0 ppm, CD₃OD: δ 49.0 ppm).

¹ Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics*, **1996**, 15, 1518.

² Perrin, D. D. and Armarego, W. L. *Purification of Laboratory Chemicals*; 3rd ed., Pergamon Press, Oxford, 1988.

tert-butyl 2-[(6S,4R)-2-(4-methoxyphenyl)-6-[(phenylmethoxy)methyl]-1,3-dioxan-4-yl]acetate (11). The diol **10**³ (310 mg, 1 mmol, 1 equiv) was dissolved in 5 mL of CH₂Cl₂. To the solution was added catalytic amount of PPTs (10 mg) and anisaldehyde dimethylacetal (273 mg, 1.5 mmol, 1.5 equiv). The resulting solution was stirred at rt overnight and then concentrated. The residue was purified by flash chromatography (5% EtOAc/hexanes) to afford the acetal **11** as a colorless oil (0.428 g, 100%): TLC R_f = 0.60 (20% EtOAc/hexanes); [α]_D²³ -6.9° (c 0.99, CHCl₃); IR (film) 2975, 2915, 2866, 1728, 1616, 1518, 1368, 1429, 1153, 1099, 1031, 828, 698 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.42 (d, *J* = 8.8 Hz, 2H), 7.35 (d, *J* = 4.4 Hz, 4H), 7.26 (m, 1H), 6.87 (d, *J* = 8.8 Hz, 2H), 5.55 (s, 1H), 4.60 (d, *J* = 4.8 Hz, 2H), 4.28 (m, 1H), 4.11 (m, 1H), 3.79 (s, 3H), 3.66 (dd, *J* = 10.3, 5.9 Hz, 1H), 3.53 (dd, *J* = 10.3, 4.4 Hz, 1H), 2.63 (dd, *J* = 15.4, 7.0 Hz, 1H), 2.43 (dd, *J* = 15.4, 6.2 Hz, 1H), 1.74 (dt, *J* = 12.8, 2.6 Hz, 1H), 1.49-1.55 (m, 1H), 1.45 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 170.0, 159.8, 138.1, 130.9, 128.4, 127.7, 127.6, 127.4, 113.5, 100.5, 80.8, 75.8, 73.5, 73.3, 72.8, 55.3, 42.2, 33.1, 28.1; Exact mass calcd. for C₂₅H₃₃O₆ [M+H]⁺: 429.2277; found: 429.2280 (ES⁺).

tert-butyl 2-[(6S,4R)-6-(hydroxymethyl)-2-(4-methoxyphenyl)-1,3-dioxan-4-yl]acetate. The benzyl ether **11** (428 mg, 1 mmol) was dissolved in 3 mL of absolute alcohol. To this solution was added two pittettes of Raney Nickel in ethanol slurry. The reaction mixture was then stirred at rt under 1 atm H₂ for 3 h. The suspension was then filtered through Celite, washed with EtOAc carefully and concentrated. The residue was then purified via flash chromatography (10%–20% EtOAc/hexanes) to afford the primary alcohol as a colorless oil (300 mg, 89%): TLC R_f = 0.55 (20% EtOAc/hexanes); [α]_D²³ -3.7 (c 0.9, CHCl₃); IR (film) 3452, 2977, 2934, 1731, 1615, 1519, 1456 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.44–7.39 (m, 2H), 6.95–6.88 (m, 2H), 5.56 (s, 1H), 4.32–4.24 (m, 1H), 4.04–3.97 (m, 1H), 3.80 (s, 3H), 3.72 (dd, *J* = 3.3, 11.9 Hz), 3.64 (dd, *J* = 11.9, 6.6 Hz, 1H), 2.64 (dd, *J* = 15.4, 7.0 Hz, 1H), 2.44 (dd, *J* = 15.6, 6.4 Hz, 1H), 2.10–1.95 (bs, 1H), 1.66–1.50 (m, 2H), 1.45 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 170.2, 160.2, 130.9, 127.6, 113.8, 100.8, 81.1, 77.13, 73.4, 65.8, 55.5, 42.4, 32.1, 28.3; Exact mass calcd. for C₁₈H₂₇O₆ [M+H]⁺: 339.1807; found: 339.1826 (ES⁺).

tert-butyl-2-[(6S,4R)-6-[[1,1-bis(methylethyl)-2-methyl-1-silapropoxy] methyl]-2-(4-methoxyphenyl)-1,3-dioxan-4-yl]acetate. The alcohol (300 mg, 0.887 mmol, 1 equiv) was dissolved in 5 mL of CH₂Cl₂. To the solution was added imidazole (121 mg, 1.774 mmol, 2 equiv) and triisopropylsilyl chloride (205 mg, 1.064 mmol, 1.2 equiv). The reaction was stirred for 16 h at rt, then concentrated and purified by flash

³ Evans, D. A. ; Kozłowski, M. C. ; Murry, J. A. ; Burgey, C. S. ; Campos, K. R. ; Connell, B. T. ; Staples, R. J. *J. Am. Chem. Soc.* **1999**, 121, 669.

chromatography (0%→5% EtOAc/hexanes), affording the silyl ether as a colorless oil (395 mg, 90%): TLC R_f = 0.83 (20% EtOAc/hexanes); $[\alpha]_D^{23}$ -7.2 (c 1.0, CHCl₃); IR (film) 2943, 2867, 1730, 1617, 1519, 1463, 1368 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.42–7.37 (m, 2H), 6.87–6.82 (m, 2H), 5.52 (s, 1H), 4.30–4.22 (m, 1H), 4.00–3.91 (m, 1H), 3.90 (dd, J = 9.5, 5.1 Hz, 1H), 3.77 (s, 3H), 3.66 (dd, J = 9.5, 6.6 Hz, 1H), 2.63 (dd, J = 15.2, 7.3 Hz, 1H), 2.43 (dd, J = 15.0, 6.2 Hz, 1H), 1.86 (dt, J = 13.1, 2.2, 2.2 Hz, 1H), 1.44 (s, 9H), 1.30–1.15 (m, 1H), 1.15–1.02 (m, 21H); ¹³C NMR (100 MHz, CDCl₃) δ 170.1, 159.7, 131.0, 127.3, 113.4, 100.4, 80.7, 77.1, 73.5, 66.2, 55.2, 42.4, 33.6, 28.0, 17.9, 17.6, 12.2, 11.8; Mass required for C₂₇H₄₇SiO₆ [M+H]⁺: 495.3142; found: 495.3140 (ES⁺).

2-((6S,4R)-6-[[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]methyl]-2-(4-methoxyphenyl)(1,3-dioxan-4-yl))-N-methoxy-N-methylacetamide (12). To a solution of ester (395 mg, 0.799 mmol, 1 equiv) in 5 mL of THF was added solid N,O-dimethyl hydroxylamine hydrochloride salt (234 mg, 2.397 mmol, 3 equiv). The resulting suspension was cooled to -30 °C immediately. Then 2 M isopropylmagnesium chloride in THF solution (2.40 mL, 4.755 mmol, 5.95 equiv) was introduced into the suspension dropwise. After stirring at -30 °C for 15 min, the reaction mixture was allowed to warm up to 0 °C slowly by turning off the cryocool over 3 h. The reaction was quenched with a mixed solution of 5 mL of satd. NaHCO₃ and 5 mL of brine and extracted with 3 × 30 mL of diethyl ether. The combined organic extracts were dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (20%→50% EtOAc/hexanes) to provide the Weinreb amide **12** as a colorless oil (333 mg, 87%): TLC R_f = 0.10 (20% EtOAc/hexanes); $[\alpha]_D^{23}$ -2.0 (c 1.0, CHCl₃); IR (film) 2942, 2866, 1666, 1615, 1519, 1463, 1384 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.40 (d, J = 8.4 Hz, 2H), 6.85 (d, J = 8.8 Hz, 2H), 5.54 (s, 1H), 4.45–4.36 (m, 1H), 4.02–3.95 (m, 1H), 3.90 (dd, J = 9.9, 5.1 Hz, 1H), 3.78 (s, 3H), 3.71–3.66 (m, 1H), 3.66 (s, 3H), 3.19 (s, 3H), 2.98 (dd, J = 15.4, 6.6 Hz, 1H), 2.54 (dd, J = 15.7, 5.5 Hz, 1H), 1.91 (dt, J = 13.1, 2.4, 2.4 Hz, 1H), 1.53–1.43 (m, 1H), 1.12–1.02 (m, 21H); ¹³C NMR (100 MHz, CDCl₃) δ 171.4, 159.7, 131.1, 127.4, 113.4, 100.6, 77.2, 73.4, 66.3, 61.4, 55.2, 38.3, 33.9, 31.9, 17.9, 12.2, 11.9, 11.6; Exact mass calcd. for C₂₅H₄₄NO₆Si [M+H]⁺: 482.2938; found: 482.2926 (ES⁺).

(5S,3R)-6-[[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-3-hydroxy-N-methoxy-5-[(4-methoxyphenyl)methoxy]-N-methylhexanamide(13). MgBr₂•Et₂O (5.90 g, 22.84 mmol, 2 equiv) was charged into a 250 mL flask containing the acetal **12** (5.50 g, 11.42 mmol, 1 equiv) in the glovebox. Then 100 mL of CH₂Cl₂ was added into the flask to make a

slurry. To the suspension was added *n*-tributyltin chloride (4.32 g, 14.84 mmol, 1.3 equiv). The resulting light yellow solution was stirred at rt for 5 h. The reaction was quenched with 60 mL of water and extracted with 3 × 200 mL of CH₂Cl₂. The combined organic extracts were dried over MgSO₄ and concentrated. The residue was purified via flash chromatography (20%–50% EtOAc/hexanes) to afford the PMB ether **13** as a colorless oil (5.10 g, 93%): TLC R_f = 0.2 (50% EtOAc/hexanes); [α]_D²³ -31.0° (c 0.98, CHCl₃); IR (film) 3474, 2942, 2866, 1651, 1614, 1514, 1464, 1386, 1302, 1249, 1174, 1098, 1036, 998, 883, 823 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.24 (d, *J* = 8.3 Hz, 2H), 6.83 (d, *J* = 8.8 Hz, 2H), 4.69 (d, *J* = 11.2 Hz, 1H), 4.49 (d, *J* = 11.2 Hz, 1H), 4.23 (m, 1H), 3.98 (d, *J* = 1.5 Hz, 1H), 3.80 (dd, *J* = 11.7, 6.8 Hz, 1H), 3.76 (s, 3H), 3.72 (dd, *J* = 11.7, 4.4 Hz, 2H), 3.61 (s, 3H), 3.14 (s, 3H), 2.54 (m, 2H), 1.71-1.80 (m, 2H), 1.03-1.08 (m, 21H); ¹³C NMR (125 MHz, CDCl₃) δ 173.0, 159.0, 130.5, 129.4, 113.6, 78.2, 71.7, 66.4, 65.9, 61.0, 55.1, 38.5, 38.2, 31.7, 17.9, 11.7; Exact mass calcd. for C₂₅H₄₅NO₆Si [M]⁺: 484.3094; found: 484.3076 (ES⁺).

(8S,6R)-9-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-6-hydroxy-8-[(4-methoxyphenyl)methoxy]non-1-en-4-one (14). The Weinreb amide **13** (5.10 g, 9.70 mmol, 1 equiv) was dissolved in 70 mL of THF. After cooling to 0 °C, the solution was treated with allylmagnesium bromide dropwise and stirred at 0 °C for 2 h. The reaction was quenched with the minimal amount of 1 N HCl solution to dissolve all the precipitates. The reaction mixture was then extracted with 3 × 200 mL of diethyl ether. The combined organic extracts were dried over MgSO₄ and concentrated in *vacuo*. Purification by flash chromatography (10%–20% EtOAc/hexanes) afforded the ketone **14** (4.00 g, 89%) as a colorless oil: TLC R_f = 0.75 (50% EtOAc); [α]_D²⁴ -38.5° (c 1.05, CHCl₃); IR (film) 3488, 2945, 2867, 1710, 1612, 1514, 1465, 1250, 1095, 1035, 908, 733 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.25 (d, J = 8.3 Hz, 2H), 6.87 (d, J = 8.3 Hz, 2H), 5.88 (m, 1H), 5.19 (d, J = 1.5 Hz, 1H), 5.17 (d, J = 1.5 Hz, 1H), 4.71 (d, J = 11.2 Hz, 1H), 4.50 (d, J = 11.2 Hz, 1H), 4.25 (m, 1H), 3.78–3.83 (m, 1H), 3.79 (s, 3H), 3.67–3.74 (m, 2H), 3.17 (d, J = 6.8 Hz, 2H), 2.60 (dd, J = 16.1, 7.8 Hz), 2.44 (dd, J = 16.6, 4.4 Hz, 1H), 1.63–1.75 (m, 2H), 1.05–1.13 (m, 21H); ¹³C NMR (125 MHz, CDCl₃) δ 201.6, 159.2, 130.3, 130.2, 129.6, 119.0, 113.8, 78.8, 71.8, 66.6, 65.9, 55.2, 49.1, 48.4, 38.3, 17.9, 11.8; Exact mass calcd. for C₂₆H₄₅O₅Si [M+H]⁺: 465.3036; found: 465.3049 (ES⁺).

(4S,6S,8S)-9-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-8-[(4-methoxyphenyl)methoxy]non-1-ene-4,6-diol. The ketone **14** (4.10 g, 8.82 mmol, 1 equiv) was dissolved in 90 mL of THF and 18 mL of MeOH. To this solution was added Et₂BOMe (1.06 g, 10.59 mmol, 1.2 equiv). After stirring at rt for 15 min, the solution was cooled down to -78 °C. NaBH₄ (0.467 g, 12.35 mmol, 1.4 equiv) was added in 4 portions with a 10 min interval. The resulting solution was stirred at -78 °C for 9 h. Then the reaction was carefully quenched by 36 mL of 30% H₂O₂ and warmed up to rt overnight. The organic solvent was then concentrated in *vacuo*. The residue was then dissolved in 600 mL of EtOAc. The resulting solution was washed with 50 mL of brine, 50 mL of satd. Na₂S₂O₃ and 50 mL of brine, dried over MgSO₄ and concentrated in *vacuo*. Purification by flash chromatography (20%–50% EtOAc/hexanes) afforded the diol as a colorless oil which was taken to the next step directly: TLC R_f = 0.6 (50% EtOAc/hexanes); [α]_D²² -40.4° (c 1.05, CHCl₃); IR (film) 3417, 2945, 2863, 1639, 1613, 1586, 1514, 1464, 1303, 1249 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.27 (d, J = 8.8 Hz, 2H), 6.89 (d, J = 8.8 Hz, 2H), 5.90–5.80 (m, 1H), 5.14–5.05 (m, 2H), 4.73 (d, J = 11.2 Hz, 1H), 4.51 (d, J = 11.2 Hz, 1H), 4.18 (s, 1H), 4.08–4.00 (m, 1H), 3.92 (s, 1H), 3.95–3.86 (m, 1H), 3.86–3.78 (m, 1H), 3.81 (s, 3H), 3.77–3.65 (m, 2H), 2.30–2.15 (m, 2H), 1.75–1.65 (m, 2H), 1.55–1.48 (m, 2H), 1.18–1.00 (m, 21H); ¹³C NMR (125 MHz, CDCl₃) δ 159.3, 134.9, 130.0, 129.6, 117.2, 113.8, 79.4, 71.8, 71.7, 71.4, 65.9, 55.2, 42.6, 42.1, 39.5, 17.9, 17.9, 11.8; Exact mass calcd. for C₂₆H₄₇O₅Si [M+H]⁺: 467.3193; found: 467.3195 (ES⁺).

1-((1S)-1-(((6S,4R)-2,2-dimethyl-6-prop-2-enyl(1,3-dioxan-4-yl))methyl)-2-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]ethoxy)methyl)-4-methoxy benzene (15). The diol from the previous step (~8.82 mmol, 1 equiv) was dissolved in 50 mL of CH₂Cl₂. To this solution was added 2,2-dimethoxypropane (22.97 g, 0.221 mol, 25 equiv) and PPTs (200 mg, catalytic amount). The resulting solution was stirred at rt for 1 h and then concentrated in *vacuo*. The residue was purified via flash chromatography to afford the acetonide **15** as a colorless oil (3.80 g, 85% over two steps): TLC R_f = 0.75 (20% EtOAc/hexanes); [α]_D²³ -29.8 (c 1.22, CHCl₃); IR (film) 2945, 2863, 1611, 1508, 1482, 1375, 1303, 1247 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.29 (d, *J* = 8.8 Hz, 2H), 6.89 (d, *J* = 8.3 Hz, 2H), 5.83–5.74 (m, 1H), 5.11–5.05 (m, 2H), 4.68 (d, *J* = 11.7 Hz, 1H), 4.51 (d, *J* = 11.7 Hz, 1H), 3.96–3.90 (m, 1H), 3.84–3.79 (m, 1H), 3.82 (s, 3H), 3.76–3.70 (m, 2H), 3.60–3.55 (m, 1H), 2.30–2.23 (m, 1H), 2.20–2.05 (m, 1H), 1.78–1.72 (m, 1H), 1.63–1.56 (m, 1H), 1.48–1.36 (m, 1H), 1.42 (s, 3H), 1.39 (s, 3H), 1.23 (dt, *J* = 12.7, 2.4, 2.4 Hz, 1H), 1.18–0.80 (m, 21H); ¹³C NMR (125 MHz, CDCl₃) δ 159.1, 134.2, 131.1, 129.7, 116.9, 113.6, 98.4, 75.4, 71.5, 68.6, 66.4, 66.0, 55.2, 40.8, 38.6, 36.2, 30.2, 19.8, 18.0, 11.9; Exact mass calcd. for C₂₉H₅₁O₅Si [M+H]⁺: 507.3506; found: 507.3521 (ES⁺).

2-((4R,6R)-6-((2S)-3-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-2-[(4-methoxyphenyl)methoxy]propyl)-2,2-dimethyl-1,3-dioxan-4-yl)ethanal (16). The olefin **15** (600 mg, 1.184 mmol, 1 equiv) was dissolved in 15 mL of CH₂Cl₂ and 0.3 mL of MeOH and a few drops of pyridine. A few particles of Sudan III dye were added into this solution to make a red solution. After cooling down to -78 °C, O₃ was bubbled through until the red color faded into colorless. PPh₃ (1.55 g, 5.920 mmol, 5 equiv) was added into the reaction mixture which was then warmed up to rt overnight. The solution was then concentrated in *vacuo*. The residue was purified by flash chromatography (10% EtOAc/hexanes) to provide the aldehyde **16** as a colorless oil (518 mg, 86%): TLC R_f = 0.4 (20% EtOAc/hexanes); ¹H NMR (500 MHz, CDCl₃) δ 9.76 (t, *J* = 2.5 Hz, 1H), 7.30 (d, *J* = 8.3 Hz, 2H), 6.90 (d, *J* = 8.3 Hz, 2H), 4.68 (d, *J* = 11.7 Hz, 1H), 4.50 (d, *J* = 11.7 Hz, 1H), 4.20 (m, 1H), 3.95–4.01 (m, 1H), 3.82 (s, 3H), 3.81 (dd, *J* = 10.7, 6.4 Hz, 1H), 3.70 (dd, *J* = 10.3, 4.9 Hz, 1H), 3.55 (m, 1H), 2.55 (ddd, *J* = 16.1, 7.3, 2.4 Hz, 1H), 2.36 (ddd, *J* = 16.6, 4.9, 1.9 Hz, 1H), 1.75 (ddd, *J* = 14.2, 7.8, 6.3 Hz, 1H), 1.61 (ddd, *J* = 10.2, 6.8, 4.9 Hz, 1H), 1.44 (s, 3H), 1.37 (s, 3H), 1.20 (dt, *J* = 12.7, 2.5 Hz, 1H), 1.04–1.15 (m, 22H).

(4S)-3-[4-(6-[(2S)-3-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-2-[(4-methoxyphenyl)methoxy]propyl)(4S,6R)-2,2-dimethyl(1,3-dioxan-4-yl))(2S,3R)-3-hydroxy-2-methylbutanoyl]-4-benzyl-1,3-oxazolidin-2-one. Propionyl oxazolidinone (0.151 g, 0.649 mmol, 1.5 equiv) in 2 mL of CH₂Cl₂ solution was cooled to -78 °C. *n*-Bu₂BOTf (0.213 g, 0.778 mmol, 1.8 mmol) and NEt₃ (0.092 g, 0.908 mmol, 2.1 equiv) were added dropwise into the solution. The resulting solution was stirred at -78 °C for 1 h, then at 0 °C for 30 min. After cooling back to -78 °C, the aldehyde **16** (0.22 g, 0.432 mmol, 1.0 equiv) in 2 mL of CH₂Cl₂ solution was added via cannula. The flask was rinsed with 2 × 1 mL of CH₂Cl₂ and added into the reaction as well. After stirring at -78 °C for 1 h, then at 0 °C for 1 h, the reaction was quenched with 2.5 mL of pH 7 buffer, 2.5 mL of MeOH and 2.5 mL of 30% H₂O₂ and stirred at 0 °C for 1 h. After diluting with 100 mL of ether, the organic solution was washed with 20 mL of satd. NaHCO₃ solution, dried over MgSO₄ and concentrated. Purification by flash chromatography (20% EtOAc/hexanes) afforded the aldol adduct as a colorless oil (0.30 g, 93%): TLC R_f = 0.10 (20% EtOAc/hexanes); [α]_D²⁵ +9.5° (c 0.52, CHCl₃); IR (film) 3503, 2934, 2853, 1781, 1695, 1609, 1512, 1461, 1385, 1350, 1248, 1207, 1167, 1111, 1035, 959, 918, 730, 699 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) 7.33 (t, 7.0 Hz, 2H), 7.25-7.29 (m, 3H), 7.21 (d, *J* = 6.6 Hz, 2H), 6.88 (d, *J* = 8.4 Hz, 2H), 4.68-4.72 (m, 1H), 4.65 (d, *J* = 11.4 Hz, 1H), 4.48 (d, *J* = 11.7 Hz, 1H), 4.16-4.24 (m, 2H), 4.09-4.13 (m, 1H), 3.92-4.01 (m, 2H), 3.85 (dd, *J* = 6.2, 3.7 Hz, 1H), 3.80 (s, 3H), 3.80 (dd, *J* = 11.7, 5.9 Hz, 1H), 3.68 (dd, *J* = 10.3, 4.4 Hz, 2H), 3.53 (m, 1H), 3.29 (dd, *J* = 13.5, 3.3 Hz, 1H), 2.87 (dd, *J* = 13.2, 9.5 Hz, 1H), 1.47-1.75 (m, 4H), 1.42 (s, 3H), 1.33 (s, 3H), 1.26 (d, *J* = 7.0 Hz, 3H), 1.01-1.20 (m, 21H); ¹³C NMR (100 MHz, CDCl₃) 175.5, 159.1, 153.1, 135.2, 130.9, 129.7, 129.3, 128.8, 127.2, 113.6, 98.5, 75.2, 71.4, 71.3, 69.5, 66.2, 66.0, 65.9, 55.3, 55.2, 55.1, 42.7, 39.9, 38.3, 37.7, 36.6, 30.0, 19.8, 17.9, 17.8, 11.8, 11.3; Exact mass calcd. for C₄₁H₆₃NO₉Si [M]⁺: 742.0; found: 742.0 (ES⁺).

4-(6-[(2S)-3-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-2-[(4-methoxyphenyl)methoxy]propyl)(4S,6R)-2,2-dimethyl(1,3-dioxan-4-yl))(2S,3R)-3-hydroxy-N-methoxy-2-methyl-N-methylbutanamide (17). A suspension of N,O-dimethyl hydroxylamine hydrochloride salt (96.6 mg, 0.991 mmol, 3.0 equiv) in 5 mL of THF was cooled to 0 °C. AlMe₃ (0.5 mL, 1.01 mmol, 3.05 equiv, 2 M in THF) was added dropwise. After stirring at rt for 30 min, the solution was cooled to -78 °C. The imide (0.245 g, 0.330 mmol, 1 equiv) in 3 mL of THF solution was added. The flask was rinsed with 2 × 3 mL of THF and the rinses were added into the reaction mixture as well. The resulting solution was then stirred at -10→0 °C for 10 h. After quenching with 15 mL of 1 M sodium potassium tartrate solution, the resulting suspension was allowed to warm up to rt over 30 min. The organic phase was separated and the aqueous phase was extracted with 3 × 20 mL CH₂Cl₂. The combined organic extracts were dried over MgSO₄ and concentrated. Purification by flash chromatography (35% EtOAc/Hexanes) afforded the Weinreb amide **17** (150 mg, 73%) together with the unreacted imide starting material (53 mg, 22%): TLC R_f = 0.55 (50% EtOAc / hexanes); [α]_D²⁵ -12.0° (c 0.87, CHCl₃); IR (film) 3504,

2942, 2866, 1655, 1638, 1612, 1513, 1459, 1382, 1248, 1200, 1171, 1112, 1036, 996, 882, 796, 682 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.25 (d, J = 8.4 Hz, 2H), 6.85 (d, J = 8.4 Hz, 2H), 4.63 (d, J = 11.7 Hz, 1H), 4.46 (d, J = 11.4 Hz, 1H), 3.92-3.98 (m, 3H), 3.79 (s, 3H), 3.76 (dd, J = 10.3, 5.8 Hz, 1H), 3.68 (s, 3H), 3.66 (m, 1H), 3.53 (m, 1H), 3.18 (s, 3H), 2.91 (bs, 1H), 1.55-1.73 (m, 3H), 1.41-1.48 (m, 1H), 1.41 (s, 3H), 1.39 (m, 1H), 1.34 (s, 3H), 1.22-1.30 (m, 2H), 1.20 (d, J = 6.9 Hz, 3H), 1.01-1.17 (m, 21H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.1, 131.0, 129.5, 113.6, 98.5, 75.6, 71.5, 71.3, 69.2, 66.2, 66.0, 61.5, 55.2, 40.2, 38.5, 36.6, 30.1, 19.8, 18.0, 12.5, 11.8; Exact mass calcd. for $\text{C}_{32}\text{H}_{56}\text{NO}_8\text{Si}$ $[\text{M}-\text{Me}]^+$: 610.4; found: 610.2 (ES^+).

4-((4R,6R)-6-((2S)-3-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-2-[(4-methoxyphenyl)methoxy]propyl)-2,2-dimethyl(1,3-dioxan-4-yl))(2S,3R)-N-methoxy-2-methyl-N-methyl-3-(tert-butyldimethylsiloxy)butanamide. A solution of the alcohol **17** (0.16 g, 0.256 mmol, 1 equiv) in 5 mL of CH_2Cl_2 was cooled to -78°C . 2,6-lutidine (0.110 g, 1.024 mmol, 4 equiv) and TBSOTf (0.135 g, 0.512 mmol, 2 equiv) were added. The resulting solution was stirred at -78°C for 3 h and then quenched with 5 mL of satd. NaHCO_3 . After warming up to rt, the reaction mixture was extracted with 3×20 mL of CH_2Cl_2 . The combined organic extracts were washed with 10 mL of 1 M NaHSO_4 solution, dried over MgSO_4 and concentrated. Purification by flash chromatography (15%–20% EtOAc/Hexanes) afforded the silyl ether (0.155 g, 82%) as a colorless oil: TLC R_f = 0.20 (20% EtOAc/hexanes); $[\alpha]_D^{25}$ -6.84° (c 1.05, CHCl_3); IR (film) 2933, 2866, 1711, 1661, 1605, 1511, 1461, 1378, 1250, 1200, 1172, 1105, 1044, 994, 883, 833, 778, 683 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.26 (d, J = 8.8 Hz, 2H), 6.85 (d, J = 8.8 Hz, 2H), 4.63 (d, J = 11.4 Hz, 1H), 4.48 (d, J = 11.4 Hz, 1H), 3.96-4.06 (m, 3H), 3.78 (s, 3H), 3.74 (dd, J = 19.4, 6.6 Hz, 1H), 3.64-3.69 (m, 1H), 3.68 (s, 3H), 3.58-3.62 (m, 1H), 3.16 (s, 3H), 3.02 (m, 1H), 1.53-1.78 (m, 5H), 1.42-1.45 (m, 1H), 1.39 (s, 3H), 1.34 (s, 3H), 1.13 (d, J = 7.0 Hz, 3H), 1.01-1.10 (m, 21H), 0.89 (s, 9H), 0.05 (s, 3H), 0.04 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 159.0, 131.1, 129.3, 113.6, 98.2, 76.3, 71.5, 70.4, 66.2, 66.1, 65.5, 61.3, 55.2, 38.6, 37.5, 30.2, 25.9, 19.8, 18.03, 17.99, 17.9, 14.1, 11.9, -4.1 , -4.6 ; Exact mass calcd. for $\text{C}_{39}\text{H}_{74}\text{NO}_8\text{Si}_2$ $[\text{M}+\text{H}]^+$: 740.4953; found: 740.4963 (ES^+).

4-((4R,6R)-6-((2S)-3-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-2-[(4-methoxyphenyl)methoxy]propyl)-2,2-dimethyl(1,3-dioxan-4-yl))(2S,3R)-2-methyl-3-(tert-butyldimethylsiloxy)butan-1-one (5). A solution of the amide (80 mg, 0.108 mmol, 1 equiv) in 5 mL of CH_2Cl_2 was cooled to -78°C . A 1 M toluene solution of DIBAL-H (0.32 mL, 0.324 mmol, 3 equiv) was added dropwise into the solution. After stirring at -78°C for 3 h, the reaction mixture was quenched with 0.5 mL of EtOAc and stirred at this temperature for 30 min. Water (0.5 mL), Celite (2 g) and MgSO_4 (0.5 g) were added into

the reaction mixture. After warming up to rt, the suspension was filtered through Celite and washed with 4×15 of mL CH_2Cl_2 and concentrated. Purification by flash chromatography (10% EtOAc/Hexanes) afforded the aldehyde **5** (65 mg, 88%) as a colorless oil: TLC R_f = 0.75 (20% EtOAc/hexanes); ^1H NMR (500 MHz, C_6D_6) δ 9.65 (s, 1H), 7.28 (d, J = 8.3 Hz, 2H), 6.78 (d, J = 8.8 Hz, 2H), 4.68 (d, J = 11.7 Hz, 1H), 4.95 (d, J = 11.7 Hz, 1H), 4.41 (quint, J = 4.9 Hz, 1H), 4.04 (m, 1H), 3.89 (dd, J = 10.3, 5.9 Hz, 1H), 3.79 (dd, J = 10.7, 4.5 Hz, 1H), 3.71 (m, 2H), 3.27 (s, 3H), 2.26 (m, 1H), 1.97 (m, 1H), 1.74-1.81 (m, 3H), 1.47-1.51 (m, 1H), 1.46 (s, 3H), 1.28 (s, 3H), 1.10 (m, 21H), 1.07 (d, J = 6.8 Hz, 3H), 0.89 (s, 9H), 0.04 (s, 3H), -0.01 (s, 3H).

2R-Methyl-5-phenyl-3R-(trimethyl-silanyloxy)-pent-4-enethioic acid S-ethyl ester:

BuLi (400 mL, 2.62M in hexanes, 1.05 mmol, 2.29 equiv) was added dropwise to a -78°C solution of ethanethiol (112 μL , 1.51 mmol, 3.29 equiv) and THF (6.0 mL). The reaction mixture was transferred to a 0°C ice bath upon completion of addition and maintained at 0°C for 15 minutes. A solution of **18**⁴ (200 mg, 0.46 mmol, 1 equiv) in THF (5.0 mL) was added slowly with rapid stirring. The resulting solution was maintained at 0°C until TLC analysis indicated complete reaction (1 hour) and then diluted with 1:1 hexanes/EtOAc (30 mL each). The mixture was washed with 0.5 M NaOH (aq, 30 mL) twice followed by one wash with brine (30 mL). The resulting organic layer was dried (Na_2SO_4) and concentrated to give 210 mg of a light yellow oil. The oil was purified by flash chromatography (10% EtOAc/hexanes) to provide 141 mg (96%) of thioester as a colorless oil: $[\alpha]_D^{23}$ = -91.8 (c 2.0, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.45–7.20 (m, 5H), 6.55 (d, J = 15.7 Hz, 1H), 6.11 (dd, J = 15.7, 7.3 Hz, 1H), 4.50–4.45 (m, 1H), 3.0–2.77 (m, 3H), 1.28 (t, J = 10.7 Hz, 3H), 1.10 (d, J = 7.0 Hz, 3H), 0.11 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 202.1, 136.5, 131.7, 129.8, 128.5, 127.7, 126.5, 75.9, 55.4, 23.1, 14.8, 14.0, 0.1; IR (film) 3027, 2965, 2932, 2875, 1686 cm^{-1} ; Exact mass calcd. for $\text{C}_{17}\text{H}_{26}\text{O}_2\text{S}_1\text{Si}_1\text{Na}_1$ ($\text{M}+\text{Na}^+$): 345.1321; found: 345.1323 (ES^+).

3R-Hydroxy-2R-methyl-5-phenyl-pent-4-enoic acid methoxy-methyl-amide (19):

$\text{Me}(\text{MeO})\text{NH}_2\text{Cl}$ (539 mg, 5.52 mmol, 3 equiv) was added to a solution of the thioester (593 mg, 1.84 mmol, 1 equiv) in THF (18 mL) at rt. The suspension was immediately transferred to a -30°C bath (cryocool), then $i\text{-PrMgCl}$ (5.5 mL of 2.0M in THF, 11.0 mmol, 5.98 equiv) was added dropwise. During the addition, the suspension turns a bright yellow and becomes opaque. After completion of addition, the reaction is allowed to slowly warm to 0°C (internal monitor) over 2 hours. During this time, the

⁴ Evans, D. A.; Tedrow, J. S.; Shaw, J. T.; Downey, C. W. J. *Am. Chem. Soc.* **2002**, 124, 392–393.

reaction becomes homogeneous then the reaction is diluted with EtOAc (400 mL) and sequentially washed with 1 M NaOH (2 × 50 mL) and brine (50 mL). The organic layer was concentrated and the residue redissolved in CH₂Cl₂/MeOH (40 mL each). Concentrated HCl (2 drops) was added and the resulting solution was stirred at rt for 30 minutes then quenched with Et₃N (4.0 mL) and concentrated. The residue was chromatographed (30% → 100% EtOAc in hexanes) to provide 390 mg (85%) of **19** as a colorless oil: $[\alpha]_D^{23} = -26.1$ (c 2.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.40–7.20 (m, 5H), 6.65 (d, *J* = 15.7 Hz, 1H), 6.23 (dd, *J* = 15.7, 6.6 Hz, 1H), 4.38 (t, 6.2 Hz, 1H), 3.71 (s, 3H), 3.50 (bs, 1H, -OH), 3.14–3.06 (m, 1H), 1.25 (d, *J* = 7.3 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 136.6, 131.5, 130.3, 128.5, 127.6, 126.5, 75.1, 61.6, 40.7, 31.9, 14.9; IR (film) cm⁻¹; Exact mass calcd. for C₁₄H₁₉N₁O₃Na₁ (M+Na⁺): 272.1263; found: 272.1259 (ES⁺)

3R-(4-Methoxy-benzyloxy)-2R-methyl-5-phenyl-pent-4-enoic acid methoxy-methylamide: NaH (101 mg of 60% wt suspension, 2.52 mmol, 1.66 equiv) was added to a 0 °C solution of **19** (378 mg, 1.52 mmol, 1 equiv), DMF (5.0 mL) and PMB-Br (1.06 mL of a 2.5M solution in PhMe, 2.65 mmol, 1.75 equiv). The suspension was stirred at 0 °C for 1 hour then poured onto H₂O (30 mL) and pentane/Ether (40 mL each). The layers were separated and the organic layer was sequentially washed with H₂O (15 mL) and brine (15 mL) then dried (Na₂SO₄) and concentrated. The residue was purified via flash chromatography (10% → 50% EtOAc/hexanes) to provide 553 mg (99%) of PMB ether as a colorless oil: $[\alpha]_D^{23} = -31.7$ (c 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.46–7.41 (m, 2H), 7.38–7.32 (m, 2H), 7.30–7.25 (m, 1H), 7.22–7.16 (m, 2H), 6.85–6.80 (m, 2H), 6.63 (d, *J* = 15.7 Hz, 1H), 6.09 (dd, *J* = 15.9, 8.6 Hz, 1H), 4.51 (d, *J* = 11.0 Hz, 1H), 4.34 (d, *J* = 11.4 Hz, 1H), 4.16 (t, *J* = 4.2 Hz, 1H), 3.78 (s, 3H), 3.68 (s, 3H), 3.25 (s, 3H), 1.06 (d, *J* = 7.0 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 175.9, 158.9, 136.3, 134.5, 130.8, 129.1, 128.6, 129.0, 127.9, 126.6, 113.5, 83.0, 70.6, 61.5, 55.2, 40.0, 32.0, 14.2; IR (film) 2936, 1655, 1612, 1513, 1460, 1420, 1387, 1301, 1247 cm⁻¹; Exact mass calcd. for C₂₂H₂₇N₁O₄, (M+H⁺): 370.2018; found: 370.2020 (ES⁺).

1-(4'R-Benzyl-2'-oxo-oxazolidin-3'-yl)-5S-hydroxy-7R-(4-methoxy-benzyloxy)-2R,4S,-6R-trimethyl-9-phenyl-non-8-ene-1,3-dione (22): Dibal-H (24.2 mL, 24.2 mmol, 3 equiv) was added to a -78 °C solution of the Weinreb amide (2.98 g, 8.06 mmol, 1 equiv) and CH₂Cl₂ (40 mL). The reaction was maintained at -78 °C for 3 h then quenched with MeOH (1.0 mL) followed after 3 minutes by H₂O (1 mL) and Na₂SO₄ (anhydrous, 20 g) and 10 g celite. The cooling bath was removed and the suspension was stirred vigorously for 1/2 h while warming to rt. The suspension was filtered through Na₂SO₄ (anhydrous) and celite with washing (CH₂Cl₂, 3 × 40 mL). The filtrate was concentrated

and purified via flash chromatography (10% EtOAc in hexanes) to provide 2.30 g (92%) of aldehyde **20** as a colorless oil.

In a separate flask, imide **21** (2.79 g, 9.63 mmol, 1.3 equiv) was dissolved in Et₂O (42 mL) then cooled to 0 °C. *c*-Hex₂BCl (2.46 g, 11.56 mmol, 1.56 equiv) and EtNMe₂ (0.85 g, 11.56 mmol, 1.56 equiv) were added sequentially and the mixture was stirred at 0 °C. After 1 h at 0 °C, the reaction was chilled to -78 °C and an ethereal solution of the aldehyde **20** (2.30 g, 7.41 mmol, 1 equiv) in 10 mL of Et₂O and 2 × 4 mL rinses were added via cannula. The reaction mixture was stirred at -78 °C for 3 h then placed a -20 °C freezer for 18 h. The reaction was placed in an ice bath (0 °C) with vigorous stirring for 5 minutes then quench with 2:1 NH₄Cl/MeOH (45 mL total volume) and stirred for an additional 5 minutes. H₂O (~4 mL) was added to make the salts soluble and the biphasic mixture was poured onto 3:1 EtOAc/hexanes (400 mL total volume). The layers were separated and the aqueous layer was extracted with 3:1 EtOAc/hexanes (200 mL total volume). The organic layers were combined, dried (Na₂SO₄), concentrated and chromatographed (10% EtOAc in hexanes) to provide 3.73 g (84%) of **22** as an amorphous white foam: $[\alpha]_D^{23} = -94.2$ (c 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.46–7.18 (m, 12H), 6.90–6.82 (m, 1H), 6.58 (d, *J* = 15.7 Hz, 1H), 6.22 (dd, *J* = 15.9, 8.2 Hz, 1H), 4.96–4.88 (m, 1H), 4.78–4.68 (m, 1H), 4.58 (d, *J* = 11.4 Hz, 1H), 4.27 (d, *J* = 11.7 Hz, 1H), 4.30–4.08 (m, 2H), 3.93 (dd, *J* = 7.9, 5.9 Hz, 1H), 3.82–3.76 (m, 1H), 3.79 (s, 3H), 3.29 (dd, *J* = 13.2, 3.3 Hz, 1H), 2.97–2.85 (m, 1H), 2.77 (dd, *J* = 13.2, 9.5 Hz, 1H), 2.72 (d, *J* = 2.6 Hz, 1H), 1.84–1.75 (m, 1H), 1.48 (d, *J* = 7.3 Hz, 3H), 1.02 (d, *J* = 7.0 Hz, 3H), 0.96 (d, *J* = 7.0 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 211.0, 170.9, 159.2, 153.3, 136.2, 135.1, 133.6, 129.9, 129.5, 129.4, 128.9, 128.6, 128.5, 127.9, 127.3, 126.5, 113.8, 83.0, 72.5, 70.2, 66.2, 55.3, 55.2, 53.1, 47.3, 39.0, 37.9, 13.6, 12.7, 9.8; IR (film) 3604, 1934, 1780, 1718, 1698, 1612, 1513, 1454, 1388, 1360 cm⁻¹; Exact mass calcd. for C₃₆H₄₁N₁O₇Na₁ (M+Na⁺): 622.2781; found: 622.2272 (ES⁺).

Hz, 1H), 1.87–1.76 (m, 1H), 1.45 (d, $J = 7.3$ Hz, 3H), 1.12 (d, $J = 7.0$ Hz, 3H), 0.96 (t, $J = 7.9$ Hz, 9H), 0.83 (d, $J = 7.3$ Hz, 3H), 0.68–0.52 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 208.9, 171.1, 158.9, 153.2, 136.5, 135.1, 133.8, 130.8, 130.0, 129.4, 129.1, 128.9, 128.6, 127.8, 127.4, 126.5, 113.7, 81.8, 72.9, 69.5, 66.2, 55.2, 52.9, 49.8, 40.5, 37.8, 14.1, 12.3, 10.4, 7.1, 5.3; IR (film) 2953, 2876, 1782, 1718, 1699, 1612, 1514, 1455, 1385, 1356 cm^{-1} ; Exact mass calcd. for $\text{C}_{42}\text{H}_{55}\text{N}_1\text{O}_7\text{Si}_1\text{Na}_1$ ($\text{M}+\text{Na}^+$): 736.3645; found: 736.3660 (FAB).

4R-Benzyl-3-(2'R-{6"S-[2"R-(4-methoxy-benzyloxy)-1"R-methyl-4"-phenyl-but-3"-enyl]-2",2",5"S-trimethyl-[1,3]dioxan-4"S-yl}-propionyl)-oxazolidin-2-one (24): Freshly prepared $\text{Zn}(\text{BH}_4)_2$ (6.7 mL of a 0.15 M solution in Et_2O , 1.0 mmol) was added dropwise to a -78°C solution of the ketone (596 mg, 0.835 mmol) and CH_2Cl_2 (9.0 mL) in a 500 mL round bottom flask then the reaction was transferred to a -35°C bath and maintained at -35°C for 30 minutes. Additional $\text{Zn}(\text{BH}_4)_2$ (3.4 mL, 0.5 mmol) was added and the reaction was allowed to warm to -25°C (reset cryocool temperature) then maintained at -25°C for 3 h. The reaction was quenched by addition of NH_4Cl (12 mL) then allowed to warm to rt [note: be careful of foaming during warming]. After the reaction reached rt, H_2O (~ 5 mL to solubilize the salts) and CH_2Cl_2 (40 mL) were added. The layers were separated and the aqueous layer was extracted with CH_2Cl_2 (3×40 mL). The combined organic layers were dried (Na_2SO_4) and concentrated to provide 632 mg of alcohol **23** as a colorless oil which was carried into the next reaction without further purification.

$\text{HF} \cdot \text{pyridine}$ (1.0 mL) was added to a solution of **23** (632 mg, 0.835), THF (16 mL) and pyridine (2.0 mL) in a HDPE plastic bottle. After 2 h, the reaction appeared to stall by TLC and additional pyridine (1.0 mL) and $\text{HF} \cdot \text{pyridine}$ (0.5 mL) was added. The reaction was quenched after 4 1/2 hours at rt by pouring onto $\text{NaHCO}_3/\text{CH}_2\text{Cl}_2$ (200 mL each). The layers were separated and the organic layer was washed with 1N NaHSO_4 (200 mL), dried (Na_2SO_4) and concentrated to provide 568 mg of the diol as a light yellow oil which was carried into the next reaction without further purification.

Camphorsulfonic acid (2 mg, cat.) was added to a solution of 2,2-dimethoxypropane (5 mL) and the diol from the previous step (568 mg, 0.835 mmol) and the mixture was stirred for 5 h at rt. The reaction was quenched with Et_3N (10 mL), concentrated and chromatographed (20% $\rightarrow$ 40% EtOAc /hexanes) to provide 340 mg (86% from **23**) of desired acetonide **24** as a colorless oil: $[\alpha]_D^{22} = -233.9$ (c 1.0, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.45–7.20 (m, 12H), 6.90–6.84 (m, 2H), 6.57 (d, $J = 16.1$ Hz, 1H), 6.08 (dd, $J = 16.1, 8.8$ Hz, 1H), 4.63–4.55 (m, 1H), 4.52 (d, $J = 11.0$ Hz, 1H), 4.26 (d, $J = 11.4$ Hz, 1H), 4.22–3.98 (m, 3H), 4.06 (dd, $J = 10.3, 1.8$ Hz, 1H), 3.95 (dd, $J = 9.9$ Hz, 3.3 Hz, 1H), 3.85 (t, $J = 10.8$ Hz, 1H), 3.80 (s, 3H), 3.39 (dd, $J = 13.2, 2.9$ Hz, 1H), 2.77 (dd, $J = 13.4, 10.1$ Hz, 1H), 1.99–1.88 (m, 1H), 1.72–1.60 (m, 1H), 1.33 (s, 3H), 1.30 (s, 3H), 1.22 (d, $J = 7.0$ Hz, 3H), 0.85–0.79 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 174.6, 158.9, 153.3, 136.7, 135.5, 133.4, 130.8, 130.1, 129.5, 129.3, 128.9, 128.6, 127.6, 127.3, 126.4, 113.7, 97.8, 80.9, 74.2, 71.7, 70.1, 66.2, 56.3, 55.2, 40.2, 38.4, 37.6, 32.7, 29.9, 19.7, 10.8, 9.2, 9.1; IR (film) 2976, 2934,

1779, 1709, 1612, 1514, 1454, 1381, 1356, 1300, 1248, 1208 cm^{-1} ; Exact mass calcd. for $\text{C}_{39}\text{H}_{47}\text{N}_1\text{O}_7\text{Na}_1$ ($\text{M}+\text{Na}^+$): 664.3250; found: 664.3253 (ES^+).

2-(6-((3E)(1R,2R)-2-[(4-methoxyphenyl)methoxy]-1-methyl-4-phenylbut-3-enyl)-(4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl))(2R)propan-1-one (25). A solution of the imide **24** (440 mg, 0.686 mmol, 1 equiv) in THF (15 mL) was cooled to 0 °C. 2 M LiBH_4 (1.72 mL, 3.428 mmol, 5 equiv) in THF solution was added dropwise to the solution. Then MeOH (109 mg, 3.428 mmol, 5 equiv) was added slowly. After 5 min at rt, the ice-water bath was removed and the solution was stirred at rt for 24 h. The reaction was then cooled to -78 °C and quenched with 15 mL of satd. NH_4Cl . After warming up to rt, the precipitate was dissolved by adding just enough 1 M NaHSO_4 solution. The resulting aqueous solution was extracted with 3 $\times$ 30 mL ether, dried over MgSO_4 and concentrated to afford the crude alcohol.

The crude alcohol from the previous step was dissolved in 10 mL of CH_2Cl_2 . Pyridine (0.542 g, 6.856 mmol, 10 equiv) and Dess-Martin periodinane (0.874 g, 2.06 mmol, 3 equiv) were added to the solution. The slightly yellow solution was stirred at rt for 3 h. The reaction mixture was then diluted with 150 mL of ether and washed with a mixed solution of 1 M NaS_2O_3 and satd. NaHCO_3 (40 mL, 1:1 v/v). The organic phase was then washed with 2 $\times$ 20 mL satd. CuSO_4 , dried over MgSO_4 and concentrated. Purification by flash chromatography (10%–20% EtOAc/hexanes) afforded the aldehyde **25** (0.306 g, 95%) as a colorless oil: TLC R_f = 0.30 (20% EtOAc/hexanes); $[\alpha]_D^{25}$ -42.7° (c 0.57, CHCl_3); IR (film) 2978, 2929, 2871, 1729, 1608, 1512, 1449, 1381, 1303, 1250, 1197, 1173, 1057, 1033, 970 cm^{-1} ; ^1H NMR (400 MHz, C_6D_6) δ 9.40 (s, 1H), 7.26 (m, 3H), 7.01–7.05 (m, 4H), 6.81 (d, J = 8.5 Hz, 2H), 6.40 (d, J = 15.7 Hz, 1H), 6.06 (dd, J = 16.1, 8.8 Hz, 1H), 4.59 (d, J = 11.0 Hz, 1H), 4.24 (d, J = 11.0 Hz, 1H), 4.20 (dd, J = 10.2, 1.8 Hz, 1H), 3.94 (t, J = 9.2 Hz, 1H), 3.84 (dd, J = 10.3, 2.6 Hz, 1H), 3.27 (s, 3H), 1.90–1.97 (m, 2H), 1.58–1.65 (m, 1H), 1.38 (s, 3H), 1.31 (s, 3H), 1.05 (d, J = 7.0 Hz, 3H), 0.89 (d, J = 7.0 Hz, 3H), 0.39 (d, J = 7.0 Hz, 3H); ^{13}C NMR (100 MHz, C_6D_6) δ 202.6, 159.7, 137.1, 134.1, 131.5, 130.4, 129.6, 128.8, 127.9, 126.8, 114.0, 98.3, 81.2, 73.9, 72.3, 70.5, 54.8, 47.3, 39.0, 31.9, 30.1, 19.9, 10.8, 9.5, 6.3; Exact mass calcd. for $\text{C}_{29}\text{H}_{42}\text{NO}_5$ [$\text{M}+\text{NH}_4$] $^+$: 484.3063; found: 484.3072 (CI^+).

6-(6-((3E)(1R,2R)-2-[(4-methoxyphenyl)methoxy]-1-methyl-4-phenylbut-3-enyl)-(6S,-4R,5R)-2,2,5-trimethyl(1,3-dioxan-4-yl))(6S,5R)-5-hydroxyheptan-3-one. A flame dried 10 mL flask was charged with (-)-DIP-Cl (0.378 g, 1.180 mmol, 1.8 equiv) in the

glovebox. The solid was then dissolved in 3 mL of diethyl ether and cooled to $-78\text{ }^{\circ}\text{C}$. NEt_3 (0.166 g, 1.639 mmol, 2.5 equiv) and 2-butanone (70.8 mg, 0.984 mmol, 1.5 equiv) were added into the solution. After stirring at $-78\text{ }^{\circ}\text{C}$ for 10 min and $0\text{ }^{\circ}\text{C}$ for 1.5 h, the suspension was cooled back to $-78\text{ }^{\circ}\text{C}$. The aldehyde **25** (0.306 g, 0.656 mmol, 1.0 equiv) in 2 mL of ether was added dropwise. The flask was rinsed with $2 \times 1.5\text{ mL}$ of ether and the rinses were added into the reaction as well. The reaction mixture was stirred at $-78\text{ }^{\circ}\text{C}$ for 7 h, then at $-20\text{ }^{\circ}\text{C}$ overnight. After placing in a $0\text{ }^{\circ}\text{C}$ bath, the reaction was quenched with a mixed solution of 3 mL of pH 7 buffer, 3 mL of MeOH and 3 mL of 30% H_2O_2 and stirred at $0\text{ }^{\circ}\text{C}$ for 1 h. The mixture was then extracted with $3 \times 25\text{ mL}$ of CH_2Cl_2 . The combined organics were washed with 20 mL of 1 M $\text{Na}_2\text{S}_2\text{O}_3$, dried over MgSO_4 and concentrated. Purification by flash chromatography (17% EtOAc /Hexanes) afforded the aldol adduct (0.273 g, 77%) as a colorless oil: TLC $R_f = 0.25$ (20% EtOAc /hexanes); $[\alpha]_D^{25} -10.6^{\circ}$ (c 0.50, CHCl_3); IR (film) 3503, 2934, 1710, 1614, 1512, 1456, 1375, 1355, 1304, 1247, 1137, 1171, 1111, 1065, 1035, 974, 821, 750, 694 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.43 (d, $J = 7.3\text{ Hz}$, 2H), 7.35 (d, $J = 7.3\text{ Hz}$, 2H), 7.25 (m, 3H), 6.87 (d, $J = 8.8\text{ Hz}$, 2H), 6.57 (d, $J = 16.1\text{ Hz}$, 1H), 6.07 (dd, $J = 16.1, 8.8\text{ Hz}$, 1H), 4.52 (d, $J = 11.2\text{ Hz}$, 1H), 4.27 (d, $J = 10.8\text{ Hz}$, 1H), 4.27 (m, 1H), 4.08-4.10 (m, 1H), 3.80-3.86 (m, 2H), 3.80 (s, 3H), 2.72 (dd, $J = 15.6, 8.3\text{ Hz}$, 1H), 2.62-2.64 (m, 1H), 2.46-2.53 (m, 2H), 2.41 (dd, $J = 15.6, 4.5\text{ Hz}$, 1H), 1.90-1.93 (m, 1H), 1.76-1.81 (m, 1H), 1.62-1.73 (m, 1H), 1.43 (s, 3H), 1.33 (s, 3H), 1.07 (t, $J = 17.3\text{ Hz}$, 3H), 0.94 (d, $J = 7.3\text{ Hz}$, 3H), 0.81 (d, $J = 6.8\text{ Hz}$, 3H), 0.70 (d, $J = 6.3\text{ Hz}$, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 210.9, 159.0, 136.6, 133.5, 130.7, 129.9, 129.3, 128.5, 127.6, 126.4, 113.7, 97.9, 80.8, 79.4, 72.5, 71.8, 70.1, 55.2, 47.0, 38.3, 37.1, 37.0, 31.8, 30.0, 19.9, 10.9, 9.2, 7.5, 5.5; Exact mass calcd. for $\text{C}_{33}\text{H}_{46}\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$: 561.3192; found: 561.3191 (ES^+).

6-(6-((3E)(1R,2R)-2-[(4-methoxyphenyl)methoxy]-1-methyl-4-phenylbut-3-enyl))(4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl))(5R,6R)-5-(tert-butyldimethylsiloxy) heptan-3-one (4). A solution of the alcohol (0.27 g, 0.501 mmol, 1 equiv) in 10 mL of CH_2Cl_2 was cooled to $-78\text{ }^{\circ}\text{C}$. 2,6-lutidine (0.215 g, 2.005 mmol, 4 equiv) and TBSOTf (0.265 g, 1.003 mmol, 2 equiv) were added. The resulting solution was stirred at $-78\text{ }^{\circ}\text{C}$ for 3 h and then quenched with 10 mL of 1 M NaHSO_4 and warmed up to rt over 30 min. The reaction mixture was extracted with $3 \times 30\text{ mL}$ of CH_2Cl_2 . The combined organics were washed with 20 mL of 1 M NaHSO_4 , dried over MgSO_4 and concentrated. Purification by flash chromatography (5% EtOAc /Hexanes) afforded the silyl ether **4** (0.24 g, 77%) as a colorless oil: TLC $R_f = 0.65$ (20% EtOAc /hexanes); $[\alpha]_D^{25} +2.61^{\circ}$ (c 0.72, CHCl_3); IR (film) 2935, 2857, 1716, 1614, 1587, 1514, 1463, 1380, 1252, 1202, 1174, 1078, 1040, 832, 777, 750 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.44 (d, $J = 7.7\text{ Hz}$, 2H), 7.35 (t, $J = 7.7\text{ Hz}$, 2H), 7.25-7.30 (m, 3H), 6.89 (d, $J = 8.4\text{ Hz}$, 2H), 6.58 (d, $J = 16.1\text{ Hz}$, 1H), 6.11 (dd, $J = 16.1, 8.4\text{ Hz}$, 1H), 4.55 (d, $J = 11.0\text{ Hz}$, 1H), 4.33 (d, $J = 11.0\text{ Hz}$, 1H), 4.14-4.20 (m, 1H), 4.07-4.13 (m, 1H), 3.98 (d, $J = 9.9\text{ Hz}$, 1H), 3.89 (t, $J = 9.1\text{ Hz}$, 1H), 3.81 (s, 3H), 2.71 (dd, $J = 15.4, 8.4\text{ Hz}$, 1H), 2.40-2.55 (m, 3H), 1.88-1.98 (m, 2H), 1.57-1.66 (m, 1H), 1.45 (s, 3H), 1.34 (s, 3H),

1.05 (t, $J = 7.3$ Hz, 3H), 0.91 (s, 9H), 0.86 (d, $J = 7.3$ Hz, 3H), 0.83 (d, $J = 7.0$ Hz, 3H), 0.72 (d, $J = 6.6$ Hz, 3H), 0.07 (s, 3H), 0.01 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 211.7, 159.0, 136.7, 133.2, 130.9, 130.1, 129.4, 128.5, 127.6, 126.4, 113.6, 97.2, 80.9, 72.2, 72.0, 71.7, 70.1, 55.2, 45.9, 38.9, 38.6, 38.0, 31.9, 30.2, 25.8, 19.9, 17.9, 11.5, 9.8, 9.3, 7.5, -4.5, -4.9; Exact mass calcd. for $\text{C}_{39}\text{H}_{60}\text{O}_6\text{SiNa}$ $[\text{M}+\text{Na}]^+$: 675.4057; found: 675.4050 (ES^+).

(6S,7S,5R)-5,7-bis(*tert*-butyldimethylsiloxy)-6,8-dimethylnonan-3-one (26).⁵ (-)-(ipc)₂BCl (DIP-chloride, 0.481 g, 1.5 mmol, 1.5 equiv) was charged into a 25 mL round bottom flask in the glove box. After adding 8 mL of diethyl ether, the solution was cooled to -78 °C. NEt_3 (0.202 g, 2.0 mmol, 2.0 equiv) and 2-butanone (86.4 mg, 1.2 mmol, 1.2 equiv) were added into the solution. The resulting solution was then warmed up and stirred at 0 °C for 1 h. After cooling back to -78 °C, the aldehyde (0.244 g, 1 mmol, 1.0 equiv, azeotroped with 2×5 mL dry benzene) in 1.5 mL of ether was added dropwise into the solution. The flask was then rinsed with 2×0.25 mL of ether and the rinses were added into the reaction mixture. After stirring at -78 °C for 10 min, the flask was placed into a -20 °C bath overnight. The reaction was quenched with a mixed solution of 3 mL of pH 7 buffer, 3 mL of MeOH and 3 mL of 30% H_2O_2 . The resulting solution was stirred at 0 °C for 1 h, diluted with 10 mL of water and extracted with 3×25 mL of ether. The combined organic phase was washed with 2×10 mL of 1 M $\text{Na}_2\text{S}_2\text{O}_3$, dried over MgSO_4 and concentrated. Purification by flash chromatography (5%–10% EtOAc/Hexanes) afforded the aldol adduct (0.424 g, 68%) as a colorless oil: TLC $R_f = 0.56$ (20% EtOAc/hexanes).

The aldol adduct (0.390 g, 1.232 mmol, 1 equiv) was dissolved in 10 mL of CH_2Cl_2 . After cooling to -78 °C, 2,6-lutidine (0.396 g, 3.697 mmol, 3.0 equiv) and TBSOTf (0.489 g, 1.849 mmol, 1.5 equiv) were added. After stirring at -78 °C for 2 h, the reaction was quenched with 5 mL of satd. NaHCO_3 and warmed up to rt. The reaction mixture was then extracted with 3×15 mL of CH_2Cl_2 , dried over MgSO_4 and concentrated. Purification by flash chromatography (4% EtOAc/Hexanes) afforded the silyl ether **26** (0.415 g, 78%) as a colorless oil. TLC $R_f = 0.86$ (20% EtOAc/hexanes); $[\alpha]^{22}_{\text{D}} +20.8^\circ$ (c 0.53, CHCl_3); IR (film) 2943, 2862, 1717, 1465, 1410, 1385, 1359, 1254, 1213, 1047, 833, 773, 674 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 4.18 (dt, $J = 5.9, 3.9$ Hz, 1H), 3.58 (dd, $J = 5.4, 2.9$ Hz, 1H), 2.57 (d, $J = 5.9$ Hz, 2H), 2.41 (ddd, $J = 14.6, 7.3, 1.5$ Hz, 2H), 1.88 (m, 1H), 1.62 (m, 1H), 1.02 (t, $J = 7.3$ Hz, 3H), 0.90 (d, $J = 6.8$ Hz, 3H), 0.899 (s, 9H), 0.86 (d, $J = 5.9$ Hz, 3H), 0.85 (s, 9H), 0.83 (d, $J = 6.8$ Hz, 3H), 0.06 (s, 3H), 0.043 (s, 3H), 0.038 (s, 3H), -0.03 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 210.2, 76.0, 70.1, 47.9, 41.0, 37.6, 32.1, 26.2, 25.9, 19.8, 18.5, 18.0, 16.9, 11.4, 7.5, -3.4, -3.6, -4.4, -4.5; Exact mass calcd. for $\text{C}_{23}\text{H}_{51}\text{O}_3\text{Si}_2$ $[\text{M}+\text{H}]^+$: 431.3377; found: 431.3380 (ES^+).

⁵ Paterson, I.; Cumming, J. G.; Smith, J.D.; Ward, R.A. *Tetrahedron Lett.* **1994**, 35, 441.

(10S,11S,3R,4R,5R,6R,9R)-5-hydroxy-2,4,6,10,12-pentamethyl-3,9,11-tris (*tert*-butyl dimethylsiloxy)tridecan-7-one (28a). A solution of ketone **26** (86.2 mg, 0.20 mmol, 1 equiv) in 2 mL of diethyl ether was cooled to $-78\text{ }^{\circ}\text{C}$. NEt_3 (30.3 mg, 0.3 mmol, 1.5 equiv) and chlorodicyclohexylborane (42.5 mg, 0.2 mmol, 1.0 equiv) were added dropwise into the solution. After stirring at $-78\text{ }^{\circ}\text{C}$ for 10 min, the solution was warmed up to $0\text{ }^{\circ}\text{C}$ and stirred at this temperature for 1.5 h. After cooling back to $-78\text{ }^{\circ}\text{C}$, the aldehyde **27** (48.9 mg, 0.2 mmol, 1.0 equiv, azeotroped with $2 \times 3\text{ mL}$ of dry benzene) in 1 mL of ether was added dropwise. The flask was rinsed with $2 \times 0.5\text{ mL}$ ether and the rinses were added into the reaction mixture as well. The resulting suspension was stirred at $-78\text{ }^{\circ}\text{C}$ for 4 h, then at $-20\text{ }^{\circ}\text{C}$ overnight. The reaction mixture was placed in $0\text{ }^{\circ}\text{C}$ bath and quenched with a mixed solution of 2 mL of pH 7 buffer, 2 mL of MeOH and 2 mL of 30% H_2O_2 . After stirring at $0\text{ }^{\circ}\text{C}$ for 1 h, the reaction mixture was diluted with 5 mL of water and extracted with $3 \times 10\text{ mL}$ of CH_2Cl_2 . The combined organic solution was washed with 10 mL of 1 M $\text{Na}_2\text{S}_2\text{O}_3$, dried over MgSO_4 and concentrated. Purification by flash chromatography (20% $\square$ 50% CH_2Cl_2 /Hexanes) afforded the aldol adduct **28a** as a colorless oil (57 mg, 42%): TLC $R_f = 0.05$ (15% CH_2Cl_2 /hexanes); $[\alpha]^{22}_{\text{D}} - 3.0^{\circ}$ ($c\ 0.94$, CHCl_3); IR (film) 3534, 2955, 2924, 1710, 1468, 1385, 1370, 1253, 1106, 1050, 1004, 837, 771, 669 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 4.21-4.24 (m, 1H), 3.75 (ddd, $J = 8.8, 3.7, 2.6\text{ Hz}$, 1H), 3.58 (t, $J = 4.0\text{ Hz}$, 1H), 3.54 (dd, $J = 6.6, 2.6\text{ Hz}$, 1H), 2.88 (dd, $J = 7.9, 6.6\text{ Hz}$, 1H), 2.56-2.68 (m, 2H), 2.50 (d, $J = 4.0\text{ Hz}$, 1H), 1.86-1.93 (m, 2H), 1.63-1.73 (m, 2H), 0.99 (d, $J = 3.3\text{ Hz}$, 3H), 0.92 (d, $J = 4.0\text{ Hz}$, 3H), 0.91 (s, 9H), 0.90 (s, 9H), 0.90 (d, $J = 3.3\text{ Hz}$, 3H), 0.89 (d, $J = 2.2\text{ Hz}$, 3H), 0.87 (s, 9H), 0.86 (d, $J = 2.6\text{ Hz}$, 3H), 0.85 (d, $J = 1.5\text{ Hz}$, 3H), 0.84 (d, $J = 2.6\text{ Hz}$, 3H), 0.07 (s, 6H), 0.06 (s, 3H), 0.05 (s, 3H), 0.04 (s, 3H), 0.02 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 214.0, 79.9, 76.5, 76.0, 68.8, 50.0, 48.7, 40.9, 36.6, 32.3, 31.7, 26.3, 26.1, 25.9, 20.3, 19.4, 18.6, 18.4, 18.0, 17.5, 16.3, 13.3, 11.0, 8.5, -3.37, -3.39, -4.0, -4.3, -4.6; Exact mass calcd. for $\text{C}_{36}\text{H}_{78}\text{O}_5\text{Si}_3$ $[\text{M}+\text{NH}_4]^+$: 675.5235; found: 675.5240 (ES^+).

2-(6-((3E)(1R,2R)-2-[(4-methoxyphenyl)methoxy]-1-methyl-4-phenylbut-3-enyl)-(4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl))(2R,3R,6R,7R,8R,9R)-10-((4R,6R)-6-((2S)-3-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-2-[(4-methoxyphenyl)methoxy]propyl)-2,2-dimethyl(1,3-dioxan-4-yl))-3,9-bis(*tert*-butyldimethylsiloxy)-7-hydroxy-6,8-di-

methyl\decan-5-one (29). A vial containing the ketone **4** (115 mg, 0.176 mmol, 2 equiv, azeotroped with 3 × 3 mL of dry benzene) was charged with 1 mL diethyl ether. The solution was cooled to -78 °C. NEt₃ (26.7 mg, 0.264 mmol, 3.0 equiv) and chlorodicyclohexylborane (37.5 mg, 0.176 mmol, 2.0 equiv) were added dropwise. After stirring at -78 °C for 10 min, then at 0 °C for 1.5 h, the suspension was cooled back to -78 °C. The aldehyde **5** (60 mg, 0.088 mmol, 1.0 equiv, azeotroped with 3 × 3 mL of dry benzene) in 1 mL of ether was added slowly into the reaction mixture. The vial was rinsed with 2 × 0.2 mL of ether and the rinses were added as well. The resulting reaction mixture was stirred at -78 °C for 4 h, then at -20 °C for 17 h. The reaction was quenched by a mixed solution of 1.5 mL of pH 7 buffer, 1.5 mL of MeOH and 1.5 mL of 30% H₂O₂ and stirred at 0 °C for 1 h. After diluting with 100 mL of ether, the reaction mixture was washed with 10 mL of 1 M Na₂S₂O₃. The ethereal solution was dried over MgSO₄ and concentrated. Purification by flash chromatography afforded the ketone starting material **4** (5 % EtOAc/Hexanes, 78 mg), the major aldol adduct **29** as a colorless oil (10% EtOAc / Hexanes, 46 mg, 40%). Further elution with 20% EtOAc / Hexanes afforded the minor diastereomer **30** (10 mg, 9%).

The major diastereomer 29: TLC R_f = 0.70 (20% EtOAc/hexanes); [α]_D²² -13.4° (c 1.25, CHCl₃); IR (film) 3506, 2940, 2865, 1717, 1610, 1514, 1461, 1381, 1247, 1173, 1108, 1039, 836, 777, 687 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.42 (d, J = 7.3 Hz, 2H), 7.34 (t, J = 7.5 Hz, 2H), 7.25-7.28 (m, 5H), 6.86 (dd, J = 8.8, 2.4 Hz, 4H), 6.55 (d, J = 16.1 Hz, 1H), 6.08 (dd, J = 16.1, 8.8 Hz, 1H), 4.65 (d, J = 11.2 Hz, 1H), 4.52 (d, J = 11.2 Hz, 1H), 4.50 (d, J = 11.7 Hz, 1H), 4.31 (d, J = 11.2 Hz, 1H), 4.20-4.26 (m, 1H), 4.02-4.09 (m, 2H), 3.92-3.98 (m, 2H), 3.82-3.92 (m, 2H), 3.80 (s, 3H), 3.78 (m, 2H), 3.70 (dd, J = 10.4, 4.5 Hz, 1H), 3.54-3.60 (m, 1H), 3.05 (bs, 1H), 2.86 (dd, J = 17.6, 7.3 Hz, 1H), 2.75 (dd, J = 17.6, 3.4 Hz, 1H), 2.63 (dd, J = 9.8, 7.3 Hz, 1H), 1.88-1.95 (m, 2H), 1.53-1.77 (m, 6H), 1.43 (s, 3H), 1.35 (s, 3H), 1.33 (s, 6H), 1.24-1.30 (m, 2H), 1.04-1.12 (m, 21H), 0.94 (d, J = 7.8 Hz, 3H), 0.88 (s, 9H), 0.87 (s, 9H), 0.85 (d, J = 4.9 Hz, 3H), 0.82 (d, J = 7.3 Hz, 3H), 0.81 (d, J = 6.8 Hz, 3H), 0.69 (d, J = 6.8 Hz, 3H), 0.08 (s, 6H), 0.07 (s, 3H), 0.05 (s, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 214.2, 159.3, 159.2, 137.1, 133.5, 131.3, 131.2, 130.5, 129.8, 129.7, 129.6, 128.8, 127.8, 126.7, 113.95, 113.91, 113.88, 98.5, 97.5, 81.3, 76.2, 76.1, 75.3, 74.5, 72.6, 72.3, 71.7, 70.9, 70.4, 66.52, 66.45, 65.9, 64.5, 55.5, 50.9, 46.1, 41.8, 41.2, 39.0, 38.8, 37.7, 37.0, 36.4, 32.2, 30.6, 30.4, 30.2, 26.3, 26.1, 26.0, 20.2, 19.8, 19.7, 18.3, 18.2, 13.3, 12.1, 11.7, 10.0, 9.6, 6.2, -3.5, -4.2, -4.36, -4.38, -4.7, -4.8; Exact mass calcd. for C₇₆H₁₂₈O₁₃Si₃Na [M+Na]⁺: 1357; found: 1357 (FAB⁺).

The minor diastereomer 30: ¹H NMR (500 MHz, C₆D₆) δ 7.26-7.30 (m, 5 H), 7.12-7.14 (m, 3H), 7.04-7.07 (M, 1H), 6.81 (d, J = 8.3 Hz, 2H), 6.80 (d, J = 8.8 Hz, 2H), 6.43 (d, J = 16.1 Hz, 1H), 6.10 (dd, J = 16.1, 8.8 Hz, 1H), 4.69 (d, J = 11.7 Hz, 1H), 4.62 (d, J = 11.2 Hz, 1H), 4.48-4.54 (m, 2H), 4.32-4.39 (m, 2H), 4.24-4.32 (m, 2H), 4.19 (d, J = 10.3 Hz, 1H), 3.98-4.05 (m, 2H), 3.90 (dd, J = 10.3, 5.9 Hz, 1H), 3.79 (dd, J = 10.3, 4.4 Hz, 1H), 3.66-3.74 (m, 2H), 3.46 (d, J = 2.0 Hz, 1H), 3.33 (s, 3H), 3.31 (s, 3H), 3.17 (dd, J = 18.1, 8.8 Hz, 1H), 2.96 (dd, J = 6.9, 4.4 Hz, 1H), 2.77 (dd, J = 18.1, 2.0 Hz, 1H), 1.85-2.08 (m, 4H), 1.70-1.78 (m, 3H), 1.59-1.66 (m, 1H), 1.58 (s, 3H), 1.56 (s, 3H), 1.50 (s, 3H), 1.44 (d, 7.3 Hz, 3H), 1.38 (s, 3H), 1.23 (d, J = 6.8 Hz, 3H), 1.24-1.50 (m, 2H), 1.08-1.14 (m, 21H), 1.00 (d, J = 6.8 Hz, 3H), 0.98 (s, 9H), 0.95 (s, 9H), 0.92 (d, J = 7.3 Hz, 3H), 0.72 (d, J = 6.3 Hz, 3H), 0.19 (s, 3H), 0.13 (s, 3H), 0.12 (s, 6H); ¹³C NMR (125 MHz, C₆D₆) δ 214.2, 159.7, 159.6, 137.2, 133.9, 131.7, 131.5, 130.6, 129.8, 129.5, 128.8, 128.3, 126.9, 114.0, 113.99, 113.95, 98.5, 97.7, 81.2, 76.0,

74.1, 73.0, 72.2, 71.7, 71.0, 70.4, 66.9, 66.34, 66.28, 54.8, 54.7, 49.3, 46.4, 42.1, 39.3, 39.1, 39.0, 38.0, 32.3, 30.6, 30.5, 26.24, 26.18, 26.1, 20.2, 19.7, 18.3, 18.2, 18.1, 12.3, 11.9, 10.7, 10.1, 9.7, 8.2, -3.3, -4.2, -4.4, -4.6.

2-[6-(3-[6-(3-(6-[(3E)(1R,2R)-2-[(4-methoxyphenyl)methoxy]-1-methyl-4-phenylbut-3-enyl)](4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl))](2R,3R)-2-(tert-butyldimethylsiloxy)butyl)](4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl)](2R,3R)-2-(tert-butyldimethylsiloxy)butyl](4R,6R)-2,2-dimethyl(1,3-dioxan-4-yl)](1S)-1-[[triisopropylsiloxy]methyl]-1-[(4-methoxyphenyl)methoxy]ethane (31). A solution of the hydroxyl ketone **29** (100 mg, 74.96 μmol , 1 equiv) was dissolved in 3 mL of CH_2Cl_2 . After cooling to 0 $^\circ\text{C}$, $\text{Zn}(\text{BH}_4)_2$ (4 mL, 0.60 mmol, 8 equiv) in ether solution (0.15 M) was added dropwise into the solution. The resulting solution was stirred at 0 $^\circ\text{C}$ for 2 h. Another batch of $\text{Zn}(\text{BH}_4)_2$ (4 mL, 0.60 mmol, 8 equiv) was added and stirred at 0 $^\circ\text{C}$ for another 2 h. After adding another batch of $\text{Zn}(\text{BH}_4)_2$ (4 mL, 0.60 mmol, 8 equiv) and stirring for 2 h, the reaction was complete and carefully quenched with 5 mL of satd. NH_4Cl and extracted with 3 $\times$ 20 mL of CH_2Cl_2 . The combined organics were dried over MgSO_4 and concentrated. Purification by flash chromatography (10% EtOAc / Hexanes) afforded the desired *syn*-diol as a colorless oil (70 mg, 70%): TLC R_f = 0.50 (20% EtOAc /hexanes).

The diol (53 mg, 39.7 μmol , 1 equiv) was dissolved in 3 mL of CH_2Cl_2 and 1 mL of 2,2-dimethoxypropane. A few particles of PPTs were added. The resulting solution was stirred at rt for 1 h before being concentrated. Purification by flash chromatography (10% EtOAc / Hexanes) afforded the acetonide **31** as a colorless oil (59 mg, 100%): TLC R_f = 0.80 (20% EtOAc /hexanes); $[\alpha]_D^{25}$ -19.3 $^\circ$ (c 1.0, CHCl_3); IR (film) 2939, 2862, 1613, 1512, 1458, 1381, 1250, 1197, 1173, 1105, 1037, 834, 771, 689 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.40 (d, J = 7.3 Hz, 2H), 7.32 (t, J = 7.3 Hz, 2H), 7.24 (m, 5H), 6.84 (d, J = 8.9 Hz, 2H), 6.83 (d, J = 8.8 Hz, 2H), 6.54 (d, J = 16.1 Hz, 1H), 6.06 (dd, J = 16.1, 8.5 Hz, 1H), 4.61 (d, J = 11.0 Hz, 1H), 4.48 (t, J = 11.8 Hz, 2H), 4.28 (d, J = 11.0 Hz, 1H), 4.05-4.13 (m, 1H), 4.03 (dd, J = 8.8, 2.0 Hz, 1H), 3.94-3.99 (m, 1H), 3.81-3.86 (m, 2H), 3.78 (s, 3H), 3.57-3.78 (m, 7H), 1.41-1.94 (m, 13H), 1.37 (s, 9H), 1.32 (s, 3H), 1.27 (s, 3H), 1.26 (s, 3H), 1.04-1.05 (m, 21H), 0.88 (s, 9H), 0.87 (s, 9H), 0.87 (d, J = 4.4 Hz, 3H), 0.84 (d, J = 7.3 Hz, 3H), 0.78 (d, J = 7.0 Hz, 3H), 0.69 (d, J = 6.6 Hz, 3H), 0.66 (d, J = 6.6 Hz, 3H), 0.028 (s, 3H), 0.024 (s, 3H), 0.018 (s, 3H), 0.010 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 159.0, 136.8, 133.1, 131.2, 131.0, 130.3, 129.3, 129.2, 128.6, 127.5, 126.4, 113.7, 113.6, 98.2, 97.3, 97.2, 81.0, 77.2, 73.2, 73.0, 72.2, 72.1, 71.8, 71.6, 71.0, 70.2, 66.4, 66.0, 65.3, 55.2, 50.9, 40.2, 38.7, 38.5, 38.3, 38.2, 38.0, 37.4, 36.2, 32.4, 30.9, 30.4, 30.3, 30.1, 29.7, 26.0, 25.9, 25.8, 20.0, 19.8, 19.6, 18.0, 12.0, 11.9, 11.1, 10.1, 10.0, 9.3, -3.9, -4.1, -4.5, -4.6; Exact mass calcd. for $\text{C}_{79}\text{H}_{134}\text{O}_{13}\text{Si}_3\text{Na}$ $[\text{M}+\text{Na}]^+$: 1399.2; found: 1399.1 (ES^+).

(1E)-4-[6-((1R,2R)-3-{6-[(1R,2R)-3-((4R,6R)-6-{(2S)-3-[1,1-bis(methylethyl)-2-methyl-1-silapropoxy]-2-hydroxypropyl)-2,2-dimethyl(1,3-dioxan-4-yl)]-1-methyl-2-(*tert*-butyldimethylsiloxy)propyl}(4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl))-1-methyl-2-(*tert*-butyldimethylsiloxy)propyl) (6S,4R, 5R)-2,2,5-trimethyl(1,3-dioxan-4-yl)](4S)-1-phenylpent-1-en-3-one (**32**). The PMB ether **31** (59 mg, 39.7 μ mol, 1 equiv) was dissolved in 4 mL of CH_2Cl_2 and 0.2 mL of water. DDQ (36 mg, 0.159 mmol, 4 equiv) was added. The resulting reaction mixture was stirred at rt for 30 min. Then 2 g of silica gel was added into the reaction mixture. The reaction was carefully concentrated. Purification by flash chromatography (10% EtOAc/Hexanes) afforded the enone **32** as a colorless oil (29 mg, 65%): TLC R_f = 0.70 (20% EtOAc/hexanes); IR (film) 2939, 2851, 1685, 1608, 1453, 1381, 1250, 1197, 1168, 1119, 1066, 998, 833, 766 cm^{-1} ; ^1H NMR (400 MHz, C_6D_6) δ 7.79 (d, J = 15.7 Hz, 1H), 7.21 (t, J = 3.7 Hz, 2H), 7.02 (m, 3H), 6.96 (d, J = 16.1 Hz, 1H), 4.28-4.36 (m, 1H), 4.24 (dd, J = 9.9, 3.3 Hz, 1H), 4.16-4.22 (m, 1H), 4.00-4.14 (m, 5H), 3.90-3.96 (m, 1H), 3.77 (dd, J = 9.9, 6.2 Hz, 1H), 3.67 (dd, J = 9.9, 5.2 Hz, 1H), 3.13 (bs, 1H), 2.72-2.80 (m, 1H), 2.16-2.24 (m, 1H), 2.02-2.14 (m, 3H), 1.72-1.90 (m, 5H), 1.60-1.70 (m, 1H), 1.50-1.58 (m, 2H), 1.48 (s, 6H), 1.45 (s, 3H), 1.40 (s, 3H), 1.39 (s, 3H), 1.33 (s, 3H), 1.15 (t, J = 7.0 Hz, 6H), 1.06-1.08 (m, 24H), 1.02 (s, 9H), 1.01 (s, 9H), 0.82 (d, J = 6.6 Hz, 3H), 0.75 (d, J = 6.6 Hz, 3H), 0.12 (s, 3H), 0.08 (s, 3H), 0.07 (s, 3H), 0.05 (s, 3H); ^{13}C NMR (100 MHz, C_6D_6) δ 199.3, 141.8, 135.4, 130.0, 129.0, 128.4, 128.1, 127.9, 124.4, 98.6, 98.1, 97.8, 75.3, 73.6, 73.5, 72.8, 72.3, 71.5, 71.0, 69.0, 68.0, 65.8, 47.9, 40.8, 40.3, 38.9, 38.5, 37.6, 36.8, 34.1, 30.7, 30.6, 30.2, 26.3, 26.2, 19.9, 19.8, 19.5, 18.3, 18.2, 12.5, 12.2, 11.8, 10.6, 9.7, -3.8, -4.0, -4.3, -4.4; Exact mass calcd. for $\text{C}_{63}\text{H}_{116}\text{O}_{11}\text{Si}_3\text{Na}$ $[\text{M}+\text{Na}]^+$: 1155.8; found: 1155.9 (ES^+).

(2R)-2-{6-[(1R,2R)-3-(6-[(1R,2R)-3-[(4R,6R)-6-(2-hydroxyethyl)-2,2-di-methyl(1,3-dioxan-4-yl)]-1-methyl-2-(*tert*-butyldimethylsiloxy)propyl}(4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl))-1-methyl-2-(1,1,2,2-tetramethyl-1-silapropoxy)propyl](4S,5S,6S)-2,2,5-trimethyl(1,3-dioxan-4-yl))} propan-1-ol (**33**). The enone **32** (29 mg, 25.6 μ mol, 1 equiv)

was dissolved in 4 mL of CH_2Cl_2 and 0.5 mL of MeOH and cooled to -78°C . O_3 was passed through until purple color persisted. O_2 was then bubbled through until the solution was colorless. NaBH_4 (20 mg, 0.52 mmol, 20 equiv) was added. The solution was slowly warmed up to rt over 3-4 h and concentrated. The residue was dissolved in 2 mL of EtOAc, passed through a pipette of silica gel column, washed with 20 mL of EtOAc and concentrated to afford a crude triol which was taken to the next step without any further purification: TLC R_f = 0.80 (50% EtOAc/hexanes).

The residue from the last step was dissolved in 3 mL of THF. TBAF (0.25 mL, 0.25 mmol, 10 equiv) as a 1 M solution in THF was added. After stirring at rt for 1 h, the reaction mixture was concentrated. Purification by flash chromatography (pipette column, 0%–10% MeOH/EtOAc) afforded the tetraol as a colorless oil which was taken to the next step: TLC R_f = 0.05 (50% EtOAc/hexanes).

The tetraol was dissolved in 3 mL of MeOH. NaIO_4 (55 mg, 0.26 mmol, 10 equiv) in 0.55 mL of water was added. After stirring at rt overnight, 30 μL of ethylene glycol was added to quench the excess NaIO_4 and stirred at rt for 2 h. The reaction mixture was concentrated, passed through a pipette column of silica gel, eluted with 15 mL of EtOAc and concentrated. The residue was dissolved in 3 mL of CH_2Cl_2 and 0.5 mL of MeOH. After cooling to -78°C , NaBH_4 (20 mg, 0.52 mmol, 20 equiv) was added. The reaction mixture was slowly warmed up to rt over 2-3 h and concentrated. Purification by flash chromatography (20%–50% EtOAc / Hexanes) afforded the diol **33** as a colorless oil (15 mg, 69% over 3 steps): TLC R_f = 0.74 (50% EtOAc/hexanes); $[\alpha]_D^{22} -9.3^\circ$ (c 0.75, CHCl_3); IR (film) 3422, 2933, 2856, 1468, 1378, 1256, 1194, 1172, 1061, 833, 767, 672 cm^{-1} ; ^1H NMR (400 MHz, C_6D_6) δ 4.26-4.33 (m, 1H), 4.12 (dd, J = 10.3, 1.8 Hz, 1H), 4.03-4.10 (m, 2H), 3.96-4.01 (m, 1H), 3.86-3.94 (m, 2H), 3.74 (dd, J = 10.2, 2.2 Hz, 1H), 3.64-3.72 (m, 1H), 3.60-3.64 (m, 2H), 3.56 (dd, J = 10.6, 4.8 Hz, 1H), 2.15-2.19 (m, 1H), 2.04-2.13 (m, 3H), 1.84-1.90 (m, 2H), 1.74-1.84 (m, 2H), 1.64-1.74 (m, 5H), 1.48 (s, 6H), 1.44 (s, 3H), 1.37 (s, 3H), 1.36 (s, 6H), 1.16 (d, J = 6.9 Hz, 3H), 1.15 (d, J = 6.9 Hz, 3H), 1.02 (s, 9H), 1.01 (s, 9H), 1.01-1.02 (d, J = 6.7 Hz, 3H), 0.83 (d, J = 6.6 Hz, 3H), 0.64 (d, J = 6.6 Hz, 3H), 0.12 (s, 3H), 0.10 (s, 3H), 0.08 (s, 3H), 0.05 (s, 3H); ^{13}C NMR (100 MHz, C_6D_6) δ 98.6, 97.8, 97.7, 76.5, 73.6, 73.5, 72.8, 72.4, 71.6, 69.0, 66.9, 65.6, 60.3, 40.7, 39.1, 38.9, 38.3, 37.7, 36.9, 35.9, 32.8, 30.7, 30.6, 30.3, 26.3, 26.2, 19.81, 19.78, 18.27, 18.26, 12.4, 11.3, 10.6, 10.5, 9.4, -3.8, -4.0, -4.3, -4.4; Exact mass calcd. for $\text{C}_{45}\text{H}_{91}\text{O}_{10}\text{Si}_2$ $[\text{M}+\text{H}]^+$: 847.6151; found: 847.6155 (ES^+).

(3*S*,4*S*,6*S*,9*S*,10*S*,12*S*,2*R*,5*R*,7*R*,11*R*,13*R*,15*R*,17*R*)-2,4,6,10,12-pentamethyl-nonadecane-1,3,5,7,9,11,13,15,17,19-decaol (**2**). The diol **33** (15 mg, 17.7 μmol) was dissolved in 4 mL of AcOH and 1 mL of water and stirred at rt for 24 h. The reaction mixture was then carefully concentrated with 4 $\times$ 5 mL toluene (35-45 torr, 35°C). The residue was further concentrated on high vacuum pump for 2 h to afford the essentially pure decaol

2 as a white solid (8.9 mg, 100%): $[\alpha]^{22}_{\text{D}} -2.47^\circ$ (c 0.59, EtOH); ^1H NMR (500 MHz, CD_3OD) δ 3.92-3.96 (m, 2H), 3.82-3.91 (m, 3H), 3.75 (dd, J = 8.3, 2.0 Hz, 1H), 3.68 (dd, J = 8.8, 2.0 Hz, 1H), 3.61 (t, J = 6.5 Hz, 2H), 3.57 (dd, J = 9.8, 1.5 Hz, 1H), 3.48 (dd, J = 10.8, 7.3 Hz, 1H), 3.39 (dd, J = 10.3, 5.9 Hz, 1H), 1.30-1.90 (m, 13H), 0.86 (d, J = 7.3 Hz, 3H), 0.85 (d, J = 7.3 Hz, 3H), 0.79 (d, J = 7.3 Hz, 3H), 0.74 (d, J = 6.8 Hz, 3H), 0.69 (d, J = 6.8 Hz, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 81.4, 78.6, 77.6, 77.0, 75.8, 73.7, 70.4, 68.8, 66.5, 60.0, 45.4, 43.1, 42.6, 41.0, 40.1, 39.9, 39.4, 38.9, 36.6, 13.4, 11.4, 9.7, 6.5, 6.2; Exact mass calcd. for $\text{C}_{24}\text{H}_{51}\text{O}_{10}$ $[\text{M}+\text{H}]^+$: 499.3482; found: 499.3481 (ES^+).

(3S,4S,6S,9S,10S,12S,2R,5R,7R,11R,13R,15R,17R)-3,5,7,9,11,13,15,17,19-nonaacetyloxy-2,4,6,10,12-pentamethylnonadecyl acetate (34). The decaol **2** (8.8 mg, 17.7 μmol) was dissolved in 4 mL of pyridine and 1 mL of Ac_2O . 2 mg of DMAP was also added. The resulting solution was stirred at rt for 28 h and then concentrated with 6×5 mL toluene (35-39 $^\circ\text{C}$, 40-45 torr). Purification by flash chromatography (20% $\square$ 50% $\square$ 100% EtOAc/Hexanes) afforded the desired polyacetate **34** as a colorless oil (11.5 mg, 70%): TLC R_f = 0.15 (50% EtOAc / hexanes); $[\alpha]^{22}_{\text{D}} +10.0^\circ$ (c 0.80, MeOH); ^1H NMR (500 MHz, CD_3OD) δ 4.88-5.10 (m, 1H), 4.80-4.88 (m, 7H), 4.06-4.10 (m, 2H), 3.94 (dd, J = 11.2, 7.3 Hz, 1H), 3.82 (dd, J = 11.2, 6.3 Hz, 1H), 2.15-2.24 (m, 2H), 2.12 (m, 1H), 2.05-2.13 (m, 3H), 1.74-1.96 (m, 8H), 0.96 (d, J = 6.8 Hz, 3H), 0.95 (d, J = 6.8 Hz, 3H), 0.94 (d, J = 6.8 Hz, 3H), 0.91 (d, J = 6.8 Hz, 3H), 0.90 (d, J = 6.8 Hz, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 172.8, 172.7, 172.5, 172.42, 172.38, 172.33, 172.30, 172.27, 172.21, 172.16, 76.5, 75.9, 75.1, 75.0, 73.4, 73.2, 70.2, 69.8, 67.9, 61.7, 40.0, 39.6, 39.3, 39.1, 38.3, 35.3, 34.2, 32.8, 21.4, 21.31, 21.28, 21.1, 20.9, 20.8, 14.6, 11.33, 11.29, 9.6, 9.0; Exact mass calcd. for $\text{C}_{44}\text{H}_{71}\text{O}_{20}$ $[\text{M}+\text{H}]^+$: 919.4538; found: 919.4531 (ES^+).

Stereochemical Proofs

Compound 28

H _n	Selected δ -values (see ref. 15b)
H ₁	-0.05
H ₂	-0.06
H ₃	-0.03
H ₄	-0.06
H ₅	+0.03
H ₆	+0.03

Compounds 29 and 30

Unfortunately, Mosher's esters could not be formed for these compounds; thus the stereochemistry of C₁₉ is assigned by analogy to compound **28**. The relative stereochemistry was assigned as follows:

Conditions: a) Zn(BH₄)₂, CH₂Cl₂; b) MeC(OMe)₂, PPTs

¹³C-NMR ketal carbon shifts: **X** --> 97.2, 97.3 and 98.2 ppm
Y --> 97.4, 98.2 and 98.4 ppm

From our data collection of more than a hundred compounds (Dart., M. J., Ph.D. Thesis, Harvard University, **1995**) we have observed the following empirical rules:

97.0–97.9 ppm = ketal of syn-acetonide derived from an anti-aldol adduct

98.0–99.0 ppm = ketal of syn-acetonide derived from a syn-aldol or acetate-aldol adduct

Thus the C₁₇–C₁₉ stereochemistries are as shown for **X** and **Y** above