Synthesis and Self-Assembly of Functionalized Donor- σ -Acceptor Molecules

H. Li, E. A. Homan, A. J. Lampkins, I. Ghiviriga, and Ronald K. Castellano*

Department of Chemistry, University of Florida, Gainesville, FL 32611

I. SYNTHESIS

General. Reagents and solvents were purchased from Acros, Aldrich, or Fluka and used without further purification unless otherwise specified. THF, ether, CH_2Cl_2 , and DMF were degassed in 20 L drums and passed through two sequential purification columns (activated alumina; molecular sieves for DMF) under a positive argon atmosphere using the GlassContour solvent system (GlassContour, Inc.). Thin layer chromatography (TLC) was performed on DURASIL TLC aluminum sheets with visualization by UV light or staining. Melting points (m.p.) were determined on a MEL-TEMP melting apparatus and are uncorrected. ^1H (300, 500 MHz) and ^{13}C NMR (75, 125 MHz) spectra were recorded on Varian Mercury 300 or Inova 500 spectrometers as specified. Chemical shifts (δ) are given in parts per million (ppm) relative to TMS and referenced to residual protonated solvent (CHCl_3 : δ_{H} 7.26 ppm, δ_{C} 77.00 ppm). Abbreviations used are s (singlet), d (doublet), t (triplet), q (quartet), quin (quintet), b (broad), and m (multiplet). MS spectra (HRMS and LRMS) were recorded on a Finnigan MAT95Q Hybrid Sector spectrometer. UV-Vis spectra were recorded using a Shimadzu UV-1700 UV-Visible spectrophotometer. IR spectra were recorded using a Perkin Elmer 1600 Series spectrometer.

1,3,5-Tris(bromomethyl)-2,4,6-trimethoxybenzene (4).¹ To a mixture of 1,3,5-trimethoxybenzene **3** (7.5 g, 45 mmol) and paraformaldehyde (5.00 g, 167 mmol) in a pressure tube was added glacial acetic acid (17 mL) and the resulting suspension was allowed to stir for 1 hour. HBr (33% in HOAc, 47.2 mL) was then added and the reaction mixture was heated to 70 °C for 3 hours in the sealed tube. After cooling to room temperature, the reaction mixture was poured into water (250 mL) and precipitated solids were dissolved in methylene chloride. The organic layer was separated and the aqueous solution was extracted with methylene chloride. All organics were combined and washed sequentially with saturated aqueous NaHCO_3 , water, and brine and concentrated in vacuo to yield an orange oil. Flash chromatography using hexanes/ethyl acetate (20:1) afforded **4** (7.37 g, 37%) as a white solid: ^1H NMR (300 MHz, CDCl_3) δ 4.14 (s, 9H), 4.60 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 160.0, 123.2, 62.6, 22.4.

1,3,5-Tribenzyl-2,4,6-trimethoxybenzene (5a): Representative Procedure.² To magnesium turnings (1.80 g, 74.1 mmol) and anhydrous diethyl ether (30 mL) in a dry three-necked flask was slowly added bromobenzene (6.00 mL, 56.9 mmol) under an inert atmosphere. After addition, the solution was heated to reflux for 30 min, cooled to room temperature, then transferred via cannulation to another dry three-necked flask equipped with a distillation apparatus and connected to an inert gas line. After adding benzene (60 mL), the ether was removed by distillation to leave a cloudy white mixture. To the remaining benzene solution (30 mL) was slowly added **4** (3.50 g, 7.83 mmol) dissolved in a small amount of benzene (ca. 0.5 h). The mixture was stirred at reflux for 24 hours. After carefully quenching the reaction, the mixture was poured into 1.0 M HCl (50 mL), and then extracted with 3 $\times$ 40 mL diethyl ether. The ether solution was dried over MgSO_4 and the organic layers were concentrated in vacuo. The resulting colorless oil was purified by flash column chromatography using hexanes/ethyl acetate (20/1). The crude solid was recrystallized twice from hexanes to give **2** as large, hemispherical, colorless crystals (3.08 g, 90%) ^1H NMR (300 MHz, CDCl_3) δ 3.48 (s, 9H), 4.06 (s, 6H), 7.15–7.27 (m, 15H); ^{13}C NMR (75 MHz, CDCl_3)

δ 157.41, 141.55, 128.17, 128.15, 125.66, 123.91, 61.51, 30.35; LRMS (EI, M^+) calcd for $C_{30}H_{30}O_3$: 438; found: 438.

1,3,5-Tris(2'-methylbenzyl)-2,4,6-trimethoxybenzene (5b). A solution of **4** (2.25 g, 5.03 mmol) in 25 mL toluene was added dropwise to a boiling benzene solution of 2-methylphenylmagnesium bromide (made from 2-bromotoluene (17.0 g, 99.6 mmol) in 130 mL of THF, and the THF was replaced by toluene). After the addition, the mixture was kept at reflux for 1 day. The reaction was

5b

quenched with dilute HCl solution, the organic layer was separated, and the aqueous layer was extracted with ether (3 $\times$ 50 mL). The organic layer was dried with anhydrous $MgSO_4$ and flash chromatography using hexane/ethyl acetate (20/1) as the eluent gave the target as a white powder (1.32 g, 55%). 1H NMR (300 MHz, $CDCl_3$) δ 2.41 (s, 9H), 3.40 (s, 9H), 3.99 (s, 6H), 6.83–6.86 (m, 3H), 7.04–7.15 (m, 9H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 157.63, 139.54, 135.82, 129.67, 127.36, 125.75, 123.43, 61.46, 27.68, 19.86; HRMS (EI, M^+) calcd for $C_{33}H_{36}O_3$: 480.2664; found: 480.2654.

1,3,5-Tris(3'-methylbenzyl)-2,4,6-trimethoxybenzene (5c). A solution of **4** (2.24 g, 5.02 mmol) in 25 mL benzene was added dropwise to a boiling benzene solution of 3-methylphenylmagnesium bromide (made from 11 g, 66 mmol of 3-bromotoluene in 100 mL of ether, and the ether was replaced by benzene). After the addition, the mixture was kept at reflux for 1 day. The reaction was quenched with dilute HCl solution, the organic layer was separated, and the aqueous layer was extracted with ether (2 $\times$ 100 mL). The organic layer was dried with anhydrous $MgSO_4$ and flash chromatography using hexane/ethyl acetate (20/1) as the eluent gave the target as a white powder (931 mg, 42%). 1H NMR (300 MHz, $CDCl_3$) δ 2.30 (s, 9H), 3.50 (s, 9H), 4.03 (s, 6H), 6.95–7.00 (m, 9H), 7.11–7.16 (m, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 157.31, 141.50, 137.61, 128.90, 128.00, 126.40, 125.15, 123.91, 61.50, 30.22, 21.48; HRMS (EI, M^+) calcd for $C_{33}H_{36}O_3$: 480.2664; found: 480.2680.

5c

1,3,5-Tris(4'-methylbenzyl)-2,4,6-trimethoxybenzene (5d). A solution of **4** (2.8 g, 6.3 mmol) in 25 mL benzene was added dropwise to a boiling benzene solution of 4-methylphenylmagnesium bromide (made from 17.1 g, 100 mmol of 4-bromotoluene in 100 mL of ether, and the ether was replaced by benzene). Standard work-up and flash chromatography using hexane/ethyl acetate (25/1) as an eluent gave the target as a white solid (2.0 g, 66%). 1H NMR (300 MHz, $CDCl_3$) δ 2.33 (s, 9H), 3.52 (s, 9H), 4.05 (s, 6H), 7.07–7.14 (m, 12H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 157.24, 138.47, 134.95, 128.84, 127.97, 124.01, 61.53, 29.87, 20.99; HRMS (EI, M^+) calcd for $C_{33}H_{36}O_3$: 480.2664; found: 480.2668.

5d

1,3,5-Tris(3',5'-dimethylbenzyl)-2,4,6-trimethoxybenzene (5e). A solution of **4** (2.24 g, 5.02 mmol) in 25 mL benzene was added dropwise to a boiling benzene solution of 3,5-dimethylphenylmagnesium bromide (made from 18 g, 95 mmol of 1-bromo-3,5-dimethylbenzene in 80 mL of ether, and the ether was replaced by benzene). After the addition, the mixture was kept at reflux for 1 day. Standard work-

up and flash chromatography using hexane/ethyl acetate (30/1) as the eluent gave the target as a white solid (705 mg, 27%). ^1H NMR (300 MHz, CDCl_3) δ 2.28 (s, 18H), 3.13 (s, 6H), 3.26 (s, 6H), 6.83 (s, 3H), 6.92 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 157.28, 141.53, 137.40, 127.29, 125.96, 123.88, 61.48, 3014, 21.35; HRMS (EI, M^+) calcd for $\text{C}_{36}\text{H}_{42}\text{O}_3$: 522.3134; found: 522.3133.

1,3,5-Tris(2'-isopropylbenzyl)-2,4,6-trimethoxybenzene (5f)

(5f). A solution of **4** (1.50 g, 3.38 mmol) in 25 mL benzene was added dropwise to a boiling benzene solution of 2-isopropylphenylmagnesium bromide (made from 5.0 g, 25 mmol of 1-bromo-2-isopropylbenzene in 50 mL of ether, and the ether was replaced by benzene). After the addition, the mixture was kept at reflux for 1 day. Standard work-up and flash chromatography using hexane/ethyl acetate (30/1) as an eluent gave the target as a white solid (1.3 g, 66%). ^1H NMR (300 MHz, CDCl_3) δ 1.32 (d, 18H, $J = 6.9$ Hz), 3.42 (s, 9H), 3.47 (septet, 3H, $J = 6.9$ Hz), 4.15 (s, 6H), 6.89 (d, $J = 8.1$ Hz), 7.07 (t, $J = 7.2$ Hz), 7.18 (t, $J = 7.2$ Hz), 7.30 (d, $J = 7.5$ Hz); ^{13}C NMR (75 MHz, CDCl_3) δ 157.64, 146.24, 137.96, 127.81, 126.04, 125.39, 124.63, 123.89, 61.45, 28.60, 26.97, 23.43; HRMS (EI, M^+) calcd for $\text{C}_{39}\text{H}_{48}\text{O}_3$: 564.3603; found: 564.3616.

1,3,5-Tris(4'-tert-butylbenzyl)-2,4,6-trimethoxybenzene (5g). A solution of **4** (2.0 g, 4.5 mmol) in 50 mL benzene was added dropwise to a boiling benzene solution of *p*-tert-butylphenylmagnesium bromide (25 mL of 2.0 M *p*-tert-butylphenylmagnesium bromide in ether solution, and the ether was replaced by benzene by distillation). After the addition, the mixture was kept at reflux for 1 day. Standard work-up and flash chromatography using hexane/ethyl acetate (30/1) as the eluent gave the product as a white solid (1.7 g, 64%). ^1H NMR (300 MHz, CDCl_3) δ 1.28 (s, 27H), 3.49 (s, 9H), 4.02 (s, 6H), 7.11 (d, 6H, $J = 8.4$ Hz), 7.25 (d, 6H, $J = 8.4$ Hz); ^{13}C NMR (75 MHz, CDCl_3) δ 157.31, 148.27, 138.38, 127.78, 125.03, 124.02, 61.55, 34.27, 31.41, 29.83; HRMS (EI, M^+) calcd for $\text{C}_{42}\text{H}_{54}\text{O}_6$: 606.4073; found: 606.4072.

1,3,5-Tris(4'-fluorobenzyl)-2,4,6-trimethoxybenzene (5h). A solution of **4** (2.24 g, 5.02 mmol) in 25 mL benzene was added dropwise to a boiling benzene solution of 4-fluorophenylmagnesium bromide (40 mL of 2.0 M ether solution, and the ether was replaced by benzene by distillation). After the addition, the mixture was kept at reflux for 1 day. Standard work-up and flash chromatography using hexane/ethyl acetate (30/1) as the eluent gave the target as a white solid (1.56 g, 63%). ^1H NMR (300 MHz, CDCl_3) δ 3.53 (s, 9H), 4.04 (s, 6H), 6.95 (t, 6H, $J = 8.4$ Hz), 7.15 (t, 6H, $J = 6.6$ Hz); ^{13}C NMR (75 MHz, CDCl_3) δ 162.79, 157.29, 136.97, 129.46, 124.09, 115.07, 61.52, 55.17, 29.55; HRMS (EI, M^+) calcd for $\text{C}_{30}\text{H}_{27}\text{F}_3\text{O}_3$: 492.1912; found: 492.1906.

1,3,5-Tris(4'-methoxybenzyl)-2,4,6-trimethoxybenzene

(5i). A solution of **4** (1.44 g, 3.22 mmol) in 25 mL benzene was added dropwise to a boiling benzene solution of 4-methoxyphenylmagnesium bromide (made from 8.0 g, 43 mmol of 4-bromoanisole in 60 mL of ether, and the ether was replaced by benzene). After the addition, the mixture was kept at reflux for 1 day. Standard work-up and flash chromatography using hexane/ethyl acetate (25/1) as the eluent gave the target as a white solid (647 mg, 63%). ¹H NMR (300 MHz, CDCl₃) δ 3.50 (s, 9H), 3.77 (s, 9H), 3.99 (s, 6H), 6.80 (d, 6H, *J* = 8.7 Hz), 7.11 (d, 6H, *J* = 8.7 Hz); ¹³C NMR (75 MHz, CDCl₃) δ 157.58, 157.14, 133.63, 129.01, 124.28, 113.56, 61.56, 55.17, 29.41; LRMS (EI, M⁺) calcd for C₃₃H₃₆O₆: 528; found: 528.

1,3,5-Tris(3'-naphthalenyl)-2,4,6-trimethoxybenzene (5j). A solution of **4** (791 mg, 1.77 mmol) in 25 mL benzene was added dropwise to a boiling toluene solution of 3-naphthalenylmagnesium bromide (made from 5.0 g, 24 mmol of 2-bromonaphthalene in 80 mL of THF, and the THF was replaced by

benzene). After the addition, the mixture was kept at reflux for 1 day. Standard work-up and flash chromatography using hexane/ethyl acetate (20/1) as the eluent gave the target as a white powder (247 mg, 24%). ¹H NMR (300 MHz, CDCl₃) δ 3.53 (s, 9H), 4.24 (s, 6H), 7.38–7.76 (m, 21H); ¹³C NMR (75 MHz, CDCl₃) δ 157.59, 139.13, 133.57, 132.03, 127.73, 127.62, 127.54, 127.25, 126.12, 125.73, 125.45, 125.06, 124.01, 61.70, 30.60; HRMS (EI, M⁺) calcd for C₄₂H₃₆O₃: 588.2664; found: 588.2681.

2,4,6-Tribenzylphloroglucinol (2a): Representative Procedure. BBr₃ (0.5 mL, 5 mmol) was added dropwise to a solution of **5a** (0.56 g, 1.3 mmol) in 40 mL of CH₂Cl₂ at –78 °C. After one hour the mixture was allowed to warm to room temperature, and stirring was continued for 10 h. The mixture was quenched with saturated NaHCO₃ and extracted with CH₂Cl₂. The organic layer was dried with anhydrous MgSO₄ and the solvent was removed in vacuo to provide a yellow solid that was used directly in the next step without purification. Compounds **2b–j** were prepared similarly.

3,5,7-Tribenzyl-1-aza-4,6,10-adamantanetrione (1a): Representative Procedure. A solution of compound **2a** (503 mg, 1.27 mmol) and hexamethylenetetramine (178 mg, 1.27 mmol) in absolute methanol (30 mL) was refluxed for 24 hours. Some solid precipitate could be detected in the solution after this time, and after cooling to room temperature, more precipitate formed. The solid was collected by filtration. The product was obtained in essentially pure form after recrystallization from benzene (371 mg, 65%). M.p. 297 °C (DSC); ¹H NMR (300 MHz, CDCl₃) δ 7.35–7.18 (m, 15 H), 3.31 (s, 6 H), 3.11 (s, 6 H); ¹³C NMR (75 MHz, CDCl₃) δ 198.06, 135.93, 131.17, 128.24, 126.63, 71.59, 68.23, 31.71; HRMS (EI, M⁺): calcd for C₃₀H₂₇NO₃: 449.1991; found: 449.1990; UV/Vis (THF, 10^{–5} M): λ_{CT} = 284 nm (ε_{CT} = 2200 M^{–1} mol^{–1}); IR (Nujol) 1735, 1692 cm^{–1}. Compounds **1b–j** were prepared similarly to **1a**.

3,5,7-Tris(2'-methylbenzyl)-1-aza-4,6,10-adamantanetrione (1b). Yield: 61%. ^1H NMR (300 MHz, CDCl_3) δ 2.34 (s, 9H), 3.13 (s, 6H), 3.50 (s, 6H), 7.08–7.11 (m, 9H), 7.21–7.24 (m, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 198.71, 137.05, 134.62, 130.89, 130.47, 126.54, 125.96, 72.58, 69.16, 26.99, 20.54; LRMS (EI, M^+) calcd for $\text{C}_{33}\text{H}_{33}\text{NO}_3$: 491; found: 491; anal. calcd for $\text{C}_{33}\text{H}_{33}\text{NO}_3$: C, 80.62; H, 6.77; N, 2.85; found: C, 80.51; H, 6.93; N, 2.83.

3,5,7-Tris(3'-methylbenzyl)-1-aza-4,6,10-adamantanetrione (1c). Yield: 62%. ^1H NMR (300 MHz, CDCl_3) δ 2.31 (s, 9H), 3.12 (s, 6H), 3.30 (s, 6H), 7.00–7.16 (m, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 198.02, 137.83, 135.90, 131.91, 128.15, 127.34, 71.48, 67.95, 31.54, 21.42; HRMS (EI, M^+) calcd for $\text{C}_{33}\text{H}_{33}\text{NO}_3$: 491.2460; found: 491.2469.

3,5,7-Tris(4'-methylbenzyl)-1-aza-4,6,10-adamantanetrione (1d). Yield: 43%. ^1H NMR (300 MHz, CDCl_3) δ 2.30 (s, 9H), 3.13 (s, 6H), 3.27 (s, 6H), 7.06 (d, 6H, $J = 7.8$ Hz), 7.17 (d, 6H, $J = 7.8$ Hz); LRMS (EI, M^+) calcd for $\text{C}_{33}\text{H}_{33}\text{NO}_3$: 491; found: 491; anal. calcd for $\text{C}_{33}\text{H}_{33}\text{NO}_3$: C, 80.62; H, 6.77; N, 2.85; found: C, 80.26; H, 6.83; N, 2.82.

3,5,7-Tris(3',5'-dimethylbenzyl)-1-aza-4,6,10-adamantanetrione (1e). Yield: 48%. ^1H NMR (300 MHz, CDCl_3) δ 2.26 (s, 18H), 3.13 (s, 6H), 3.26 (s, 6H), 6.83 (s, 3H), 6.92 (s, 6H); ^{13}C NMR (75 MHz,

CDCl_3) δ 197.97, 137.64, 135.91, 131.23, 128.96, 128.18, 71.38, 67.85, 31.36, 21.29; LRMS (EI, M^+) calcd for $\text{C}_{36}\text{H}_{39}\text{NO}_3$: 533; found: 533.

1f

3,5,7-Tris(2'-isopropylbenzyl)-1-aza-4,6,10-adamantanetrione (1f). Yield: 64%. ^1H NMR (300 MHz, CDCl_3) δ 1.23 (d, 18H, $J = 6.9$ Hz), 3.06 (s, 6H), 3.20 (septet, 3H, $J = 6.9$ Hz), 3.60 (s, 6H), 7.06–7.11 (m, 3H), 7.16–7.21 (m, 3H), 7.24–7.26 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 198.55, 147.73, 132.88, 131.37, 127.00, 125.60, 125.41, 72.26, 68.93, 29.10, 25.92, 23.90; LRMS (EI, M^+) calcd for $\text{C}_{39}\text{H}_{45}\text{NO}_3$: 575; found: 575; anal. calcd for $\text{C}_{39}\text{H}_{45}\text{NO}_3$: C, 81.35; H, 7.88; N, 2.43; Found: C, 80.90; H, 8.31; N, 2.46.

1g

3,5,7-Tris(4'-tert-butylbenzyl)-1-aza-4,6,10-adamantanetrione (1g). Yield: 33%. ^1H NMR (300 MHz, CDCl_3) δ 1.19 (s, 27H), 3.06 (s, 6H), 3.18 (s, 6H), 7.11–7.18 (m, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 198.30, 149.35, 147.20, 132.81, 130.86, 128.32, 125.08, 71.64, 68.32, 34.37, 31.32; LRMS (EI, M^+) calcd for $\text{C}_{42}\text{H}_{51}\text{NO}_3$: 617; found: 617; anal. calcd for $\text{C}_{42}\text{H}_{51}\text{NO}_3$: C, 81.64; H, 8.32; N, 2.27; Found: C, 82.12; H, 8.68; N, 2.15.

1h

3,5,7-Tris(4'-fluorobenzyl)-1-aza-4,6,10-adamantanetrione (1h). Yield: 44%. ^1H NMR (300 MHz, CDCl_3) δ 1.19 (s, 27H), 3.14 (s, 6H), 3.28 (s, 6H), 6.91–6.98 (m, 6H), 7.26–7.30 (m, 6H); HRMS (EI, M^+) calcd for $\text{C}_{30}\text{H}_{24}\text{F}_3\text{NO}_3$: 503.1708; found: 503.1719; anal. calcd for $\text{C}_{30}\text{H}_{24}\text{F}_3\text{NO}_3$: C, 71.56; H, 4.80; N, 2.78; Found: C, 71.25; H, 4.73; N, 2.76.

1i

3,5,7-Tris(4'-hydroxybenzyl)-1-aza-4,6,10-adamantanetrione (1i). Yield: 58%. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 3.03 (s, 6H), 3.12 (s, 6H), 6.61 (d, 6H, $J = 8.4\text{Hz}$), 7.06 (d, 6H, $J = 8.4\text{Hz}$), 9.18 (s, 3H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 198.58, 155.60, 131.86, 129.58, 125.96, 114.61, 70.87, 69.62, 30.56; HRMS (EI, M^+) calcd for $\text{C}_{30}\text{H}_{27}\text{NO}_6$: 497.1838; found: 497.1831.

1j

3,5,7-Tris(3'-naphthalenylbenzyl)-1-aza-4,6,10-adamantanetrione (1j). Yield: 40%. ^1H NMR (300 MHz, CDCl_3) δ 3.14 (s, 6H), 3.53 (s, 6H), 7.42–7.47 (s, 9H), 7.72–7.79 (s, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 198.06, 133.48, 132.20, 129.87, 129.32, 127.80, 126.05, 125.70, 71.85, 68.26, 31.84; HRMS (EI, M^+) calcd for $\text{C}_{42}\text{H}_{33}\text{NO}_3$: 599.2460; found: 599.2474.

1,3,5-Tris(allyloxy)benzene. To phloroglucinol dihydrate (15.0 g, 119 mmol) and anhydrous K_2CO_3 (65.76 g, 457.8 mmol) in a round-bottomed flask was added dry DMF (125 mL) and the resulting suspension was heated to 50 °C for 1 hour. Allyl bromide (57.56 g, 475.8 mmol) was then added dropwise and the reaction mixture was warmed to 65 °C for 36 hours. After cooling to room temperature, the reaction mixture was diluted with water (1000 mL) and extracted with ether (3 $\times$ 150 mL). The combined ethereal extracts were washed with water (5 $\times$ 150 mL), dried over MgSO_4 , and concentrated in vacuo to a crude orange oil. Flash chromatography using hexanes/ethyl acetate (15:1) afforded the suitably pure target (20.0 g, 88%) as a yellow oil: ^1H NMR (300 MHz, CDCl_3) δ 4.50 (m, 6H), 5.35 (m, 6H), 6.04 (m, 3H), 6.13 (s, 3H); ^{13}C NMR (CDCl_3) δ 68.8, 94.4, 117.7, 133.1, 160.3.

7

2,4,6-Triallyl-benzene-1,3,5-triol (7). To 1,3,5-tris(allyloxy)benzene (11.0 g, 44.7 mmol) in a round-bottomed flask was added *N,N*-diethylaniline (55 mL) and the solution was allowed to reflux for 4 hours. After cooling to room temperature, the reaction mixture was poured into ice-cold dilute HCl (300 mL) and extracted with ether (4 $\times$ 100 mL). These combined ethereal extracts were washed with 1.0 M HCl (2 $\times$ 150 mL), water (1 $\times$ 50 mL), 2% aqueous NaHCO_3 (1 $\times$ 50 mL), brine (1 $\times$ 50 mL), dried over MgSO_4 . After concentration, crude

7 (8.71g, 79%) as a thick brown oil, was used without further purification.

3,5,7-Triallyl-1-aza-adamantane-4,6,10-trione (1k). To crude **7** (2.46 g, 10.0 mmol) and HMTA (1.46 g, 10.0 mmol) in a dry flask was added absolute MeOH (50 mL) and the reaction mixture was allowed to reflux for 15 hours. After cooling to room temperature, the solid precipitate was collected by suction filtration and purified via recrystallization from EtOH/EtOAc to give pure **1k** (2.2 g, 74%) as a white solid: ^1H NMR (CDCl_3) δ 2.58 (d, 6H, $J = 7.2$ Hz), 3.42 (s, 6H), 5.14 (m, 6H), 5.91 (m, 3H); ^{13}C NMR (CDCl_3) δ 31.6, 69.6, 70.7, 119.1, 132.6, 198.5; HRMS (EI, M^+) calcd for $\text{C}_{18}\text{H}_{21}\text{NO}_3$: 299.1521; found: 299.1519; anal. calcd for $\text{C}_{18}\text{H}_{21}\text{NO}_3$: C, 72.22; H, 7.07; N, 4.68; Found: C, 72.24; H, 7.22; N, 4.54.

3,5,7-Tripropyl-1-aza-adamantane-4,6,10-trione (1l). To **1k** (0.500 g, 1.67 mmol) and a spatula tip-full of 10% Pd/C in a clean flask under a H_2 atmosphere was added THF (50 mL) and EtOH (10 mL). This resulting suspension was allowed to stir for 12 hours. The reaction mixture was then filtered through a pad of celite and the filtrate concentrated in vacuo to give **1l** (0.300 g, 98%) as a white, flaky solid: ^1H NMR (CDCl_3) δ 0.98 (t, 9H, $J = 7.2$ Hz), 1.39 (sextet, 6H, $J = 7.2$ Hz), 1.70 (t, 6H, $J = 7.2$ Hz), 3.40 (s, 6H); ^{13}C NMR (CDCl_3) δ 15.0, 16.9, 29.4, 71.4, 72.3, 200.5; HRMS (EI, M^+) calcd for $\text{C}_{18}\text{H}_{27}\text{NO}_3$: 305.1991; found: 305.1992.

II. NMR EXPERIMENTS AND DATA

General. The NOESY experiment and the measurement of the temperature dependence of the chemical shifts were run on a Varian Inova spectrometer equipped with a 5 mm indirect detection probe, operating at 500 MHz. Chemical shifts are reported in ppm relative to TMS.

^1H NMR Spectra. Routine ^1H NMR spectra (300 MHz, CDCl_3) are attached (pages S12–S43) for all new compounds which have not been subjected to combustion analysis.

NOESY Spectrum of 1a (Figure S1a). A sample of **1a** was prepared in CDCl_3 (~5 mM). The NOESY experiment was run at 25 °C, in phase-sensitive mode, on a spectral window of 1453 Hz, collecting 2048 points in f2 and 1024 increments in f1, with 32 transients per increment. The preacquisition delay and the mixing time were both 1 s.

Figure S1. NMR data related to the conformational analysis of **1a**. a) NOESY spectrum showing evidence for the “all-arms-up” conformer in CDCl_3 . b) ^{13}C CPMAS NMR spectrum revealing multiple non-equivalent carbons.

Temperature Coefficient Determinations. The temperature dependence of the chemical shifts was followed in a series of 11 spectra in which the temperature was increased in 10 °C steps from –80

°C to 20 °C in CD₂Cl₂. Since the variation of chemical shifts with the temperature displayed good linearity, the temperature coefficient of the chemical shifts was measured as $[\delta(20\text{ °C}) - \delta(-80\text{ °C})]/100 \times 1000$, in ppb/K. The digital resolution in the proton spectra was 0.2 Hz/point, and the temperature was the one indicated by the thermocouple in the probe. Under these conditions, the temperature coefficients of the chemical shifts were measured with a precision of 95%.

Solid-State ¹³C CPMAS (Charge Polarized Magic Angle Spinning) NMR Spectrum of **1a (Figure S1b).** Performed with the dry monomer **1a** (full rotor, no insert) at 10 kHz MAS, D1 = 10 s, 16384 scans. We are grateful to Prof. Joanna Long for assistance with the spectrum.

III. GEL PROPERTIES

Gel Formation. 1-Aza-adamantanetrione **1a** (10.5 mg, 0.5% by weight) and DMSO (2.1 g) were combined in a sealed vial. The vial was heated to 180 °C using an oil bath until an isotropic solution was formed. The vial was then placed in a water bath at room temperature (~ 25 °C) and slowly cooled. The gel gradually formed over the course of several hours.

Xerogel Preparation. *Freeze drying (FD):* A vial containing the organogel of **1a** in DMSO was frozen in liquid nitrogen and transferred to a conventional freeze dry system (Labconco FreeZone 4.5 liter) for two days. A fibrous solid was obtained. *Critical point drying (CPD):* The DMSO was exchanged for liquid CO₂ using an SPI-DRY™ CPD (Model #13200J, Structure Probe, Inc.).

Xerogel Analysis by SEM/TEM. Some of the xerogel was flaked onto a 01801 Formvar/Carbon 200 mesh copper grid (Ted Pella, Inc.). The sample was then sputter-coated with a thin layer of amorphous carbon and imaged on a JEOL 6335f Field Emission Scanning Electron Microscope and at 200 kV on a JEOL 200cx Transmission Electron Microscope.

DSC Analysis of 1a. Differential scanning calorimetry (DSC) analysis was performed using a Perkin Elmer DSC 7 at a heating rate of 10 °C/min. The sample of **1a** (1.5 mg) was heated in a hermetically sealed pan.

a)

10/1/03 10:15:55 AM

1) Hold for 2.0 min at 100.00°C	3) Hold for 1.0 min at 320.00°C
2) Heat from 100.00°C to 320.00°C at 10.00°C/min	4) Cool from 320.00°C to 240.00°C at 10.00°C/min

b)

10/1/03 10:17:49 AM

1) Hold for 2.0 min at 200.00°C	3) Hold for 1.0 min at 320.00°C
2) Heat from 200.00°C to 320.00°C at 10.00°C/min	4) Cool from 320.00°C to 200.00°C at 10.00°C/min

IV. COMPUTATIONAL DETAILS

Monte Carlo Conformational Searching and Geometry Minimizations. All calculations were performed on a Dell PC (2.4 GHz) running Red Hat Linux. Monte Carlo conformational searching was done using MacroModel v. 8.6 (Schrodinger, LLC) and the MCMM method. For **1a**, relevant parameters include: steps = 500, iterations = 5000, solvent (GB/SA) = CHCl₃, force field = Amber*. The lowest energy “all-arms-up” conformer and the lowest energy “all-arms-down” conformer identified were further refined at the HF/3-21G* level using Gaussian 98 (version A.7)³ as implemented on the Linux machine. Both the relative energies and the dipole moments are reported based on these calculations.

V. COPIES OF ¹H NMR AND ¹³C NMR SPECTRA (SEE THE FOLLOWING PAGES)

¹H NMR of **1a** (300 MHz, CDCl₃)

¹³C NMR of **1a** (75 MHz, CDCl₃)

¹H NMR of **1c** (300 MHz, CDCl₃)

¹³C NMR of **1c** (75 MHz, CDCl₃)

¹H NMR of **1e** (300 MHz, CDCl₃)

1H Standard Parameters
expt1 szpu1

SAMPLE		SPECIAL	
date	Jan 11 2004	temp	25.0
solvent	CDCl ₃	gain	not used
file/export/name	1e	spin	2.0
hengfeng/azcl1.fid	9.830	ps30	9.830
ACQUISITION	4803.1	atfa	20.000
SW	4.000	ii	n
at	38426	in	n
np	not used	dp	v
bs	16	hs	mn
d1	1.000	PROCESSING	not used
nt	16	fn	not used
ct	16	DISPLAY	130.3
TRANSMITTER	H1	sp	300.161
stf	300.161	rf1	2482.3
tof	597.6	rfp	2179.2
tpwr	59	fp	6.0
pw	4.700	lp	-63.5
DECOUPLER	C13	WC	PLOT
dn	0	sc	250
dof	0	vs	0
dm	nmn	th	52
dpm	c	at	12
dwt	12300	ph	

¹³C NMR of **1e** (75 MHz, CDCl₃)

¹H NMR of **1i** (300 MHz, DMSO-*d*₆)

^{13}C NMR of **1i** (75 MHz, DMSO- d_6)

^1H NMR of **1j** (300 MHz, CDCl_3)

^{13}C NMR of **1j** (75 MHz, CDCl_3)

C13 Standard Parameters
expt s2pu1

SAMPLE		SPECIAL	
date	Jan 11 2004	temp	25.0
solvent	CDCl_3	gtn	not used
file	/export/home/~hengfeng/aza-NAT-C~	spn	20
hengfeng/aza-NAT-C~	bst	hst	0.008
13 fid	pw90	9.250	
ACQUISITION	alpha	20.000	
sw	18367.9	FLAGS	
at	1.815	i1	n
np	68492	in	y
fb	10400	dp	nn
bs	64	hs	nn
ss	2	PROCESSING	
d1	1.000	j0	1.00
nt	2008	tn	not used
ct	840	DISPLAY	
TRANSMITTER	SP	-1149.4	
to	C13	WD	18867.9
sfrq	75.483	rfl	6961.0
tof	739.8	rff	5811.6
tdwr	58	fp	73.0
pw	4.900	tp	-192.0
DECOUPLER		PLOT	
dn	H1	WC	250
dm	0	SC	0
dmm	yyv	VS	267
dmm	w	tn	13
dpwr	36	at	
dnt	7547	no	ph

¹H NMR of **5b** (300 MHz, CDCl₃)

^{13}C NMR of **5b** (75 MHz, CDCl_3)

STANDARD HI PARAMETERS

exp2	std1h
------	-------

SAMPLE	DATE	JUL 11 2003	DEC. & VI
SOLVENT	CDCl3	dn	299.948
FILE /EXPORT/HOME/~		dn	H1
hengfeng/t-5-h.tid		not used	-75.2
ACQUISITION		dot	nmr
CFREQ	299.949	dm	200
AT		dm	200
SW	3.741	dsen	1.0
FW	4199.9	dres	nmr
TS		hom.	1.0
BS	2300		PROCESSING
TPWR	16	gf	2.228
DI		59	not used
TOF	800.0	Proc	ft
CT	16	math	not used
ALOCK		werr	f
GAIN	not used	wexp	
IN	n	wts	
IN	n		
DISPLAY	218.1	nm	
SP	3509.7	y	
WD	359.870		
SC	0		
WC	250		
HZMAX	13.60		
TS	4838.50		
RF1	2428.7		
RF2	2177.6		
TH	11		
INS	9.000		
AT	cdc	ph

exp2 std13c

SAMPLE	date	DEC. & VI
solvent	11 2003	dfrq 299.948
file/export/home/~	CDCl3	dn H1
hengfeng/i-5-C.tif	dfrq	not used
ACQUISITION	75.428	dm 0
sfreq	75.428	dm yyy
at	1.815	dres 9900
np	55904	dres 1.0
sw	16501.7	homo
fb	9100	PROCESSING 1.00
bs	16	1b
twpr	59	wtfile
pw	4.0	proc
dl	0	ft
rf	0	not used f
tofr	1020	math
ct	320	wexp
alock	n	wbs
gain	not used	wnt
1l	FLAGS	
in	n	
in	n	
dp	n	
hs	nn	
DISPLAY	-848.2	
sp	16500.7	
sc	0522	
sc	0	
wc	250	
hzm	66.01	
is	500.00	
rfl	6655.7	
rffp	5807.5	
th	13	
ins	1.000	
at	no ph

¹H NMR of **5d** (300 MHz, CDCl₃)

¹³C NMR of **5d** (75 MHz, CDCl₃)

¹H NMR of **5e** (300 MHz, CDCl₃)

^{13}C NMR of **5e** (75 MHz, CDCl_3)

^{13}C NMR of **5f** (75 MHz, CDCl_3)

¹H NMR of **5g** (300 MHz, CDCl₃)

SAMPL

[illegible]

Supporting Information

^{13}C NMR of **5h** (75 MHz, CDCl_3)

¹H NMR of **5i** (300 MHz, CDCl₃)

^{13}C NMR of **5i** (75 MHz, CDCl_3)

^1H NMR of **5j** (300 MHz, CDCl_3)

^{13}C NMR of **5j** (75 MHz, CDCl_3)

¹H NMR of tris(allyloxy)benzene (300 MHz, CDCl₃)

^{13}C NMR of tris(allyloxy)benzene (75 MHz, CDCl_3)

^1H NMR of **11** (300 MHz, CDCl_3)

^{13}C NMR of **11** (75 MHz, CDCl_3)

VI. REFERENCES

- ¹ Cho, B. R.; Chajara, K.; Oh, H. J.; Son, K. H.; Jeon, S. J. *Org. Lett.* **2002**, *4*, 1703-1706.
- ² Simaan, S.; Siegel, J. S.; Biali, S. E. *J. Org. Chem.* **2003**, *68*, 3699-3701.
- ³ Gaussian 98, Revision A.7, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, Jr., J. A.; Vreven, T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Strain, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; and Pople, J. A.; Gaussian, Inc., Wallingford CT, 2004.