Synthesis of 2,3-Substituted Thienylboronic Acids and Esters

Claus Christophersen^{,†,‡}, Mikael Begtrup[‡], Søren Ebdrup[§], Henning Petersen[§] and Per Vedsø[§]*

Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd, Denmark and Department of Medicinal Chemistry, The Danish University for Pharmaceutical Sciences, Universitetsparken 2, DK-2200 Copenhagen N, Denmark

clcp@novonordisk.com

Table of contents.

General Experimental Procedure	S3
3-Iodothiophene	S3-S4
Typical Procedure for the Preparation of 3-Bromo-2-thienylmagnesium Chloride (2)	S4
Typical Procedure for the Preparation of 2-Bromo-3-thienylmagnesium Chloride (5)	S4
Ethyl 3-Bromo-2-thiophenecarboxylate (7a)	S4
<i>N</i> -tert-Butyl 3-Bromothieryl-2-carboxamide (7b)	S5
3-Bromo-2-trimethylsilylthiophene (7c)	S5
3-Acetyl-2-bromothiophene (8a)	S5-S6
2-Bromo-3-cyanothiophene (8b)	S6
2-Bromo-3-thienylcarbaldehyde (8c)	S6-S7
Ethyl 2-Bromo-3-thiophenecarboxylate (8d)	S7
General Procedure for the Borylation of 2,3-Substituted Bromothiophenes with Pinacolborane Catalyzed by Pd(P(<i>t</i> -Bu) ₃) ₂	S7

2-Carboxyethyl-3-thienylboronic Acid (9a)	S7-S8
<i>N-tert</i> -Butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-thiophene Carboxamide (9b)	S8
3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3-trimethylsilylthiophene (9c)	S8
2-Nitro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-thiophene (9d)	S9
3-Acetyl-2-thienylboronic Acid (10a)	S9
2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3-thienylcarbaldehyde (10c)	S9
3-Carboxyethyl-2-thiopheneboronic Acid (10d)	S10
<i>N-tert</i> -Butyl 3-(4-(<i>N',N'</i> -dimethylamino)phenyl)-2-thiophene Carboxamide (11)	S10-S11
References	S11-S12
¹ H- and ¹³ C-NMR spectra for Ethyl 3-Bromo-2-thiophenecarboxylate (7a)	S13-S14
¹ H- and ¹³ C-NMR spectra for <i>N-tert</i> -Butyl 3-Bromothieryl-2-carboxamide (7b)	S15-S16
¹ H- and ¹³ C-NMR spectra for 3-Bromo-2-trimethylsilylthiophene (7c)	S17-S18
¹ H- and ¹³ C-NMR spectra for 3-Acetyl-2-bromothiophene (8a)	S19-S20
¹ H- and ¹³ C-NMR spectra for 2-Bromo-3-cyanothiophene (8b)	S21-S22
¹ H- and ¹³ C-NMR spectra for 2-Bromo-3-thienylcarbaldehyde (8c)	S23-S24
¹ H- and ¹³ C-NMR spectra for Ethyl 2-Bromo-3-thiophenecarboxylate (8d)	S25-S26
¹ H- and ¹³ C-NMR spectra for 2-Carboxyethyl-3-thienylboronic Acid (9a)	S27-S28
¹ H- and ¹³ C-NMR spectra for <i>N-tert</i> -Butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-thiophene Carboxamide (9b)	S29-S30
¹ H- and ¹³ C-NMR spectra for 3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3-trimethylsilylthiophene (9c)	S31-S32
¹ H- and ¹³ C-NMR spectra for 2-Nitro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-thiophene (9d)	S33-S34
¹ H- and ¹³ C-NMR spectra for 3-Acetyl-2-thienylboronic Acid (10a)	S35-S36
¹ H- and ¹³ C-NMR spectra for 2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3-thienylcarbaldehyde (10c)	S37-S38
¹ H- and ¹³ C-NMR spectra for 3-Carboxyethyl-2-thiopheneboronic Acid (10d)	S39-S40
¹ H- and ¹³ C-NMR spectra for <i>N-tert</i> -Butyl 3-(4-(<i>N',N'</i> -dimethylamino)phenyl)-2-thiophene Carboxamide (11)	S41-S42

Experimental Section

General Experimental Procedure: All reactions were performed in flame-dried equipment under nitrogen and magnetic stirring. Anhydrous THF, DMF, $(\text{CH}_3)_3\text{SiCl}$ and pinacolborane were purchased in Sure/Seal bottles. All other solvents were dried over molecular sieves ($3\text{\AA}$). Ethylmagnesium chloride in THF was purchased in Sure/Seal bottles and titrated using the procedure described by Lin and Paquette.¹ Commercially available reagents and catalysts were used without further purification. Tri-*tert*-butylphosphine was purchased in ampoules and transferred to vials equipped with a septum under N_2 . The vial was stored at ambient temperature and gently heated ($\text{P}(t\text{-Bu})_3$ is a gel at ambient temperature) with a heat gun prior to taking out aliquots with a syringe. Z-HSM-5 zeolite was purchased on H^+ -form from Zeolyst Inc. 2-Bromo-3-iodothiophene was prepared by bromination of 3-iodothiophene as described by Gronowitz *et al.*² and 3-bromo-2-nitrothiophene was prepared as described by Bacon *et al.*³ HPLC analyses were performed using a YMC $5\text{ }\mu\text{m}$ $120\text{\AA}$ C-18 reversed phase column. A gradient system using acetonitrile and water (pH 2.3) over 35 min was used. UV-detection at 226 nm was applied. ^1H -NMR and ^{13}C -NMR spectra were recorded at 400 MHz and 100 MHz, respectively. Mass spectra were recorded in the EI mode. Melting points were obtained on a differential scanning calorimeter. Melting intervals were determined as onset and endset temperatures.

3-Iodothiophene. A mixture of 2-iodothiophene (157.4 g, 0.75 mol) and H-ZSM-5 zeolite (7.86 g) was stirred for 3 hours at 100°C . After cooling to 60°C acetic acid (160 mL) and water (80 mL) were added. Zinc dust (24.5 g, 0.375 mol, $< 10\text{ }\mu\text{m}$) was added under vigorous stirring in small portions at such a rate that the temperature was kept below 40°C . After addition of the Zn-dust the mixture was stirred at ambient temperature for further 16 hours. Water (100 mL) was added to the reaction mixture followed by extraction with MTBE (3 x 250 mL). The combined organic phases were concentrated to approx. 300 mL under reduced

pressure and washed with aq. NaHCO_3 (500 mL). The aqueous phase was extracted with MTBE (100 mL) and the combined organic phases were dried over MgSO_4 . The solvent was removed under reduced pressure. Distillation of the remaining oil afforded 3-iodothiophene (93.4 g, 59%) as a colorless oil. bp 46-49°C (6 Torr). ^1H NMR data were in accordance with literature data.⁴

Typical Procedure for the Preparation of 3-Bromo-2-thienylmagnesium Chloride (2).

A solution of 2,3-dibromothiophene (**1**) (8.46 g, 35 mmol) in THF (30 mL) was cooled to 15°C under magnetic stirring. Ethylmagnesium chloride in THF (19.0 mL, 38 mmol) was added over 15 min. The solution was stirred for further 15 min at 15°C.

Typical procedure for the Preparation of 2-Bromo-3-thienylmagnesium Chloride (5).

A solution of 2-bromo-3-iodothiophene (**4**) (31.8 g, 0.11 mol) in THF (50 mL) was cooled to 15°C under magnetic stirring. Ethylmagnesium chloride in THF (58.5 mL, 0.112 mol) was added over 15 min. The solution was stirred for further 15 min at 15°C.

Ethyl 3-Bromo-2-thiophenecarboxylate (7a). A THF-solution of **2** (63.3 mmol) cooled to 15°C was added to ethyl cyanoformate (6.54 mL, 66 mmol) in THF (20 mL) at 25-30°C over 30 min. The reaction mixture was stirred for further 30 min at 20-25°C followed by addition of sat. NaHCO_3 (100 mL). The mixture was vigorously stirred until all solid material had dissolved and the solution was extracted with MTBE (2 x 100 mL). The combined organic phases were dried over MgSO_4 and the solvents were removed under reduced pressure. The resulting brown oil was distilled on a 30 cm Vigreux column affording **7a** as a colorless oil (8.4 g, 59%): bp 77-80°C (0.8 Torr) (lit.⁵ 63-66°C (0.8 Torr)); ^1H -NMR (CDCl_3)⁶ δ 1.39 (t, J = 7.3 Hz, 3 H), 4.37 (q, J = 7.3 Hz, 2 H), 7.08 (d, J = 5.3 Hz, 1 H), 7.45 (d, J = 5.3 Hz, 1 H); ^{13}C -NMR (CDCl_3) δ 14.2, 61.3, 116.4, 127.7, 131.0, 132.9, 160.7; MS (EI, m/z) 236/234 (29/30), 208/206 (40/41), 191/189 (99/100), 82 (62), 45 (49).

***N*-tert-Butyl 3-Bromothieryl-2-carboxamide (7b).** To a THF-solution of **2** (35 mmol) was added *tert*-butyl isocyanate (8.0 mL, 70 mmol) at 25-30°C over 15 min. The reaction mixture was stirred for further 45 min at 20-25°C followed by addition of 1 M HCl (50 mL). The mixture was extracted with MTBE (2 x 50 mL). The combined organic phases were washed with water (50 mL) and dried over MgSO₄. The solvents were removed under reduced pressure and the crude product was re-crystallized from heptane to give **7b** as white needles (6.98 g, 76%): mp 72.2-76.0°C; ¹H-NMR (CDCl₃) δ 1.47 (s, 9 H), 6.90-6.98 (broad s, 1 H), 7.00 (d, *J* = 5.3 Hz, 1 H), 7.40 (d, *J* = 5.3 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 28.9, 52.3, 107.6, 129.4, 131.7, 136.5, 159.3; MS (EI, *m/z*) 263/261 (9/8), 248/246 (25/24), 191/189 (100/99), 182 (10), 163/161 (3/3), 82 (29). Anal. Calcd. for C₉H₁₂BrNOS C 41.23, H 4.61, N 5.34, Br 30.48, found C 41.20, H 4.60, N 5.30, Br 30.67.

3-Bromo-2-trimethylsilylthiophene (7c). To a THF-solution of **2** (25 mmol) was added trimethylsilyl chloride (3.5 mL, 28 mmol) at 20-25°C over 5 min. The reaction mixture was stirred for 3 hours at 22-25°C followed by addition of sat. NH₄Cl (50 mL). The mixture was extracted with heptane (2 x 25 mL). The combined organic phases were dried over MgSO₄ and the solvents were removed under reduced pressure. Kugelrohr distillation gave **7c** as a colorless oil (4.8 g, 81%): bp 100-103°C (1 Torr) (lit.⁷ 110°C (2 Torr)). ¹H NMR and ¹³C NMR data were in accordance with published data.⁸

3-Acetyl-2-bromothiophene (8a).⁹ A THF-solution of **5** (0.10 mol) cooled to 15°C was added to acetic anhydride (14.2 mL, 0.15 mol) in THF (40 mL) at 25-30°C over 30 min. The reaction mixture was stirred for further 45 min at 20-25°C followed by addition of 1 M HCl (150 mL). The mixture was extracted with MTBE (2 x 50 mL). The combined organic phases were washed with 4 M KOH (100 mL), water (50 mL) and dried over MgSO₄. The solvents were removed under reduced pressure. Distillation on a 30 cm Vigreux column afforded **8a** as

a slight yellow oil (14.2 g, 69%; purity: > 95% (HPLC)); bp 56-61°C (0.4 Torr). Crystallization from heptane gave a product containing approx. 3% of 2-acetyl-3-bromothiophene:¹⁰ mp 32.7-36.7°C; ¹H-NMR (CDCl₃) δ 2.60 (s, 3 H), 7.21 (d, *J* = 5.8 Hz, 1 H), 7.33 (d, *J* = 5.8 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 29.9, 117.9, 126.0, 129.0, 139.3, 192.1; MS(EI, *m/z*): 206/204 (34/32), 191/189 (100/95), 163/161 (8/6), 82, (43), 45 (43). Anal. Calcd. for C₆H₅BrOS C 35.14, H 2.46, Br 38.96, found C 35.00, H 2.40, Br 38.81.

2-Bromo-3-cyanothiophene (8b). A THF-solution of **5** (0.11 mol) cooled to 1°C was added to *p*-toluenesulfonyl cyanide (20.3 g, 0.11 mol) in THF (100 mL) at 2-7°C over 30 min. The reaction mixture was stirred for further 1 hour at 20-25°C followed by addition of sat. NaHCO₃ (600 mL) and MTBE (100 mL). The mixture was extracted with CH₂Cl₂ (3 x 100 mL) and the combined organic phases were dried over MgSO₄. The solvents were removed under reduced pressure. Distillation on a 30 cm Vigreux column afforded **8b** as a colorless oil (11.5 g, 64%; purity: ≈ 90% (¹H-NMR)); bp 82-85°C (3 Torr) (lit.¹¹ bp 130°C (22 Torr)). Crystallization of the oil from heptane gave a product containing 4% of 3-bromo-2-cyanothiophene:¹² mp 20.2-27.4°C; ¹H-NMR (CDCl₃) δ 7.12 (d, *J* = 5.6 Hz, 1 H), 7.34 (d, *J* = 5.6 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 113.7, 114.0, 123.4, 128.1, 129.0; MS (EI, *m/z*) 189/187 (100), 108 (73), 82 (37), 64 (91), 45 (85). Anal. Calcd. for C₅H₂BrNS C 31.94, H 1.07, N 7.45, found C 31.91, H 1.04, N 7.66.

2-Bromo-3-thienylcarbaldehyde (8c). To a cooled (11°C) THF-solution of **5** (35 mmol) was added DMF (5.4 mL, 70 mmol) at 20-30°C over 5 min. The reaction mixture was stirred for further 1 hour at 20-25°C followed by addition of 1 M HCl (100 mL). The mixture was extracted with MTBE (2 x 50 mL). The combined organic phases were washed with sat. NaHCO₃ (50 mL) and dried over MgSO₄. The solvents were removed under reduced pressure resulting in a brown oil. The oil was dissolved in heptane (50 mL) and kept at 2-4°C for 3

weeks which gave **8c** as pale yellow needles (3.2 g, 48%; purity: 97% (GC)): mp 28.5-34.2°C (lit.¹¹ mp 34-35°C). Further crystallization from heptane gave the analytically pure product: mp 34.1-39.3°C; ¹H-NMR (CDCl₃)³² δ 7.28 (dd, *J* = 5.9, 0.7 Hz, 1 H), 7.36 (d, *J* = 5.9 Hz, 1 H), 9.94 (d, *J* = 0.7 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 125.2, 126.2, 126.9, 138.5, 184.6; MS (EI, *m/z*) 192/190 (73/73), 191/189 (100/99), 163/161 (9/8), 119/117 (6/6), 82 (51), 57 (23), 45 (47).

Ethyl 2-Bromo-3-thiophenecarboxylate (8d). A THF-solution of **5** (40 mmol) cooled to 1°C was added to ethyl cyanofomate (4.4 mL, 44 mmol) in THF (20 mL) at 4-7°C over 20 min. The reaction mixture was stirred for further 30 min at 20-25°C followed by addition of sat. NaHCO₃ (100 mL). The mixture was extracted with MTBE (2 x 100 mL) and the combined organic phases were dried over MgSO₄. The solvents were removed under reduced pressure. Distillation on Podbielniak column afforded **8d** as a colorless oil (4.6 g, 49%) (The product contained 4% of **7a**): bp 79-81°C (0.7 Torr); ¹H-NMR (CDCl₃) δ 1.39 (t, *J* = 7.3 Hz, 3 H), 4.35 (q, *J* = 7.3 Hz, 2 H), 7.21 (d, *J* = 5.8 Hz, 1 H), 7.36 (d, *J* = 5.8 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 14.2, 60.9, 119.6, 125.7, 129.3, 131.3, 161.9; MS (EI, *m/z*) 236/234 (25/23), 208/206 (33/31), 191/189 (100/99), 82 (74), 45 (60). Anal. Calcd. for C₇H₇BrO₂S C 35.76, H 3.00, found C 35.30, H 3.10.

General Procedure for the Borylation of 2,3-Substituted Bromothiophenes with Pinacolborane Catalyzed by Pd(P(*t*-Bu)₃)₂. The specified amount of Pd(P(*t*-Bu)₃)₂ (weighed in the air) was added to 2,3-substituted bromothiophenes **7a-d** or **8a-d** (1 eqv.) dissolved in dry THF (2 ml/mmol). Triethylamine (3 eqv.) and pinacolborane (1.5 eqv.) were added to the mixture. The reaction mixture was stirred at 40°C under N₂ for the specified time followed by cooling to room temperature and filtration. The solid was extracted with MTBE and the filtrates were combined. The solvents were removed under reduced pressure.

2-Carboxyethyl-3-thienylboronic Acid (9a). This was prepared from **7a** (4.70 g, 20 mmol). The evaporation residue was re-dissolved in THF (20 mL) and water (20 mL). The mixture was adjusted to pH 3.5 by acetic acid and stirred vigorously for 50 min, this resulted in precipitation of a light brown solid. The mixture was filtered and the solid was washed with heptane (2 x 5 mL) and cold water (2 x 5 mL) giving **9a** (3.51 g, 88%; purity: $\approx$ 95% (HPLC)). Re-crystallization from toluene gave the pure product: mp 138.3-150.7°C; $^1\text{H-NMR}$ (DMSO- d_6) δ 1.34 (t, J = 6.8 Hz, 3 H), 4.34 (q, J = 6.8 Hz, 2 H), 7.34 (d, J = 5.0 Hz, 1 H), 7.89 (d, J = 5.0 Hz, 1 H), 8.48 (s, 2 H); $^{13}\text{C-NMR}$ (DMSO- d_6) δ 14.2, 61.6, 132.6, 133.9, 136.0, 163.8. Anal. Calcd. for $\text{C}_7\text{H}_9\text{BO}_4\text{S}$ C 42.03, H 4.54, found C 42.25, H 4.53.

***N-tert*-Butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2-thiophene Carboxamide (9b).** This was prepared from **7b** (787 mg, 3 mmol). The evaporation residue was re-dissolved in heptane (5 mL) and washed with water (3 x 10 mL) adjusted to pH 4. The combined organic phases were dried over MgSO_4 and the solvent was removed under reduced pressure. The brown oil was distilled at 130-138°C (0.5 Torr) and **9b** (505 mg, 54%) was collected as white crystals by condensation on a water cooled “cold finger”: mp 100.6-105.0°C; $^1\text{H-NMR}$ (CDCl_3) δ 1.38 (s, 12 H), 1.48 (s, 9 H) 7.38 (d, J = 5.7 Hz, 1 H), 7.42 (d, J = 5.7 Hz, 1 H), 8.6-8.68 (broad s, 1 H); $^{13}\text{C-NMR}$ (CDCl_3) δ 24.8, 28.8, 51.9, 84.9, 128.4, 135.86, 153.4, 161.5. MS (EI, m/z): 309 (5), 294 (5), 276 (3), 251 (29), 237 (18), 195 (56), 154 (38), 137 (54), 83 (100), 59 (25). Anal. Calcd. for $\text{C}_{15}\text{H}_{24}\text{BNO}_3\text{S}$ C 58.26, H 7.82, N 4.53, found C 58.17, H 7.94, N 4.48.

3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3-trimethylsilylthiophene (9c). This was prepared from **7c** (247 mg, 1.05 mmol). The evaporation residue was re-dissolved in CH_2Cl_2 and subjected to column chromatography on silica gel (heptane/ CH_2Cl_2 (1:1), R_f =0.68) to give **9c** (171 mg, 58%): mp 64.4-68.5°C; $^1\text{H-NMR}$ (CDCl_3) δ 0.39 (s, 9 H), 1.33 (s, 12 H),

7.52 (d, $J = 4.6$ Hz, 1 H), 7.59 (d, $J = 4.6$ Hz, 1 H); ^{13}C -NMR (CDCl_3) δ 0.2, 24.8, 83.5, 128.8, 136.1, 152.6; MS(EI, m/z) 282 (12), 267 (40), 185 (36), 167 (15), 125 (13), 83 (100), 55 (40). Anal. Calcd. for $\text{C}_{13}\text{H}_{23}\text{SiBO}_2\text{S}$ C 55.31, H 8.21, found C 55.46, H 8.32.

2-Nitro-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-thiophene (9d). This was prepared from **7d** (625 mg, 3 mmol). The evaporation residue was distilled at 95-105°C (0.3 Torr) and **9d** (329 mg, 43%) was collected as pale orange crystals by condensation on a water cooled “cold finger”: mp 80.9-84.0°C; ^1H -NMR (CDCl_3) δ 1.41 (s, 12 H), 7.08 (d, $J = 5.2$ Hz, 1 H), 7.52 (d, $J = 5.2$ Hz, 1 H); ^{13}C -NMR (CDCl_3) δ 24.7, 85.1, 130.7, 132.3; MS (EI, m/z) 255 (<5), 240 (11), 154 (56), 139 (100). Anal. Calcd. for $\text{C}_{10}\text{H}_{14}\text{BNO}_4\text{S}$ C 47.08, H 5.53, N 5.49, found C 47.20, H 5.57, N 5.43.

3-Acetyl-2-thienylboronic Acid (10a).¹³ This was prepared from **8a** (2.99 g, 14.6 mmol). The evaporation residue was re-dissolved in THF (5 mL) and water (3 mL). The mixture was adjusted to pH 4 by acetic acid and stirred vigorously for 45 min resulting in precipitation of a blue-green solid. The mixture was filtered and the solid was washed with heptane (2 x 10 mL) and cold water (4 mL) resulting in **10a** (527 mg, 24%; purity: $\approx 95\%$ (^1H -NMR)). Re-crystallization from toluene gave the pure product: mp dec. $\approx 125^\circ\text{C}$ (lit.¹³ dec. 120-140°C); ^1H -NMR ($\text{DMSO}-d_6$)¹³ δ 2.68 (s, 3 H), 7.84 (d, $J = 5.2$ Hz, 1 H), 7.89 (d, $J = 5.2$ Hz, 1 H), 9.06 (s, 2 H); ^{13}C -NMR ($\text{DMSO}-d_6$) δ 29.5, 131.0, 131.9, 145.3, 197.9.

2-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-3-thienylcarbaldehyde (10c). This was prepared from **8c** (1.91 g, 10 mmol). The evaporation residue was re-dissolved in heptane (20 mL) and washed with water (3 x 10 mL) adjusted to pH 4. The combined organic phases were dried over MgSO_4 and the solvent was removed under reduced pressure. Re-crystallization from heptane gave **10c** (1.17 g, 49%; purity: $\approx 93\%$ (HPLC)). A further re-crystallization from ethyl acetate gave the pure product: mp 65.3-68.3°C; ^1H -NMR (CDCl_3) δ 1.38 (s, 12 H),

7.57 (d, $J = 4.9$ Hz, 1 H), 7.68 (d, $J = 4.9$ Hz, 1 H), 10.41 (s, 1 H); ^{13}C -NMR (CDCl_3) δ 24.8, 84.8, 127.0, 132.3, 150.2, 187.9; MS(EI, m/z) 238 (1), 223 (5), 182 (81), 180 (100), 165 (28), 139 (70), 111 (23), 84 (60), 57 (55). Anal. Calcd. for $\text{C}_{11}\text{H}_{15}\text{BO}_3\text{S}$ C 55.49, H 6.35, found: C 55.40, H 6.30.

3-Carboxyethyl-2-thiopheneboronic Acid (10d). This was prepared from **8d** (2.82 g, 12 mmol). The evaporation residue was re-dissolved in THF (10 mL) and water (10 mL). The mixture was adjusted to pH 4 by acetic acid and vigorously stirred for 25 min resulting in precipitation of a light brown solid. The slurry was filtered and the solid was washed with heptane (2 x 10 mL) and cold water (2 x 10 mL) resulting in **10d** (1.53 g, 64%; purity: $\approx 95\%$ (NMR)). Re-crystallization from toluene gave the title compound containing 5% of **9a**. The product was purified by preparative HPLC: mp 124.4-134.6°C. Dec. $\approx 156^\circ\text{C}$; ^1H -NMR (DMSO-d_6) δ 1.35 (t, $J = 6.6$ Hz, 3 H), 4.35 (q, $J = 6.6$ Hz, 2 H), 7.59 (d, $J = 4.6$ Hz, 1 H), 7.82 (d, $J = 4.6$ Hz, 1 H), 8.86 (s, 2 H); ^{13}C -NMR (DMSO-d_6) δ 14.9, 62.3, 131.0, 131.7, 137.6, 166.2. Anal Calcd. for $\text{C}_7\text{H}_9\text{BO}_4\text{S}$ C 42.03, H 4.54, found: C 42.01, H 4.46.

***N*-tert-Butyl 3-(4-(*N,N'*-dimethylamino)phenyl)-2-thiophene Carboxamide (11).** Crude **9b** was prepared from **7b** (262 mg, 1.0 mmol) according to the general borylation procedure described above. The evaporation residue was dissolved in DMF (2 mL) and 4-bromo-*N,N*-dimethylaniline (304 mg, 1.5 mmol), $\text{Pd}(\text{P}(t\text{-Bu})_3)_2$ (9.7 mg, 0.019 mmol) and anhydrous K_3PO_4 (flame dried) (636 mg, 3.0 mmol) were added to the solution. The reaction mixture was stirred at 50°C for 18 hours and at 100°C for 2 hours. 1 M HCl (6 mL) was added to the reaction mixture followed by extraction with MTBE (6 mL). The organic phase was extracted with 1 M HCl (2 x 5 mL). The combined aqueous phases were basified with 4 M KOH and extracted with MTBE (3 x 5 mL). The solvent was removed under reduced pressure and the crude product was purified by flash chromatography on silica gel (ethyl acetate/heptane (3:7), $R_f = 0.33$) affording **11** (179 mg, 59% overall yield) of white crystals: mp 151.1-154.8 $^\circ\text{C}$; ^1H -

NMR (CDCl₃) δ 1.20 (s, 9 H), 3.00 (s, 6 H), 6.78 (dt, J = 8.5, 2.7 Hz, 2 H), 6.95 (d, J = 5.1 Hz, 1 H), 7.27 (dt, J = 8.5, 2.7 Hz, 2 H), 7.37 (d, J = 5.1 Hz, 1 H); ¹³C-NMR (CDCl₃) δ 28.6, 40.4, 51.2, 112.4, 122.7, 127.7, 130.1, 130.8, 135.7, 141.8, 150.6, 161.8; MS(EI, m/z) 302 (100), 246 (49), 245 (48), 230 (92), 202 (14), 187 (14), 186 (14), 114 (43), 79 (27). Anal. Calcd. for C₁₇H₂₂N₂O₂ C 67.51, H 7.33, N 9.26, found C 67.36, H 7.48, N 9.13.

References

- (1) Lin, S.-H.; Paquette, L. A. *Synth. Commun.* **1994**, *24*, 2503-2506.
- (2) Gronowitz, S.; Petterson, K. *J. Heterocycl. Chem.* **1976**, *13*, 1099-1100.
- (3) Bacon, R. G. R.; Hamilton, S. D. *J. Chem. Soc., Perkin Trans. I* **1974**, 1970-1975.
- (4) Satonaka, H. *Bull. Chem. Soc. Jpn.* **1983**, *56*, 2463-2468.
- (5) Chadwick, D. J.; Chambers, J.; Meakins, G. D.; Snowden, R. L. *J. Chem. Soc., Perkin Trans. I* **1973**, 1766-1773.
- (6) Gromova, G. P.; Belin'kii, L. I.; Krayushkin, M. M. *Chem. Heterocycl. Compd. (Engl. Transl.)* **1993**, *29*, 889-891.
- (7) Seconi, G.; Eaborn, C.; Stamper, J. G. *J. Organometal. Chem.* **1981**, *204*, 153-168.
- (8) Fröhlich, J.; Kalt, W. *J. Org. Chem.* **1990**, *55*, 2993-2995.
- (9) The compound was indicated but no characterization data were presented in: Noto, R.; Frenna, V.; Consiglio, G.; Spinelli, D. *J. Chem. Res., Miniprint* **1991**, 2701-2710.
- (10) Muratake, H.; Hayakawa, A.; Natsume, M. *Chem. Pharm. Bull.* **2000**, *48*, 1558-1566.
- (11) Fournai, P.; Guillard, R.; Person, M. *Bull. Soc. Chim. Fr.* **1967**, 4115-4120.

- (12) Crude 3-bromo-2-cyanothiophene was prepared from 3-bromo-2-thienylmagnesium chloride and *p*-toluenesulfonyl cyanide according to the procedures described in the Experimental section. ¹H-NMR data for this product: ¹H-NMR (CDCl₃) δ 7.11 (d, *J* = 4.9 Hz, 1 H), 7.54 (d, *J* = 4.9 Hz, 1 H). These data are in accordance with the data observed for the by-product in **8b**.
- (13) Gronowitz, S.; Dahlgren, T.; Namtvedt, J.; Roos, C.; Rosén, G.; Sjöberg, B.; Forsgren, U. *Acta Pharm. Suecica* **1971**, 8, 623-638.

10d

11