SUPPLEMENTARY MATERIAL

Synthesis of 3-arylpiperidines by a radical 1,4-aryl migration

Alexandru George, Béatrice Quiclet-Sire, Xavier Vila and Samir Z. Zard

Laboratoire de Synthèse Organique, Département de Chimie. École Polytechnique. Route de Saclay. 91128 Palaiseau cedex (France)

zard@poly.polytechnique.fr

General procedures

All reactions were carried out under an inert atmosphere. Commercial reagents were used as received without further purification. All products were purified by using silica gel (SDS, Silice 60 A. C. C. 40-63 μm) or by crystallization. Analytical TLC was carried out on Merck silica gel plates using short wave (254 nm) UV light, 1% aq. KMnO_4 solution to visualise components. NMR spectra were recorded in CDCl_3 using a Bruker AMX400 operating at 400 MHz for ^1H and 100 MHz for ^{13}C . The chemical shifts are expressed in parts per million (ppm) referenced to residual chloroform. ^1H NMR data are reported as follows: δ , chemical shift; multiplicity (recorded as: s, singlet; bs, broad singlet; d, doublet; t, triplet; q, quadruplet; qt, quintuplet; ht, heptuplet; dd, double doublet; ddd, double double doublet; dddd, double double double doublet; dt, double triplet; ddt, double double triplet; dq, double quadruplet; tt, triple triplet; td, triple doublet; tdd, triple double doublet; m, multiplet), coupling constants (J are given in Hertz, Hz) and integration. Infrared Absorption spectra were recorded as a solution in CCl_4 with a Perkin-Elmer 1600 Fourier Transform Spectrophotometer. Mass spectra were recorded with an HP 5989B mass spectrometer via direct introduction for chemical positive ionization (CI) using ammonia as the reagent gas. Melting points were determined by Reichert microscope apparatus and were uncorrected. Microanalyses were performed by the Service de Microanalyse, Institut de Chimie des Substances Naturelles, CNRS, F-91198, Gif-sur-Yvette.

General procedure for the synthesis of *N*-allylbenzenesulfonamides 1

To a solution of NaOH (1.2 equiv) in water (0.5 mL/mmol allylamine) and Et_2O (0.5 mL/mmol allylamine) was added allylamine (1 equiv) and the arenesulfonyl chloride (1.1 equiv) at room temperature. After 5 h, the reaction mixture was extracted with CH_2Cl_2 . The organic extracts were dried and evaporated, and the residue was purified by flash column chromatography.

N-Allyl-*p*-toluenesulfonamide (1a)

White solid, 99% (petroleum ether: AcOEt 6:4).

Tehrani, K. A.; Van, T. N.; Karikomi, M.; Rottiers, M.; De Limpe, N. *Tetrahedron* **2002**, 58, 7145.

N-Allyl-4-*tert*-butylbenzenesulfonamide (1b)

White solid, 85% (petroleum ether: AcOEt 7:3).

mp: 65-68 $^{\circ}\text{C}$.

IR (CCl_4): ν 3395, 3276, 1596 (NH), 1343, 1167 (SO_2) cm^{-1} .

^1H -NMR (CDCl_3 , 400 MHz): δ 1.34 (s, 9 H, CH_3), 3.59 (t, J = 5.6 Hz, 2 H, CH_2), 4.75 (t, J = 5.8 Hz, 1 H, NH), 5.09 (d, J = 10.4 Hz, 1 H, CH=), 5.17 (d, J = 17.2 Hz, 1 H, CH=), 5.67-5.80 (m, 1 H, CH=), 7.52 (d, J = 8 Hz, 2 H, ArH), 7.79 (d, J = 8 Hz, 2 H, ArH).

^{13}C -NMR (CDCl_3 , 100 MHz): δ 31.1 (CH_3), 35.1 (C), 45.7 (CH_2), 117.6 ($\text{CH}_2=$), 126.0 (CH Ar), 126.9 (CH Ar), 133.1 (CH=), 136.9 (C-*ipso*), 156.4 (C-*p*).

MS (Cl, NH₃): 254 (MH⁺), 271 (MNH₄⁺).

N-Allyl-2-chlorobenzenesulfonamide (1c)

White solid, 55% (petroleum ether: AcOEt 9:1).

mp: 67-70 °C.

IR (CCl₄): ν 3392, 1572 (NH), 1352, 1172 (SO₂) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 3.58 (t, J = 6 Hz, 2 H, CH₂), 5.07 (d, J = 10.4 Hz, 1 H, CH=), 5.10 (bs, 1 H, NH), 5.16 (d, J = 16.8 Hz, 1 H, CH=), 5.69 (ddt, J = 6, 10.4, 16.8 Hz, 1 H, CH=), 7.41 (ddd, J = 2.5, 5.7, 7.9 Hz, 1 H, ArH), 7.45-7.55 (m, 2 H, ArH), 8.08 (d, J = 7.2 Hz, 1 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 45.9 (CH₂), 118.0 (CH₂=), 127.2 (CH Ar), 131.26 (CH Ar), 131.33 (CH Ar), 131.5 (CH Ar), 132.5 (CH=), 133.7 (CH Ar), 137.4 (C-*ipso*).

MS (Cl, NH₃): 232 (MH⁺), 249 (MNH₄⁺).

N-Allyl-4-bromobenzenesulfonamide (1d)

White solid, 76% (petroleum ether: AcOEt 8:2).

For NMR description, see: Klapars, A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2002**, 124, 14844.

N-Allyl-4-iodobenzenesulfonamide (1e)

White solid, 80% (petroleum ether: AcOEt 8:2).

For NMR description, see: Klapars, A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2002**, 124, 14844.

N-Allyl-4-trifluoromethylbenzenesulfonamide (1f)

White solid, 73% (petroleum ether: AcOEt 9:1).

mp: 100-102 °C.

IR (CCl₄): ν 3399 (NH), 1323, 1174 (SO₂) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 3.65 (d, J = 5.6 Hz, 2 H, CH₂), 4.68 (bs, 1 H, NH), 5.13 (dd, J = 0.8, 10.4 Hz, 1 H, CH=), 5.18 (dd, J = 0.8, 16.8 Hz, 1 H, CH=), 5.72 (ddt, J = 6, 10.4, 16.8 Hz, 1 H, CH=), 7.80 (d, J = 8.4 Hz, 2 H, ArH), 8.02 (d, J = 8 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 45.6 (CH₂), 117.9 (CH₂=), 123.1 (q, J = 271.4 Hz, CF₃), 126.2 (C-*m*), 127.6 (C-*o*), 132.4 (CH=), 134.2 (q, J = 32.5 Hz, C-*p*), 143.5 (C-*ipso*).

MS (Cl, NH₃): 266 (MH⁺), 283 (MNH₄⁺).

N-Allyl-4-cyanobenzenesulfonamide (1g)

White solid, 93% (petroleum ether: AcOEt 7:3).

mp: 80-81 °C.

IR (CCl₄): ν 3400, 3263 (NH), 2233 (CN), 1350, 1165 (SO₂) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 3.65 (t, *J* = 5.4 Hz, 2 H, CH₂), 4.93 (t, *J* = 5.4 Hz, 2 H, NH), 5.11 (d, *J* = 10.4 Hz, 1 H, CH=), 5.16 (d, *J* = 17.2 Hz, 1 H, CH=), 5.69 (ddt, *J* = 5.6, 11.2, 16.8 Hz, 1 H, CH=), 7.82 (d, *J* = 8.4 Hz, 2 H, ArH), 7.99 (d, *J* = 8.4 Hz, 2 H, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 45.7 (CH₂), 116.4 (C-*p*), 117.2 (CN), 118.2 (CH₂=), 127.6 (CH Ar), 132.4 (CH=), 132.9 (CH Ar), 144.5 (C-*ipso*).

MS (CI, NH₃): 223 (MH⁺), 240 (MNH₄⁺).

***N*-Acetyl-*N*-allyl-*p*-toluenesulfonamide (1h)**

To a solution of sulfonamide **1a** (5 g, 23.7 mmol) in toluene (47 mL) was added acetyl chloride (25.5 mL, 356 mmol) and was heated at 110 °C for 2 days. After that time the reaction mixture was filtered and crystallized from ethanol, obtaining the desired sulfonamide as a white solid (3.6 g, 60%).

For NMR description, see: Ozaki, S.; Matsushita, H.; Ohmori, H.; *J. Chem. Soc., Perkin Trans 1*, **1993**, 2339.

Ethoxythiocarbonylsulfanylacetic acid methyl ester (2a)

For preparation details and NMR description, see: Boutillier, P. Ph. D. Thesis, École Polytechnique, Palaiseau, 2001.

Dithiocarbonic acid *O*-ethyl ester *S*-(2-oxo-3-phenylpropyl) ester (2b)

To a solution of phenylacetone (10 mL, 74.8 mmol) in glacial AcOH (25 mL) was added concentrated HBr (12.5 mL) and a solution of Br₂ (8.5 mL, 165 mmol) in glacial AcOH (41 mL), and the resulting mixture was stirred at room temperature for 6 h. Acetone (132 mL) was then added, and stirring continued for a further 18 h. The reaction mixture was then concentrated, and extracted with CH₂Cl₂. The organic extracts were dried and concentrated. The residue was purified by flash column chromatography (petroleum ether: AcOEt 95:5), which afforded 1-bromo-3-phenylacetone as a dark yellow oil, as well as other minority isomers due to non-regioselective bromination. A mixture of the intermediate bromide and potassium ethylxanthogenate (15.6 g, 97.2 mmol) in acetone (375 mL) was stirred at room temperature for 4 h. The reaction mixture was then concentrated, the residue taken up in Et₂O and filtrated through Celite™. The solvent was evaporated, and the residue was purified by flash column chromatography (petroleum ether: AcOEt 9:1), and the desired xanthate was obtained as a brown solid (16.9 g, 89%, two steps).

mp: 55-57 °C.

IR (CCl₄): ν 1730 (CO), 1224 (COS), 1052 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.39 (t, *J* = 7.2 Hz, 3 H, CH₃), 3.90 (s, 2 H, CH₂), 4.00 (s, 2 H, CH₂), 4.60 (q, *J* = 7.2 Hz, 2 H, OCH₂), 7.20-7.40 (m, 5 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.6 (CH₃), 44.8 (CH₂), 44.9 (CH₂), 70.8 (OCH₂), 127.2, 128.7, 129.5 (CH Ar), 133.2 (C-*ipso*), 200.6 (CO), 213.0 (CS).

MS (CI, NH₃): 255 (MH⁺), 272 (MNH₄⁺).

Dithiocarbonic acid *S*-(2-cyclohexyl-2-oxoethyl) ester *O*-ethyl ester (2c)

To a solution of cyclohexane carboxylic acid (2.45 g, 19.1 mmol) in THF (143 mL) at 0 °C was added MeLi (48 mL, 1.6 M solution in cyclohexane, 76.3 mmol) and the reaction mixture stirred at that temperature for 2 h. TMSCl (48.5 mL, 382 mmol) was then added, and the ice bath was removed. 1 M aq. HCl (7.5 mL) was then added, and the reaction mixture was stirred for 30 minutes. The two layers were then separated, and the aqueous phase was further extracted with Et₂O. The organic extracts

were dried and evaporated. The residue was purified by flash column chromatography (petroleum ether: AcOEt 9:1), and acetylcyclohexane was obtained as a yellow oil (1.20 g, 50%).

Acetylcyclohexane (1.2 g, 9.52 mmol) was dissolved in MeOH (7.5 mL), cooled to 0 °C, and a single portion Br₂ (0.5 mL) added, and the resulting mixture stirred at a temperature below 15 °C for 3 h. Water (8 mL) was then added and stirring continued for a further 15 h at room temperature. The reaction mixture was then extracted with AcOEt, and the organic extracts were dried and evaporated. The residue was purified by flash column chromatography (petroleum ether: AcOEt 9:1), and the desired bromoacetylcyclohexane was obtained as a yellow oil (1.35 g, 70%).

To a solution of bromoacetylcyclohexane (1.35 g, 6.59 g) in acetone (33 mL) was added potassium ethylxanthogenate (1.37 g, 8.56 g), and the mixture was stirred at room temperature for 1.5 h. The reaction mixture was then concentrated, the residue taken up in Et₂O, and filtered through Celite™. The solvent was evaporated, and the residue was purified by flash column chromatography (petroleum ether: AcOEt 97:3), and the desired xanthate was obtained as a yellow semi-solid (1.03 g, 64%).

IR (CCl₄): ν 1713 (CO), 1224 (COS), 1052 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.25-1.50 (m, 5 H), 1.40 (t, J = 7.2 Hz, 3 H, CH₃), 1.67 (d, J = 11.2 Hz, 3 H, CH₃), 1.79 (d, J = 12.4 Hz, 2 H), 1.90 (d, J = 12.8 Hz, 2 H), 2.59 (tt, J = 3.3, 10.9 Hz, 1 H, H-1 cyclohexyl), 4.07 (s, 2 H, CH₂S), 4.62 (q, J = 7.1 Hz, 2 H, OCH₂).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃), 25.5 (C-4 cyclohexyl), 25.7 (C-3 and C-5 cyclohexyl), 28.5 (C-2 and C-6 cyclohexyl), 44.5 (CH₂S), 49.8 (C-1 cyclohexyl), 70.6 (OCH₂), 205.7 (CO), 213.5 (CS).

MS (Cl, NH₃): 247 (MH⁺), 264 (MNH₄⁺).

Dithiocarbonic acid O-ethyl ester S-(2-oxotetrahydrofuran-3-yl) ester (2d)

For preparation details and NMR description, see: Tanaka, K.; Unema, H.; Yamagishi, N.; Tanikaga, R.; Kaji, A.; *Bull. Chem. Soc. Jpn.* **1980**, 53, 2910.

Dithiocarbonic acid O-ethyl ester S-(2-oxo-2-piperidin-1-yl-ethyl) ester (2e)

To a suspension of K₂CO₃ (55.8 g, 404 mmol) and chloroacetyl chloride (8.1 mL, 101 mmol) in THF (116 mL) was added dropwise a solution of piperidine (10 mL, 101 mmol) in THF (232 mL), and the resulting mixture was stirred at room temperature for 16 h. The reaction mixture was then filtered and concentrated. The residue was purified by flash column chromatography (petroleum ether: AcOEt 7:3), and the desired chloroacetylpiperidine was obtained as a yellow oil (13.1 g, 80%).

To a solution of chloroacetylpiperidine (13.1 g, 81.1 mmol) in acetone (81 mL) at 0 °C was added dropwise a suspension of potassium ethylxanthogenate (14.3 g, 89.2 mmol) in acetone (357 mL), and the resulting mixture was stirred at room temperature for 3 h. The reaction mixture was then concentrated, the residue taken up in Et₂O and filtered through Celite™. The solvent was evaporated, and the residue purified by flash column chromatography (petroleum ether: AcOEt 7:3) and the desired xanthate was obtained as a yellow oil (16 g, 80%).

IR (CCl₄): ν 1651 (CO), 1220 (COS), 1054 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.41 (t, J = 7.2 Hz, 3 H, CH₃), 1.50-1.70 (m, 6 H), 3.48 (t, J = 5.2 Hz, 2 H, CH₂N), 3.56 (t, J = 5.2 Hz, 2 H, CH₂N), 4.11 (s, 2 H, CH₂CO), 4.64 (q, J = 7.2 Hz, 2 H, OCH₂).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃), 24.3, 25.4, 26.5 (CH₂ ring), 39.6 (CH₂CO), 43.4 (CH₂N), 47.3 (CH₂N), 70.4 (OCH₂), 164.7 (CO), 213.9 (CS).

MS (Cl, NH₃): 248 (MH⁺), 265 (MNH₄⁺).

Dithiocarbonic acid O-ethyl ester S-(2-oxo-2-thiophen-2-yl-ethyl) ester (2f)

For preparation details and NMR description, see: Boutillier, P. Ph. D. Thesis, École Polytechnique, Palaiseau, 2001.

Dithiocarbonic acid S-(3,3-dimethyl-2-oxobutyl) ester O-ethyl ester (2g)

For preparation details and NMR description, see: Batanero, B.; Picazo, O.; Barba, F. *J. Org. Chem.* **2001**, 66, 320.

Dithiocarbonic acid O-ethyl ester S-(2-oxo-2-phenylethyl) ester (2h)

For preparation details and NMR description, see: Batanero, B.; Picazo, O.; Barba, F. *J. Org. Chem.* **2001**, 66, 320.

2-Ethoxythiocarbonylsulfanyl-3,3-dimethoxypropionic acid methyl ester (2i)

For preparation details and NMR description, see: Sortais, B. Ph. D. Thesis, École Polytechnique, Palaiseau, 2002.

Dithiocarbonic acid O-ethyl ester S-[2-(4-methoxyphenyl)-2-oxoethyl] ester (2j)

For preparation details and NMR description, see: Batanero, B.; Picazo, O.; Barba, F. *J. Org. Chem.* **2001**, 66, 320.

General procedure for the synthesis of addition products 3

To a solution of xanthate **2** (1.5 equiv) in 1,2-dichloroethane (1 mL/mmol of **2**) was added sulfonamide **1** (1 equiv), and the mixture heated at reflux temperature. DLP (5% molar) was then added every 1.5 h until disparition of starting material. Then the reaction mixture was evaporated, and the residue was purified by flash column chromatography.

4-Ethoxythiocarbonylsulfanyl-5-(toluene-4-sulfonylamino)pentanoic acid methyl ester (3a)

Yellow oil, 85% (13% DLP, petroleum ether: AcOEt 6:4).

IR (film): ν 1741 (CO), 1598 (NH), 1495, 1437, 1337, 1163 (SO₂), 1222 (COS), 1052 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.41 (t, J = 8 Hz, 3 H, CH₃), 1.85-1.95 (m, 1 H, CH), 2.13-2.19 (m, 1 H, CH), 2.44 (s, 3 H, CH₃ tolyl), 2.46-2.50 (m, 2 H, CH₂CO), 3.21-3.24 (m, 2 H, CH₂N), 3.68 (s, 3 H, OCH₃), 3.81-3.85 (m, 1 H, CHS), 4.60 (q, J = 7.2 Hz, 2 H, OCH₂), 7.30 (d, J = 8 Hz, 2 H, CH Ar), 7.71 (d, J = 8 Hz, 2 H, CH Ar).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.8 (CH₃), 21.7 (CH₃ tolyl), 26.2 (CH₂), 31.1 (CH₂CO), 44.6 (OCH₃), 46.3 (CH₂N), 50.1 (CHS), 70.6 (OCH₂), 127.2, 127.5, 129.9, 131.2 (CH Ar), 136.7 (C-*p*), 143.7 (C-*ipso*), 176.0 (CO), 212.6 (CS).

MS (CI, NH₃): 406 (MH⁺), 423 (MNH₄⁺).

Dithiocarbonic acid O-ethyl ester S-{4-oxo-5-phenyl-1-[(toluene-4-sulfonylamino)methyl]pentyl} ester (3b)

Yellow oil, 70% (80% DLP, petroleum ether: AcOEt 8:2).

IR (CCl₄): ν 3250 (NH), 1715 (CO), 1343, 1165 (SO₂), 1222 (COS), 1052 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.39 (t, J = 7.2 Hz, 3 H, CH₃), 1.78 (ddd, J = 6.8, 8.8, 21.6 Hz, 1 H, CH), 2.06 (ddd, J = 7.2, 12.2, 14.4 Hz, 1 H, CH), 2.42 (s, 3 H, CH₃ tolyl), 2.61 (t, J = 7 Hz, 2 H, CH₂CO), 3.15 (t, J = 6.2 Hz, 2 H, CH₂N), 3.67 (s, 2 H, CH₂Ph), 3.66-3.78 (m, 1 H, CHS), 4.52-4.66 (m, 2 H, OCH₂), 5.14 (t, J = 6.4 Hz, 1 H, NH), 7.17 (d, J = 6.8 Hz, 2 H, ArH), 7.23-7.34 (m, 5 H, ArH), 7.73 (d, J = 8.4 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.6 (CH₃), 21.5 (CH₃ tolyl), 24.6 (CH₂), 38.6 (CH₂CO), 46.2 (CH₂N), 49.8 (CHS), 50.0 (CH₂Ph), 70.3 (OCH₂), 127.0, 128.7, 129.3, 129.7 (CH Ar), 133.8 (C Ar), 136.8 (C Ar), 143.5 (C Ar), 207.1 (CO), 212.5 (CS).

MS (Cl, NH₃): 466 (MH⁺), 483 (MNH₄⁺).

Dithiocarbonic acid S-{1-[(4-*tert*-butyl-benzenesulfonylamino)methyl]-4-cyclohexyl-4-oxobutyl} ester O-ethyl ester (3c)

Yellow oil, 80% (15% DLP, petroleum ether: AcOEt 8:2).

IR (CCl₄): ν 3274, 1596 (NH), 1710 (CO), 1342, 1167 (SO₂), 1221, 1113 (COS), 1053 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.10-1.30 (m, 5 H, CH cyclohexyl), 1.32 (s, 9 H, CH₃ ^{*t*}Bu), 1.37 (t, J = 7 Hz, 3 H, CH₃), 1.63 (d, J = 10.4 Hz, 1 H, CH cyclohexyl), 1.67-1.75 (m, 5 H, 4 CH cyclohexyl and CH), 1.98-2.12 (m, 1 H, CH), 2.22-2.35 (m, 1 H, H-1 cyclohexyl), 2.56 (t, J = 6.8 Hz, 2 H, CH₂CO), 3.15 (dd, J = 6.8, 13.6 Hz, 1 H, CHN), 3.71 (dd, J = 6.2, 13.4 Hz, 1 H, CHN), 3.70-3.80 (m, 1 H, CHS), 4.58 (q, J = 6.3 Hz, 2 H, OCH₂), 5.29 (t, J = 6.2 Hz, 1 H, NH), 7.49 (d, J = 8 Hz, 2 H, ArH), 7.76 (d, J = 8 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.6 (CH₃), 24.5 (CH₂ cyclohexyl), 25.5 (CH₂), 25.7 (CH₂ cyclohexyl), 28.4 (CH₂ cyclohexyl), 31.0 (CH₃ ^{*t*}Bu), 35.0 (C ^{*t*}Bu), 37.1 (CH₂CO), 46.3 (CH₂N), 50.0 (CHS), 50.7 (C-1 cyclohexyl), 70.2 (OCH₂), 126.0 (CH Ar), 126.8 (CH Ar), 136.8 (C Ar), 156.4 (C Ar), 212.6 and 212.9 (CO and CS).

MS (Cl, NH₃): 500 (MH⁺), 517 (MNH₄⁺).

Dithiocarbonic acid S-[2-(2-chlorobenzenesulfonylamino)-1-(2-oxotetrahydrofuran-3-yl-methyl)ethyl] ester O-ethyl ester (3d) 2 diastereomers, D1 and D2

Yellow oil, 92% (15% DLP, petroleum ether: AcOEt 8:2).

IR (CCl₄): ν 3392, 1572 (NH), 1352, 1172 (SO₂) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.39 and 1.41 (2 t, J = 6.4 Hz, 6 H, CH₃), 1.78 (ddd, J = 7, 8.4, 15 Hz, 1 H, CH D2), 1.87-2.07 (m, 3 H, CH D1, CH ring D1 and CH ring D2), 2.16 (ddd, J = 4.3, 10.5, 14.7 Hz, 1 H, CH D1), 2.27 (dt, J = 6.9, 14.8 Hz, 1 H, CH D2), 2.36-2.50 (m, 2 H, CH ring D1 and CH ring D2), 2.70 (ddd, J = 7, 10.8, 14.8 Hz, 1 H, CHCO D2), 2.78 (ddd, J = 4.4, 9.8, 19.4 Hz, 1 H, CHCO D1), 3.16-3.36 (m, 4 H, CH₂N), 3.92 (td, J = 5.6, 10.8 Hz, 1 H, CHS D1), 4.08 (dt, J = 5.8, 15.2 Hz, 1 H, CHS D2), 4.18 (dd, J = 9.2, 16.4 Hz, 2 H, CHO ring D1 and CHO ring D2), 4.36 (dd, J = 7.8, 15.8 Hz, 1 H, CHO ring D1 and CHO ring D2), 4.57-4.65 (m, 4 H, OCH₂), 5.52 (t, J = 7.2 Hz, 1 H, NH), 5.56 (t, J = 6.8 Hz, 1 H, NH), 7.40-7.45 (m, 2 H, ArH), 7.50-7.55 (m, 4 H, ArH), 8.08 (t, J = 7.6 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃), 29.2 (CH₂ ring D1), 29.4 (CH₂ ring D2), 31.5 (CH₂ D1), 32.2 (CH₂ D2), 36.96 (CHCO D1), 37.00 (CHCO D2), 46.4 (CH₂N D2), 46.7 (CH₂N D1), 48.8 (CHS D2), 49.0 (CHS D1), 66.45 (CH₂O ring D2), 66.51 (CH₂O ring D1), 70.5 (OCH₂ D2), 70.7 (OCH₂ D1), 127.3 (CH Ar), 131.1 (CH Ar D2), 131.3 (CH Ar D1), 131.39 (CCl D1), 131.42 (CCl D2), 131.6 (CH Ar D1), 131.7 (CH Ar D2), 133.8 (CH Ar D2), 133.9 (CH Ar D1), 136.9 (C-*ipso* D1), 137.0 (C-*ipso* D2), 178.4 (CO D2), 178.5 (CO D1), 211.9 (CS D2), 212.2 (CS D1).

MS (Cl, NH₃): 438 (MH⁺), 455 (MNH₄⁺).

Dithiocarbonic acid S-{1-[(4-bromobenzenesulfonylamino)methyl]-4-oxo-4-piperidin-1-yl-butyl} ester O-ethyl ester (3e)

Yellow oil, 66% (35% DLP, petroleum ether: AcOEt 1:1).

IR (CCl₄): ν 3197, 1576 (NH), 1632 (CO), 1347, 1168 (SO₂), 1221 (COS), 1054 (CS).

¹H-NMR (CDCl₃, 400 MHz): δ 1.40 (t, J = 7.2 Hz, 3 H, CH₃), 1.45-1.65 (m, 6 H, CH₂ piperidine), 1.80-1.96 (m, 1 H, CH), 2.07-2.19 (m, 1 H, CH), 2.32-2.48 (m, 2 H, CH₂CO), 3.13 (dt, J = 6.7, 13.4 Hz, 1 H, CHN), 3.26 (ddd, J = 5.6, 6.4, 13.2 Hz, 1 H, CHN), 3.34 (t, J = 5.4 Hz, 2 H, 2 CH piperidine), 3.50 (t, J = 5.6 Hz, 2 H, 2 CH piperidine), 3.74-3.83 (m, 1 H, CHS), 4.53-4.66 (m, 2 H, OCH₂), 6.13 (t, J = 6.2 Hz, 1 H, NH), 7.62 (d, J = 8.8 Hz, 2 H, ArH), 7.74 (d, J = 8.4 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃), 24.4 (CH₂ piperidine), 25.5 (CH₂ piperidine), 25.7 (CH₂), 26.3 (CH₂ piperidine), 29.6 (CH₂CO), 43.0 (NCH₂ piperidine), 46.0 (NCH₂), 46.5 (NCH₂ piperidine), 49.9 (CHS), 70.4 (OCH₂), 127.3 (C Ar), 128.7 (CH Ar), 132.2 (CH Ar), 139.3 (C Ar), 169.9 (CO), 212.7 (CS).

MS (Cl, NH₃): 525 (MH⁺).

Dithiocarbonic acid O-ethyl ester S-{4-oxo-4-thiophen-2-yl-1-[(4-trifluoromethylbenzenesulfonylamino)methyl]butyl} ester (3f)

Yellow oil, 49% (1.2 equiv DLP, petroleum ether: AcOEt 7:3).

IR (CCl₄): ν 3283, 1517 (NH), 1668 (CO), 1322, 1173 (SO₂), 1226 (COS), 1063 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.37 (t, J = 7.2 Hz, 3 H, CH₃), 1.94-2.04 (m, 1 H, CH), 2.22-2.32 (m, 1 H, CH), 3.07 (t, J = 6.8 Hz, 2 H, CH₂CO), 3.31 (td, J = 2.1, 6.4 Hz, 2 H, CH₂N), 3.82-3.90 (m, 1 H, CHS), 4.50-4.63 (m, 2 H, OCH₂), 5.47 (t, J = 6.4 Hz, 1 H, NH), 7.12 (dd, J = 4, 4.8 Hz, 1 H, H-4 thiophene), 7.64 (dd, J = 1, 5 Hz, 1 H, CH thiophene), 7.70 (dd, J = 1, 5 Hz, 1 H, CH thiophene), 7.77 (d, J = 8.4 Hz, 2 H, ArH), 8.00 (d, J = 8 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.6 (CH₃), 25.1 (CH₂), 35.9 (CH₂CO), 46.4 (CH₂N), 50.0 (CHS), 70.5 (OCH₂), 123.2 (q, J = 271.1 Hz, CF₃), 126.3 (q, J = 3.7 Hz, C-*m*), 127.6 (C-*o*), 128.2 (C-4 thiophene), 132.2 (CH thiophene), 134.0 (CH thiophene), 134.4 (q, J = 32.9 Hz, C-*p*), 143.56 (C-*ipso*), 143.61 (C-*ipso*), 191.8 (CO), 212.4 (CS).

MS (Cl, NH₃): 512 (MH⁺), 529 (MNH₄⁺).

Dithiocarbonic acid S-{1-[(4-cyanobenzenesulfonylamino)methyl]-5,5-dimethyl-4-oxohexyl} ester O-ethyl ester (3g)

Yellow oil, 81% (30% DLP, petroleum ether: AcOEt 8:2).

IR (CCl₄): ν 3225 (NH), 2234 (CN), 1705 (CO), 1352, 1166 (SO₂), 1225 (COS), 1052 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.10 (s, 9 H, CH₃ ^{*t*}Bu), 1.40 (t, J = 7 Hz, 3 H, CH₃), 1.69-1.81 (m, 1 H, CH), 2.00-2.10 (m, 1 H, CH), 2.55-2.72 (m, 2 H, CH₂CO), 3.23 (dd, J = 6.4, 13.2 Hz, 1 H, CHN), 3.29 (dd, J = 6, 13.2 Hz, 1 H, CHN), 3.63-3.76 (m, 1 H, CHS), 4.53-4.66 (m, 2 H, OCH₂), 5.46 (t, J = 6 Hz, 1 H, NH), 7.81 (d, J = 8 Hz, 2 H, H-*m*), 7.99 (d, J = 8 Hz, 2 H, H-*o*).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃), 24.5 (CH₂), 26.4 (CH₃ ^{*t*}Bu), 33.3 (CH₂CO), 44.1 (C ^{*t*}Bu), 46.6 (CH₂N), 50.1 (CHS), 70.6 (OCH₂), 116.4 (C-*p*), 117.3 (CN), 127.7 (C-*o*), 132.9 (C-*m*), 144.4 (C-*ipso*), 212.8 and 215.2 (CO and CS).

MS (Cl, NH₃): 443 (MH⁺), 460 (MNH₄⁺).

Dithiocarbonic acid O-ethyl ester S-{1-[(4-iodobenzenesulfonylamino)methyl]-5,5-dimethoxy-4-oxopentyl} ester (3h)

Yellow oil, 95% (15% DLP, petroleum ether: AcOEt 8:2).

IR (CCl₄): ν 3279 (NH), 1740 (CO), 1316, 1166 (SO₂), 1221 (COS), 1054 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.40 (t, 3 H, J = 7.1 Hz, OCH₃), 1.75-1.86 (m, 1H, CH), 2.00-2.13 (m, 1 H, CH), 2.83-2.88 (m, 0.5 H, CHCO), 2.93-2.98 (m, 0.5 H, CHCO), 3.23-3.37 (m+s, 8 H, CH₂N, 2 OCH₃), 3.64-3.88 (m+s, 4 H, CH, CO₂CH₃), 4.49-4.52 (m, 1 H, CHS), 4.58-4.65 (m, 2 H, OCH₂), 5.09 (m, 1 H, NH), 7.57 (d, 2 H, J = 7.7 Hz, ArH), 7.87 (d, 2 H, J = 8.3 Hz, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.8 (CH₃) 28.4, 28.9 (CH₂), 45.8 (CH₂N), 46.5, 46.7 (CH), 47.2 (CH₂N), 48.9, 49.1 (CHCO), 52.3, 53.9, 54.1, 54.9, 55.0 (OCH₃), 70.6, 70.7 (OCH₂), 100.2 (Cl), 104.5, 104.8 (CHOCH₃), 128.6, 138.5, (CH Ar), 172.5, 172.6 (CO), 212.8 (CS).

MS (Cl, NH₃): 609 (MNH₄⁺).

General procedure for the preparation of acetylated products 4

Method A: To a solution of sulfonamide **3** (1 equiv) in pyridine (53 equiv) was added acetic anhydride (17 equiv), and the reaction mixture stirred until disparition of starting material. The mixture was then evaporated, and CH₂Cl₂ was added. The organic phase was washed with 1 M aq. HCl, sat. aq. NaHCO₃ solution and sat. aq. NaCl solution, dried and evaporated. The residue was purified by flash column chromatography.

Method B: To a solution of sulfonamide **3** (1 equiv) in CH₂Cl₂ (2 mL/mmol of **3**) was added DMAP (4 equiv), and the solution was cooled to 0 °C. Acetyl chloride was then added dropwise (3 equiv), and the resulting mixture was stirred at room temperature for 2 h. CH₂Cl₂ was then added, and the organic phase was washed with 1 M aq. HCl, sat. aq. NaHCO₃ solution and sat. aq. NaCl solution, dried and evaporated. The residue was purified by flash column chromatography.

5-[Acetyl(toluene-4-sulfonyl)amino]-4-ethoxythiocarbonylsulfanylpentanoic acid methyl ester (4a)

Yellow oil, 95% (method A, petroleum ether: AcOEt 7:3).

IR (film): ν 1741 (CO), 1707 (CO), 1366, 1169 (SO₂), 1227 (CSO), 1051 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.45 (t, J = 8 Hz, CH₃ xanthate), 1.85-1.95 (m, 1 H, CH), 2.13-2.19 (m, 1 H, CH), 2.35 (s, 3 H, CH₃ acetyl), 2.46 (s, 3 H, CH₃ tolyl), 2.51-2.59 (m, 2 H, CH₂CO), 3.68 (s, 3 H, OCH₃), 4.10-4.15 (m, 2 H, CH₂N), 4.18-4.23 (m, 1 H, CHS), 4.65 (q, J = 7.2 Hz, 2 H, OCH₂), 7.36 (d, J = 8 Hz, 2 H, ArH), 7.74 (d, J = 8 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.8 (CH₃ xanthate), 21.7 (CH₃ acetyl), 25.2 (CH₂), 26.6 (CH₃ tolyl), 31.4 (CH₂CO), 49.06 (OCH₃), 49.1 (CH₂N), 49.5 (CHS), 70.5 (OCH₂), 126.3, 127.6, 130.2, 130.3 (CH Ar), 136.6 (C Ar), 145.3 (C Ar), 170.2 (NCO), 174.0 (CO), 212.8 (CS).

MS (Cl, NH₃): 448 (MH⁺), 465 (MNH₄⁺).

Dithiocarbonic acid S-(1-{[acetyl(toluene-4-sulfonyl)amino]methyl}-4-oxo-5-phenylpentyl) ester O-ethyl ester (4b)

Brown oil, 94% (method B, petroleum ether: AcOEt 7:3).

IR (CCl₄): ν 1759 (CO), 1710 (CO), 1367, 1170 (SO₂), 1224 (COS), 1049 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.42 (t, J = 7.1 Hz, 3 H, CH₃ xanthate), 1.75-1.90 (m, 1 H, CH), 2.05-2.15 (m, 1 H, CH), 2.33 (s, 3 H, CH₃ acetyl), 2.44 (s, 3 H, CH₃ tolyl), 2.60-2.75 (m, 2 H, CH₂CO), 3.70 (s, 2 H, CH₂Ph), 4.00-4.30 (m, 3 H, CH₂N and CHS), 4.63 (q, J = 7.1 Hz, 2 H, OCH₂), 7.20 (d, J = 7.1 Hz, 2 H, ArH), 7.75 (dd, J = 2.8, 8.1 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃ xanthate), 21.6 (CH₃ tolyl), 25.0 (CH₃ acetyl), 25.1 (CH₂), 38.9 (CH₂CO), 48.9 (CH₂N), 49.4 (CHS), 50.1 (CH₂Ph), 70.3 (OCH₂), 127.4, 128.7, 129.4, 130.0 (CH Ar), 134.0 (C Ar), 136.6 (C Ar), 145.1 (C Ar), 170.4 (CO acetyl), 206.8 (CO), 212.7 (CS).
MS (Cl, NH₃): 509 (MH⁺), 525 (MNH₄⁺).

Dithiocarbonic acid S-(1-[[acetyl-(4-tert-butylbenzenesulfonyl)amino]methyl]-4-cyclohexyl-4-oxobutyl) ester O-ethyl ester (4c)

Yellow oil, 93% (method B, petroleum ether: AcOEt 8:2).

IR (CCl₄): ν 1709 (CO), 1366, 1172 (SO₂), 1227, 1113 (COS), 1049 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.10-1.40 (m, 5 H, CH ring), 1.34 (s, 9 H, CH₃^tBu), 1.43 (t, *J* = 6.8 Hz, 3 H, CH₃ xanthate), 1.65 (d, *J* = 13.2 Hz, 1 H, CH cyclohexyl), 1.70-1.85 (m, 5 H, CH and 4 CH cyclohexyl), 2.05-2.15 (m, 1 H, CH), 2.25-2.40 (m, 1 H, H-1 cyclohexyl), 2.35 (s, 3 H, CH₃ acetyl), 2.55-2.75 (m, 2 H, CH₂CO), 4.02-4.20 (m, 3 H, CH₂N and CHS), 4.58-4.70 (m, 2 H, OCH₂ xanthate), 7.54 (d, *J* = 8.4 Hz, 2 H, ArH), 7.78 (d, *J* = 8.4 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃ xanthate), 25.0 (CH₂), 25.1 (CH₃ acetyl), 25.62, 25.64, 25.8, 28.46, 28.52 (CH₂ cyclohexyl), 31.0 (CH₃^tBu), 35.3 (C^tBu), 37.5 (CH₂CO), 49.0 (CH₂N), 49.6 (CHS), 50.9 (C-1 cyclohexyl), 70.3 (OCH₂ xanthate), 126.4 (CH Ar), 127.3 (CH Ar), 136.5 (C Ar), 158.0 (C Ar), 170.3 (CO acetyl), 212.5 and 212.8 (CO and CS).

MS (Cl, NH₃): 542 (MH⁺), 559 (MNH₄⁺).

Dithiocarbonic acid S-[2-[acetyl(2-chlorobenzenesulfonyl)amino]-1-(2-oxotetrahydrofuran-3-ylmethyl)ethyl] ester O-ethyl ester (4d)
2 diastereomers, D1 and D2

Yellow oil, 89% (method B, petroleum ether: AcOEt 1:1).

IR (CCl₄): ν 1780 (CO ester), 1709 (CO amide), 1368, 1168 (SO₂), 1227 (COS), 1045 (CS).

¹H-NMR (CDCl₃, 400 MHz): δ 1.42 (t, *J* = 3.6 Hz, 3 H, CH₃), 1.45 (t, *J* = 3.6 Hz, 3 H, CH₃), 1.80-1.92 (m, 2 H, CH D1 and CH D2), 1.94-2.17 (m, 2 H, CH ring D1 and CH ring D2), 2.19-2.39 (m, 2 H, CH D1 and CH D2), 2.25 (s, 3 H, CH₃), 2.28 (s, 3 H, CH₃), 2.48 (dddd, *J* = 2, 6.4, 8.6, 14.8 Hz, 1 H, CH ring D1), 2.56 (dddd, *J* = 2, 6.4, 8.4, 14.8 Hz, 1 H, CH ring D2), 2.75-2.91 (m, 2 H, CHCO D1 and CHCO D2), 4.14-4.27 (m, 6 H, CH₂N D1, CH₂N D2, CHO ring D1 and CHO ring D2), 4.32-4.46 (m, 4 H, CHS D1, CHS D2, CHO ring D1 and CHO ring D2), 4.59-4.73 (m, 4 H, OCH₂ D1 and D2), 7.44-7.62 (m, 6 H, ArH), 8.19 (d, *J* = 8 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃), 24.7 (CH₃ acetyl D2), 24.8 (CH₃ acetyl D1), 29.1 (CH₂ ring D1), 29.4 (CH₂ ring D2), 31.9 (CH₂ D1), 32.8 (CH₂ D2), 37.2 (CH₂ ring D2), 37.4 (CH₂ ring D1), 48.1 (CHS D2), 48.6 (CHS D1), 49.4 (CH₂N D2), 50.4 (CH₂N D1), 66.4 (CHO ring D1), 66.5 (CHO ring D2), 70.4 (OCH₂ D2), 70.7 (OCH₂ D1), 127.3 (CH Ar), 131.8 (C Ar D2), 131.9 (C Ar D1), 132.1 (CH Ar D2), 132.2 (CH Ar D1), 132.36 (CH Ar D1), 132.41 (CH Ar D2), 134.8 (CH Ar D2), 134.9 (CH Ar D1), 136.7 (C Ar D1), 136.9 (C Ar D2), 169.8 (CO acetyl D1), 169.9 (CO acetyl D2), 178.38 (CO), 178.41 (CO), 212.4 (CS D2), 212.6 (CS D1).

MS (Cl, NH₃): 480 (MH⁺), 497 (MNH₄⁺).

Dithiocarbonic acid S-(1-[[acetyl(4-bromobenzenesulfonyl)amino]methyl]-4-oxo-4-piperidin-1-yl-butyl) ester O-ethyl ester (4e)

Yellow oil, 91% (method B, petroleum ether: AcOEt 1:1).

IR (CCl₄): ν 1710 (CO), 1645 (CO), 1367, 1170 (SO₂), 1224 (COS), 1054 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.42 (t, *J* = 7 Hz, 3 H, CH₃), 1.45-1.65 (m, 6 H), 1.80-1.95 (m, 1 H, CH), 2.15-2.25 (m, 1 H, CH), 2.33 (s, 3 H, CH₃ acetyl), 2.42-2.58 (m, 2 H, CH₂CO), 3.36 (bs, 2 H, 2 NCH

piperidine), 3.52 (ddd, $J = 5.2, 13.2, 24$ Hz, 2 H, 2 NCH piperidine), 4.05-4.25 (m, 3 H, CHS and CH₂N), 4.62 (q, $J = 7$ Hz, 2 H, OCH₂), 7.66 (d, $J = 8.4$ Hz, 2 H, ArH), 7.78 (d, $J = 8.4$ Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃), 24.5 (CH₂ piperidine), 25.0 (CH₃ acetyl), 25.5, 26.4 (CH₂ piperidine), 27.0 (CH₂), 30.2 (CH₂CO), 42.8 (NCH₂ piperidine), 46.5 (NCH₂ piperidine), 49.3 (NCH₂), 50.1 (CHS), 70.4 (OCH₂), 129.0 (C Ar), 129.3 (CH Ar), 132.5 (CH Ar), 138.5 (C Ar), 169.6 (CO), 170.3 (CO acetyl), 212.8 (CS).

MS (Cl, NH₃): 566 (MH⁺).

Dithiocarbonic acid S-(1-{[acetyl(4-trifluoromethylbenzenesulfonyl)amino]methyl}-4-oxo-4-thiophen-2-yl-butyl) ester O-ethyl ester (4f)

Yellow oil, 86% (method B, petroleum ether: AcOEt 7:3).

IR (CCl₄): ν 1712 (CO), 1669 (CO), 1368, 1173 (SO₂), 1226 (COS), 1063 (CS) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.40 (t, $J = 7$ Hz, 3 H, CH₃), 1.92-2.05 (m, 1 H, CH), 2.30-2.42 (m, 1 H, CH), 2.38 (s, 3 H, CH₃ acetyl), 3.16 (t, $J = 7$ Hz, 2 H, CH₂CO), 4.18 (d, $J = 6.8$ Hz, 2 H, NCH), 4.22-4.31 (m, 1 H, CHS), 4.52-4.66 (m, 2 H, OCH₂), 7.13 (t, $J = 4.0$ Hz, 1 H, H-4 thiophene), 7.64 (d, $J = 4.8$ Hz, 1 H, CH thiophene), 7.73 (d, $J = 3.2$ Hz, 1 H, CH thiophene), 7.81 (d, $J = 8.4$ Hz, 2 H, H-*m*), 8.07 (d, $J = 8.4$ Hz, 2 H, H-*o*).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃ xanthate), 25.0 (CH₃ acetyl), 25.8 (CH₂), 36.0 (CH₂CO), 49.5 (NCH₂), 49.7 (CHS), 70.6 (OCH₂), 122.9 (q, $J = 271.4$ Hz, CF₃), 126.4 (q, $J = 3.6$ Hz, C-*m*), 128.1 (C-4 thiophene), 128.4 (C-*o*), 132.0 (CH thiophene), 133.8 (CH thiophene), 135.4 (q, $J = 33.2$ Hz, C-*p*), 142.8 (C-*ipso*), 143.8 (C-*ipso*), 170.3 (CO acetyl), 191.6 (CO), 212.4 (CS).

MS (Cl, NH₃): 554 (MH⁺), 571 (MNH₄⁺).

Dithiocarbonic acid S-(1-{[acetyl(4-cyanobenzenesulfonyl)amino]methyl}-5,5-dimethyl-4-oxohexyl) ester O-ethyl ester (4g)

Yellow oil, 93% (method B, petroleum ether: AcOEt 3:1).

IR (CCl₄): ν 1709 (CO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.12 (s, 9 H, CH₃), 1.42 (t, $J = 7.2$ Hz, 3 H, CH₃ xanthate), 1.70-1.82 (m, 1 H, CH), 2.07-2.20 (m, 1 H, CH), 2.35 (s, 3 H, CH₃ acetyl), 2.60-2.80 (m, 2 H, CH₂CO), 4.05-4.20 (m, 3 H, CH₂N and CHS), 4.61 (q, $J = 7.2$ Hz, 2 H, OCH₂), 7.83 (d, $J = 8.8$ Hz, 2 H, ArH), 8.07 (d, $J = 8.4$ Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.7 (CH₃ xanthate), 24.9 (CH₃ acetyl), 25.1 (CH₂), 26.5 (CH₃ ^tBu), 33.3 (CH₂CO), 44.1 (C ^tBu), 49.7 (CHS), 49.8 (CH₂N), 70.6 (OCH₂), 116.9 and 117.4 (C Ar and CN), 128.7 (CH Ar), 132.8 (CH Ar), 143.4 (C Ar), 170.3 (CO acetyl), 212.6 and 214.5 (CO and CS).

MS (Cl, NH₃): 485 (MH⁺), 502 (MNH₄⁺).

Dithiocarbonic acid S-(1-{[acetyl(toluenesulfonyl)amino]methyl}-4-oxo-4-phenylbutyl) ester O-ethyl ester (4h)

To a solution of olefin **1h** (1.5 g, 5.92 mmol) in 1,2-dichloroethane (9 mL) was added xanthate **2h** (2.13 g, 8.87 mmol) and the mixture was heated at reflux temperature. 5% molar DLP was then added every 1.5 h until disparition of the starting olefin (15%). The mixture was concentrated, and the residue was purified by flash column chromatography (petroleum ether: AcOEt 7:3), to afford the desired addition product as a yellow oil (1.2 g, 42%).

IR (film): ν 1706 (NCO), 1691 (CO), 1494, 1448, 1367, 1160 (SO₂), 1224 (COS), 1048 (CS).

¹H-NMR (CDCl₃, 400 MHz): δ 1.43 (t, $J = 8$ Hz, 3 H, CH₃ xanthate), 1.98-2.03 (m, 1 H, CH), 2.27-2.34 (m, 1 H, CH), 2.36 (s, 3 H, CH₃ acetyl), 2.46 (s, 3 H, CH₃ tolyl), 3.18-3.30 (m, 2 H, CH₂CO), 4.15-4.23 (m, 2 H, NCH₂), 4.24-4.33 (m, 1 H, CHS), 4.64 (q, $J = 8$ Hz, 2 H, OCH₂), 7.35-7.58 (m, 5 H, CH Ar), 7.79 (d, $J = 8$ Hz, 2 H, ArH), 7.88 (d, $J = 8$ Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.8 (CH₃ xanthate), 21.7 (CH₃ acetyl), 25.2 (CH₂), 25.7 (CH₃ tolyl), 35.7 (CH₂CO), 49.1 (NCH₂), 49.7 (CHS), 70.5 (OCH₂), 127.5-133.2 (CH Ar), 136.6 (C Ar), 136.7 (C Ar), 145.2 (C Ar), 170.6 (NCO), 198.8 (CO), 212.8 (CS).
MS (Cl, NH₃): 494 (MH⁺), 511 (MNH₄⁺).

Dithiocarbonic acid S-(1-([acetyl(4-iodobenzenesulfonyl)amino]methyl)-5,5-dimethoxy-4-oxopentyl) ester O-ethyl ester (4i)

Yellow oil, 91% (method B, petroleum ether: AcOEt 7:3).

IR (CCl₄): ν 1739 (CO), 1711 (NCO), 1367, 1170 (SO₂), 1226 (CS), 1055 (COS).

¹H-NMR (CDCl₃, 400 MHz): δ 1.43 (td, 3 H, *J* = 7.2 Hz, OCH₃). 1.81-1.88 and 2.02-2.17 (2 m, 2 H, CH₂), 2.34 and 2.35 (2 s, 3H, CH₃), 2.92 (m, 0.5 H, CH), 3.07 (m, 0.5 H, CH), 3.32, 3.36, 3.37 (s, bs, s, 6 H, 2 OCH₃), 3.71, 3.73 (2 s, 3 H, CO₂CH₃), 4.15-4.03 (m, 2.5 H, CHS, CH₂N), 4.18-4.25 (m, 0.5 H, CHS), 4.52 (t, 1 H, *J* = 7 Hz, CHOCH₃), 4.63 (dq, 2 H, *J* = 7.2 Hz, OCH₂), 7.60 and 7.61 (2 d, 2 H, *J* = 8.4 Hz, ArH), 7.89 and 7.90 (2 d, 2 H, *J* = 8.4Hz, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 13.8 (CH₃ xanthate), 25.1 (CH₃ acetyl), 28.5, 30.3 (CH₂), 46.6, 47.1, 48.1, 48.6 (CH), 48.9, 49.8 (CH₂N), 52.3, 53.4, 54.2, 54.9 (OCH₃), 70.5, 70.6 (OCH₂), 101.8 (Cl), 104.3, 105.0 (CHOCH₃), 129.1, 138.6 (CH Ar), 170.4, 172.3, 172.5 (CO), 212.4 (CS).

General procedure for the synthesis of the transfer products 5

Method A: To a solution of acetylsulfonamide **4** (1 equiv) in isopropanol (4 mL/mmol of **4**) at reflux temperature, was added DLP (15% molar) every 1.5 h, until disparition of starting material. The reaction mixture was then evaporated, and the residue was purified by flash column chromatography.

Method B: To a solution of acetylsulfonamide **4** (1 equiv) in a mixture of 1,2-dichloroethane:isopropanol 1:1 (0.02 M) at reflux temperature, was added DLP (0.5 equiv) every hour, until 4 equiv. had been added. The reaction mixture was then evaporated, and the residue purified by column chromatography.

5-Acetylamino-4-*p*-tolylpentanoic acid methyl ester (5a)

Yellow oil, 71% (method A, 1.35 equiv DLP, petroleum ether: AcOEt 3:7).

IR (film): ν 3451, 3350, 1436 (NH), 1739 (CO ester), 1685 (CO amide), 1513 (OMe) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.88 (s, 3 H, CH₃ acetyl), 2.02-2.04 (m, 1 H, CH), 2.16-2.20 (m, 1 H, CH), 2.33 (s, 3 H, CH₃ tolyl), 2.69-2.77 (m, 1 H, CHN), 3.16-3.24 (m, 1 H, CHN), 3.61 (s, 3 H, OCH₃), 3.66-3.71 (m, 1 H, CHAr), 5.44 (bs, 1 H, NH), 7.05 (d, *J* = 8.4 Hz, 2 H, ArH), 7.14 (d, *J* = 7.6 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 23.3 (CH₃ acetyl), 28.8 (CH₂), 30.0 (CH₃ tolyl), 36.2 (CH₂CO), 44.6 (CHAr), 45.2 (CH₂N), 52.0 (OCH₃), 127.7, 129.7, 129.9 (4 CH Ar), 136.8 and 138.3 (C-*ipso* and C-*p*), 170.2 (CO amide), 174.0 (CO ester).

MS (Cl, NH₃): 264 (MH⁺), 281 (MNH₄⁺).

N-(5-Oxo-6-phenyl-2-*p*-tolylhexyl)acetamide (5b)

Yellow oil, 60% (method B, petroleum ether: AcOEt 3:7).

IR (CCl₄): ν 3451 (NH), 1714 (CO ketone), 1684 (CO amide) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.70-1.80 (m, 1 H, CH), 1.87 (s, 3 H, CH₃ acetyl), 1.95-2.05 (m, 1 H, CH), 2.35 (s, 3 H, CH₃ tolyl), 2.25-2.45 (m, 2 H, CH₂CO), 2.65-2.75 (m, 1 H, CHAr), 3.18 (ddd, J = 4.7, 8.5, 13.3 Hz, 1 H, CH₂N), 3.59 (s, 2 H, CH₂Ph), 3.55-3.66 (m, 1 H, CH₂N), 5.64 (bs, 1 H, NH), 6.97 (d, J = 7.9 Hz, 2 H, ArH), 7.05-7.20 (m, 4 H, ArH), 7.20-7.40 (m, 3 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 20.9 (CH₃ tolyl), 23.0 (CH₃ acetyl), 27.1 (CH₂), 39.3 (CH₂CO), 44.2 (CHAr), 44.8 (CH₂N), 49.9 (CH₂Ph), 126.8, 127.5, 128.5, 129.2, 129.4 (CH Ar), 133.9 (C Ar), 136.4 (C Ar), 138.4 (C Ar), 170.0 (CO amide), 207.9 (CO).

MS (CI, NH₃): 324 (MH⁺), 341 (MNH₄⁺).

Elmt. Anal.: calculated for C₂₁H₂₅NO₂·1/2H₂O C, 75.87; H, 7.88; N, 4.21. Found C, 75.39; H, 7.84; N, 4.07.

***N*-[2-(4-*tert*-Butylphenyl)-5-cyclohexyl-5-oxopentyl]acetamide (5c)**

White foam, 70% (method B, AcOEt).

IR (CCl₄): ν 3453 (NH), 1708 (CO ketone), 1684 (CO amide) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.00-1.40 (m, 5 H), 1.31 (s, 9 H, CH₃ ^{*t*}Bu), 1.55-1.85 (m, 7 H), 1.88 (s, 3 H, CH₃ acetyl), 1.90-2.05 (m, 1 H, CH), 2.15-2.45 (m, 3 H, H-1 cyclohexyl and CH₂CO), 2.71 (t, J = 14.6 Hz, 1 H, CHAr), 3.21 (ddd, J = 4.8, 8.2, 13.1 Hz, 1 H, CHN), 3.61 (td, J = 6.6, 13.3 Hz, 1 H, CHN), 5.79 (bs, 1 H, NH), 7.06 (d, J = 8.2 Hz, 2 H, ArH), 7.32 (d, J = 8.3 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 23.1 (CH₃ acetyl), 25.4, 25.5, 25.6 (CH₂ cyclohexyl), 26.9 (CH₂), 28.2, 28.4 (CH₂ cyclohexyl), 31.2 (CH₃ ^{*t*}Bu), 34.3 (C ^{*t*}Bu), 38.0 (CH₂CO), 44.3 (CHAr), 44.9 (CH₂N), 50.6 (C-1 cyclohexyl), 125.4 (CH Ar), 127.2 (CH Ar), 138.6 (C Ar), 149.5 (C Ar), 170.0 (CO acetyl), 213.7 (CO).

MS (CI, NH₃): 358 (MH⁺).

***N*-[2-(2-Chlorophenyl)-3-(2-oxotetrahydrofuran-3-yl)propyl]acetamide (5d)**

2 diastereomers, D1 and D2

White foam, 72% (method B, petroleum ether: AcOEt 3:7).

IR (CCl₄) ν 1780 (CO ester), 1686 (CO amide).

¹H-NMR (CDCl₃, 400 MHz): δ 1.62-1.85 (m, 3 H, CH D1, CH D2, CH ring D2), 1.90-1.91 (2 s, 6 H, CH₃ acetyl), 1.90-2.05 (m, 1 H, CH ring D1), 2.12-2.25 (m, 2 H, CH ring D1, CHCO D2), 2.27-2.41 (m, 3 H, CH ring D1, CH D1, CH D2), 2.52-2.62 (m, 1 H, CH ring D1), 3.20-3.40 (m, 2 H, CHN D1, CHN D2), 3.57-3.84 (m, 4 H, CHN D1, CHN D2, CHAr D1, CHAr D2), 4.09 (ddd, J = 6.4, 10.1, 19.4 Hz, 2 H, CHO) 4.27 (dt, J = 1.6, 8.9 Hz, 1 H, CHO), 4.34 (dt, J = 2, 8.8 Hz, 1 H, CHO), 5.39 and 5.44 (2 bs, 2 H, NH), 7.15-7.45 (m, 8 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 23.0 (CH₃), 28.8 (CH₂ ring D2), 29.3 (CH₂ ring D1), 33.3 (CH₂ D1), 33.8 (CH₂ D2), 37.1 (CH₂ ring D1), 37.7 (CH₂ ring D2), 38.7 (CHAr D1), 39.6 (CHAr D2), 43.3 (CH₂N D2), 44.8 (CH₂N D1), 66.5 (CHO D1 and D2), 127.3, 127.7, 127.8, 128.2, 128.3, 129.7, 129.8 (CH Ar), 134.4, 134.8, 138.1, 139.3 (C-*ipso* and CCl), 170.35 and 170.42 (NCO D1 and D2), 179.3 (CO D1 and D2).

MS (CI, NH₃): 296 (MH⁺), 313 (MNH₄⁺).

***N*-[2-(4-Bromophenyl)-5-oxo-5-piperidin-1-yl-pentyl]acetamide (5e)**

White foam, 70% (method B, AcOEt: MeOH 95:5).

IR (CCl₄): ν 3465 (NH), 1683 (CO), 1641 (CO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.40-1.60 (m, 6 H), 1.72-1.82 (m, 1 H, CH), 1.87 (s, 3 H, CH₃ acetyl), 2.00-2.22 (m, 3 H, CH and CH₂CO), 2.74-2.84 (m, 1 H, CHAr), 3.14-3.28 (m, 3 H, NCH₂ piperidine and NCH), 3.40-3.58 (m, 3 H, NCH₂ piperidine and NCH), 5.99 (bs, 1 H, NH), 7.03 (d, J = 8.4 Hz, 2 H, ArH), 7.41 (d, J = 8 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 23.1 (CH₃), 24.4, 25.5, 26.3 (3 CH₂ piperidine), 28.6 (CH₂), 30.3 (CH₂CO), 42.7 (NCH₂ piperidine), 44.4 (CHAr), 44.8 (NCH₂), 46.4 (NCH₂ piperidine), 120.6 (C Ar), 129.4 (CH Ar), 131.8 (CH Ar), 141.3 (C Ar), 170.2 (CO), 170.5 (CO amide).

MS (Cl, NH₃): 380 (MH⁺).

Elmt. Anal.: calculated for C₁₈H₂₅BrN₂O₂·1/2H₂O C, 55.39; H, 6.71; N, 7.18. Found C, 54.99; H, 6.53; N, 6.91.

***N*-[5-Oxon-5-thiophen-2-yl-2-(4-trifluoromethylphenyl)pentyl]acetamide (5f)**

Yellow oil, 33% (method B, AcOEt).

IR (CCl₄): ν 3458 (NH), 1685 (CO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.90 (s, 3 H, CH₃ acetyl), 1.92-2.03 (m, 1 H, CH), 2.15-2.27 (m, 1 H, CH), 2.70-2.87 (m, 1 H, CH₂CO), 2.93-3.03 (m, 1 H, CHAr), 3.32 (ddd, J = 5.4, 8.2, 13.6 Hz, 1 H, NCH), 3.65 (qt, J = 6.7 Hz, 1 H, NCH), 5.63 (bs, 1 H, NH), 7.07 (dd, J = 4, 4.8 Hz, 1 H, H-4 thiophene), 7.31 (d, J = 8 Hz, 2 H, H-*o*), 7.54-7.64 (m, 4 H, H-*m* and 2 CH thiophene).

¹³C-NMR (CDCl₃, 100 MHz): δ 23.1 (CH₃ acetyl), 27.7 (CH₂), 36.5 (CH₂CO), 44.8 (NCH), 44.9 (CHAr), 124.1 (q, J = 270.4 Hz, CF₃), 125.8 (q, J = 3.65 Hz, C-*m*), 128.1 (C-4 thiophene), 128.2 (C-*o*), 129.4 (q, J = 32.2 Hz, C-*p*), 131.9 and 133.7 (C-3 and C-5 thiophene), 143.9 (C-*ipso*), 146.1 (C-*ipso*), 170.2 (CO acetyl), 192.4 (CO).

MS (Cl, NH₃): 370 (MH⁺), 387 (MNH₄⁺).

***N*-[2-(4-Cyanophenyl)-6,6-dimethyl-5-oxoheptyl]acetamide (5g)**

Transparent oil, 81% (method B, petroleum ether: AcOEt 3:7).

IR (CCl₄): ν 3458 (NH), 2231 (CN), 1705 (CO ketone), 1687 (CO amide) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.07 (s, 9 H, CH₃ ^tBu), 1.74-1.87 (m, 1 H, CH), 1.92 (s, 3 H, CH₃ acetyl), 1.98-2.10 (m, 1 H, CH), 2.34 (ddd, J = 5.5, 7.9, 18 Hz, 1 H, CHCO), 2.44 (td, J = 7.6, 17.8 Hz, 1 H, CHCO), 2.92 (t, J = 14.6 Hz, 1 H, CHAr), 3.31 (ddd, J = 5.6, 8.1, 13.6 Hz, 1 H, NCH), 3.60 (td, J = 6.5, 13.2 Hz, 1 H, NCH), 5.73 (bs, 1 H, NH), 7.32 (d, J = 8.4 Hz, 2 H, ArH), 7.65 (d, J = 8 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): 23.1 (CH₃ acetyl), 26.3 (CH₃ ^tBu), 26.9 (CH₂), 33.6 (CH₂CO), 44.0 (C ^tBu), 44.7 (NCH), 45.1 (CHAr), 110.8 and 118.6 (C Ar and CN), 128.6 (CH Ar), 132.5 (CH Ar), 148.0 (C Ar), 170.1 (CO acetyl), 215.3 (CO).

MS (Cl, NH₃): 301 (MH⁺), 318 (MNH₄⁺).

Elmt. Anal.: calculated for C₁₈H₂₄N₂O₂·1/2H₂O C, 69.87; H, 8.14; N, 9.05. Found C, 69.81; H, 8.13; N, 8.99.

N-(5-Oxo-5-phenyl-2-p-tolylpentyl)acetamide (5h)

Yellow oil, 53% (method A, petroleum ether: AcOEt 1:9).

IR (film): ν 3453, 1549 (NH), 1706 (CO), 1691 (NCO).

¹H-NMR (CDCl₃, 400 MHz): δ 1.90 (s, 3 H, CH₃ acetyl), 1.92-2.00 (m, 1 H, CH), 2.10-2.18 (m, 1 H, CH), 2.34 (s, 3 H, CH₃ tolyl), 2.81-2.88 (m, 2 H, CH₂CO), 2.90-2.98 (m, 1 H, NCH), 3.24-3.30 (m, 1 H, NCH), 3.68-3.76 (m, 1 H, CHAr), 5.51 (bs, 1 H, NH), 7.09 (d, J = 8 Hz, 2 H, ArH), 7.15 (d, J = 8 Hz, 2 H, ArH), 7.40-7.55 (m, 5 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 23.3 (CH₃ acetyl), 27.9 (CH₂), 30.0 (CH₃ tolyl), 36.2 (CH₂CO), 44.6 (CHAr), 45.2 (NCH₂), 127.7-133.1 (CH Ar), 136.7 (C Ar), 136.8 (C Ar), 138.8 (C Ar), 170.2 (NCO), 200.2 (CO).

MS (Cl, NH₃): 310 (MH⁺).

5-Acetylamino-2-dimethoxymethyl-4-(4-iodophenyl)pentanoic acid methyl ester (5i)

White foam, 60% (method A, AcOEt).

IR (CCl₄): ν 3453 (NH), 1739 (CO), 1687 (NCO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.84 (s, 1 H, CH₃ acetyl), 1.85 (s, 2 H, CH₃ acetyl), 1.78-2.07 (m, 2 H, CH₂), 2.61-2.84 and 2.40-2.46 (2 m, 2 H), 3.22 and 3.14 (2 s, 6 H, 2 OCH₃), 3.16-3.26 (m, 1 H, NCH), 3.28 and 3.29 (2 s, 2 H, CO₂CH₃), 3.40 (s, 1 H, CO₂CH₃), 3.54-3.66 (m, 1 H, CHAr), 3.68 (s, 2 H, CO₂CH₃), 4.39 (d, 0.33 H, J = 7.6Hz, CHOCH₃), 4.43 (d, 0.67 H, J = 7.6Hz, CHOCH₃), 5.43 (m, 1 H, NH), 6.90 (d, 2 H, J = 8.3Hz, ArH), 7.64 (d, 2 H, J = 8.4 Hz, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 23.3 (CH₃CO), 31.4, 31.7 (CH₂), 43.3, 43.9 (CH), 44.6, 45.1 (CH₂N), 46.6, 47.6 (CH), 52.6, 53.6, 54.5, 54.7 (OCH₃), 92.4, 92.6 (Cl), 104.4, 104.9 (CHOCH₃), 130.0, 130.2, 137.8, 137.9, 141.4, 140.9 (CH Ar), 170.1 (CO amide), 173.0 (CO ester).

MS (Cl, NH₃): 418 (MH⁺-MeOH), 450 (MH⁺), 467 (MNH₄⁺).

Elmt. Anal.: calculated for C₁₇H₂₄INO₅ C, 45.45; H, 5.38. Found C, 45.89; H, 5.58.

General procedure for the synthesis of imines 6

Method A: To a solution of acetamide **5** (1 equiv) in dioxane (0.4 mL/mmol of **5**) was added concentrated HCl (4 mL/mmol of **5**), and the mixture was heated at reflux temperature for 1.5 days. The reaction mixture was then evaporated, the residue dissolved in CH₂Cl₂, and washed with sat. aq. NaHCO₃ solution. The organic phase was dried and concentrated, and the imine was obtained without further purification.

Method B: To a solution of acetamide **5** (1 equiv) in dioxane (0.4 mL/mmol of **5**) was added concentrated HCl (4 mL/mmol of **5**), and the mixture was heated at reflux temperature for 1.5 days. The reaction mixture was then evaporated, and the imine was obtained as the hydrochloride, which was used in the next reaction without further purification.

5-*p*-Tolylpiperidin-2-one (7)

A solution of acetamide **5a** (0.51 g, 1.94 mmol) in concentrated HCl (9.7 mL) was heated at reflux temperature for 4 h. The reaction mixture was then concentrated, and the residue purified by flash column chromatography (AcOEt), yielding the desired piperidone as a white solid (0.295 g, 88%).

mp: 95-97 °C.

IR (film): ν 3451, 3305, 1496 (NH), 1685 (CO) cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 2.02-2.08 (m, 2 H, H-4), 2.34 (s, 3 H, CH_3), 2.49-2.56 (m, 2 H, H-3), 2.99-3.05 (m, 1 H, H-5), 3.32-3.38 (m, 1 H, H-6), 3.46-3.50 (m, 1 H, H-6), 6.33 (bs, 1 H, H-1), 7.13-7.14 (m, 4 H, ArH).

$^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz): δ 21.1 (CH_3), 27.9 (C-3), 31.4 (C-4), 39.2 (C-5), 48.8 (C-6), 126.9 (CH Ar), 129.5 (CH Ar), 136.8 (C Ar), 138.8 (C Ar), 172.2 (C-2).

MS (CI, NH_3): 190 (MH^+), 207 (MNH_4^+).

General procedure for the synthesis of piperidines 8

To a solution of imine **6** (1 equiv) in methanol (8 mL/mmol of **6**) at 0 °C was added NaBH_3CN (2.5 equiv), a few drops of glacial AcOH, and the resulting mixture was stirred at room temperature for 3 h. After that time, the reaction mixture was concentrated, CH_2Cl_2 was added, and the organic phase washed with sat. aq. NaHCO_3 solution. The organic phase was dried and concentrated, and the residue was purified by flash column chromatography.

(2*SR*,5*SR*)-2-Benzyl-5-*p*-tolylpiperidine (8a)

mp: 118-119 °C.

IR (CCl_4): ν 3100 (NH) cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.60 (ddd, J = 2.6, 12.4, 24.6 Hz, 1 H, H-4_{ax}), 1.75 (dt, J = 2.7, 13.9 Hz, 1 H, H-3_{ax}), 1.92-2.06 (m, 2 H, H-4_{eq} and H-3_{ax}), 2.29 (s, 3 H, CH_3), 2.88 (dd, J = 10.7, 15.0 Hz, 1 H, CHPh), 2.92 (t, J = 12.6 Hz, 1 H, H-6_{ax}), 3.09 (tt, J = 3.3, 12.3 Hz, 1 H, H-5_{ax}), 3.17-3.26 (m, 2 H, CHPh and H-2), 3.45 (dd, J = 1.8, 12.4 Hz, 1 H, H-6_{eq}), 7.05 (d, J = 8.3 Hz, 2 H, ArH), 7.09 (d, J = 8.2 Hz, 2 H, ArH), 7.20-7.33 (m, 5 H, ArH).

$^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz): δ 21.0 (CH_3), 28.2 (C-3), 30.0 (C-4), 39.1 (C-5), 39.7 (CH_2Ph), 50.6 (C-6), 58.7 (C-2), 126.8, 127.4, 129.0, 129.4, 129.6 (CH Ar), 134.9, 137.0, 137.3 (C Ar).

MS (CI, NH_3): 266 (MH^+).

(2*SR*,5*SR*)-5-(4-*tert*-Butylphenyl)-2-cyclohexylpiperidine (8b)

White solid, 77% (petroleum ether: AcOEt 3:7).

mp: 201-204 °C.

IR (CCl_4): ν 3100 (NH) cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.00-1.40 (m, 5 H), 1.29 (s, 9 H, CH_3), 1.55-1.75 (m, 6 H, H-4_{ax}, H-3_{ax}, H-1 cyclohexyl and 3 CH cyclohexyl), 1.78-1.85 (m, 2 H, CH_2 cyclohexyl), 2.00-2.15 (m, 2 H, H-4_{eq} and H-3_{eq}), 2.82-2.91 (m, 1 H, H-2_{ax}), 2.97 (t, J = 12.5 Hz, 1 H, H-6_{ax}), 3.07-3.21 (m, 1 H, H-5_{ax}), 3.54 (dd, J = 1.2, 10.8 Hz, 1 H, H-6_{eq}), 7.15 (d, J = 8.3 Hz, 2 H, ArH), 7.33 (d, J = 8.3 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 25.65, 25.7, 25.8, 26.0, 28.2 (CH₂), 29.1 and 30.1 (C-3 and C-4), 31.3 (CH₃ ^{*t*}Bu), 34.4 (C ^{*t*}Bu), 39.0 (C-5), 40.1 (C-1 cyclohexyl), 51.0 (C-6), 62.4 (C-2), 125.7 (CH Ar), 126.7 (CH Ar), 137.1 (C Ar), 150.4 (C Ar).
MS (CI, NH₃): 299 (MH⁺).

(2*SR*,5*SR*)-2-(thiophene-2-yl)-5-(4-trifluorophenyl)piperidine (8c)

Yellow solid, 64% (petroleum ether: AcOEt 3:7).

mp: 103-105 °C.

IR (CCl₄): ν 3073 (NH) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.78 (m, 2 H, H-4_{ax} and H-3_{ax}), 1.92 (bs, 1 H, NH), 2.07-2.18 (m, 2 H, H-4_{eq} and H-3_{eq}), 2.89 (dd, *J* = 10.6, 17 Hz, 2 H, H-2_{ax} and H-6_{ax}), 3.31 (dd, *J* = 11, 17.4 Hz, 1 H, H-5_{ax}), 4.04 (dd, *J* = 2.4, 10 Hz, 1 H, H-6_{eq}), 6.96-7.02 (dd, *J* = 0.9, 5.2 Hz, 1 H, ArH), 7.22 (dd, *J* = 0.9, 5.2 Hz, 2 H, ArH), 7.36 (d, *J* = 8.1 Hz, 2 H, ArH), 7.57 (d, *J* = 8.1 Hz, 2 H, ArH).

¹³C-NMR (CDCl₃, 100 MHz): δ 31.9 and 35.7 (C-3 and C-4), 43.1 (C-5), 54.0 (C-6), 56.8 (C-2), 123.0, 123.8, 125.4, 126.4, 127.5 (CH Ar), 148.3 (C Ar), 148.6 (C Ar).

MS (CI, NH₃): 311 (MH⁺).

Elmt. Anal.: calculated for C₁₆H₁₆F₃NS C, 61.72; H, 5.18. Found C, 61.41; H, 5.11.

(2*SR*,5*SR*)-2-Phenyl-5-*p*-tolylpiperidine (8d)

White solid, 89% (AcOEt).

mp: 119-121 °C.

IR (film): ν 3050 (NH) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.68-1.87 (m, 2 H, H-4_{ax} and H-3_{ax}), 1.96-2.15 (m, 2 H, H-4_{eq} and H-3_{eq}), 2.38 (s, 3 H, CH₃), 2.85 (tt, *J* = 3.2, 11.2 Hz, 1 H, H-5_{ax}), 2.91 (t, *J* = 11.2 Hz, 1 H, H-6_{ax}), 3.35 (d, *J* = 9.6 Hz, 1 H, H-2_{ax}), 3.73 (dd, *J* = 2.4, 10.8 Hz, 1 H, H-6_{eq}), 7.18 (d, *J* = 8 Hz, 2 H), 7.22 (d, *J* = 8 Hz, 2 H), 7.31 (d, *J* = 7.2 Hz, 1 H), 7.38 (t, *J* = 7.6 Hz, 2 H), 7.45 (d, *J* = 7.2 Hz, 2 H).

¹³C-NMR (CDCl₃, 100 MHz): δ 21.0 (CH₃), 32.5 and 35.0 (C-3 and C-4), 42.7 (C-5), 54.7 (C-6), 61.8 (C-2), 126.6, 127.0, 127.1, 128.4, 129.1 (CH Ar), 135.8, 141.5, 145.0 (C Ar).

MS (CI, NH₃): 252 (MH⁺).

1-Acetyl-5-(4-iodo-phenyl)-1,4,5,6-tetrahydropyridine-3-carboxylic acid ethyl ester (8e)

To a solution of acetamide **5i** (96 mg, 0.21 mmol) in CH₂Cl₂ (2 mL) was added *p*TsOH·H₂O (45 mg, 0.24 mmol) and heated at reflux temperature for 4.5 h. After that time, the reaction mixture was cooled and washed with sat. aq. NaHCO₃ solution and sat. aq. NaCl solution. The organic phase was dried and concentrated, the residue was purified by flash column chromatography (petroleum ether: AcOEt 1:1), and the desired piperidine was obtained as a white solid (72 mg, 89%).

mp: 111-113 °C.

IR (CCl₄): ν 1694 (CO), 1627 (CO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 2.32 (s, 3 H, CH₃ acetyl), 2.38 (ddd, *J* = 1.6, 14.8, 17.2 Hz, 1 H, H-4_{ax}), 2.74 (dd, *J* = 3.6, 17.2 Hz, 1 H, H-4_{eq}), 2.87 (tt, *J* = 4, 10.8 Hz, 1 H, H-5), 3.10 (t, *J* = 12 Hz, 1 H, H-6_{ax}), 3.75 (s, 3 H, OCH₃), 4.39 (d, *J* = 12.4 Hz, 1 H, H-6_{eq}), 6.96 (d, *J* = 8.4 Hz, 2 H, ArH), 7.64 (d, *J* = 8.4 Hz, 2 H, ArH), 7.88 (bs, 1 H, H-2).

¹³C-NMR (CDCl₃, 100 MHz): δ 21.6 (CH₃ acetyl), 28.5 (C-4), 36.7 (C-5), 45.3 (C-6), 51.7 (OCH₃), 92.4 (C), 109.1 (C), 129.0 (CH Ar), 135.4 (C-2), 137.7 (CH Ar), 141.1 (C-*ipso*), 167.2 (CO), 169.1 (CO).

2-Phenyl-5-*p*-tolylpyridine (9)

To a solution of imine **6h** (100 mg, 0.60 mmol) in xylenes (15 mL) was added 10% Pd/C (100 mg) and heated at reflux temperature for 2.5 h. The reaction mixture was then filtered over Celite™ and concentrated. The residue was recrystallized from EtOH, and the desired pyridine was obtained as a white solid (83%).

mp: 141-143 °C.

IR (film): ν 2925, 2855, 1594, 1472, 1446 cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 2.43 (s, 3 H, CH₃), 7.30-7.55 (m, 7 H, CH Ar), 7.80 (d, *J* = 8.4 Hz, 1 H, H-3), 7.94 (dd, *J* = 1.4, 8.4 Hz, 1 H, H-4), 8.03 (dd, *J* = 2.2, 8.2 Hz, 2 H, CH Ar), 8.92 (d, *J* = 1.4 Hz, 1 H, H-6).

¹³C-NMR (CDCl₃, 100 MHz): δ 21.2 (CH₃), 120.3 (C-3), 126.77 (CH Ar), 126.80 (CH Ar), 128.8 (CH Ar), 128.9 (CH Ar), 129.8 (CH Ar), 134.7 (C-5), 134.81 (C Ar), 134.84 (C-4), 138.0 (C Ar), 139.0 (C Ar), 147.9 (C-6), 155.8 (C-2).

MS (CI, NH₃): 246 (MH⁺).

2-Ethanesulfonyl-3-(4-trifluoromethylphenyl)pent-4-enoic acid ethyl ester (10)

For preparation details and NMR description, see: Ouvry, G. Ph. D. Thesis, École Polytechnique, Palaiseau, 2002.

(*E*)-7-(4-Methoxyphenyl)-7-oxo-4-(4-trifluoromethylphenyl)hept-2-enoic acid ethyl ester (11)

For preparation details and NMR description, see: Ouvry, G. Ph. D. Thesis, École Polytechnique, Palaiseau, 2002.

General procedure for the synthesis of amines 12

To a solution of ketone **10** (1 equiv) in absolute EtOH (2.5 mL/mmol) were added 4Å molecular sieves, the amine (as its acetate salt, 10 equiv), NaBH₃CN (1.1 equiv) and the resulting mixture was heated at reflux temperature for 7 h. The reaction mixture was then concentrated, dissolved in CH₂Cl₂ and washed with sat. aq. NaHCO₃ solution. The organic layer was dried and evaporated, and the residue purified by flash column chromatography.

(2*SR*,3*SR*,6*SR*)-[6-(4-Methoxyphenyl)-3-(4-trifluoromethylphenyl)piperidin-2-yl]acetic acid ethyl ester (12a)

Yellow solid, 60% (petroleum ether: AcOEt 8:2).

mp: 80-83 °C.

IR (CCl₄): ν 1731 (CO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.16 (t, J = 7.2 Hz, 3 H, CH₃), 1.55-1.75 (m, 2 H, H-5_{ax} and H-4_{ax}), 1.91 (dd, J = 2.2, 12.2 Hz, 1 H, H-5_{eq}), 2.03 (dd, J = 2.8, 12.8 Hz, 1 H, H-4_{eq}), 2.23 (dd, J = 3.2, 16.8 Hz, 1 H, CHCO), 2.30 (dd, J = 8.8, 16.8 Hz, 1 H, CHCO), 2.62 (td, J = 3.6, 11.6 Hz, 1 H, H-3_{ax}), 3.39 (td, J = 2.7, 10 Hz, 1 H, H-2_{ax}), 3.59 (bs, 1 H, NH), 3.80 (s, 3 H, OCH₃), 3.86 (dd, J = 1.6, 10.8 Hz, 1 H, H-6_{ax}), 3.95-4.05 (m, 2 H, OCH₂), 6.89 (d, J = 8.4 Hz, 2 H, ArH methoxyphenyl), 7.35 (d, J = 8.4 Hz, 4 H, 2 ArH methoxyphenyl and 2 ArH trifluoromethylphenyl), 7.58 (d, J = 8 Hz, 2 H, ArH trifluoromethylphenyl).

¹³C-NMR (CDCl₃, 100 MHz): δ 14.0 (CH₃), 34.0 (C-4), 34.2 (C-5), 38.6 (CH₂CO), 48.3 (C-3), 55.2 (OCH₃), 58.4 (C-2), 60.4 (OCH₂), 61.0 (C-6), 113.8 (CH Ar methoxyphenyl), 124.2 (q, J = 270 Hz, CF₃), 125.5 (q, J = 2.7 Hz, C-*m* trifluoromethylphenyl), 127.8 (CH Ar methoxyphenyl), 128.2 (C-*o* trifluoromethylphenyl), 129.0 (q, J = 32.1 Hz, C-*p* trifluoromethylphenyl), 136.6 (C Ar), 147.6 (C Ar), 158.8 (C Ar), 172.3 (CO).

MS (Cl, NH₃): 422 (MH⁺).

Elmt. Anal.: calculated for C₂₃H₂₆F₃NO₃ C, 65.55; H, 6.22. Found C, 65.39; H, 6.52.

(2*SR*,3*SR*,6*RS*)-[6-(4-Methoxyphenyl)-3-(4-trifluoromethylphenyl)piperidin-2-yl]acetic acid ethyl ester (12b)

Yellow solid, 11% (petroleum ether: AcOEt 1:1).

mp: 85-86 °C.

IR (CCl₄): ν 1729 (CO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 1.24 (t, J = 7.2 Hz, 3 H, CH₃), 1.67-1.76 (m, 1 H, H-4_{ax}), 1.81 (ddd, J = 4, 9, 18 Hz, 1 H, H-4_{eq}), 2.00 (tt, J = 4.2, 12.5 Hz, 1 H, H-5_{ax}), 2.26 (dd, J = 3.8, 15.8 Hz, 1 H, CHCO), 2.32 (dd, J = 9, 16.2 Hz, 1 H, CHCO), 2.20-2.35 (m, 1 H, H-5_{eq}), 2.50 (bs, 1 H, NH), 2.60 (td, J = 4, 9.6 Hz, 1 H, H-3_{ax}), 3.47 (t, J = 4, 8.8 Hz, 1 H, H-2_{ax}), 4.05-4.15 (bs, 1 H, H-6_{eq}), 6.95 (d, J = 8.4 Hz, 2 H, ArH methoxyphenyl), 7.28 (d, J = 8 Hz, 2 H, H-*o* trifluoromethylphenyl), 7.46 (d, J = 8.4 Hz, 2 H, ArH methoxyphenyl), 7.50 (d, J = 8 Hz, 2 H, H-*m* trifluoromethylphenyl).

¹³C-NMR (CDCl₃, 100 MHz): δ 14.2 (CH₃), 28.4 (C-4), 29.2 (C-5), 39.0 (CH₂CO), 48.2 (C-3), 51.9 (C-2), 53.3 (C-6), 55.3 (OCH₃), 60.5 (OCH₂), 114.0 (CH Ar methoxyphenyl), 124.2 (q, J = 270 Hz, CF₃), 125.4 (q, J = 3.7 Hz, C-*m* trifluorophenyl), 127.9 (CH Ar methoxyphenyl), 128.2 (C-*o* trifluoromethylphenyl), 128.8 (q, J = 32.4 Hz, C-*p* trifluorophenyl), 134.5 (C Ar), 148.2 (C Ar), 158.2 (C Ar), 172.4 (CO).

MS (Cl, NH₃): 423 (MH⁺).

Elmt. Anal.: calculated for C₂₃H₂₆F₃NO₃·3/4H₂O: C, 63.51; H, 6.37, N, 3.22. Found C, 63.41; H, 6.18; N, 2.81.

(2*SR*,3*SR*,6*SR*)-[1-Cyclopropyl-6-(4-methoxyphenyl)-3-(4-trifluoromethylphenyl)piperidin-2-yl]acetic acid ethyl ester (13)

Yellow solid, 86% (petroleum ether: AcOEt 9:1).

mp: 78-80 °C.

IR (CCl₄): ν 1737 (CO) cm⁻¹.

¹H-NMR (CDCl₃, 400 MHz): δ 0.02 (q, J = 3.2 Hz, 2 H, CH₂ cyclopropyl), 0.13 (qt, J = 3.4 Hz, 2 H, CH₂ cyclopropyl), 1.11 (t, J = 7.2 Hz, 3 H, CH₃), 1.82 (ht, J = 3.6 Hz, 1 H, CH cyclopropyl), 1.87 (ddd, J = 2.8, 5.6, 12.4 Hz, 1 H, H-5_{ax}), 1.94-2.06 (m, 2 H, H-4), 2.02 (dd, J = 5.2, 14.8 Hz, 1 H, CHCO), 2.07-2.18 (m, 1 H, H-5_{eq}), 2.77 (dd, J = 8.4, 14.8 Hz, CHCO), 3.38-3.44 (m, 1 H, H-3_{ax}), 3.81 (s, 3 H, OCH₃), 3.85 (q, J = 6.8 Hz, 2 H, OCH₂), 3.86-3.92 (m, 1 H, H-2_{ax}), 3.93-3.97 (m, 1 H, H-6_{ax}), 6.84 (d, J = 8.4

Hz, 2 H, ArH methoxyphenyl), 7.24 (d, $J = 8.4$ Hz, 2 H, ArH methoxyphenyl), 7.38 (d, $J = 8$ Hz, 2 H, ArH trifluoromethylphenyl), 7.58 (d, $J = 8.4$ Hz, 2 H, ArH trifluoromethylphenyl).

$^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz): δ 7.4 and 8.9 (CH_2 cyclopropyl), 14.0 (CH_3), 23.9 (C-4), 29.7 (CH_2CO), 29.9 (C-5), 32.7 (CH isopropyl), 42.5 (C-3), 55.2 (OCH_3), 59.4 (C-2), 60.2 (OCH_2), 62.4 (C-6), 113.1 (CH Ar methoxyphenyl), 124.2 (q, $J = 270$ Hz, CF_3), 125.2 (q, $J = 3.7$ Hz, C-*m* trifluoromethylphenyl), 128.0 (C-*o* trifluoromethylphenyl), 128.6 (q, $J = 32.2$ Hz, C-*p* trifluoromethylphenyl), 129.8 (CH Ar methoxyphenyl), 134.8 (C Ar), 146.8 (C Ar), 158.6 (C Ar), 172.6 (CO).

MS (Cl, NH_3): 462 (MH^+).

Elmt. Anal.: calculated for $\text{C}_{26}\text{H}_{30}\text{F}_3\text{NO}_3$ C, 67.66; H, 6.55; N, 3.03. Found C, 67.59; H, 6.59; N, 2.88.

(1*SR*,7*SR*,10*SR*)-7-(4-Methoxyphenyl)-10-(4-trifluoromethylphenyl)octahydropyrido[1,2-*d*][1,4]diazepin-2-one (14a)

White solid, 56% (AcOEt).

mp: 220-222 °C.

IR (CCl_4): ν 1679 cm^{-1} .

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.60 (ddd, $J = 3.3, 12.3, 24.3$ Hz, 1 H, H-9_{ax}), 1.72 (ddd, $J = 2.8, 13, 24.2$ Hz, 1 H, H-8_{ax}), 1.79 (tdd, $J = 3.2, 6.4, 13.2$ Hz, 1 H, H-8_{eq}), 1.87 (ddd, $J = 3.2, 6, 12.8$ Hz, 1 H, H-9_{eq}), 2.07 (dd, $J = 9.6, 13.6$ Hz, 1 H, H-5), 2.30 (d, $J = 14.4$ Hz, 1 H, H-1), 2.62 (dd, $J = 8.4, 14$ Hz, 1 H, H-1), 2.65-2.78 (m, 2 H, H-10 and H-11), 2.93 (qt, $J = 7.1$ Hz, 1 H, H-4), 3.03 (dd, $J = 6.2, 13.4$ Hz, 1 H, H-5), 3.23 (dd, $J = 3, 11$ Hz, 1 H, H-7), 3.27 (dd, $J = 4, 9.6$ Hz, 1 H, H-4), 6.80 (bs, 1 H, H-3), 6.87 (d, $J = 8.8$ Hz, 2 H, CH Ar methoxyphenyl), 7.26 (d, $J = 8.4$ Hz, 2 H, H-*o* trifluoromethylphenyl), 7.32 (d, $J = 8$ Hz, 2 H, CH Ar methoxyphenyl), 7.57 (d, $J = 8.4$ Hz, 2 H, H-*m* trifluoromethylphenyl).

$^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz): δ 34.0 and 35.6 (C-8 and C-9), 42.3 (C-1 and C-5), 49.5 (C-10), 55.2 (OCH_3), 56.2 (C-4), 63.4 (C-7), 68.5 (C-11), 114.0 (CH Ar methoxyphenyl), 124.1 (q, $J = 271$ Hz, CF_3), 125.8 (C-*m* trifluoromethylphenyl), 128.1 (CH Ar methoxyphenyl and C-*o* trifluoromethylphenyl), 128.8 (q, $J = 32$ Hz, C-*p* trifluoromethylphenyl), 136.8 (C Ar), 148.2 (C Ar), 158.6 (C Ar), 176.6 (C-2).

MS (Cl, NH_3): 419 (MH^+).

(1*SR*,7*RS*,10*SR*)-7-(4-Methoxyphenyl)-10-(4-trifluoromethylphenyl)octahydropyrido[1,2-*d*][1,4]diazepin-2-one (14b)

White foam, 16% (AcOEt: MeOH 99:1).

IR (CCl_4): 3429 (NH), 1676 (CO).

$^1\text{H-NMR}$ (CDCl_3 , 400 MHz): δ 1.64-1.77 (m, 2 H, H-10), 1.78-1.87 (m, 1 H, H-9_{ax}), 1.98 (dt, $J = 4.1, 10.6$ Hz, 1 H, H-9_{eq}), 2.25 (d, $J = 14.8$ Hz, H-6), 2.72-2.84 (m, 1 H, H-3), 2.84 (dd, $J = 9.6, 14.8$ Hz, H-2), 2.97 (q, $J = 4.8$ Hz, H-8), 3.10 (dd, $J = 5.6, 14.8$ Hz, 1 H, H-2), 3.38 (dd, $J = 9.8, 13$ Hz, 1 H, H-6), 3.59 (ddd, $J = 4.8, 9.6, 14.8$ Hz, 1 H, H-3), 3.77-3.83 (m, 1 H, H-7), 3.81 (s, 3 H, OCH_3), 4.08 (dd, $J = 4, 8$ Hz, 1 H, H-11_{eq}), 6.28 (bs, 1 H, NH), 6.88 (d, $J = 8.4$ Hz, 2 H, ArH methoxyphenyl), 7.27 (d, $J = 8.4$ Hz, 2 H, ArH methoxyphenyl), 7.57 (dd, $J = 8.4$ Hz, 2 H, ArH trifluoromethylphenyl), 7.61 (d, $J = 8.4$ Hz, 2 H, ArH trifluoromethylphenyl).

$^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz): δ 27.4 and 28.6 (C-9 and C-10), 38.7 (C-6), 39.4 (C-2), 44.3 (C-8), 52.9 (C-3), 55.2 (OCH_3), 56.0 and 59.1 (C-11 and C-7), 114.0 (CH Ar methoxyphenyl), 125.1 (q, $J = 3$ Hz, C-*m* trifluoromethylphenyl), 128.2 (q, $J = 32.3$ Hz, C-*p* trifluoromethylphenyl), 128.5 (CH Ar methoxyphenyl), 128.6 (C-*o* trifluoromethylphenyl), 134.6 (C Ar), 148.6 (C Ar), 158.6 (C Ar), 177.0 (C-5).

MS (Cl, NH_3): 419 (MH^+).