Synthesis of [60]Fullerene Adducts Bearing Carbazole Moieties by Bingel Reaction and Their Properties

Yosuke Nakamura, Masato Suzuki, Yumi Imai, and Jun Nishimura*

Department of Nano-Material Systems, Graduate School of Engineering,, Gunma University, Tenjin-cho, Kiryu, Gunma 376-8515, Japan

nisimura@chem.gunma-u.ac.jp

Supporting information

Contents

- 1) ^1H NMR spectrum of **4** in CDCl_3 .
- 2) ^{13}C NMR spectrum of **4** in CDCl_3 .
- 3) UV-Vis spectrum of **4** in toluene.
- 4) ^1H NMR spectrum of **5** in CDCl_3 .
- 5) ^{13}C NMR spectrum of **5** in CDCl_3 .
- 6) UV-Vis spectrum of **5** in toluene.
- 7) ^1H NMR spectrum of **7** in CDCl_3 .
- 8) ^{13}C NMR spectrum of **7** in CDCl_3 .
- 9) UV-Vis spectrum of **7** in toluene.

1) ^1H NMR spectrum of **4** in CDCl_3 .

2) ^{13}C NMR spectrum of **4** in CDCl_3 .

3) UV-Vis spectrum of **4** in toluene.

4) ^1H NMR spectrum of **5** in CDCl_3 .

5) ^{13}C NMR spectrum of **5** in CDCl_3 .

6) UV-Vis spectrum of **5** in toluene.

7) ^1H NMR spectrum of 7 in CDCl_3 .

8) ^{13}C NMR spectrum of 7 in CDCl_3 .

9) UV-Vis spectrum of 7 in toluene.