Synthesis of Biaryls via Decarboxylative Pd-Catalyzed Cross-Coupling Reaction

Jean-Michel Becht,^{†*} Cédric Catala,[‡] Claude Le Drian[†] and Alain Wagner^{†*}

[†] Université de Haute-Alsace, Ecole Nationale Supérieure de Chimie de Mulhouse,
Laboratoire de Chimie Organique et Bioorganique, UMR-CNRS 7015,
3 rue Alfred Werner, 68093 Mulhouse Cedex, France

[‡] Novalyst Discovery, Bioparc, Boulevard Sébastien Brant BP 30170, 67405 Illkirch Cedex, France

SUPPORTING INFORMATION

Table of contents

General Remarks	S2
General Procedure for the Synthesis and Characterizations of Biaryls 3a-t	S2-S5
NMR Spectra of Biaryls 3a-t	S6-S33

General Remarks.

The reagents were obtained from commercial sources and were used without further purifications. DMSO was dried on molecular sieves 4 Å. The syntheses of biaryls **3a-r** were carried out in anhydrous glassware under an atmosphere of argon. Purifications of compounds **3a-r** were performed by flash-chromatography on silica gel (40-63 µm, Merck). ¹H and ¹³C NMR spectra were recorded using a 400 MHz instrument in CDCl₃. Chemical shifts are reported in parts per million (δ) downfield from TMS. Spin multiplicities are indicated by the following symbol: s (singlet), d (doublet), t (triplet), q (quadruplet), hept (heptuplet) and m (multiplet). IR spectra were recorded on a Nicolet 205 FT-IR spectrometer. 2,6-Diisopropoxybenzoic acid was prepared according to a previous literature report.¹ Compounds **3a**, **3b**, **3f**, **3g**, **3i**, **3j**, **3k**, **3m**, **3n**, **3o**, **3p**, **3s** and **3t** were previously described in the literature.²

General Procedure for the Syntheses of Biaryls **3a-t**.

DMSO (6 mL) was added to a mixture of the aryl iodide (0.50 mmol, 1.0 equiv), the arene carboxylic acid (0.65 mmol, 1.3 equiv), Ag₂CO₃ (1.50 mmol, 414 mg, 3.0 equiv), AsPh₃ (0.30 mmol, 91.9 mg, 0.6 equiv) and PdCl₂ (0.15 mmol, 26.6 mg, 0.3 equiv). The reaction mixture was degassed twice with argon and directly heated at 150 °C for 6 h. After cooling to rt, the reaction mixture was filtered with Celite and AcOEt (100 mL) was added to the filtrate. The organic phase was washed with saturated NH₄Cl (70 mL), dried with MgSO₄, filtered and concentrated under vacuum. The residue was purified by flash-chromatography on silica gel to afford pure biaryls after drying under vacuum (0.1 mbar).

2,4',6-Trimethoxybiphenyl (3a): Elution with AcOEt / Cyclohexane 4:96 afforded 110 mg (90% yield) of a white solid; mp 114-116 °C (lit. mp 116-117 °C).^{2a} ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.77 (s, 6H), 3.87 (s, 3H), 6.68 (d, ³J(H,H) = 8.3 Hz, 2H), 6.98 (d, ³J(H,H) = 8.8 Hz, 2H), 7.31 (m, 3H).

2,4,4',6-Tetramethoxybiphenyl (3b): Elution with AcOEt / Cyclohexane 10:90 afforded 103 mg (75% yield) of a white solid; mp 101-103 °C (lit. mp 102-103 °C).^{2b} ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.74 (s, 6H), 3.84 (s, 3H), 3.87 (s, 3H), 6.24 (s, 2H), 6.94 (d, ³J(H,H) = 8.8 Hz, 2H), 7.27 (d, ³J(H,H) = 8.8 Hz, 2H).

3-Bromo-2,4',6-trimethoxybiphenyl (3c): Elution with AcOEt / Cyclohexane 5:95 afforded 105 mg (65% yield) of a white solid; mp 78-80 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.40 (s, 3H), 3.74 (s, 3H), 3.87 (s, 3H), 6.67 (d, ³J(H,H) = 9.1 Hz, 1H), 6.97 (d, ³J(H,H) = 8.9 Hz, 2H), 7.34 (d, ³J(H,H) = 8.9 Hz, 2H), 7.47 (d, ³J(H,H) = 9.1 Hz, 1H). ¹³C NMR (80 MHz, CDCl₃) δ (ppm): 55.2, 56.1, 60.2, 108.3, 108.7, 113.3, 114.1, 125.3, 127.7, 131.5, 131.6, 155.3, 157.3, 158.7. IR (CHCl₃) ν

(cm⁻¹): 3002, 2999, 2935, 1610, 1579, 1516, 1463, 1396, 1286, 1244, 1221, 1088, 1043, 812. Anal. Calcd for C₁₅H₁₅BrO₃: C 55.75, H 4.68, Br 24.72, found: C 55.86, H 4.64, Br 24.79.

2,6-Diisopropoxy-4'-methoxybiphenyl (3d): Elution with AcOEt / Cyclohexane 2:98 afforded 90 mg (62% yield) of a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 1.16 (d, ³J(H,H) = 6.1 Hz, 12H), 3.86 (s, 3H), 4.29 (hept, ³J(H,H) = 6.1 Hz, 2H), 6.63 (d, ³J(H,H) = 8.1 Hz, 2H), 6.90 (d, ³J(H,H) = 8.8 Hz, 2H), 7.16 (t, ³J(H,H) = 8.1 Hz, 1H), 7.29 (d, ³J(H,H) = 8.8 Hz, 2H). ¹³C NMR (80 MHz, CDCl₃) δ (ppm): 22.1, 55.1, 71.2, 108.7, 112.5, 122.9, 126.8, 127.7, 132.3, 156.4, 157.8. IR (CHCl₃) ν (cm⁻¹): 3006, 2976, 2936, 1590, 1517, 1459, 1383, 1243, 1176, 1115, 1061, 830. HRMS: m/z calcd for C₁₉H₂₄O₃ (M⁺): 300.1725, found: 300.1719.

4,4',5'-Trimethoxy-2-nitrobiphenyl (3e): Elution with AcOEt / Cyclohexane 15:85 afforded 114 mg (79% yield) of a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.86 (s, 3H), 3.96 (s, 3H), 3.98 (s, 3H), 6.78 (s, 1H), 6.95 (d, ³J(H,H) = 8.8 Hz, 2H), 7.23 (d, ³J(H,H) = 8.8 Hz, 2H), 7.52 (s, 1H). ¹³C NMR (80 MHz, CDCl₃) δ (ppm): 55.3, 56.3, 56.4, 107.8, 113.6, 114.0, 129.2, 130.4, 131.0, 141.1, 147.8, 152.1, 159.4. IR (CHCl₃) ν (cm⁻¹): 3002, 2937, 2922, 1610, 1582, 1504, 1460, 1439, 1337, 1284, 1237, 1216, 1180, 1022. HRMS: m/z calcd for C₁₅H₁₅NO₅ (M⁺): 289.0950, found: 289.0944.

4'-Methoxy-2-nitrobiphenyl (3f): Elution with AcOEt / Cyclohexane 2:98 afforded 72 mg (63% yield) of a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.86 (s, 3H), 6.96 (d, ³J(H,H) = 8.6 Hz, 2H), 7.26 (m, 2H), 7.45 (m, 2H), 7.60 (t, ³J(H,H) = 8.2 Hz, 1H), 7.81 (d, ³J(H,H) = 8.1 Hz, 1H).

2,3,4,5,6-Pentafluoro-4'-methoxybiphenyl (3g): Elution with AcOEt / Cyclohexane 4:96 afforded 126 mg (92% yield) of a white solid; mp 116-118 °C. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.88 (s, 3H), 7.02 (d, ³J(H,H) = 8.8 Hz, 2H), 7.37 (d, ³J(H,H) = 8.8 Hz, 2H).

2-Chloro-6-fluoro-4'-methoxybiphenyl (3h): Elution with AcOEt / Cyclohexane 2:98 afforded 97 mg (82% yield) of a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.87 (s, 3H), 7.00 (d, ³J(H,H) = 8.8 Hz, 2H), 7.07 (t, ³J(H,H) = 9.1 Hz, 1H), 7.35 (m, 4H). ¹³C NMR (80 MHz, CDCl₃) δ (ppm): 55.2, 113.6, 114.2 (d, ²J(C,F) = 19.1 Hz), 124.7, 125.3 (d, ⁴J(C,F) = 2.8 Hz), 128.7 (d, ²J(C,F) = 7.9 Hz), 131.4, 134.6 (d, ³J(C,F) = 2.8 Hz), 160.5 (d, ¹J(C,F) = 196.6 Hz), 159.4. IR (CHCl₃) ν (cm⁻¹): 3003, 2936, 2914, 1612, 1569, 1519, 1446, 1294, 1249, 1179, 1037, 891, 830, 782. HRMS: m/z calcd for C₁₃H₁₀ClFO (M⁺): 236.0404, found: 236.0401.

2,3',6-Trimethoxybiphenyl (3i): Elution with AcOEt / Cyclohexane 4:96 afforded 109 mg (89% yield) of a yellowish oil. ¹H NMR (400 MHz, CDCl₃) δ (ppm): 3.76 (s, 6H), 3.84 (s, 3H), 6.68 (d, ³J(H,H) = 8.3 Hz, 2H), 6.94 (m, 3H), 7.32 (m, 2H).

2,2',6-Trimethoxybiphenyl (3j): Elution with AcOEt / Cyclohexane 10:90 afforded 76 mg (62% yield) of a white solid; mp 138-140 °C (lit. mp 141 °C).^{2c} ¹H NMR (400 MHz, CDCl₃) δ (ppm):

3.74 (s, 6H), 3.77 (s, 3H), 6.67 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.02 (m, 2H), 7.19 (d, $^3J(\text{H,H}) = 7.6$ Hz, 1H), 7.30 (m, 2H).

2,6-Dimethoxy-4'-methylbiphenyl (3k): Elution with AcOEt / Cyclohexane 2:98 afforded 87 mg (76% yield) of a white solid; mp 83-85 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 2.42 (s, 3H), 3.76 (s, 6H), 6.67 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.30 (m, 5H).

4'-Chloro-2,6-dimethoxybiphenyl (3l): Elution with AcOEt / Cyclohexane 4:96 afforded 97 mg (78% yield) of a white solid; mp 118-120 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 3.75 (s, 6H), 6.65 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.29 (m, 3H), 7.37 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H). ^{13}C NMR (80 MHz, CDCl_3) δ (ppm): 55.8, 104.1, 118.2, 127.9, 129.0, 132.3, 132.4, 132.5, 157.5. IR (CHCl_3) ν (cm^{-1}): 3060, 2941, 2908, 1588, 1564, 1472, 1432, 1396, 1268, 1247, 1117, 1102, 730. Anal. Calcd for $\text{C}_{14}\text{H}_{13}\text{ClO}_2$: C 67.61, H 5.27, Cl 14.26, found: C 67.68, H 5.21, Cl 14.29.

2'-Bromo-2,6-dimethoxybiphenyl (3m): Elution with AcOEt / Cyclohexane 3:97 afforded 85 mg (58% yield) of a white solid; mp 139-141 °C (lit. mp 141-142 °C).^{2a} ^1H NMR (400 MHz, CDCl_3) δ (ppm): 3.77 (s, 6H), 6.68 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.24 (m, 2H), 7.37 (m, 2H), 7.68 (d, $^3J(\text{H,H}) = 8.1$ Hz, 1H).

2,6-Dimethoxybiphenyl (3n): Elution with AcOEt / Cyclohexane 2:98 afforded 89 mg (84% yield) of a white solid; mp 84-86 °C (lit. mp 83-85 °C).^{2e} ^1H NMR (400 MHz, CDCl_3) δ (ppm): 3.77 (s, 6H), 6.70 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.36 (m, 6H).

2,6-Dimethoxy-3'-(trifluoromethyl)biphenyl (3o): Elution with AcOEt / Cyclohexane 5:95 afforded 109 mg (77% yield) of a white solid; mp 77-79 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 3.76 (s, 6H), 6.67 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.32 (t, $^3J(\text{H,H}) = 8.3$ Hz, 1H), 7.58 (m, 4H).

2,6-Dimethoxy-4'-acetylbiphenyl (3p): Elution with AcOEt / Cyclohexane 15:85 afforded 89 mg (70% yield) of a white solid; mp 137-139 °C. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 2.64 (s, 3H), 3.75 (s, 6H), 6.67 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.32 (t, $^3J(\text{H,H}) = 8.3$ Hz, 1H), 7.47 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 8.01 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H).

2,6-Dimethoxy-2'-nitrobiphenyl (3q): Elution with AcOEt / Cyclohexane 10:90 afforded 92 mg (71% yield) of a yellow oil. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 3.72 (s, 6H), 6.65 (d, $^3J(\text{H,H}) = 8.6$ Hz, 2H), 7.33 (t, $^3J(\text{H,H}) = 8.6$ Hz, 1H), 7.48 (m, 2H), 7.64 (t, $^3J(\text{H,H}) = 8.1$ Hz, 1H), 7.99 (d, $^3J(\text{H,H}) = 8.1$ Hz, 1H). ^{13}C NMR (80 MHz, CDCl_3) δ (ppm): 55.8, 104.2, 115.2, 123.9, 127.8, 129.2, 129.7, 132.1, 133.7, 149.8, 156.9. IR (CHCl_3) ν (cm^{-1}): 3002, 2928, 2362, 1612, 1591, 1525, 1474, 1432, 1355, 1253, 1110, 782. HRMS: m/z calcd for $\text{C}_{14}\text{H}_{13}\text{NO}_4$ (M^+): 259.0845, found: 259.0838.

Ethyl 2',6'-Dimethoxybiphenyl-2-carboxylate (3r): Elution with AcOEt / Cyclohexane 5:95 afforded 84 mg (59% yield) of a yellowish oil. ^1H NMR (400 MHz, CDCl_3) δ (ppm): 1.01 (t,

$^3J(\text{H,H}) = 7.1$ Hz, 3H), 3.70 (s, 6H), 4.07 (q, $^3J(\text{H,H}) = 7.1$ Hz, 2H), 6.63 (d, $^3J(\text{H,H}) = 8.3$ Hz, 2H), 7.31 (m, 2H), 7.40 (t, $^3J(\text{H,H}) = 8.3$ Hz, 1H), 7.54 (d, $^3J(\text{H,H}) = 7.8$ Hz, 1H), 7.96 (d, $^3J(\text{H,H}) = 7.8$ Hz, 1H). ^{13}C NMR (80 MHz, CDCl_3) δ (ppm): 13.7, 55.8, 60.3, 104.0, 119.3, 126.9, 128.6, 129.7, 131.1, 132.0, 132.3, 134.8, 157.1, 167.9. IR (CHCl_3) ν (cm^{-1}): 3061, 2958, 1724, 1591, 1471, 1432, 1365, 1289, 1250, 1110, 1045, 785. HRMS: m/z calcd for $\text{C}_{17}\text{H}_{18}\text{O}_4$ (M^+): 286.1205, found: 286.1200.

2-(4-Methoxyphenyl)-3-methylbenzofurane (3s): Elution with AcOEt / Cyclohexane 1:99 afforded 77 mg (65% yield) of a yellow solid; mp 86-88 °C (lit. mp 87 °C).^{2g} ^1H NMR (400 MHz, CDCl_3) δ (ppm): 2.40 (s, 3H), 3.82 (s, 3H), 6.96 (d, $^3J(\text{H,H}) = 8.8$ Hz, 2H), 7.22 (m, 2H), 7.44 (m, 2H), 7.69 (d, $^3J(\text{H,H}) = 8.8$ Hz, 2H).

1-(2,6-Dimethoxyphenyl)naphthalene (3t): Elution with AcOEt / Cyclohexane 3:97 afforded 96 mg (73% yield) of a yellowish solid; mp 144-146 °C (lit. mp 147 °C).^{2h} ^1H NMR (400 MHz, CDCl_3) δ (ppm): 3.66 (s, 6H), 6.74 (d, $^3J(\text{H,H}) = 8.0$ Hz, 2H), 7.42 (m, 5H), 7.56 (t, $^3J(\text{H,H}) = 8.0$ Hz, 1H), 7.88 (m, 2H).

3a

¹H NMR, 400 MHz, CDCl₃

3c

¹H NMR, 400 MHz, CDCl₃

3c
¹³C NMR, 80 MHz, CDCl₃

3d

¹H NMR, 400 MHz, CDCl₃

3d

^{13}C NMR, 80 MHz, CDCl_3

3e

¹H NMR, 400 MHz, CDCl₃

3e

¹³C NMR, 80 MHz, CDCl₃

3f

¹H NMR, 400 MHz, CDCl₃

3h
¹H NMR, 400 MHz, CDCl₃

3h

^{13}C NMR, 80 MHz, CDCl_3

3i

¹H NMR, 400 MHz, CDCl₃

3j
¹H NMR, 400 MHz, CDCl₃

3k
¹H NMR, 400 MHz, CDCl₃

3I

^1H NMR, 400 MHz, CDCl_3

3I

^{13}C NMR, 80 MHz, CDCl_3

3m

¹H NMR, 400 MHz, CDCl₃

3n
¹H NMR, 400 MHz, CDCl₃

3o
¹H NMR, 400 MHz, CDCl₃

3p
¹H NMR, 400 MHz, CDCl₃

3q
¹H NMR, 400 MHz, CDCl₃

3q
¹³C NMR, 80 MHz, CDCl₃

3r

¹H NMR, 400 MHz, CDCl₃

3r

^{13}C NMR, 80 MHz, CDCl_3

3s
¹H NMR, 400 MHz, CDCl₃

3t

¹H NMR, 400 MHz, CDCl₃

References

- (1) Florvall, L.; Ögren, S.-O. *J. Med. Chem.* **1982**, 25, 1280.
- (2) (a) For compounds **3a** and **3m** see: Milne, J. E.; Buchwald, S. L. *J. Am. Chem. Soc.* **2004**, 126, 13028. (b) For compound **3b** see: Kawano, N. *Chem. & Pharm. Bull.* **1959**, 7, 698. (c) For compounds **3i**, **3j**, **3k**, and **3o** see: Becht, J.-M.; Gissot, A.; Wagner, A.; Mioskowski, C. *Chem. Eur. J.* **2003**, 9, 3209. (c) For compound **3f** see: Wang, L.; Zhang, Y.; Liu, L.; Wang, Y. *J. Org. Chem.* **2006**, 71, 1284. (d) For compound **3g** see: Korenaga, T.; Kosaki, T.; Fukumura, R.; Ema, T.; Sakai, T. *Org. Lett.* **2005**, 7, 4915. (e) For compound **3n** see: Saa, J. M.; Martorell, G.; Garcia-Raso, A. *J. Org. Chem.* **1992**, 57, 678. (f) For compound **3p** see: Song, C.; Ma, Y.; Chai, Q.; Ma, C.; Jiang, W.; Andrus, M. B. *Tetrahedron* **2005**, 61, 7438. (g) For compound **3s** see: Bach, T.; Bartels, M. *Synthesis* **2003**, 6, 925. (h) For compound **3t** see: Kamikawa, T.; Hayashi, T. *Tetrahedron*, 55, **1999**, 3455.