Synthesis of Bisubstrate and Donor Analogs of Sialyltransferase and Their Inhibitory Activities

Masayuki Izumi,* Katsuhiro Wada, Hideya Yuasa, and Hironobu Hashimoto

Table of contents

General procedures	S2
¹ H and ¹³ C NMR spectra of 10	S3,4
¹ H and ¹³ C NMR spectra of 12	S5,6
¹ H and ¹³ C NMR spectra of 13	S7,8
¹ H and ¹³ C NMR spectra of 14	S9,10
¹ H and ¹³ C NMR spectra of 15	S11,12
¹ H and ¹³ C NMR spectra of 16	S13,14
¹ H and ¹³ C NMR spectra of 17	S15,16
¹ H and ¹³ C NMR spectra of 18	S17,17
¹ H and ¹³ C NMR spectra of 19	S19,20
¹ H and ¹³ C NMR spectra of 21	S21,22
¹ H and ¹³ C NMR spectra of 4	S23,24
¹ H and ¹³ C NMR spectra of 5	S25,26
¹ H and ¹³ C NMR spectra of 22	S27,28
¹ H NMR spectrum of 23	S29
¹ H and ¹³ C NMR spectra of 6	S30,31
¹ H NMR spectrum of 24	S32
¹ H and ¹³ C NMR spectra of 7	S33,34
¹ H NMR spectrum of 26	S35
¹ H and ¹³ C NMR spectra of 8	S36,37

General procedures. ^1H NMR spectra were recorded at 270 or 400 MHz. ^{13}C NMR spectra were recorded at 67.8 or 100 MHz, and ^{31}P NMR spectra were recorded at 109 MHz. Where indicated, NMR peak assignments were made using COSY; all others are subjective. All chemical shifts are quoted on the δ -scale and were referenced to tetramethylsilane ($\delta = 0$ in CDCl_3) or HDO ($\delta = 4.80$ in D_2O) for ^1H NMR, and CDCl_3 ($\delta = 77.0$ in CDCl_3) or NAc ($\delta = 23.1$ in D_2O) for ^{13}C NMR as an internal standard, and 85% H_3PO_4 ($\delta = 0$) for ^{31}P NMR as an external standard. Combinations of the following abbreviations are used to describe NMR spectra: s, singlet; d, doublet; t, triplet; q, quartet; qu, quintet; m, multiplet; br, broad. Low and high-resolution mass spectra were recorded using electrospray ionization techniques with time-of-flight detector. Thin layer chromatography (TLC) was carried out on glass plates coated with silica gel 60F254 (E. Merck). Plates were visualized by irradiation with UV lamp, or dipping in 1% $\text{Ce}(\text{SO}_4)_2$ -1.5% $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$ -10% H_2SO_4 , or orcinol in 10% H_2SO_4 -EtOH and charred. Silica gel column chromatography was performed using Wakogel C-300 (40-75 μm , Wako Chemical). Spherical silica gel 60N (0.1-0.2 mm, neutral) was purchased from Kanto Kagaku. Rat recombinant α -2,6-sialyltransferase, α -2,3-(*N*)-sialyltransferase and CMP-Neu5Ac were purchased from Calbiochem. Inertsil-ODS-3 (5 μm , ϕ 4.6 mm $\times$ 100 mm) HPLC column was purchased from GL Science. Dehydrated THF was purchased from Kanto Kagaku. Dehydrated toluene was purchased from Wako Chemical.

270 MHz ^1H NMR spectrum of compound **10**

67.8 MHz ^{13}C NMR spectrum of compound **10**

270 MHz ^1H NMR spectrum of compound **12**

67.8 MHz ^{13}C NMR spectrum of compound **12**

270 MHz ^1H NMR spectrum of compound **13**

67.8 MHz ^{13}C NMR spectrum of compound **13**

270 MHz ^1H NMR spectrum of compound **14**

67.8 MHz ^{13}C NMR spectrum of compound **14**

270 MHz ^1H NMR spectrum of compound **15**

67.8 MHz ^{13}C NMR spectrum of compound **15**

270 MHz ^1H NMR spectrum of compound **16**

67.8 MHz ^{13}C NMR spectrum of compound **16**

270 MHz ^1H NMR spectrum of compound **17**

67.8 MHz ^{13}C NMR spectrum of compound **17**

270 MHz ^1H NMR spectrum of compound **18**

67.8 MHz ^{13}C NMR spectrum of compound **18**

270 MHz ^1H NMR spectrum of compound **19**

67.8 MHz ^{13}C NMR spectrum of compound **19**

270 MHz ^1H NMR spectrum of compound **21**

67.8 MHz ^{13}C NMR spectrum of compound **21**

400 MHz ^1H NMR spectrum of compound 4

100 MHz ^{13}C NMR spectrum of compound **4**

400 MHz ^1H NMR spectrum of compound **5**

100 MHz ^{13}C NMR spectrum of compound **5**

270 MHz ^1H NMR spectrum of compound **22**

67.8 MHz ^{13}C NMR spectrum of compound **22**

270 MHz ^1H NMR spectrum of compound **23**

400 MHz ^1H NMR spectrum of compound **6**

100 MHz ^{13}C NMR spectrum of compound **6**

270 MHz ^1H NMR of compound **24**

400 MHz ^1H NMR spectrum of compound **7**

100 MHz ^{13}C NMR spectrum of compound 7

270 MHz ^1H NMR of compound **26**

400 MHz ^1H NMR spectrum of compound **8**

100 MHz ^{13}C NMR spectrum of compound **8**