The Synthesis Of Chiral Chromans By The Pd-catalyzed Asymmetric Allylic Alkylation (AAA): Scope, Mechanism And Applications

Barry M. Trost, Hong C. Shen, Li Dong, Jean-Philippe Surivet and Catherine Sylvain

Department of Chemistry

Stanford University

Stanford, CA 94305-5080

Supporting information

General

All reactions were performed under a nitrogen atmosphere unless otherwise indicated. Some solvents were freshly purified by column before use: acetonitrile, methylene chloride, toluene and diethyl ether. Other solvents were purified by distillation: benzene, triethylamine, and pyridine from calcium hydride; tetrahydrofuran, dioxane from sodium benzophenone ketyl; and acetone from calcium sulfate; and dimethylformamide from barium oxide and stored over 4Å molecular sieves. Methanol was distilled from magnesium methoxide. Tris(dibenzylideneacetone)palladium(0) monochloroform complex, $\text{Pd}_2\text{dba}_3\cdot\text{CHCl}_3$ was prepared by the procedure of Ibers¹. All other reagents were used as obtained unless otherwise noted.

Flash Chromatography was performed with EM Science silica gel (0.040-0.063 μm grade). Analytical thin-layer chromatography was performed with 0.2mm coated commercial silica gel plates (E. Merck, DC-Plasrikfolien, kieselgel 60 F₂₅₄). Melting points were obtained on a Thomas-Hoover apparatus in open capillary tubes and are uncorrected. Kugelrohr distillations were performed on a Büchi GKR-50 glass tube oven.

Proton nuclear magnetic resonance (¹H-NMR) data were acquired at 300 MHz on a Varian GEM-300 and at 500 MHz on a Varian GEM-500 spectrometer. Chemical shifts are reported in delta (δ) units, in parts per million (ppm) downfield from tetramethylsilane, or in ppm relative to the singlet at 7.26ppm from chloroform-*d*. Splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; p, pentet; m, multiplet; br, broad. Carbon-13 nuclear

¹ Ukai, T.; Kawazura, H.; Iishii, Y.; Bonnett, J.J.; Ibers, J.A. *J. Organomet. Chem.* **1974**, 65, 263.

magnetic resonance (^{13}C -NMR) data were acquired at 75 MHz on a Varian GEM-300 and at 125 MHz on a Varian GEM-500 spectrometer. Chemical shifts are reported in ppm relative to the center line of a triplet at 77.0 ppm for chloroform-*d* and the center line of a septet at 39.5 ppm in dimethylsulfoxide-*d*₆.

Infrared (IR) data were recorded as films on sodium chloride plates or a potassium bromide (KBr) pellets on a Nicolet 205 and Perkin-Elmer Paragon 500 FT-IR spectrometers. Absorbance frequencies are reported in reciprocal centimeters (cm^{-1}). High resolution mass spectral data (HRMS) were obtained from Mass Spectrometry Resource, School of Pharmacy, University of California-San Francisco, on a Kratos MS9 spectrometer at an ionizing current of 98 mA and an ionizing voltage of 70 eV. Elemental analyses (Anal.) were performed by M.-H.-W. Laboratories of Phoenix, AZ. Chiral HPLC analyses were performed on a Spectra Series P-100 chromatograph using the chiral column and heptane:*iso*-propanol mixtures indicated. Optical rotations were measured on a Jasco DIP-1000 digital polarimeter using 5cm cells and the sodium D line (589 nm) at ambient temperature in the solvent and concentration indicated.

2-(3-Methoxy-5-methylphenoxy)-tetrahydropyran (19): To an orcinol monomethyl ether (10 g, 72 mmol) in distilled THF (105 mL) were added *p*-TsOH (0.69 g, 3.6 mmol) and 3,4-dihydro-2*H*-pyran (15.1 g, 16.4 mL, 180 mmol) at 0°C. The solution was stirred at rt overnight. The reaction mixture was then diluted with an aqueous saturated solution of sodium bicarbonate (80 mL). The aqueous layer was extracted with diethyl ether (2x100 mL). The combined ethereal layers were washed with brine and dried over magnesium sulfate. The removal of solvent *in vacuo* gave an oily residue, which was purified by silica gel flash chromatography eluting with 5% diethyl ether in petroleum ether to afford 2-(3-methoxy-5-methylphenoxy)-tetrahydropyran **19** (15.5 g, 69.8 mmol, 97%) as a colorless oil.

IR (neat): 2943, 1596, 1469, 1196 cm^{-1} ; ^1H -NMR (500 MHz, CDCl_3): δ 6.52 (s, 1H), 6.47 (s, 1H), 6.40 (s, 1H), 5.42 (t, $J=2.9$ Hz, 1H), 3.94 (ddd, $J=2.9, 9.8, 11.8$ Hz, 1H), 3.79 (s, 3H), 3.63 (dt, $J=2.9, 11.8$ Hz, 1H), 2.32 (s, 3H), 2.03 (m, 1H), 1.86 (m, 2H), 1.74-1.60 (m, 3H); ^{13}C -NMR

(125 MHz, CDCl₃): δ 160.4, 158.0, 140.0, 109.4, 108.1, 99.7, 96.2, 55.1, 30.3, 25.1, 21.7, 18.7.
Anal. Calc'd for C₁₃H₁₈O₃: C, 70.24, H, 8.16. Found: C, 70.40; H, 7.93.

2-Methoxy-4-methyl-6-tetrahydropyran-2-yloxybenzaldehyde (20): To a solution of 2-(3-methoxy-5-methylphenoxy)-tetrahydropyran **19** (6.30 g, 28.4 mmol) in distilled THF (100 mL) at 0°C were added tetramethylethylenediamine (8.6 mL, 57.0 mmol) and *n*-butyllithium (23.4 mL, 37.4 mmol, 1.6 M in hexanes) at 0 °C. The resulting dark red mixture was stirred at rt for 5h. To this solution was added DMF (12.3 mL, 159 mmol) and stirring continued for an additional 12h. To this mixture was added water (60 mL) and dichloromethane (120 mL). The aqueous layer was extracted with dichloromethane (2 x 100 mL). The combined organic extracts were washed with brine (60 mL) and dried over magnesium sulfate. The solution was concentrated *in vacuo* and purified by silica gel flash chromatography eluting with diethyl ether:petroleum ether (20:1) to petroleum ether/dichloromethane/diethyl ether (2:1:1) to afford 2-methoxy-4-methyl-6-tetrahydropyran-2-yloxybenzaldehyde **20** (7.10 g, 28.4 mmol, 100%) as a colorless oil.

IR (neat): 2944, 1684, 1606, 1573, 1463, 1238, 1125, 1098 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃): δ 10.5 (s, 1H), 6.64 (s, 1H), 6.42 (s, 1H), 5.57 (t, *J*=2.9 Hz, 1H), 3.90 (s, 3H), 3.89 (m, 1H), 3.64 (m, 1H), 2.37 (s, 1H), 2.05 (m, 1H), 1.91 (m, 2H), 1.74 (m, 2H), 1.65 (m, 1H); ¹³C-NMR (125 MHz, CDCl₃): δ 189.0, 188.8, 161.2, 160.2, 147.4, 108.2, 105.4, 96.2, 61.8, 55.9, 30.0, 28.2, 25.0, 18.1. HRMS: Calc'd for C₁₄H₁₈O₄ [M⁺]: 250.1205. Found: 250.1192.

4-(2-Methoxy-4-methyl-6-tetrahydropyran-2-yloxyphenyl)-but-3-en-2-one (21): A solution of 2-methoxy-4-methyl-6-tetrahydropyran-2-yloxybenzaldehyde **20** (7.4 g, 29.6 mmol) and 1-triphenylphosphanylidene-propan-2-one (14.1 g, 44.4 mmol) in toluene (100 mL) was refluxed for 12 h. The solvent was removed under reduced pressure and the residue was directly chromatographed (petroleum ether/diethyl ether/dichloromethane 2:1:1) to afford 4-(2-methoxy-

4-methyl-6-tetrahydropyran-2-yloxyphenylbut-3-en-2-one **21** (6.9 g, 23.8 mmol, 80%) as white flakes.

m. p.: 90-92 °C; IR (KBr): 2944, 1356, 1250, 1199, 1121, 1094 cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3): δ 7.98 (d, $J=16.6$ Hz, 1H), 7.13 (d, $J=16.6$ Hz, 1H), 6.61 (s, 1H), 6.37 (s, 1H), 5.50 (t, $J=2.7$ Hz, 1H), 3.84 (s, 3H), 3.84 (m, 1H), 2.33 (s, 3H), 2.31 (s, 3H), 2.04-1.84 (m, 3H), 1.80-1.60 (m, 3H); ^{13}C -NMR (50 MHz, CDCl_3): δ 200.0, 159.7, 157.0, 142.4, 135.1, 112.9, 110.2, 108.0, 105.1, 96.2, 61.7, 55.5, 30.1, 26.7, 25.0, 22.3, 18.5. HRMS: Calc'd for $\text{C}_{17}\text{H}_{22}\text{O}_4$ [M^+]: 290.1674. Found: 290.1675.

22

4-(2-Methoxy-4-methyl-6-tetrahydropyran-2-yloxyphenyl)-butan-2-one (22): To 4-(2-methoxy-4-methyl-6-tetrahydropyran-2-yloxyphenyl)-but-3-en-2-one **21** (4.70 g, 16.2 mmol) in methanol (200 mL) was added sodium borohydride (1.75 g, 46.3 mmol, 98%) at rt. The solution was stirred for 30 min and water (100 mL) was added. The methanol was removed under reduced pressure. The residue was diluted with dichloromethane (200 mL) and aqueous layer was extracted with dichloromethane (2 x 100 mL). The combined organic layers were washed with brine, dried over magnesium sulfate. After the removal of solvent, the allylic alcohol was dissolved in ethyl acetate (HPLC grade, 100 mL). To this solution was added platinum dioxide (175 mg, 0.77 mmol) and the reaction mixture was stirred under 1 atm of hydrogen gas for 40 min. The suspension was filtered over a pad of celite and washed with ethyl acetate. The filtrate was concentrated *in vacuo*. The residue was purified over silica gel chromatography eluting with 1:1 to 3:1 diethyl ether/petroleum ether to afford alcohol. The alcohol was resuspended in dichloromethane (20 mL) and to this solution were added 4Å molecular sieves (8.0 g) and *N*-methylmorpholine oxide (3.0 g, 26.8 mmol). The reaction mixture was stirred at rt for 40 min before the reaction mixture was cooled to 0 °C. To this mixture was added tetrapropylammonium perruthenate (0.21 g, 0.61 mmol) in portions (caution: exothermic reaction occurred). After 15 min, the mixture was warmed to rt and stirred for an additional 30 min. The reaction mixture was filtered through a pad of silica gel and the filtrate was concentrated *in vacuo*. The residue was then chromatographed eluting with 1:1 diethyl ether in

petroleum ether to afford 4-(2-methoxy-4-methyl-6-tetrahydropyran-2-yloxyphenyl)-butan-2-one **22** (4.70 g, 16.1 mmol, 99%) as a light yellow oil.

IR (neat): 2942, 1712, 1609, 1587, 1122, 1096 cm^{-1} ; ^1H -NMR (500 MHz, CDCl_3): δ 6.61 (s, 1H), 6.39 (s, 1H), 5.42 (t, $J=3.2$ Hz, 1H), 3.90 (ddd, $J=3.2, 10.0, 11.7$ Hz, 1H), 3.80 (s, 3H), 3.63 (dt, $J=3.2, 11.7$ Hz, 1H), 2.95 (m, 2H), 2.64 (m, 2H), 2.33 (s, 3H), 2.17 (s, 3H), 1.98 (m, 1H), 1.87 (m, 2H), 1.76-1.60 (m, 3H); ^{13}C -NMR (125 MHz, CDCl_3): δ 209.5, 157.8, 155.2, 137.1, 114.8, 107.8, 105.5, 96.1, 61.9, 55.4, 43.5, 30.5, 29.6, 25.2, 21.9, 18.8, 17.8. HRMS: Calc'd for $\text{C}_{17}\text{H}_{24}\text{O}_4$ [M^+]: 292.1674. Found: 292.1675.

Methyl *E*-5-(2-methoxy-4-methyl-6-(tetrahydropyran-2-yloxy))-phenyl-3-methyl-pentanoate (23) Method A: To a suspension of sodium hydride (1 g, 25.6 mmol, 60% oil dispersion, washed twice with distilled hexane) in THF (200 mL), was added trimethylphosphonoacetate (4.3 mL, 26.5 mmol) at 0°C . The resulting mixture was stirred at rt for 1h, and a solution of 4-(2-methoxy-4-methyl-6-tetrahydropyran-2-yloxyphenyl)-butan-2-one **22** (3.7 g, 12.6 mmol) in THF (50 mL) was added. The resulting mixture was stirred at 50°C for 10 h. Water (100 mL) was added and the aqueous layer was extracted with dichloromethane (3 x 100 mL). The combined organic layers were washed with brine (100 mL) and dried over magnesium sulfate. After evaporation under reduced pressure, the residue was purified over silica gel eluting with 6/1 to 4/1 petroleum ether /diethyl ether to afford *E*-5-[2-methoxy-4-methyl-6-(tetrahydropyran-2-yloxy)-phenyl]-3-methyl-pent-2-enoic methyl ester **23** (4.2 g, 12.0 mmol) as a 7:1 (*E*:*Z*) mixture.

Method B: Following the Still's procedure², to a mixture of bis(2,2,2-trifluoroethyl)-methoxycarbonylmethyl phosphonate (36 mg, 25 μL , 0.11 mmol) and 18-crown-6 (79 mg, 0.30 mmol) in distilled THF (0.5 mL) at -78°C was added potassium bis(trimethylsilyl)amide (0.23 mL, 0.5 M in toluenes). After 15 min, ketone **22** (30 mg, 0.10 mmol) was added. The reaction mixture was stirred at -78°C for 30 min and 50°C for 16 h. This mixture was chromatographed

² Still, W. C.; Gennari, C. *Tetrahedron Lett.* **1983**, 24, 4405.

directly eluting with 5% to 20% diethyl ether in petroleum ether to afford the ester **23** (23 mg, 0.066 mmol, 66%, *E/Z* = 4.6:1).

Method C: To a solution of trimethylphosphonoacetate (5.86 g, 5.21 mL, 32.2 mmol) in distilled THF (100 mL) was added *n*-butyllithium (22 mL, 35.4 mmol, 1.6 M in hexanes) at 0°C. After 30 min, to this solution was added 4-[2-methoxy-4-methyl-6-(tetrahydropyran-2-yloxy)-phenyl]-butan-2-one **22** (4.70 g, 16.1 mmol) in distilled THF (50 mL). The solution was stirred at 50°C for 10 h and quenched with water (100 mL). The aqueous layer was extracted with dichloromethane (3 x 100 mL). The combined organic extracts were washed with brine (100 mL) and dried over magnesium sulfate. After removing the solvent, the residue was purified by silica gel chromatography eluting with 6/1 to 4/1 petroleum ether/diethyl ether to afford *E*-5-[2-methoxy-4-methyl-6-(tetrahydropyran-2-yloxy)-phenyl]-3-methylpent-2-enoic methyl ester **23** (5.53 g, 15.9 mmol, 99%) as a colorless oil.

IR (neat): 2944, 1720, 1650, 1609, 1587, 1225, 1167 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃): δ 6.62 (s, 1H), 6.39 (s, 1H), 5.73 (t, *J*=1.2 Hz, 1H), 5.44 (t, *J*=3.1 Hz, 1H), 3.91 (ddd, *J*=3.1, 10.3, 11.5 Hz, 1H), 3.81 (s, 3H), 3.71 (s, 3H), 3.65 (m, 1H), 2.83 (m, 2H), 2.35 (m, 2H), 2.33 (s, 3H), 2.02 (m, 1H), 1.89 (m, 2H), 1.72 (m, 2H), 1.64 (m, 1H); ¹³C-NMR (125 MHz, CDCl₃): δ 167.4, 161.2, 157.8, 155.2, 137.0, 115.1, 114.7, 107.7, 105.0, 95.9, 61.8, 55.4, 50.6, 40.7, 30.5, 25.2, 21.9, 21.6, 18.8, 18.7. HRMS: Calc'd for C₂₀H₂₈O₅ [M⁺]: 348.1936. Found: 348.1936.

2-(5-Hydroxy-3-methyl-pent-3-enyl)-3-methoxy-5-methylphenol (24): Method A: To a solution of α,β -unsaturated ester **23** (4.2 g, 12.0 mmol) in diethyl ether (60 mL), cooled at -78°C, was added DIBAL-H (30 mL, 1M in hexanes, 30 mmol). The reaction was stirred for 20 min., and water (3 mL) was added. The mixture was then warmed up at room temperature and stirred for 1h. The resulting heterogeneous mixture was filtered through celite (washing with diethyl ether) and the filtrate was concentrated *in vacuo*. The residue was taken up in methanol (200 mL) and *p*-TsOH (0.300 g, 1.5 mmol) was added. The solution was stirred at room temperature for 1h. A saturated solution of NaHCO₃ (100 mL) was added and the volatiles were removed under reduced pressure. The residue was diluted with CH₂Cl₂ (100 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 100 mL). The combined organic layers were dried over magnesium

sulfate. After evaporation to dryness, the resulting solid crude material was recrystallized from ethyl acetate/petroleum ether to afford directly allylic alcohol **24** (1.9 g, 8.04 mmol) as a crystalline white solid.

Method B: To allylic alcohol **33** (50 mg, 0.2 mmol) was added *n*-propyl lithium thiolate (2 mL, 1.0 mmol, 0.5 M in HMPA). The resulting orange solution was heated at 120°C for overnight. Without workup, the mixture was purified by flash chromatography eluting with 5% to 60% diethyl ether in petroleum ether to afford allylic alcohol **24** (41 mg, 0.17 mmol, 87%) as a white solid.

m.p.: 92-94°C; IR (neat): 3329, 2935, 1672, 1616, 1595, 1463, 1418, 1109 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ: 6.31 (s, 1H), 6.28 (s, 1H), 5.42 (qt, *J* = 1.2, 5.8 Hz, 1H), 5.37 (s, 1H), 4.16 (br t, *J* = 5.8 Hz, 2H), 3.80 (s, 3H), 2.75 (dd, *J* = 7.6 Hz, 2H), 2.28 (s, 3H), 2.21 (dd, *J* = 8.3 Hz, 2H), 1.80 (s, 1H), 1.77 (d, *J* = 1.2 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 158.2, 1154.2, 140.9, 136.9, 122.8, 113.4, 108.9, 104.0, 59.3, 55.5, 38.6, 21.6, 21.5, 16.4. HRMS: Calc'd for C₁₄H₁₉O₂ [M⁺-OH]: 219.1385. Found: 219.1343.

***E*-5-(2-Hydroxy-6-methoxy-4-methylphenyl)-3-methylpent-2-enyl methyl carbonate (**15c**):**

To a solution of *E*-5-[2-methoxy-4-methyl-6-(tetrahydropyran-2-yloxy)-phenyl]-3-methyl-pent-2-enoic methyl ester **23** (4.2 g, 12.0 mmol) in diethyl ether (60 mL) was added Dibal-H (30 mL, 1 M in hexanes, 30 mmol) at -78 °C. The reaction was stirred for 20 min., and water (3 mL) was added. The mixture was warmed to rt and stirred for 1h. The resulting heterogeneous mixture was filtered through celite and washed with diethyl ether. The filtrate was concentrated *in vacuo*. The residue was taken up in methanol (200 mL) and *p*-TsOH (0.3 g, 1.5 mmol) was added. The solution was stirred at rt for 1 h. A saturated aqueous solution of sodium bicarbonate (100 mL) was added and the volatiles were removed under reduced pressure. The residue was diluted with dichloromethane (100 mL). The aqueous layer was extracted with dichloromethane (3 x 100 mL). The combined organic layers were dried over magnesium sulfate. After evaporating to dryness, the resulting solid crude material was crystallized in ethyl acetate/petroleum ether to afford *E*-2-(5-hydroxy-3-methylpent-3-enyl)-3-methoxy-5-

methylphenol **24** (1.9 g, 8.04 mmol, 67%) as a crystalline white solid. To a solution of *E*-2-(5-hydroxy-3-methylpent-3-enyl)-3-methoxy-5-methylphenol (0.70 g, 2.96 mmol) in dichloromethane (17.5 mL) was added pyridine (0.75 mL, 9.27 mmol). The solution was cooled to 0 °C, and methyl chloroformate (0.275 mL, 3.55 mmol) was added dropwise. After 15 min, water (10 mL) and ethyl acetate (100 mL) were added. The organic layer was washed with an aqueous saturated solution of CuSO₄ (3 x 50 mL), water (50 mL) and brine (50 mL). The organic layer was dried over magnesium sulfate and concentrated to dryness to afford carbonate **15c** (0.87 g, 2.96 mmol, 100%) as a pale yellow oil.

IR (neat): 3445, 2957, 2859, 1650, 1731, 1614, 1594, 1513, 1446, 1383, 1278, 1168, 1092, 934 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃): δ 6.28 (s, 1H), 6.25 (s, 1H), 5.38 (t, *J*=7.2 Hz, 2H), 4.66 (d, *J*=6.9 Hz, 2H), 3.79 (s, 3H), 3.78 (s, 3H), 2.73 (t, *J*=8.1 Hz, 2H), 2.26 (s, 3H), 2.21 (t, *J*=7.8 Hz, 2H), 1.79 (s, 3H); ¹³C-NMR (75 MHz, CDCl₃): δ 158.4, 156.0, 154.2, 144.0, 136.8, 117.3, 113.4, 108.9, 104.0, 64.9, 55.5, 54.6, 38.7, 21.5, 21.4, 16.6. HRMS: Calc'd for C₁₆H₂₂O₅ [M⁺]: 294.1467. Found: 294.1463.

26

3,6-Dimethoxy-4-methylbenzaldehyde (26): To a solution of **25** (2.05 g, 13.5 mmol) in 20 mL of diethyl ether at 0°C was added TMEDA (3.14 g, 4.07 mL, 27 mmol) and *n*-butyllithium (10.1 mL, 16.2 mmol). The solution was heated at reflux for 3h to result in a yellow slurry. The mixture was cooled to 0°C before the addition of DMF (2.35 g, 2.5 mL, 27 mmol). The mixture was slowly warmed to rt and stirred for an additional 2 h before the mixture was quenched with water (10 mL). The mixture was extracted with ethyl acetate (3 x 50 mL). The combined organic extracts were dried and concentrated *in vacuo* before purification by flash chromatography eluting with 5% to 60% diethyl ether in petroleum ether afforded aldehyde **26** as white oily solid (2.35 g, 13.1 mmol, 97%).

IR (neat): 2923m, 2851w, 1689m, 1608s, 1581m, 1464s, 1413m, 1320w, 1237m, 1126s, 970w, 813m cm⁻¹; ¹H-NMR (400 MHz, CDCl₃): δ 10.2 (s, 1H), 6.16 (s, 2H), 3.64 (s, 6H), 2.14 (s, 3H); ¹³C-NMR (100 MHz, CDCl₃): δ 188.0, 161.5, 147.2, 111.2, 104.0, 55.3, 22.2. Anal. Calc'd for C₁₀H₁₂O₃: C, 66.65, H, 6.71. Found: C, 66.70; H, 6.69.

1,3-Dimethoxy-5-methyl-2-vinylbenzene (27): To methyl triphenylphosphonium bromide (5.3 g, 14.8 mmol) in distilled THF (40 mL) at rt was added sodium bis(trimethylsilyl)amide (16.5 mL, 16.5 mmol, 1 M in THF). The resulting yellow slurry was stirred for 3h then cooled to -78 °C. To the mixture was then added a solution of aldehyde **26** (1.90 g, 10.6 mmol) in distilled THF (40 mL) and the resulting mixture was warmed to rt overnight. Reaction was quenched with acetone and water, and extracted with diethyl ether (4 x 50 mL). The combined organic extracts were dried over magnesium sulfate and concentrated *in vacuo*. Flash chromatography eluting with 5% diethyl ether in petroleum ether afford the styrene derivative **27** (1.85 g, 10.4 mmol, 98%) as a colorless oil.

IR (neat): 2938s, 2838s, 2219w, 1820w, 1608s, 1573s, 1464s, 1407s, 1313m, 1241s, 1198s, 1161s, 1123s, 1046m, 1005m, 972m, 909s, 815s, 775w cm⁻¹; ¹H-NMR (300 MHz, CDCl₃): δ 6.93 (dd, *J*=12.3, 18 Hz, 1H), 6.38 (s, 2H), 6.00 (dd, *J*=2.7, 18 Hz, 1H), 5.38 (dd, *J*=3.0, 12.0 Hz, 1H), 3.83 (s, 6H), 2.34 (s, 3H); ¹³C-NMR (75 MHz, CDCl₃): δ 158.4, 138.6, 127.4, 117.4, 112.1, 104.8, 55.6, 22.1. HRMS: Calc'd for C₁₁H₁₄O₂: 178.0993. Found: 178.0999.

2-(2,6-Dimethoxy-4-methylphenyl)-ethanol (28): To a solution of olefin **27** (86 mg, 0.48 mmol) in distilled THF (2 mL) was added pinacolborane (0.97 mL, 0.97 mmol, 1 M in THF) and rhodium tris(triphenylphosphine) chloride (8 mg). The mixture was heated to 50°C for 4.5 h and then stirred rt overnight. Flash chromatography eluting with 5% diethyl ether in petroleum ether afforded pinacol borane (125 mg, 0.41 mmol, 85%) as an oily white solid. To the resulting borane (101 mg, 0.33 mmol) in 4 mL of THF was added sodium hydroxide aqueous solution (1 M, 1.0 mL) and 0.5 mL of hydrogen peroxide (0.5 mL, 30% in water). The mixture was heated to 50°C for 1.5 h. Without workup, the mixture was purified by flash chromatography eluting with 5% to 25% diethyl ether in petroleum ether to afford alcohol **28** as a white oily solid (65 mg, 0.33 mmol, 100%).

IR (neat): 3314b, 2999w, 2942m, 2837w, 1589s, 1463m, 1453m, 1415m, 1317w, 1279w, 1246s, 1187s, 1174s, 1133s, 1049w, 1010m, 972w, 806m cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 6.38 (s, 2H), 3.79 (s, 6H), 3.74 (t, $J=4.8$ Hz, 2H), 2.94 (t, $J=4.8$ Hz, 2H), 2.33 (s, 3H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 158.1, 137.5, 111.9, 104.6, 62.8, 55.5, 26.0, 21.9;

HRMS: Calc'd for $\text{C}_{11}\text{H}_{16}\text{O}_3$: 196.1099. Found: 196.1096.

IR (neat): 2977m, 2929m, 1608w, 1586m, 1464m, 1042m, 1371s, 1314s, 1237m, 1148s, 1112m, 968w, 848w, 813w cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 6.35 (s, 2H), 2.79 (s, 6H), 2.70 (t, $J=8.4$ Hz, 2H), 2.33 (s, 3H), 1.26 (s, 12H), 1.00 (t, $J=8.4$ Hz, 2H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 157.8, 136.2, 118.2, 104.8, 82.7, 55.5, 29.7, 24.8, 21.9, 17.1. Anal. Calc'd for $\text{C}_{17}\text{H}_{27}\text{O}_4\text{B}$: C, 66.68, H, 8.89. Found: C, 66.64; H, 8.91.

For the experimental leading to compounds **15d** and **15d'**:

1-(2-Iodoethyl)-2-methoxybenzene: To a stirred solution of 2-(2-hydroxy)phenyl ethanol (2.0 g, 14.47 mmol) in 10% NaOH (8.0 mL, 20.0 mmol) was added dimethylsulfate dropwise at room temperature. The mixture was stirred at room temperature for 4 h and was partitioned between ether (60 mL) and water (10 mL). After separation of the layers, the aqueous phase was further exacted with ether (2×15 mL). The combined organic layer was washed with brine (10 mL), dried (Na_2SO_4), filtered, concentrated under reduced pressure to give a light yellow oil (2.18 g). The crude methyl ether was used directly for the next step without further purification. To a solution of the above oil (2.18 g) in CH_3CN (12 mL) and ether (20 mL) was added PPh_3 (5.172 g, 19.17 mmol), imidazole (1.547 g, 22.76 mmol), and iodine (5.39 g, 21.24 mmol) at 0°C . The suspension was stirred at room temperature for 2.5 h and was treated with ether (200 mL). The mixture was washed with saturated NaHSO_3 (100 mL), saturated CuSO_4 (20 mL), brine (20 mL), dried (MgSO_4), filtered, and concentrated under reduced pressure. The residue was purified by column chromatography (petroleum ether/ether, 100:3) to produce 1-(2-iodoethyl)-2-methoxybenzene as a light sensitive colorless oil (3.42 g, 86% for two steps).

IR (neat): 2957w, 2834w, 1600w, 1493s, 1464m, 1439w, 1288w, 1244s, 1174m, 1134m, 1050w, 1030m, 752s cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.26 (m, 1H), 7.13 (d, $J=7.8$ Hz, 1H), 6.91 (m, 2H), 3.83 (s, 3H), 3.37 (dd, $J=\text{td}$, $J=1.2, 8.4$ Hz, 2H), 3.22 (t, $J=7.2$ Hz, 2H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 130.2, 128.9, 128.2, 120.4, 111.0, 110.4, 55.2, 35.6.

2-(2-Iodoethyl)-phenol: To a solution of 1-(2-iodoethyl)-2-methoxybenzene (470 mg, 1.81 mmol) in dichloromethane (5 mL) at -78°C was added boron tribromide (1.36 g, 0.52 mL, distilled). The resulting bright yellow solution was slowly warmed to rt and stirred for 2h to result in a brown solution. To this solution was added water (0.5 mL) at rt. After the bubbling stops, the mixture was loaded to silica gel column. The flash chromatography eluting with 30% diethyl ether in petroleum ether afforded 2-(2-iodoethyl)-phenol (421 mg, 1.72 mmol, 95%) as a colorless oil, which turned slightly pink after stored at rt overnight.

IR (neat): 3529bs, 3033w, 2963w, 1608m, 1591m, 1500m, 1455s, 1328m, 1268m, 1236s, 1172s, 1124s, 1079w, 1040w, 935w, 753s cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.17 (m, 2H), 6.95 (t, $J=7.2$ Hz, 1H), 6.76 (d, $J=7.8$ Hz, 1H), 4.94 (s, 1H), 3.43 (t, $J=8.1$ Hz, 2H), 3.24 (t, $J=7.5$ Hz, 2H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 153.1, 130.6, 128.2, 127.0, 121.0, 115.5, 35.0, 4.95. Anal. Calc'd for $\text{C}_8\text{H}_9\text{IO}$: C, 38.73, H, 3.66. Found: C, 38.67; H, 3.65.

2-(2-Iodoethyl)-phenol *tert*-butyldimethylsilyl ether: To a solution of 2-(2-iodoethyl)-phenol (421 mg, 1.72 mmol) in dichloromethane (5 mL) at rt were added imidazole (234 mg, 3.44 mmol) and TBSCl (390 mg, 2.58 mmol). The mixture was stirred for 2h and the reaction mixture was purified by silica gel chromatography eluting with 5% diethyl ether in petroleum ether to afford 2-(2-iodoethyl)-phenol *tert*-butyldimethylsilyl ether (607 mg, 1.69 mmol, 98%) as a colorless oil.

IR (neat): 2932s, 2859s, 1599m, 1561m, 1490s, 1449m, 1255s, 1170m, 1131m, 1043w, 916m, 831m, 780m, 756m cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.11 (m, 2H), 6.86 (t, $J=7.5$ Hz, 1H), 6.76 (d, $J=8.1$ Hz, 1H), 3.23 (td, $J=1.2, 8.4$ Hz, 2H), 3.13 (dd, $J=7.2, 8.4$ Hz, 2H), 1.02 (s, 9H), 0.23 (s, 6H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 153.5, 131.2, 130.3, 128., 121.1, 118.4, 36.0, 25.8, 18.1, 4.37, -4.16. Anal. Calc'd for $\text{C}_{14}\text{H}_{23}\text{IOSi}$: C, 46.41, H, 6.40. Found: C, 46.35; H, 6.37.

***E*-2-(5-Hydroxy-3-methyl-3-pentenyl)-phenol:** To a colorless solution of iodide 2-(2-iodoethyl)-phenol *tert*-butyldimethylsilyl ether (240 mg, 0.67 mmol) and zinc dichloride (200 mg, 1.47 mmol) in THF (4 mL) at -78°C was added *t*-butyllithium (1.0 mL, 1.5 mmol, 1.5 M in pentane). The solution remained colorless. After 30 min, the solution was warmed to rt and stirred for 2 h. To another flask containing a solution of Pd(dppf)Cl₂ (24 mg, 0.055 mmol) in THF (1 mL) was added *n*-butyllithium (44 μ L, 0.066 mmol, 1.5 M in hexane) to give a purple red solution. After 3 min, this solution was transferred to a third flask containing a solution of vinyl iodide **32** (209 mg, 0.67 mmol) in THF (3 mL) via cannula. The resulting dark red solution was stirred for 1 min. To this solution was then added alkyl zinc solution prepared as above via cannula. The solution turned dark brown. After 12 h, the mixture was purified eluting with 2% diethyl ether in petroleum ether to give a colorless oil, which was directly submitted to the next reaction without further purification. To a solution of the oil in THF (3 mL) at rt was added TBAF (2.4 mL, 1 M in THF). The resulting brown solution was stirred for 1 h and purified by silica gel chromatography eluting with 10% to 50% diethyl ether in petroleum ether to afford *E*-2-(5-hydroxy-3-methyl-3-pentenyl)-phenol (20 mg, 0.10 mmol, 16%) as a slightly yellow oil.

IR (neat): 3336b, 2925s, 2854s, 1593w, 1458s, 1379w, 1241m, 1180w, 1104w, 1044w, 990m, 840w, 752s cm⁻¹; ¹H-NMR (300 MHz, CDCl₃): δ 7.09 (m, 2H), 6.86 (t, *J*=6.6, 7.5 Hz, 1H), 6.74 (d, *J*=8.1 Hz, 1H), 5.43 (t, *J*=6.6 Hz, 1H), 4.15 (d, *J*=6.9 Hz, 2H), 2.75 (dd, *J*=7.8, 13.2 Hz, 2H), 2.32 (t, *J*=7.8 Hz, 2H), 1.75 (s, 3H); ¹³C-NMR (75 MHz, CDCl₃): δ 153.6, 139.8, 130.2, 128.1, 127.2, 123.6, 120.7, 115.3, 59.3, 39.5, 29.7, 28.6. HRMS: Calc'd for C₁₂H₁₆O₂: 192.1150. Found: 192.1148.

***Z*-2-(5-Hydroxy-3-methyl-3-pentenyl)-phenol:** To a colorless solution of 2-(2-iodoethyl)-phenol *tert*-butyldimethylsilyl ether (390 mg, 1.09 mmol) and zinc dichloride (325 mg, 2.39 mmol) in THF (4 mL) at -78°C was added *t*-butyllithium (1.6 mL, 2.39 mmol, 1.5 M in pentane). The solution remained colorless. After 30 min, the solution was warmed to rt and stirred for 2 h. To another flask containing a solution of Pd(dppf)Cl₂ (40 mg, 0.055 mmol) in THF (1 mL) was added *n*-butyllithium (84 μ L, 0.11 mmol, 1.5 M in hexane) to give a purple red solution. After 3 min, this solution was transferred to a third flask containing a solution of vinyl iodide **30** (340

mg, 1.09 mmol) in THF (3 mL) via cannula. The resulting dark red solution was stirred for 1 min. To this solution was then added alkyl zinc solution prepared as above via cannula. The solution turned dark brown. After 18 h at rt, the mixture was purified eluting with 2% diethyl ether in petroleum ether to give a colorless oil, which was directly submitted to the next reaction without further purification. To a solution of the oil in THF (3 mL) at rt was added TBAF (4 mL, 1 M in THF). The resulting brown solution was stirred for 1 h and purified by silica gel chromatography eluting with 10% to 50% diethyl ether in petroleum ether to afford *Z*-2-(5-hydroxy-3-methyl-3-pentenyl)-phenol (74 mg, 0.43 mmol, 39%) as a slightly yellow oil.

IR (neat): 3322b, 2929s, 2859m, 1593w, 1457s, 1378m, 1242m, 1176w, 1108w, 982m, 836w, 752s cm⁻¹; ¹H-NMR (300 MHz, CDCl₃): δ 7.08 (m, 2H), 6.81 (m, 2H), 5.49 (t, *J*=6.9 Hz, 1H), 3.99 (d, *J*=7.2 Hz, 2H), 2.75 (t, *J*=7.5 Hz, 2H), 2.39 (t, *J*=7.8 Hz, 2H), 1.81 (s, 3H); ¹³C-NMR (75 MHz, CDCl₃): δ 154.2, 141.8, 130.5, 127.7, 127.5, 123.8, 120.2, 115.5, 58.7, 32.3, 29.8, 23.6. HRMS: Calc'd for C₁₂H₁₄O ([M-H₂O]⁺): 174.1045. Found: 174.1035.

For the experimental leading to compounds **15e** and **15e'**:

2-Iodoethyl-4-methoxyphenol *tert*-butyldimethylsilyl ether: To a stirred solution of (2-allyl-4-methoxy-phenoxy)-*tert*-butyldimethylsilane (2.1 g, 7.54 mmol) and NMO (1.764 g, 15.08 mmol) in CH₂Cl₂ (20 mL) was added OsO₄ (4 w/w% aqueous solution, 0.958 mL, 0.151 mmol) at 0 °C. The brown solution was warmed to room temperature and stirred for 1.5 h. The mixture was treated with EtOAc (60 mL) and washed with 0.5 N HCl (15 mL), brine (5 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to give the crude diol as a brown oil.

To a stirred solution of the above residue in acetone/water (4:1, 25 mL) was added NaIO₄ (3.2 g, 15.08 mmol) at room temperature. The white suspension was stirred for 1 h and treated with ether (60 mL). The mixture was washed with water (20 mL), brine (10 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure to give the crude aldehyde. The crude material was purified by column chromatography (petroleum ether/ether, 100:6) to produce the aldehyde as a clear liquid.

To a stirred solution of the above aldehyde in MeOH (15 mL) was added NaBH₄ (270 mg, 7.14 mmol) at 0 °C. The mixture was warmed up to room temperature and stirred for 10 min before treated with ether (60 mL). The mixture was washed with water (20 mL), brine (10 mL),

dried (MgSO₄), filtered, and concentrated under reduced pressure to give the crude alcohol. The crude material was purified by column chromatography (petroleum ether/ether, 100:6) to produce compound the alcohol as a clear liquid (2.1 g, 99%).

Employing the similar procedure for the preparation of 1-(2-iodoethyl)-2-methoxybenzene, the title compound was prepared from the above alcohol (93%). IR (neat) 2930, 1497, 1426, 1263, 1223, 1043 cm⁻¹; ¹H-NMR (300 MHz, CDCl₃) δ 6.62 (m, 3 H), 3.66(s, 3 H), 3.24 (t, *J* = 7.8 Hz, 2 H), 3.05 (t, *J* = 7.8 Hz, 2 H), 0.96 (s, 9 H), 0.16 (s, 6 H); ¹³C-NMR (75 MHz, CDCl₃) δ 153.6, 147.2, 131.9, 118.9, 115.7, 112.6, 55.5, 36.2, 25.7, 18.0, 4.1, -4.2.

***E*-2-(5-Hydroxy-3-methyl-pent-2-enyl)-4-methoxyphenol:** To a colorless solution of (2-iodoethyl)-4-methoxy-phenol *tert*-butyldimethylsilyl ether (490 mg, 1.30 mmol) and zinc dichloride (350 mg, 2.60 mmol) in THF (5 mL) at -78°C was added *t*-butyllithium (1.73 mL, 2.6 mmol, 1.5 M in pentane). The solution remained colorless. After 30 min, the solution was warmed to rt and stirred for 2 h. To another flask containing a solution of Pd(dppf)Cl₂ (38 mg, 0.052 mmol) in THF (1.5 mL) was added was added *n*-butyllithium (42 uL, 0.10 mmol, 2.5 M in hexane) to give a purple red solution. After 3 min, this solution was transferred to a third flask containing a solution of vinyl iodide **32** (324 mg, 1.0 mmol) in THF (3.5 mL) via cannula. The resulting dark red solution was stirred for 1 min. To this solution was then added alkyl zinc solution prepared as above via cannula. The solution turned dark brown. After 12 h, the mixture was purified eluting with 2% diethyl ether in petroleum ether to give a colorless oil, which was directly submitted to the next reaction without further purification. To a solution of the oil in THF (3 mL) at rt was added TBAF (2.5 mL, 1 M in THF). The resulting brown solution was stirred for 1 h, and without workup directly purified by silica gel chromatography eluting with 10% to 50% diethyl ether in petroleum ether to afford *E*-2-(5-hydroxy-3-methyl-pent-2-enyl)-4-methoxyphenol (25 mg, 0.11 mmol, 11%) as a slightly yellow oil.

IR (neat): 3364b, 2926s, 2860m, 1506s, 1451m, 1434s, 1209s, 1153w, 1046w, 989w, 805w cm⁻¹; ¹H-NMR (300 MHz, CDCl₃): δ 6.67 (m, 3H), 5.42 (t, *J*=6.6 Hz, 1H), 4.14 (d, *J*=6.9 Hz, 2H), 3.75 (s, 3H), 2.72 (t, *J*=7.8 Hz, 2H), 2.31 (t, *J*=7.8 Hz, 2H), 1.74 (s, 3H); ¹³C-NMR (75 MHz, CDCl₃): δ 153.6, 147.6, 139.7, 129.4, 123.6, 116.0, 115.9, 111.7, 59.3, 55.7, 39.5, 29.7, 28.9; HRMS: Calc'd for C₁₃H₁₈O₃: 222.1256. Found: 222.1258.

Z-2-(5-Hydroxy-3-methyl-pent-2-enyl)-4-methoxyphenol: To a colorless solution of (2-iodoethyl)-4-methoxy-phenol *tert*-butyldimethylsilyl ether (128 mg, 0.34 mmol) and zinc dichloride (92 mg, 0.68 mmol) in THF (3 mL) at -78°C was added *t*-butyllithium (0.26 mL, 0.39 mmol, 1.5 M in pentane). The solution remained colorless. After 30 min, the solution was warmed to rt and stirred for 2 h. To another flask containing a solution of Pd(dppf)Cl₂ (10 mg, 0.014 mmol) in THF (1 mL) was added was added *n*-butyllithium (11 uL, 0.028 mmol, 2.5 M in hexane) to give a purple red solution. After 3 min, this solution was transferred to a third flask containing a solution of vinyl iodide **30** (88 mg, 0.28 mmol) in THF (2 mL) via cannula. The resulting dark red solution was stirred for 1 min. To this solution was then added alkyl zinc solution prepared as above via cannula. The solution turned dark brown. After 12 h, the mixture was purified eluting with 2% diethyl ether in petroleum ether to give a colorless oil, which was directly submitted to the next reaction without further purification. To a solution of the oil in THF (3 mL) at rt was added TBAF (0.5 mL, 1 M in THF). The resulting brown solution was stirred for 1 h, and without workup directly purified by silica gel chromatography eluting with 10% to 50% diethyl ether in petroleum ether to afford Z-2-(5-hydroxy-3-methyl-pent-2-enyl)-4-methoxyphenol (14 mg, 0.063 mmol, 22%) as a slightly yellow oil.

IR (neat): 3375b, 2928s, 1508s, 1433s, 1377w, 1209s, 1154m, 1032m, 984w, 808w, 713w cm⁻¹; ¹H-NMR (400 MHz, CDCl₃): δ 6.67 (m, 3H), 5.50 (t, *J*=6.8 Hz, 1H), 4.01 (d, *J*=7.6 Hz, 2H), 3.75 (s, 3H), 2.71 (t, *J*=8.4 Hz, 2H), 2.38 (t, *J*=8.4 Hz, 2H), 1.76 (s, 3H); ¹³C-NMR (100 MHz, CDCl₃): δ 153.3, 147.9, 141.8, 128.7, 124.0, 116.2, 116.1, 112.0, 58.8, 55.7, 32.5, 30.2, 29.7; HRMS: Calc'd for C₁₃H₁₈O₃: 222.1256. Found: 222.1255.

For the preparation of compound **15f**:

1,4-Dimethoxy-2,3-dimethylbenzene.³ To DMSO (75 ml), was added powdered KOH (16.2 g, 288 mmol). After stirring for 5 min., 2,3-dimethyl hydroquinone (5.0 g, 36,1 mmol) was added

³ Johnston, R. A. W.; Rose, M. E. *Tetrahedron*, **1979**, 35, 2169.

followed immediately by iodomethane (9 ml, 744 mmol). After stirring at for 1 h, H₂O was added (700 ml), and the mixture was extracted with dichloromethane (3 x 500 ml). The combined organic extracts were washed with water (5 x 250 ml). The combined layers were dried over MgSO₄. After evaporation of the solvent under reduced pressure, chromatographic purification on silica gel (P.E./Et₂O 95/5) gave the title compound (5.6 g, 94 % yield) as a white solid, *R_f* = 0.29 (P.E./Et₂O 98/2), mp = 73-74°C.

IR (KBr): 2955, 2838, 1483, 1259, 1116, 1097, 799, 719 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.17 (s, 6H), 3.78 (s, 6H), 6.66 (s, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 11.9, 55.9, 107.6, 126.0, 151.8.

2,5-Dimethoxy-3,4-dimethoxybromobenzene.⁴ To a solution of the above compound (0.375, 2.2 mmol) in acetic acid (27 ml) was added at bromine (0.113 ml, 2.2 mmol). After stirring for 10 min. at , the mixture was quenched by aqueous bisulfite (50 ml) and extracted with ether (3x30 ml). After drying the ethereal layer on MgSO₄, the solvent was evaporated under reduced pressure. Filtration on silica gel (P.E./Et₂O 95/5) gave the title compound (0.535 g, 97% yield) as a colorless oil, *R_f* = 0.31 (P.E./Et₂O 98/2).

IR (neat): 2996, 2936, 2833, 1558, 1469, 1400, 1301, 1231, 1113, 1014, 774 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 6.86 (s, 1H), 3.77 (s, 3H), 3.73 (s, 3H), 2.24 (s, 3H), 2.09 (s, 3H); ¹³C NMR (75 MHz, CDCl₃): δ 154.1, 149.0, 132.3, 125.9, 113.3, 112.2, 60.5, 55.8, 13.3, 12.1.

2,5 Dimethoxy-3,4-dimethoxyphenol. Method 1:⁵ To a stirred solution of the above bromide (0.2 g, 0.81 mmol) in THF (1.6 ml) under argon at -78°C, was added drop wise *n*-BuLi (2.15 M) (0.376 ml, 0.81 mmol). In a separate flask, *n*-BuLi (2.15 M) (0.753 ml, 1.6 mmol) was added to a solution of *tert*-butylhydroperoxide in toluene (3.27 M) (0.495 ml, 1.6 mmol) at -78°C under argon. After stirring for 30 min., the hydroperoxide anion solution was added by cannula to the

⁴ Ebersson, L.; Hashorn, M.P.; Radner, F.; Persson, O. *J. Chem. Soc., Perkin Trans. 2*, **1998**, 59.

⁵ Julia, M.; Pfeuty-Saint Jalmes V.; Plé, K.; Verpeaux, J.-N. *Bull. Soc. Chim. Fr.* **1996**, 133, 15.

solution of the aromatic anion, and stirring was continued for 2h at 0°C. The reaction was quenched by pouring into a cold 10% HCl solution (10 ml). The aqueous layer was extracted with dichloromethane (3 x 20 ml). The combined layers were dried over MgSO₄ and the solvent was evaporated under reduced pressure. Chromatographic purification on silica gel (P.E./Et₂O 8/2) gave the title compound (108 mg, 74 % yield) as a white solid.

Method 2:⁶

Freshly cut potassium (1.78 g, 45.6 mmol), anhydrous magnesium chloride (2.42 g, 25.4 mmol) and THF (60 ml) were placed in an oven-dried 250 ml three-necked flask under argon. The mixture was stirred and heated to reflux for 3 h. to give a dark gray mixture. Then, after cooling at rt, the above bromide (2.62 g, 10.6 mmol) in solution into THF (2 ml) was added followed by BH₃.THF (1M) (13.5 ml, 13.4 mmol). The mixture was refluxed for 1 h., then cooled at rt . Water was carefully added drop wise until H₂ gas evolution ceased. Next, (1M) aqueous solution of NaOH (58.3 ml) was added followed by 30% H₂O₂ (5.87 ml). After 62h the reaction mixture was poured into water and extracted with AcOEt (3 x 100 ml). The combined layers were dried over MgSO₄ and the solvent was evaporated under reduced pressure. Chromatographic purification on silica gel (P.E./Et₂O 8/2) gave the title compound (1.1 g, 60 % yield) as a white solid; R_f = 0.36 (P.E./ Et₂O 7/3), mp = 70-71°C.

IR (neat): 3412, 2932, 1603, 1484, 1467, 1344, 1314, 1246, 1192, 1128, 1092, 1037, 994, 828 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 6,41 (s, 1H), 5,50 (s, 1H), 3,76 (s, 3H), 3,71 (s, 3H), 2,20 (s, 3H), 2,06 (s, 3H); ¹³C NMR (75 MHz, CDCl₃): δ 154.1, 146.5, 138.9, 13.1, 116.8, 96.4, 61.0, 55.6, 12.6, 11.9.

2,5,8-Trimethoxy-2,6,7-dimethyl chroman.⁷ To a degassed solution of the above compound (1.73 g, 9.5 mmol) in dry methanol (16 ml) and trimethyl orthoformate (4 ml) under argon and cooled in an ice bath, was added H₂SO₄ (0.023 ml) followed by methyl vinyl ketone (1.6 ml, 19 mmol). The mixture was stirred without cooling for 18 h. Then methyl vinyl ketone (1.6 ml, 19 mmol) was added at 0°C, and stirred overnight at rt. This operation was repeated two more times.

⁶ Garofalo, A.W.; Litvak, J.; Wang, L.; Dubenko, L.G.; Cooper, R.; Bierer, D.E. *J. Org. Chem.* **1999**, *64*, 3369.

⁷ Mizuguchi, E.; Achiwa, K. *Chem.Pharm. Bull.* **1997**, *45*, 1209.

Then, H₂O was added to the flask and the mixture was extracted with ether (3 x 20 ml). After drying the ethereal layer over MgSO₄, the solvent was removed under reduced pressure. Chromatographic purification on silica gel (P.E./Et₂O 9/1) afforded the title compound (1.76 g, 70% yield) as a colorless oil, *R_f* = 0.32 (P.E./Et₂O 9/1).

IR (neat): 3416, 2939, 1610, 1463, 1415, 1379, 1330, 1267, 1234, 1150, 1088, 1061 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 3.78 (s, 3H), 3.68 (s, 3H), 3.32 (s, 3H), 2.82-2.74 (m, 2H), 2.16 (s, 3H), 2.31 (s, 3H), 2.15-2.03 (m, 1H), 1.81-1.65 (m, 1H), 1.59 (s, 3H); ¹³C NMR (50 MHz, CDCl₃): δ 151.5, 143.7, 142.9, 128.5, 121.1, 114.4, 97.9, 60.3, 59.9, 49.1, 31.3, 23.1, 16.3, 12.1, 11.9. HRMS calcd for C₁₅H₂₂O₄ (M) = 266.1518, found = 266.1515.

5,8-Dimethoxy-2,6,7-dimethyl chroman-2-ol. To a solution of the above acetal (2.4 g, 9.1 mmol) in acetone (10 ml), was added a solution of (0.1 N) HCl (7 ml). The stirred reaction was heated at reflux as the acetone was removed by distillation over a 15 min. period. An additional 5 ml of acetone was added and the distillation process repeated over 5 min. The reaction mixture was cooled and extracted with ether. After drying ethereal layer over MgSO₄, the solvent was removed under reduced pressure. Crystallization from pentane afforded the title compound (1.93 g, 90% yield) as a white solid, *R_f* = 0.25 (P.E./Et₂O 6/4), mp = 74-75°C.

IR (KBr): 3480, 2936, 1457, 1410, 1330, 1259, 1238, 1150, 1083, 1022, 948 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 3.74 (s, 3H), 3.68 (s, 3H), 2.85-2.81 (m, 2H), 2.66 (s, 1H), 2.16 (s, 3H), 2.13 (s, 3H), 2.07 (m, 1H), 1.76 (m, 1H), 1.68 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 152.9, 151.5, 145.3, 142.3, 128.7, 128.2, 121.2, 118.4, 113.6, 100.2, 96.0, 60.9, 60.8, 60.3, 60.0, 43.8, 31.0, 29.6, 28.6, 18.3, 16.6, 12.5, 12.2, 12.0. HRMS calcd for C₁₄H₂₀O₄ (M) = 252.1361, found = 252.1373.

Acetic acid 2,5-dimethoxy-3,4-dimethyl-6-(3-oxo-butyl)-phenyl ester. A solution of acetyl chloride (1.7 ml, 23.7 mmol) in THF (7 ml) was added at 0°C to a solution of the above acetal (2 g, 7.9 mmol) and triethylamine (5.2 ml, 37.6 mmol) in THF (43 ml). After stirring for 20 h. at rt,

the mixture was quenched with H₂O and extracted with EtOAc (3 x 50 ml). The combined layers were dried over MgSO₄, and the solvent was evaporated under reduced pressure. Filtration on silica gel (P.E./EtOAc 7/3) gave the title ketone (2.3 g, 98%) as a white solid, mp = 63-64°C, *R_f* = 0.3 (P.E./EtOAc 8/2).

IR (KBr): 2937, 1764, 1716, 1462, 1405, 1369, 1250, 1205, 1077, 1014 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.12 (s, 3H), 2.16 (s, 6H), 2.32 (s, 3H), 2.6–2.8 (m, 4H), 3.67 (s, 6H); ¹³C NMR (75 MHz, CDCl₃): δ 208.2, 169.3, 152.7, 146.3, 140.4, 129.7, 128.6, 124.8, 60.9, 60.6, 43.5, 29.7, 20.5, 18.8, 12.7, 12.5. HRMS calcd for C₁₆H₂₂O₅ (M) = 294.1467, found = 294.1464.

5-(2-Acetoxy-3,6-dimethoxy-4,5-dimethylphenyl)-3-methylpent-2-enoic acid. A solution of trimethyl phosphonoacetate (2.94 g, 16.1 mmol) in THF (140 ml) under an atmosphere of argon, was treated with 60% NaH in oil (0.586 g, 14.66 mmol) at 0°C. The mixture was stirred for 2 h. at rt., which was followed by the addition of a solution of the above ketone (2.16 g, 7.33 mmol) in THF (40 ml). The reaction mixture was stirred for 18 h. at 40°C, then quenched with H₂O and extracted with EtOAc (3 x 150 ml). The combined layers were dried over MgSO₄, and the solvent was evaporated under reduced pressure. Chromatographic purification on silica gel (P.E./EtOAc 85/15), provided the title ester (2 g, 80% yield) as a colorless oil, *R_f* = 0.56 (P.E./EtOAc 7/3).

IR (neat): 2947, 1765, 1716, 1651, 1463, 1408, 1368, 1250, 1204, 1149, 1087 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.17 (s, 6H), 2.22 (s, 3H), 2.28–2.36 (m, 2H), 2.34 (s, 3H), 2.60–2.66 (m, 2H), 3.67 (s, 3H), 3.69 (s, 6H), 5.7 (s, 1H); ¹³C NMR (75 MHz, CDCl₃): δ 169.1, 167.1, 159.7, 152.7, 146.3, 140.5, 129.8, 128.6, 124.9, 115.1, 61.0, 60.7, 50.8, 41.0, 23.5, 20.6, 18.8, 12.8, 12.6. HRMS calcd for C₁₉H₂₆O₆ (M) = 350.1729, found = 350.1730.

2-((E)-5-Hydroxy-3-methylpent-3-enyl)-3,6-dimethoxy-4,5-dimethyl phenol. A solution of the above ester (0.957 g, 2.7 mmol) in toluene (23 ml) was treated with (1 M) DIBAL-H (11.4 ml, 11.4 mmol) at 0°C. The mixture was stirred for 4 h. at rt., then carefully treated with a dilute

aqueous solution of HCl, and the grainy precipitate was removed by filtration. The filtrate was washed with H₂O, then the organic layer was dried over MgSO₄ and the solvent was evaporated under reduced pressure. Chromatographic purification on silica gel (P.E./EtOAc 7/3 then 6/4) afforded the title alcohol (0.711 g, 93% yield) as a mixture of two isomers *E/Z* in the ratio 3/1. Several purifications were necessary to isolate the *E* isomer as a white solid, mp = 124 - 125°C, *R*_f = 0.32 (P.E./EtOAc 6/4).

IR (KBr): 3447, 3286, 2940, 1607, 1462, 1413, 1236, 1078, 975, 848 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 5.57 (s, 1H), 5.43 (t, *J* = 7 Hz, 1H), 4.13 (t, *J* = 6 Hz, 1H), 3.72 (s, 3H), 3.67 (s, 3H), 2.75 (t, *J* = 8.4 Hz, 2H), 2.25 (t, *J* = 8.4 Hz, 2H), 2.17 (s, 3H), 2.11 (s, 3H), 1.77 (s, 3H), 1.2 (t, *J* = 5.4 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃): δ 153.1, 145.1, 141.7, 140.4, 127.3, 123.0, 121.0, 119.1, 60.99, 60.91, 59.3, 39.4, 23.1, 16.2, 12.5, 12.1. HRMS calcd for C₁₆H₂₄O₄ (M) = 280.1674, found = 280.1674.

Carbonic acid (*E*)-5-(2-hydroxy-3,6-dimethoxy-4,5-dimethylphenyl)-3-methylpent-2-enyl ester methyl ester (15f**).** The above alcohol (0.1 g, 0.35 mmol) was dissolved in CH₂Cl₂ (2.5 ml) and pyridine (0.1 ml), and the solution was cooled to 0°C. Methyl chloroformate (0.0355 ml, 0.46 mmol) was added and the mixture stirred for 2h. at rt. The reaction mixture was then diluted with H₂O (10 ml) and extracted with EtOAc (3 x 15 ml). The combined layers were dried over MgSO₄, and the solvent was evaporated under reduced pressure. Chromatographic purification on silica gel (P.E./EtOAc 9/1 then 8/2) following by a recrystallization in ether, afford **15f** (0.111 g, 93% yield) as a white solid, mp = 59 - 60°C, *R*_f = 0.35 (P.E./EtOAc 8.5/1.5).

IR (neat): 3452, 2941, 1747, 1462, 1414, 1264, 1076, 1014, 936 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 5.72 (s, 1H), 5.44 (t, *J* = 7 Hz, 1H), 4.67 (d, *J* = 7 Hz, 1H), 3.77 (s, 3H), 3.71 (s, 3H), 3.66 (s, 3H), 2.74 (t, *J* = 8 Hz, 2H), 2.28 (t, *J* = 8 Hz), 2.17 (s, 3H), 2.11 (s, 3H), 1.81 (s, 3H); ¹³C NMR (75 MHz, CDCl₃): δ 155.8, 153.0, 145.1, 143.6, 141.6, 127.3, 120.9, 120.9, 119.0, 117.3, 64.7, 60.9, 60.8, 54.4, 39.3, 22.9, 16.4, 12.4, 12.1. HRMS calcd for C₁₈H₂₆O₆ (M) = 338.1729, found = 338.1732.

***E*-5-(5-Fluoro-2-hydroxyphenyl)-pent-2-enyl methyl carbonate (15g):** To allylic alcohol **38**⁸ (1.32 g, 4.26 mmol) in dichloromethane (5 mL) were added pyridine (1.01 g, 1.03 mL, 12.8 mmol) and then methyl chloroformate (604 mg, 0.50 mL, 6.39 mmol) at 0 °C. The reaction mixture was warmed to rt over 1h. After the addition of water (5 mL), the mixture was extracted with diethyl ether (3 x 50 mL). The combined organic extracts were washed with saturated copper sulfate aqueous solution (2 x 20 mL), brine, and dried over magnesium sulfate. After the removal of solvent, silyl phenol ether carbonate was obtained as a slightly yellow oil (1.44 g, 3.91 mmol, 92%). To a solution of silyl phenol ether carbonate (1.34 g, 3.64 mmol) in THF (5 mL) was added TBAF (7.2 mL, 1 M in THF) at rt. After 1h the mixture was directly purified by silica gel chromatography eluting with 5% to 50% diethyl ether in petroleum ether to afford phenol **15g** (0.81 g, 3.19 mmol, 88%) as a brown oil.

IR (neat): 3446bs, 2960s, 2859w, 1746s, 1728s, 1621w, 1510s, 1443s, 1383m, 1264s, 1186s, 941m, 873w, 793m, 712w cm⁻¹; ¹H-NMR (400 MHz, CDCl₃): δ 6.77 (m, 3H), 5.84 (dt, *J*=6.8, 15.2 Hz, 1H), 5.61 (dt, *J*=6.8, 15.2 Hz, 1H), 4.56 (d, *J*=6.4 Hz, 2H), 3.78 (s, 3H), 2.63 (dd, *J*=6.8, 8.0 Hz, 2H), 2.32 (q, *J*=8.0 Hz, 2H); ¹³C-NMR (100 MHz, CDCl₃): δ 157.9, 155.6 (d, *J*=17.5 Hz), 149.8, 136.2, 129.5 (d, *J*=6.8 Hz), 123.8, 116.5, 116.2 (d, *J*=11.9 Hz), 113.0 (d, *J*=12.7 Hz), 68.6, 54.8, 32.0, 29.4. HRMS: Calc'd for C₁₃H₁₅O₄F: 254.0954. Found: 254.0955.

For the preparation of compounds **15b'**, **15h**, **15h'**, **15i**, **15i'**, **15j**, and **15j'**:

(*E*)-Ethyl 5-(2-*tert*-butyldimethylsilyloxy-5-methylphenyl)-pent-2-enoate. To a stirred solution of 6-methylchroman-2-one⁹ (500 mg, 3.086 mmol) in THF (8 mL) was added DIBAL-H (1.5 M in toluene, 2.2 mL, 3.27 mmol) in 40 min at -90 °C. The mixture was stirred at this temperature for 1 h and quenched with water (20 mL). The suspension was extracted with ether (3 x 15 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure to afford a light yellow oil. The lactol was purified by column chromatography to provide a clear oil (349 mg, 69%).

⁸ For preparation of **38**, see: Chandrasekhar, S.; Reddy, M. V. *Tetrahedron*, **2000**, 56, 6339.

⁹ Yates, P.; Macas, T.S. *Can. J. Chem.* **1988**, 66, 1.

To a stirred solution of the above lactol (196 mg, 1.195 mmol) in benzene (3 mL) was added carboethoxymethylenetriphenylphosphorane (458 mg, 1.315 mmol) at room temperature. The mixture was stirred overnight and the volatiles were removed under reduced pressure. The crude product was used without purification for the next step.

To a stirred solution of the above residue in DMF (2 mL) was added TBSCl (216.1 mg, 1.434 mmol) and imidazole (203.1 mg, 2.979 mmol) at room temperature. The mixture was stirred for 20 h before treated with ether (20 mL) and water (10 mL). After separation of the layers, the organic phase was washed with water (3 × 15 mL), brine (10 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure to afford the title compound as a clear oil (257.2 mg, 62%). IR (neat) 1725, 1499, 1270 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.21 (s, 6 H), 1.00 (s, 9 H), 1.28 (t, *J* = 6.9 Hz, 3 H), 2.24 (s, 3 H), 2.46 (t, *J* = 7.2 Hz, 2 H), 2.67 (t, *J* = 7.2 Hz, 2 H), 4.17 (q, *J* = 6.9 Hz, 2 H), 5.83 (d, *J* = 15.6 Hz, 1 H), 6.67 (d, *J* = 8.1 Hz, 1 H), 7.03-6.86 (m, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 166.7, 151.2, 148.7, 131.0, 130.7, 130.2, 127.5, 121.4, 118.1, 60.1, 32.7, 29.3, 25.7, 20.5, 18.2, 14.2, -4.2; ESIMS, *m/z* calcd for C₂₀H₃₂O₃Si[MH]⁺: 349, found: 349.

General Procedure for the Preparation of the Carbonate: Preparation of (*E*)-5-(2-*tert*-Butyldimethylsilyloxy-5-methylphenyl)-pent-2-enyl Methyl Carbonate. To a stirred solution of compound **7** (257 mg, 0.738 mmol) in THF (2 mL) was added DIBAL (1.5 M in toluene, 1.9 mL, 1.846 mmol) at -78 °C. After being stirred at this temperature for 1 h, the solution was warmed up to room temperature and stirred for an additional 1 h before quenched with the addition of water (10 mL) and ether (20 mL). The mixture was filtered through a pad of celite and the layers were separated. The aqueous phase was extracted with ether (2 × 10 mL). The combined organic layer was washed with brine (5 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure to afford (*E*)-5-(2-*tert*-butyldimethylsilyloxy-5-methylphenyl)-pent-2-en-1-ol as a clear oil (194.4 mg, 86%). IR (neat) 3345, 1498, 1250 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.22 (s, 6 H), 1.01 (s, 9 H), 2.25 (s, 3 H), 2.33 (t, *J* = 8.4 Hz, 2 H), 2.62 (t, *J* = 7.8 Hz, 2 H), 4.07 (m, 2 H), 5.74-5.66 (m, 2 H), 6.67 (d, *J* = 8.1 Hz, 1 H), 6.90-6.85

(m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 151.2, 132.8, 131.9, 130.8, 130.0, 129.1, 127.2, 118.1, 63.7, 32.8, 30.4, 25.7, 20.5, 18.2, 14.2, -4.2.

To a stirred solution of the above alcohol (117 mg, 0.382 mmol), DMAP (9.3 mg, 0.077 mmol), and pyridine (120.8 mg, 1.528 mmol) in CH_2Cl_2 (2 mL) was added dropwise methyl chloroformate (144.4 mg, 1.528 mmol) at 0 °C. After being stirred at room temperature for 24 h, the mixture was treated with ether (15 mL) and water (5 mL). The organic layer was washed with CuSO_4 (10 mL), brine (5 mL), dried (MgSO_4), filtered, and concentrated under reduced pressure to afford the title compound as a clear oil (128.5 mg, 91%). IR (neat) 1751, 1500, 1266 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.22 (s, 6 H), 1.01 (s, 9 H), 2.25 (s, 3 H), 2.35 (t, $J = 7.2$ Hz, 2 H), 2.63 (t, $J = 7.2$ Hz, 2 H), 3.78 (s, 3 H), 4.57 (d, $J = 6.3$ Hz, 2 H), 5.64 (m, 1 H), 5.84 (m, 1 H), 6.87 (d, $J = 8.1$ Hz, 1 H), 6.91-6.85 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 155.6, 151.1, 136.9, 131.6, 130.8, 130.0, 127.3, 123.4, 118.1, 68.6, 54.6, 32.7, 30.1, 25.7, 20.5, 18.1, 14.2, -4.2.

General Procedure for the TBS Deprotection: Preparation of (*E*)-5-(2-Hydroxy-5-methylphenyl)-pent-2-enyl Methyl Carbonate (15h**).** To a stirred solution of the previous carbonate (128 mg) in THF (2 mL) was added TBAF (1 M in THF, 0.7 mL, 0.7 mmol) at 0 °C. After being stirred at room temperature for 1 h, the mixture was treated with ether (15 mL) and water (5 mL). The layers were separated and the organic phase was washed with brine (5 mL), dried (MgSO_4), filtered, and concentrated under reduced pressure to afford compound **15h** as a clear oil (80.6 mg, 93%). IR (neat) 3638, 1511, 1444 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.24 (s, 3 H), 2.33 (t, $J = 7.2$ Hz, 2 H), 2.66 (t, $J = 7.2$ Hz, 2 H), 3.78 (s, 3 H), 4.57 (d, $J = 6.3$ Hz, 2 H), 5.21 (s, br, 1 H), 5.60 (dt, $J = 15.3, 6.3$ Hz, 1 H), 5.86 (dt, $J = 15.3, 6.3$ Hz, 1 H), 6.64 (d, $J = 7.8$ Hz, 1 H), 6.89-6.85 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 155.8, 151.2, 136.7, 130.8, 129.7, 127.4, 123.5, 115.1, 68.6, 54.7, 32.5, 29.4, 20.4. HRMS: Calc'd for $\text{C}_{14}\text{H}_{18}\text{O}_4$: 250.1245. Found: 250.1254.

General Procedure for the Preparation of Allyl Phenol: Preparation of 2-(*tert*-Butyldimethylsilyloxy)-3-allyl-5-methylanisole. To a stirred solution of 2-methoxy-4-methylphenol (1.658 g, 11.87 mmol) and allyl bromide (1.723 g, 14.20 mmol) in acetone (20 mL) was added potassium carbonate (2.867 g, 20.77 mmol). The mixture was refluxed for 12 h before it was treated with ether (80 mL) and water (20 mL). After separation of the layers, the organic phase was washed with water (15 mL), brine (10 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure. The crude material was purified by column chromatography (2% ether in petroleum ether) to afford the allyl ether as a clear oil (2.125 g, 100%).

The mixture of the above allyl ether (2.125 g, 11.87 mmol) and *N,N*-dimethylaniline (2.5 mL) was heated at 190-200 °C for 6 h. After it was cooled to room temperature, ether (50 mL) and 1 N HCl (10 mL) were added. With separation of the layers, the organic phase was further washed with 1 N HCl (2 × 10 mL), brine (10 mL), dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude material was purified by column chromatography (4% EtOAc in petroleum ether) to afford 2-allyl-4-methyl-6-methoxyphenol as a clear oil (1.6747 g, 79%).

To a stirred solution of the above phenol (1.6747 g, 9.41 mmol) in DMF (5 mL) was added TBSCl (1.702 g, 12.90 mmol), imidazole (1.536 g, 22.58 mmol) at room temperature. The mixture was stirred overnight and partitioned between ether (40 mL) and water (20 mL). The organic layer was further washed with water (2 × 20 mL), brine (7 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure. The crude material was purified by column chromatography (3% ether in petroleum ether) to afford the title compound (2.534 g, 92%) as a clear syrup. IR (neat) 1589, 1492, 1466, 1333, 1251 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.15 (s, 6 H), 0.98 (s, 9 H), 2.23 (s, 3 H), 3.95 (d, *J* = 6.3 Hz, 2 H), 3.72 (s, 3 H), 4.99 (m, 1 H), 5.04 (m, 1 H), 5.92 (m, 1 H), 6.50 (s, 1 H), 6.52 (s, 1 H), ¹³C NMR (75 MHz, CDCl₃) δ 149.6, 140.2, 137.2, 130.8, 130.1, 122.2, 115.4, 110.3, 54.7, 34.4, 26.1, 21.1, -3.9.

General Procedure for the Hydroboration Reaction: Preparation of 3-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-propan-1-ol. To a stirred solution of 2-(*tert*-Butyldimethylsilyloxy)-3-allyl-5-methylanisole (840.5 mg, 3.02 mmol) in THF (5 mL) was added $\text{BH}_3 \cdot \text{THF}$ (1 M in THF, 3.0 mL, 3.0 mmol) at 0 $^\circ\text{C}$. After being stirred at room temperature for 1.5 h, a NaOH solution (1 M, 2.6 mL) was added dropwise at 0 $^\circ\text{C}$. The mixture was stirred at 0 $^\circ\text{C}$ for 30 min followed by the addition of H_2O_2 (30% solution, 1.3 mL). After 2 h at 0 $^\circ\text{C}$, the mixture was treated with ether (50 mL) and water (10 mL). After separation of the layers, the aqueous phase was extracted with ether (15 mL). The combined organic layers were washed with brine (5 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure. The residue was purified by column chromatography (petroleum ether/EtOAc, 3:1) to afford the title compound as a clear oil (640.3 mg, 72%). IR (neat) 3358, 1498, 1472, 1255, 1225 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.19 (s, 6 H), 0.99 (s, 9 H), 1.82 (m, 2 H), 1.88 (m, 1 H), 2.64 (t, J = 7.5 Hz, 2 H), 3.60 (m, 2 H), 3.73 (s, 3 H), 6.71-6.57 (m, 3 H); ^{13}C NMR (75 MHz, CDCl_3) δ 153.8, 147.3, 133.0, 119.1, 115.7, 111.4, 62.0, 55.5, 32.9, 26.6, 25.8, 18.2, -4.3.

3-(2-*tert*-Butyldimethylsilyloxy-3-methoxy-5-methylphenyl)-propan-1-ol. A clear oil (58%). IR (neat) 3355, 1588, 1493, 1249 1156 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.18 (s, 6 H), 1.00 (s, 9 H), 1.83 (t, J = 7.5 Hz, 2 H), 2.03 (m, 1 H), 2.27 (s, 3 H), 2.69 (t, J = 7.5 Hz, 2 H), 3.59 (t, J = 6.3 Hz, 2 H), 3.74 (s, 3 H), 6.52 (m, 1 H), 6.56 (m, 1 H), ^{13}C NMR (75 MHz, CDCl_3) δ 149.5, 140.3, 132.4, 130.3, 122.3, 110.1, 62.0, 54.6, 33.0, 26.2, 26.1, 21.0, 18.8, -4.0.

General Procedure for the Preparation of the Aldehyde by Swern Oxidation:

Preparation of 3-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-propan-1-al. To a stirred solution of (COCl)₂ (554.0 mg, 4.36 mmol) in CH₂Cl₂ (15 mL) was added a solution of DMSO (795.7 mg, 10.19 mmol) in CH₂Cl₂ (2 mL) at -78 °C. After being stirred for 5 min, a solution of 3-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-propan-1-ol (861.9 mg, 2.91 mmol) in CH₂Cl₂ (8 mL) was added dropwise. The mixture was stirred at -78 °C for 15 min before TEA (1.23 g, 12.22 mmol) was added slowly. The suspension was stirred at this temperature for 10 min and was allowed to warm up to room temperature. The reaction was quenched with addition of water (10 mL) and ether (60 mL). After separation of the layers, the organic phase was washed with water (10 mL), brine (5 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure to afford the crude material. The title compound was obtained after column chromatography (petroleum ether/EtOAc, 100:6) as a clear oil (838.9 mg, 98%). IR (neat) 1726, 1499, 1256, 1226 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.19 (s, 6 H), 0.97 (s, 9 H), 2.71 (t, *J* = 7.2 Hz, 2 H), 2.86 (t, *J* = 7.5 Hz, 2 H), 3.72 (s, 3 H), 6.70-6.60 (m, 3 H), 9.78 (s, 1 H); ¹³C NMR (75 MHz, CDCl₃) δ 201.9, 153.7, 147.3, 131.6, 118.9, 115.7, 111.9, 55.5, 43.9, 25.7, 23.8, 18.1, -4.3.

3-(2-*tert*-Butyldimethylsilyloxy-3-methoxy-5-methylphenyl)-propan-1-al. A clear oil (98%). IR (neat) 1727, 1493, 1329, 1249 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.17 (s, 6 H), 0.98 (s, 9 H), 2.25 (s, 3 H), 2.71 (t, *J* = 7.5 Hz, 2 H), 2.91 (t, *J* = 7.5 Hz, 2 H), 3.75 (s, 3 H), 6.54 (m, 2 H), 9.78 (s, 1 H), ¹³C NMR (75 MHz, CDCl₃) δ 202.1, 149.5, 140.3, 130.9, 130.3, 122.0, 110.5, 77.2, 54.6, 44.0, 26.0, 23.4, 21.0, 18.8, -4.0.

General Procedure for the Wittig Reaction: Preparation of (*E*)-Ethyl 5-(2-*tert*-butyldimethylsilyloxy-5-methoxyphenyl)-pent-2-enoate. To a stirred solution of 3-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-propan-1-al (200.0 mg, 0.68 mmol) in benzene (5 mL)

was added carboethoxymethylenetriphenylphosphorane (237.0 mg, 1.315 mg) at room temperature. The mixture was stirred overnight and the volatiles were removed under reduced pressure. The residue was purified by column chromatography (petroleum ether/EtOAc, 25:1) to afford the title compound as a clear oil (232.7 mg, 94%). IR (neat) 1722, 1655, 1499, 1472, 1269, 1226 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.19 (s, 6 H), 0.98 (s, 9 H), 1.26 (t, $J = 7.2$ Hz, 3 H), 2.47 (m, 2 H), 2.69 (t, $J = 7.2$ Hz, 2 H), 3.72 (s, 3 H), 4.16 (q, $J = 7.2$ Hz, 2 H), 5.82 (d, $J = 15.9$ Hz, 1 H), 6.70-6.59 (m, 3 H), 7.00 (dt, $J = 15.6, 7.2$ Hz, 1 H); ^{13}C NMR (75 MHz, CDCl_3) δ 166.5, 153.6, 148.4, 147.2, 132.1, 121.5, 118.8, 115.6, 111.7, 60.0, 32.5, 29.4, 25.7, 18.2, 14.2, -4.2; EIHRMS m/z calcd for $\text{C}_{20}\text{H}_{32}\text{O}_4\text{Si}[\text{M}]^+$: 364.2069, found: 364.2069.

(E)-Ethyl 5-(2-*tert*-butyldimethylsilyloxy-3-methoxy-5-methylphenyl)-pent-2-enoate.

A clear oil (82%). IR (neat) 1722, 1493, 1250, 911 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.16 (s, 6 H), 0.97 (s, 9 H), 1.27 (t, $J = 7.5$ Hz, 3 H), 2.25 (s, 3 H), 2.48 (m, 2 H), 2.72 (t, $J = 7.5$ Hz, 2 H), 3.75 (s, 3 H), 4.18 (q, $J = 7.5$ Hz, 2 H), 5.83 (dd, $J = 15.3, 1.0$ Hz, 1 H), 6.53 (s, 1 H), 7.01 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 166.6, 149.5, 148.7, 140.3, 131.6, 130.1, 122.1, 121.3, 110.4, 77.2, 60.0, 54.6, 32.7, 29.1, 26.1, 21.0, 18.8, 14.2, -3.9.

(E)-5-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-pent-2-en-1-yl Methyl Carbonate. Employing the general procedure for the DIBAL reduction, compound, (*E*)-5-(2-*tert*-butyldimethylsilyloxy-5-methoxyphenyl)-pent-2-en-1-ol was prepared as a clear oil in 76% yield. IR (neat) 3355, 1499, 1471, 1255, 1224 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.18 (s, 6 H), 0.99 (s, 9 H), 2.31 (m, 2 H), 2.62 (t, $J = 7.5$ Hz, 2 H), 3.73 (s, 3 H), 4.05 (m, 2 H), 5.67 (m, 2 H), 6.70-6.57 (m, 3 H); ^{13}C NMR (75 MHz, CDCl_3) δ 153.6, 147.3, 133.1, 132.5, 129.3, 118.1, 115.8, 111.3, 63.6, 55.5, 32.5, 30.5, 25.8, 18.1, -4.3.

Employing the general procedure for the preparation of the carbonate, the title compound was prepared in 91 % yield as a clear oil. IR (neat) 1750, 1599, 1268, 1225 cm^{-1} ; ^1H NMR (300

MHz, CDCl₃) δ 0.11 (s, 6 H), 0.92 (s, 9 H), 2.26 (t, J = 7.2 Hz, 2 H), 2.55 (t, J = 7.2 Hz, 2 H), 3.65 (s, 3 H), 3.68 (s, 3 H), 4.48 (d, J = 6.6 Hz, 2 H), 5.54 (ddd, J = 15.3, 6.6, 6.6 Hz, 1 H), 5.75 (ddd, J = 15.6, 6.6, 6.6 Hz, 1 H), 6.63-6.50 (m, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 155.6, 153.6, 147.2, 136.5, 132.8, 123.5, 118.1, 115.7, 111.4, 68.5, 55.4, 54.5, 32.5, 30.2, 25.7, 18.1, -4.3.

(*E*)-5-(2-*tert*-Butyldimethylsilyloxy-3-methoxy-5-methylphenyl)-pent-2-enyl Methyl Carbonate. A clear oil (92%). IR (neat) 1750, 1493, 1466, 1258, 1156 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.18 (s, 6 H), 1.00 (s, 9 H), 2.26 (s, 3 H), 2.35 (quintet, J = 7.2 Hz, 2 H), 2.67 (t, J = 7.5 Hz, 2 H), 3.75 (s, 3 H), 3.77 (s, 3 H), 4.57 (d, J = 6.3 Hz, 2 H), 5.65 (m, 1 H), 5.87 (m, 1 H), 6.53 (m, 2 H); ¹³C NMR (75 MHz, CDCl₃) δ 149.4, 140.2, 136.8, 136.6, 132.2, 123.4, 122.2, 110.1, 68.5, 54.5, 32.7, 29.9, 26.0, 21.0, 18.8, -4.0; HRMS. (*E*)-5-(2-*tert*-butyldimethylsilyloxy-3-methoxy-5-methylphenyl)-pent-2-en-1-ol. IR (neat) 3334, 1492, 1466, 1250, 922 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.16 (s, 6 H), 0.99 (s, 9 H), 2.25 (s, 3 H), 2.31 (t, J = 7.5 Hz, 2 H), 2.66 (t, J = 7.5 Hz, 2 H), 3.75 (s, 3 H), 4.05 (dd, J = 6.0, 1.0 Hz, 2 H), 5.64 (m, 1 H), 5.71 (m, 1 H), 6.51 (s, 1 H), 6.53 (s, 1 H); ¹³C NMR (75 MHz, CDCl₃) δ 149.5, 140.2, 132.7, 132.4, 129.9, 129.1, 122.2, 110.1, 63.6, 54.6, 32.8, 30.2, 26.1, 21.0, 18.8, -3.9; EIHRMS m/z calcd for C₁₉H₃₂O₃Si[M]⁺: 336.2121, found: 336.2125.

(*E*)-5-(2-Hydroxy-5-methoxyphenyl)-pent-2-enyl Methyl Carbonate (15i). Employing the general procedure for the TBS deprotection, the title compound was prepared in 93% yield from (*E*)-5-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-pent-2-enyl Methyl Carbonate as a clear oil. IR (neat) 3436, 1748, 1509, 1448, 1272, 1205 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.35 (t, J = 7.2 Hz, 2 H), 2.67 (t, J = 7.2 Hz, 2 H), 3.74 (s, 3 H), 3.77 (s, 3 H), 4.56 (d, J = 6.6 Hz, 2 H), 5.60 (dt, J = 15.6, 6.6 Hz, 1 H), 5.86 (dt, J = 15.3, 6.6 Hz, 1 H), 6.70-6.59 (m, 3 H); ¹³C NMR (75 MHz, CDCl₃) δ 155.8, 153.3, 147.6, 136.5, 128.9, 123.5, 115.9,

115.8, 111.8, 68.6, 55.7, 32.2, 29.6, 20.4; EIHRMS, m/z calcd for $C_{14}H_{18}O_5[MH+1]^+$: 268.1224, found: 268.1230.

(E)-5-(2-Hydroxy-3-methoxy-5-methylphenyl)-pent-2-enyl Methyl Carbonate (15j).

A clear oil (79%). IR (neat) 3521, 1747, 1499, 1447, 1268 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$) δ 2.25 (s, 3 H), 2.36 (quintet, $J = 7.5$ Hz, 2 H), 2.67 (t, $J = 7.5$ Hz, 2 H), 3.76 (s, 3 H), 3.82 (s, 3 H), 4.57 (dd, $J = 6.0, 1.0$ Hz, 2 H), 5.63 (m, 2 H), 5.86 (m, 1 H), 6.53 (m, 2 H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 155.6, 146.0, 141.0, 136.8, 128.4, 126.8, 123.3, 122.3, 109.3, 68.5, 55.7, 32.3, 29.1, 25.5, 20.9; EIHRMS m/z calcd for $C_{15}H_{20}O_5[M]^+$: 280.1311, found: 280.1310.

3-(2-*tert*-Butyldimethylsilyloxy-5-methylphenyl)-propanol. Employing the general procedure for hydroboration reaction, the title compound was prepared from 2-allyl-4-methyl-*tert*-butyldimethylsilylphenol in 80% yield as a clear oil. IR (neat) 3332, 1500, 1472, 1265, 1227, 1044 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 0.24 (s, 6 H), 1.03 (s, 9 H), 1.85 (quintet, $J = 7.5$ Hz, 2 H), 2.25 (s, 3 H), 2.66 (t, $J = 7.2$ Hz, 2 H), 3.62 (t, $J = 6.3$ Hz, 2 H), 6.70 (d, $J = 8.1$ Hz, 1 H), 6.88 (m, 1 H), 6.96 (m, 1 H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 151.1, 131.8, 130.9, 130.3, 127.2, 118.3, 62.0, 33.0, 26.4, 25.7, 20.4, 18.1, -4.3.

General Procedure for the Alkyne Formation: Preparation of 4-(2-*tert*-Butyldimethylsilyloxyphenyl)-1-butyne. Employing the general procedure for the Swern oxidation, 3-(2-*tert*-Butyldimethylsilyloxyphenyl)-propanal was prepared from 3-(2-*tert*-butyldimethylsilyloxyphenyl)-propanol in 76% yield as a clear oil. IR (neat) 1727, 1492, 1254, 924, 839 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 0.25 (s, 6 H), 1.00 (s, 9 H), 2.72 (t, $J = 7.5$ Hz, 2

H), 2.92 (t, $J = 7.2$ Hz, 2 H), 6.90-6.78 (m, 2 H), 7.24-7.06 (m, 2 H), 9.80 (s, 1 H); ^{13}C NMR (75 MHz, CDCl_3) δ 202.0, 153.5, 130.7, 130.1, 127.3, 121.1, 118.3, 43.9, 25.7, 23.5, 18.1, -4.2.

To a stirred solution of the above aldehyde (578.0 mg, 2.20 mmol) in MeOH (22 mL) was added Bestman reagent (506.2 mg, 2.634 mmol) and K_2CO_3 (607.2 mg, 4.40 mmol) at room temperature. After being stirred overnight, the reaction mixture was treated with ether (60 mL) and water (10 mL). The aqueous layer was further extracted with ether (2×15 mL). The combined organic phase was washed with brine (10 mL), dried (Na_2SO_4), filtered, and concentrated under reduced pressure to afford a yellow residue. The crude material was purified by column chromatography (petroleum ether/EtOAc, 10:1) to generate 4-(2-hydroxyphenyl)-1-butyne (238.7 mg, 74%) as a clear oil.

To a stirred solution of the above compound in DMF (3 mL) was added TBSCl (295.7 mg, 1.96 mmol) and imidazole (266.8 mg, 3.92 mmol) at room temperature. The mixture was stirred for 16 h and was treated with ether (30 mL) and water (20 mL). After separation of the layers, the organic phase was washed with water (2×15 mL), brine (5 mL), dried (MgSO_4), filtered, and concentrated under reduced pressure to afford the crude material which was purified by column chromatography (petroleum ether/EtOAc, 100:2) to yield the title compound as a clear oil (376.8 mg, 89%). IR (neat) 3313, 1601, 1492, 1255, 925 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.22 (s, 6 H), 1.00 (s, 9 H), 1.92 (d, $J = 2.7$ Hz, 1 H), 2.43 (dt, $J = 7.8, 3.0$ Hz, 2 H), 2.81 (t, $J = 7.8$ Hz, 2 H), 6.88-6.75 (m, 2 H), 7.16-7.04 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 153.6, 130.9, 130.3, 127.4, 120.9, 118.3, 84.2, 68.5, 30.2, 25.8, 18.9, -4.2.

4-(2-*tert*-Butyldimethylsilyloxy-5-methylphenyl)-1-butyne. A clear oil (61%). (IR (neat) 3313, 1501, 1472, 1265, 1228, 1120 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.20 (s, 6 H), 1.00 (s, 9 H), 1.92 (d, $J = 2.7$ Hz, 1 H), 2.23 (s, 3 H), 2.40 (dt, $J = 7.5, 2.7$ Hz, 2 H), 2.77 (t, $J = 7.5$ Hz, 2 H), 6.65 (d, $J = 8.1$ Hz, 1 H), 6.87 (m, 1 H), 6.95 (m, 1 H); ^{13}C NMR (75 MHz, CDCl_3) δ 151.2, 130.9, 130.6, 130.0, 127.8, 118.0, 84.3, 68.3, 30.2, 25.8, 20.5, 18.2, -4.2.

4-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-1-butyne. A clear oil (54%). (IR (neat) 3311, 1500, 1272, 1227 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 0.21 (s, 6 H), 1.01 (s, 9 H), 1.96 (t, $J = 1.5$ Hz, 1 H), 2.45 (dt, $J = 7.5, 3.0$ Hz, 2 H), 2.80 (t, $J = 7.5$ Hz, 2 H), 3.75 (s, 3 H), 6.65 (m, 1 H), 6.70 (m, 1 H), 6.76 (m, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 153.6, 147.3, 131.7, 118.7, 115.9, 111.9, 84.1, 68.5, 55.5, 30.3, 25.8, 18.9, 18.1, -4.2.

General Procedure for the Preparation of (*Z*)-Allylic Alcohol: Preparation of (*Z*)-5-(2-*tert*-Butyldimethylsilyloxyphenyl)-pent-2-en-1-ol. To a stirred solution of 4-(2-*tert*-Butyldimethylsilyloxyphenyl)-1-butyne (376.2 mg, 1.45 mmol) in THF (7 mL) was added *n*-BuLi (1.6 M in hexanes, 1.8 mL, 2.89 mmol) dropwise at -78°C . After being stirred at this temperature for 1 h, paraformaldehyde (86.7 mg, 2.89 mmol) was added in one portion. The mixture was allowed to warm up to room temperature and stirred for 1 h. The reaction was quenched with the addition of saturated NH_4Cl solution (8 mL) and ether (30 mL). After separation of the layers, the organic phase was washed with water (2 $\leftrightarrow$ 5 mL), brine (5 mL), dried (MgSO_4), filtered, and concentrated under reduced pressure to afford the crude material which was purified by column chromatography (petroleum ether/EtOAc, 10:1) to yield 5-(2-*tert*-butyldimethylsilyloxyphenyl)-pent-2-yn-1-ol as a clear oil (212.7 mg, 51%). IR (neat) 3347, 1493, 1254, 923 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.23 (s, 6 H), 1.01 (s, 9 H), 2.47 (t, $J = 7.5$ Hz, 2 H), 2.80 (t, $J = 7.8$ Hz, 2 H), 3.61 (t, $J = 6.6$ Hz, 1 H), 4.20 (s, br, 2 H), 6.90-6.76 (m, 2 H), 7.17-7.06 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 153.5, 131.0, 130.3, 127.4, 120.9, 118.3, 85.9, 78.4, 51.2, 30.1, 25.7, 19.2, -4.2; EIHRMS m/z calcd for $\text{C}_{17}\text{H}_{26}\text{O}_2\text{Si}[\text{M}]^+$: 294.1858, found: 294.1856.

To a degassed solution of the above compound (104.0 mg, 0.329 mmol) in MeOH (3 mL) was added Lindlar's catalyst (40 mg). The mixture was stirred under 1 atm H_2 for 4 h and was filtered through a pad of celite[®]. The filtrate was concentrated under reduced pressure and was purified by column chromatography (petroleum ether/EtOAc, 10:1) to afford compound **31** as a

clear syrup (92.1 mg, 88%). IR (neat) 3428, 1491, 1253, 924 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.15 (s, 6 H), 0.93 (s, 9 H), 2.28 (t, $J = 6.6$ Hz, 2 H), 2.57 (t, $J = 6.9$ Hz, 2 H), 3.90 (t, $J = 4.8$ Hz, 1 H), 5.50 (m, 2 H), 6.78-6.69 (m, 2 H), 7.01-6.99 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 153.8, 131.9 (2), 130.6, 129.1, 127.0, 120.8, 118.3, 60.4, 58.3, 30.4, 27.6, 25.7, 18.2, -4.2.

(Z)-5-(2-*tert*-Butyldimethylsilyloxy-5-methylphenyl)-pent-2-en-1-ol (33). A clear oil. IR (neat) 3356, 1654, 1501, 1265 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.12 (s, 6 H), 0.91 (s, 9 H), 2.14 (s, 3 H), 2.26 (quintet, $J = 7.5$ Hz, 2 H), 2.51 (t, $J = 6.9$ Hz, 2 H), 3.92 (t, $J = 6.9$ Hz, 1 H), 5.47 (m, 2 H), 6.59 (d, $J = 8.1$ Hz, 1 H), 6.78 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 151.2, 132.0, 131.6, 131.1, 129.9, 129.0, 127.4, 118.1, 58.3, 30.5, 27.7, 25.7, 20.5, 18.2, -4.2.

(Z)-5-(2-*tert*-Butyldimethylsilyloxy-5-methoxyphenyl)-pent-2-en-1-ol. A clear oil. IR (neat) 3384, 1502, 1232 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.13 (s, 6 H), 0.93 (s, 9 H), 2.31 (quintet, $J = 7.5$ Hz, 2 H), 2.52 (t, $J = 6.9$ Hz, 2 H), 3.74 (s, 3 H), 5.47 (m, 2 H), 6.61 (m, 1 H), 6.68 (m, 1 H), 6.74 (m, 1 H); ^{13}C NMR (75 MHz, CDCl_3) δ 153.4, 147.3, 131.8, 129.8, 127.6, 118.1, 116.0, 111.8, 57.3, 51.2, 30.2, 25.7, 19.2, 18.1, -4.3.

(Z)-5-(2-Hydroxyphenyl)-pent-2-en-1-yl Methyl Carbonate (15b'). Employing the general procedure for the preparation of the carbonate, (Z)-5-(2-*tert*-butyldimethylsilyloxyphenyl)-pent-2-en-1-yl methyl carbonate was obtained as a clear oil in 91% yield. IR (neat) 1751, 1491, 1292 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.28 (s, 6 H), 1.06 (s, 9 H), 2.45 (t, $J = 7.5$ Hz, 2 H), 2.73 (t, $J = 7.5$ Hz, 2 H), 3.82 (s, 3 H), 4.62 (t, $J = 3.5$ Hz, 2 H), 5.61 (m, 1 H), 5.75 (m, 1 H), 6.92-6.82 (m, 2 H), 7.11-7.09 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 155.7, 153.5, 134.9, 131.7, 130.3, 127.0, 123.2, 120.9, 118.3, 63.6, 54.6, 30.3, 27.9, 25.7, 18.1, -4.2.

Employing the general TBS deprotection procedure, the title compound was obtained as a clear oil in 93% yield. IR (neat) 3438, 1748, 1726, 1456, 1284 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.44 (t, $J = 8.5$ Hz, 2 H), 2.67 (m, 2 H), 3.77 (s, 3 H), 4.61 (dd, $J = 7.5, 1.0$ Hz, 2 H), 5.57 (s, br, 1 H), 5.75 (dt, $J = 15.3, 6.3$ Hz, 1 H), 5.81 (s, 1 H), 6.85-6.77 (m, 2 H), 7.08-7.05 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 155.9, 153.8, 135.3, 130.4, 127.4, 127.2, 123.2, 120.5, 115.4, 63.5, 54.8, 30.2, 27.9; EIHRMS m/z calcd for $\text{C}_{13}\text{H}_{16}\text{O}_4[\text{MH}+1]^+$: 238.1067, found: 238.1056.

Z-5-(5-Fluoro-2-hydroxyphenyl)-pent-2-enyl methyl carbonate (15g'): To phenol silyl ether **40** (0.60 g, 1.63 mmol) in 5 mL of THF was added TBAF (3.26 mL, 3.26 mmol, 1M in THF) at rt. The reaction mixture was stirred at rt for 2 h and purified by silica gel chromatography eluting with 5% to 20% diethyl ether in petroleum ether to afford phenol **15g** (328 mg, 1.29 mmol, 79%) as a pale yellow oil.

IR (neat): 3450b, 3022w, 2960s, 2868m, 1734s, 1619w, 1508s, 1438s, 1365m, 1264s, 1184s, 1143m, 1103m, 944m, 872w, 813w, 736w, 713w, 668m cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3): δ 6.76 (m, 3H), 5.73 (m, 1H), 5.59 (m, 1H), 4.63 (d, $J=7.2$ Hz, 2H), 3.78 (s, 3H), 2.65 (dd, $J=5.7, 9.6$ Hz, 2H), 2.45 (q, $J=7.2$ Hz, 2H); ^{13}C -NMR (75 MHz, CDCl_3): δ 158.3, 155.9, 155.2, 149.9, 134.9, 128.8 (d, $J=7.3$ Hz), 123.4, 116.4 (d, $J=23.0$ Hz), 116.1 (d, $J=7.4$ Hz), 113.3 (d, $J=23.0$ Hz), 63.5, 54.9, 30.3, 27.6, 27.3. HRMS: Calc'd for $\text{C}_{13}\text{H}_{15}\text{O}_4\text{F}$: 254.0954. Found: 254.0951.

(Z)-5-(2-Hydroxy-5-methylphenyl)-pent-2-enyl Methyl Carbonate (15h'). A clear oil (79%). IR (neat) 3444, 1749, 1727, 1511, 1264 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.33 (s, 3 H), 2.40 (quintet, $J = 7.5$ Hz, 2 H), 2.62 (t, $J = 7.2$ Hz, 2 H), 3.76 (s, 3 H), 4.62 (dd, $J = 6.9, 1.0$ Hz, 2 H), 5.55 (m, 1 H), 5.76 (m, 1 H), 5.84 (s, 1 H), 6.65 (d, $J = 8.1$ Hz, 1 H), 6.86 (m, 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ 155.8, 151.6, 135.3, 130.8, 129.4, 127.7, 127.0, 123.1, 115.2, 63.5, 60.5, 54.7, 30.2, 28.0, 20.4.

(Z)-5-(2-tert-Butyldimethylsilyloxyphenyl)-pent-2-enyl Methyl Carbonate, IR (neat) 1751, 1501, 1443, 1266 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.22 (s, 6 H), 1.00 (s, 9 H), 2.25 (s, 3 H), 2.37 (quintet, $J = 7.8$ Hz, 2 H), 2.62 (t, $J = 7.5$ Hz, 2 H), 3.76 (s, 3 H), 4.59 (t, $J = 6.9$ Hz, 2 H), 5.54 (m, 1 H), 5.69 (m, 1 H), 6.66 (m, 1 H), 6.88 (m 2 H); ^{13}C NMR (75 MHz, CDCl_3) δ

155.7, 151.2, 135.1, 131.4, 130.9, 127.4, 123.1, 118.1, 63.6, 54.6, 30.3, 28.5, 25.7, 20.5, 18.2, -
 4.2. EIMS m/z calcd for $C_{14}H_{18}O_4[MH]^+$: 251, found: 251.

(Z)-5-(2-Hydroxy-5-methoxyphenyl)-pent-2-enyl Methyl Carbonate (15i'). A clear oil (88%). IR (neat) 3541, 1751, 1519 cm^{-1} ; 1H NMR (300 MHz, $CDCl_3$) δ 2.39 (s, 3 H), 2.61 (quintet, $J = 7.5$ Hz, 2 H), 2.61 (t, $J = 7.2$ Hz, 2 H), 3.74 (s, 3 H), 3.77 (s, 3 H), 4.62 (d, $J = 6.9$ Hz, 2 H), 5.54 (m, 1 H), 5.76 (m, 1 H), 6.72-6.61 (m, 3 H); ^{13}C NMR (75 MHz, $CDCl_3$) δ 155.8, 153.2, 147.6, 136.1, 129.3, 127.4, 115.9, 111.7, 64.2, 55.6, 32.0, 28.0, 20.4; EIHRMS m/z calcd for $C_{14}H_{18}O_5[MH]^+$: 267.1236, found: 268.1240.

(R)-5-Methoxy-2,7-dimethylchroman-2-carboxylic acid (42): To a solution of (*R*)-5-methoxy-2,7-dimethyl-2-vinylchroman **41** (0.65 g, 2.97 mmol) in 10 mL of dichloromethane was added *N*-methylmorpholine-*N*-oxide (0.9 g, 7.68 mmol) and aqueous osmium tetroxide (0.8 mL, 4% in water, 0.13 mmol). The solution was stirred for 14 h. at rt and diluted with water (30 mL) and dichloromethane (50 mL). The organic layer was dried over magnesium sulfate and evaporated to dryness. The residue was chromatographed with 1:1 to 1:5 petroleum ether in diethyl ether. The resulting brown oil (contaminated with osmium residue) was resuspended in acetone (12 mL) and a solution of sodium periodate (1.2 g, 5.61 mmol) in water (3 mL) was added. After a white precipitate was formed the reaction mixture was stirred at rt for an additional 20 min. The residue was partitioned between water (10 mL) and dichloromethane (20 mL). The aqueous layer was extracted with dichloromethane (2 x 20 mL). The combined organic layers were washed with brine (20 mL) and dried over magnesium sulfate. After evaporating the solvent, the residue was diluted with *t*-butanol (30 mL). Sodium dihydrophosphate (1.5 g) and 2-methyl-2-butene (16 mL) were successively added, the solution was cooled to 0°C. A freshly prepared solution of sodium hypochlorite ($NaClO_2$, 20% in water, 14 mL) was added to the mixture. The resulting solution was stirred for 2 h at 0°C. Water (20 mL) was added and the aqueous layer was extracted with ethyl acetate (50 mLx3). The combined organic layers were washed with

brine and dried over magnesium sulfate. After concentrated *in vacuo*, the residue was chromatographed with acetic acid in ethyl acetate (1/100) to afford 5-methoxy-2,7-dimethylchroman-2-carboxylic acid **42** (0.628 g, 2.65 mmol, 89%) as a sticky oil.

$[\alpha]_D^{25} +67.8^\circ$ (*c* 0.94, CHCl₃); IR (neat): 3500-3300b, 1716s, 1621s, 1588s, 1463s, 1143s cm⁻¹; ¹H-NMR (500 MHz, CDCl₃): δ 9.8-7.9 (br, 1H), 6.42 (s, 1H), 6.29 (s, 1H), 3.81 (s, 3H), 2.73 (m, 1H), 2.49 (m, 1H), 2.36 (m, 1H), 2.30 (s, 3H), 1.89 (m, 1H), 1.64 (s, 3H); ¹³C-NMR (125 MHz, CDCl₃): δ 179.9, 157.3, 153.5, 137.4, 109.7, 106.6, 103.5, 77.2, 55.3, 29.6, 24.7, 21.5, 16.6. HRMS: Calc'd for C₁₃H₁₆O₄ [M⁺]: 236.1048. Found: 236.1050.

(S)-[2-(R)-(5-Methoxy-2,7-dimethyl-chroman-2-carbonyl)-amino]-phenylacetic acid methyl ester (43). To a mixture of (*S*)-PGME·HCl (0.236 g, 1.17 mmol), PyBOP (0.610 g, 1.17 mmol), and HOBT (0.158 g, 1.17 mmol) was added at 0°C a solution of 5-methoxy-2,7-dimethyl-chroman-2-carboxylic acid **42** (0.22 g, 0.93 mmol) in DMF (4 mL) and *N*-methylmorpholine (0.647 mL, 5.86 mmol). The reaction mixture was stirred at rt for 5 h. To the reaction was then added ethyl acetate (50 mL) and the resulting mixture was successively washed with 1N HCl (25 mL), saturated sodium bicarbonate (25 mL) and brine (25 mL). The organic layer was dried over magnesium sulfate and concentrated *in vacuo*. The residue was chromatographed eluting with 3/1 petroleum ether/diethyl ether to afford (*S*)-[2-(*R*)-(5-methoxy-2,7-dimethyl-chroman-2-carbonyl)-amino]-phenyl-acetic acid methyl ester **43** (0.225 g, 0.583 mmol) as a sticky oil.

$[\alpha]_D^{25} +67.4^\circ$ (*c* 1.44, CHCl₃); IR (neat): 3420, 2953, 1746, 1681, 1503, 1224, 1112 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃): δ 7.48 (d, *J*=6.8 Hz, 1H), 7.40-7.32 (m, 5H), 6.44 (s, 1H), 6.32 (s, 1H), 5.46 (d, *J*=6.8 Hz, 1H), 3.81 (s, 3H), 3.65 (s, 3H), 2.66 (td, *J*=6.6, 17.6 Hz, 1H), 2.59 (td, *J*=6.6, 17.6 Hz, 1H), 2.32 (s, 3H), 2.29 (td, *J*=6.6, 13.2 Hz, 1H), 1.93 (ddd, *J*=6.1, 7.3, 13.6 Hz, 1H), 1.48 (s, 3H); ¹³C-NMR (125 MHz, CDCl₃): δ 174.3, 170.5, 157.5, 152.7, 137.2, 136.1, 128.9, 128.4, 127.0, 109.7, 107.2, 103.7, 78.6, 56.3, 55.3, 52.5, 28.7, 23.5, 21.5, 16.1. HRMS: Calc'd for C₂₂H₂₅O₅N [M⁺]: 383.1732. Found: 383.1730.

(R)-[2-(R)-(5-Methoxy-2,7-dimethyl-chroman-2-carbonyl)-amino]-phenylacetic acid methyl ester (44): The exactly same procedure was followed as described in the synthesis of **42** except that (*R*)-PGME·HCl was used instead of (*S*)-PGME·HCl. (*R*)-[2-(*R*)-(5-methoxy-2,7-dimethyl-chroman-2-carbonyl)-amino]-phenyl-acetic acid methyl ester **44** (0.194 g, 0.50 mmol, 53%) was obtained as a sticky oil.

$[\alpha]_D^{25} = -9.5^\circ$ (*c* 1.24, CHCl₃); IR (neat): 3421, 2934, 1743, 1683, 1500, 1224, 1112 cm⁻¹; ¹H-NMR (500 MHz, CDCl₃): δ 7.56 (d, *J*=7.3 Hz, 1H), 7.19-7.26 (m, 3H), 7.05 (d, *J*=7.3 Hz, 2H), 6.50 (s, 1H), 6.31 (s, 1H), 5.53 (d, *J*=7.3 Hz, 1H), 3.79 (s, 3H), 3.72 (s, 3H), 2.57 (td, *J*=5.5, 17.4 Hz, 1H), 2.24-2.36 (m, 2H), 2.33 (s, 3H), 1.84 (ddd, *J*=5.3, 8.3, 13.1 Hz, 1H), 1.57 (s, 3H); ¹³C-NMR (125 MHz, CDCl₃): δ 173.2, 170.8, 157.6, 153.0, 137.5, 136.3, 128.6, 128.1, 126.6, 109.7, 107.8, 103.7, 78.8, 55.8, 55.3, 52.7, 29.0, 24.0, 21.5, 16.2. HRMS: Calc'd for C₂₂H₂₅O₅N [M⁺]: 383.1732. Found: 383.1740.

(R)-2,7-Dimethyl-2-vinyl-chroman-5-ol (65): To a flame-dried flask was added sodium hydride (119 mg, 2.98 mmol, 60% in mineral oil). This flask was then cooled in the ice bath and to it was added DMF (anhydrous, 1 mL). To this slurry at 0°C was added ethanethiol (171 mg, 0.20 mL, 2.75 mmol). After the effervescence, the mixture was warmed to rt and stirred for 30 min followed by the addition of chroman **17c** (50 mg, 0.23 mmol) in 1.5 mL of DMF. The solution was heated at 120°C for 24 h. Without workup, the mixture was directly chromatographed eluting with 5% to 50% diethyl ether in petroleum ether to afford phenol **65** (45 mg, 0.22 mmol, 94%) as a yellow oil.

$[\alpha]_D^{25} = +56.7^\circ$ (*c* 4.5, Et₂O); IR (film): 3418b, 2977m, 2929s, 2854w, 1629s, 1586s, 1515m, 1454s, 1417s, 1373w, 1352m, 1318m, 1255m, 1211w, 1138s, 1118s, 1099m, 1053s, 996s, 924w, 889w, 821m cm⁻¹; ¹H-NMR (500 MHz, CDCl₃): δ 6.34 (s, 1H), 6.17 (s, 1H), 5.86 (ddd, *J*=1.0, 10.5,

17.0 Hz, 1H), 5.18 (dd, $J=1.0$, 17.5 Hz, 1H), 5.06 (dd, $J=1.5$, 11.0 Hz, 1H), 4.90 (s, 1H), 2.64 (dt, $J=5.0$, 16.5 Hz, 1H), 2.48 (ddd, $J=6.5$, 10.0, 16.5 Hz, 1H), 2.22 (s, 3H), 1.94 (dt, $J=5.0$, 13.5 Hz, 1H), 1.80 (ddd, $J=6.0$, 10.0, 16.0 Hz, 1H), 1.42 (s, 3H); ^{13}C -NMR (125 MHz, CDCl_3): δ 154.7, 153.5, 141.1, 137.3, 113.8, 110.0, 107.3, 105.8, 76.3, 31.2, 26.8, 21.1, 16.6. Anal. Calc'd for $\text{C}_{13}\text{H}_{16}\text{O}_2$: C, 76.44; H, 7.90. Found: C, 76.39; H, 7.93.

(R)-5-Methoxymethoxy-2,7-dimethyl-2-vinylchroman (66): To phenol **65** (20 mg, 0.098 mmol) in 1 mL of THF was added *n*-butyllithium (74 μL , 0.12 mmol) at 0°C . After 30 min, to this mixture was added MOMCl (9.5 mg, 9 μL , 0.12 mmol). The reaction mixture was warmed to rt and stirred for 5 h. Without workup, this mixture was directly chromatographed eluting with 5% diethyl ether in petroleum ether to afford **66** (23 mg, 0.093 mmol, 95 %) as a pale yellow oil.

$[\alpha]_D^{25} = +39.9^\circ$ (c 1.3, CHCl_3) IR (film): 2927s, 2853m, 1621m, 1585s, 1496w, 1454m, 1404m, 1372w, 1346m, 1319w, 1290w, 1260w, 1234w, 1210w, 1151s, 1140s, 1122s, 1106s, 1086s, 1013s, 947w, 923m, 824w cm^{-1} ; ^1H -NMR (300 MHz, CDCl_3): δ 6.41 (d, $J=5.7$ Hz, 1H), 5.85 (dd, $J=10.8$, 17.4 Hz, 1H), 5.19 (d, $J=17.4$ Hz, 1H), 5.17 (d, $J=1.2$ Hz, 2H), 5.06 (d, $J=10.8$ Hz, 1H), 3.48 (s, 3H), 2.69 (dt, $J=5.1$, 17.1 Hz, 1H), 2.48 (ddd, $J=6.3$, 9.9, 16.8 Hz, 1H), 2.27 (s, 3H), 1.92 (dt, $J=5.7$, 13.5 Hz, 1H), 1.78 (ddd, $J=5.7$, 9.6, 15.6 Hz, 1H), 1.41 (s, 3H); ^{13}C -NMR (75 MHz, CDCl_3): δ 155.1, 154.4, 141.3, 137.1, 113.7, 111.0, 108.2, 106.0, 94.4, 76.2, 56.1, 31.3, 29.7, 26.8, 21.5. HRMS: Calc'd for $\text{C}_{15}\text{H}_{20}\text{O}_3$ [M^+]: 248.1412. Found: 248.1409.

(R)-2-(5-Methoxymethoxy-2,7-dimethyl-chroman-2-yl)-ethanol (67): To a solution of alkene **66** (14 mg, 0.056 mmol) in THF (1.5 mL) at 0°C was added borane THF complex (56 μL , 0.056 mmol, 1 M in THF). The reaction mixture was warmed to rt. After 1 h, to this mixture was added sodium hydroxide (0.05 mL, 1 N in water) and then hydrogen peroxide (0.03 mL, 30% in water). The reaction mixture was then heated at 50°C for 1 h. Without workup, the mixture was

purified by flash chromatography eluting with 10% diethyl ether in petroleum ether to afford alcohol **67** (14 mg, 0.052 mmol, 94%, regioselectivity=3:1) as a colorless oil.

$[\alpha]_D^{25} = +1.6^\circ$ (c 1.0, CHCl_3); IR (film): 3374b, 2926s, 1619m, 1585m, 1452m, 1403m, 1347w, 1232w, 1153s, 1105s, 1056s, 1013s, 923w, 824w cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 6.36 (s, 1H), 6.22 (s, 1H), 5.09 (s, 2H), 3.83 (m, 2H), 3.40 (s, 3H), 2.67 (bs, 1H), 2.61 (m, 2H), 2.16 (s, 3H), 1.90 (m, 1H), 1.77 (m, 1H), 1.66 (m, 1H), 1.56 (m, 1H), 1.23 (s, 3H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 155.2, 153.4, 137.3, 111.2, 107.9, 106.3, 94.3, 76.3, 59.1, 56.0, 41.6, 31.1, 23.3, 21.4, 16.5. HRMS: Calc'd for $\text{C}_{15}\text{H}_{22}\text{O}_4$ [M^+]: 266.1518. Found: 266.1520.

(R)-2-(5-Methoxymethoxy-2,7-dimethyl-chroman-2-yl)-ethyl-*tert*-butyldiphenylsilyl ether (68): To a solution of alcohol **67** (1.2 g, 4.5 mmol), 0.96 mL of triethylamine in 10 mL of dichloromethane was added *tert*-butyldiphenylsilyl chloride (1.42 mL) followed by the addition of DMAP (65 mg). The mixture was stirred at rt for 2.5 h, and without workup, purified by flash chromatography eluting with 5% diethyl ether in petroleum ether to afford silyl ether **68** (1.87g, 3.70 mmol, 82%) as a yellow oil.

$[\alpha]_D^{25} = -13.7^\circ$ (c 1.5, CDCl_3); IR (neat): 2932m, 2857m, 1619m, 1584m, 1470w, 1401w, 1428m, 1346w, 1152s, 1106s, 1058s, 1014m, 823m, 738m, 702s, 668s cm^{-1} ; $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.58 (m, 4H), 7.31 (m, 6H), 6.33 (s, 1H), 6.19 (s, 1H), 5.08 (s, 2H), 3.76 (m, 2H), 3.40 (s, 3H), 2.52 (m, 2H), 2.15 (s, 3H), 1.83 (t, $J=6.3$ Hz, 2H), 1.69 (m, 2H), 1.18 (s, 3H), 0.96 (s, 9H); $^{13}\text{C-NMR}$ (75 MHz, CDCl_3): δ 155.2, 154.2, 137.1, 135.5, 133.7, 129.6, 127.6, 111.4, 111.3, 107.9, 105.8, 94.3, 74.9, 60.0, 56.0, 41.7, 31.0, 26.8, 24.5, 21.5, 19.1, 16.6. HRMS: Calc'd for $\text{C}_{31}\text{H}_{40}\text{O}_4\text{Si}$ [M^+]: 504.2696. Found: 504.2695.