Synthesis of β,β -Disubstituted Vinyl Boronates via the Ruthenium-Catalyzed Alder Ene Reaction of Borylated Alkynes and Alkenes

Eric C. Hansen and Daesung Lee*

Department of Chemistry, University of Wisconsin, Madison, WI 53706

Supporting Information

Materials and Methods

All reagents were obtained from Aldrich Chemical Company except for the catalyst, $\text{RuCp}(\text{CH}_3\text{CN})_3\text{PF}_6$, which was purchased from Strem Chemicals or can be prepared from known procedures.¹ Alkynyl boronates **1a-d** were prepared from the cooresponding alkynyl lithium reagents and 2-isopropoxy-4,4,5,5-tetramethyl-1,3,2-dioxaborolane according to the known procedure.² Flash chromatography was performed using silica gel 60 Å (32-63 mesh) from Sorbent Technologies. Reactions were monitored by thin layer chromatography (TLC) using 0.25 mm E. Merk pre-coated silica gel 60 (particle size 0.040 – 0.063 mm). ¹H NMR and ¹³C NMR spectra were recorded on a Bruker AC-300 spectrometer. Chemical shifts are reported in δ values relative to the internal standard tetramethylsilane. Chemical shifts for ¹H NMR are referenced directly to tetramethylsilane (TMS, 0.0 ppm) and for ¹³C NMR relative to the carbon peak of the solvent (CDCl_3 , 77.2 ppm). Multiplicities are indicated as follows: s (singlet); d (doublet); t (triplet); q (quartet); p (pentet). nOe experiments were performed on a Varian INOVA-500.

¹ (a) Trost, B. M.; Older, C. M. *Organometallics* **2002**, 21, 2544. (b) Kündig, E. P.; Monnier, F. R. *Adv. Synth. Catal.* **2004**, 346, 901.

² Brown, H. C.; Bhat, N. G.; Srebnik, M. *Tetrahedron Lett.* **1988**, 29, 2631.

Spectroscopic data for **1a-d**:

1a: ^1H NMR (300 MHz, CDCl_3) δ 1.91 (s, 3H), 1.28 (s, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 84.22, 24.84, 4.74.

1b: ^1H NMR (300 MHz, CDCl_3) δ 2.52 (t, $J = 7.1$ Hz, 2H), 1.53 (m, 2H), 1.38 (m, 2H), 1.27 (s, 12H), 1.25 (m, 4H), 0.88 (t, $J = 7.0$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 84.19, 83.37, 31.49, 28.74, 28.28, 24.85, 22.67, 19.73, 14.22; **HRMS** (ESI) calcd for $\text{C}_{14}\text{H}_{25}\text{BO}_2$ $[\text{M}+\text{Na}]^+$: 259.1845, found 259.1848.

1c: ^1H NMR (300 MHz, CDCl_3) δ 4.15 (s, 2H), 3.40 (s, 3H), 1.28 (s, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 84.61, 83.34, 60.19, 57.91, 24.81; **HRMS** (ESI) calcd for $\text{C}_{10}\text{H}_{17}\text{BO}_3$ $[\text{M}-\text{CH}_3]^+$: 181.1036, found 181.1035.

1d: ^1H NMR (300 MHz, CDCl_3) δ 4.53 (q, $J = 6.6$ Hz, 1H), 1.42 (d, $J = 6.6$ Hz, 3H), 1.27 (s, 12H), 0.89 (s, 9H), 0.13 (s, 3H), 0.11 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 84.38, 59.29, 25.97, 25.20, 24.83, 18.36, -4.37, -4.85.

Representative procedure for the Alder ene reaction: To a solution of borylated alkyne **1a** (21 mg, 0.12 mmol) and alkene **2a** (14 mg, 0.10 mmol) in acetone (0.2 mL) under a nitrogen atmosphere was added $\text{RuCp}(\text{CH}_3\text{CN})_3\text{PF}_6$ (4 mg, 0.010 mmol). The reaction was stirred at room temperature for approximately 2 hours, after which time the solvent was removed in vacuo and the residue was purified by silica gel chromatography (hexane:ether 5:1) giving **3a** (21.5 mg, 70%) as a colorless oil.

3a: 8:1 mixture of cis:trans isomers ^1H NMR (300 MHz, CDCl_3) δ 5.45 (m, 2H), 5.14 (s, 1H), 4.06 (t, $J = 6.8$ Hz, 2H), 3.08 (d, $J = 5.4$ Hz, 2H), 2.08 (m, 2H), 2.05 (s, 3H), 1.83 (d, $J = 1.2$ Hz, 3H), 1.70 (p, $J = 7.9$ Hz, 2H), 1.26 (s, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.36, 161.57, 130.40, 129.53, 82.80, 64.20, 39.76, 29.02, 28.57, 26.41, 25.00, 21.18; **HRMS** (ESI) calcd for $\text{C}_{17}\text{H}_{29}\text{BO}_4$ $[\text{M}+\text{Na}]^+$: 331.2057, found 331.2042.

3b: ^1H NMR (300 MHz, CDCl_3) δ 5.57 (m, 2H), 5.17 (s, 1H), 3.15 (d, $J = 6.4$ Hz, 2H), 3.13 (d, $J = 6.4$ Hz, 2H), 2.15 (s, 3H), 1.84 (s, 3H), 1.25 (s, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ

207.59, 160.80, 133.38, 82.86, 47.84, 39.80, 29.51, 26.45, 24.99; **HRMS** (ESI) calcd for $C_{15}H_{25}BO_3$ $[M+Na]^+$: 287.1794, found 287.1791.

3c: 8:1 mixture of cis:trans isomers **1H NMR** (300 MHz, $CDCl_3$) δ 7.40 (m, 3H), 7.29 (m, 2H), 5.66 (m, 3H), 5.16 (s, 1H), 4.76 (p, $J = 6.6$ Hz, 1H), 3.70 (m, 2H), 3.17 (d, $J = 6.0$ Hz, 2H), 1.85 (s, 3H), 1.26 (s, 12H), 0.89 (d, $J = 6.7$ Hz, 3H); **^{13}C NMR** (75 MHz, $CDCl_3$) δ 171.61, 153.12, 133.54, 133.45, 128.89, 125.81, 122.64, 82.84, 79.20, 54.96, 39.83, 39.42, 26.43, 25.00, 14.72; **HRMS** (ESI) calcd for $C_{24}H_{32}BNO_5$ $[M+Na]^+$: 448.2271, found 448.2268.

3d: 9:1 mixtures of cis:trans isomers **1H NMR** (300 MHz, $CDCl_3$) δ 5.73 (dt, $J = 15.3, 6.8$ Hz, 1H), 5.55 (dd, $J = 15.3, 6.8$ Hz, 1H), 5.16 (s, 1H), 5.15 (p, $J = 6.8$ Hz, 1H), 4.16 (q, $J = 7.0$ Hz, 2H), 3.12 (d, $J = 6.8$ Hz, 2H), 1.82 (s, 3H), 1.34 (d, $J = 6.8$ Hz, 3H), 1.29 (t, $J = 7.0$ Hz, 3H), 1.24 (s, 12H); **^{13}C NMR** (75 MHz, $CDCl_3$) δ 160.35, 154.66, 132.10, 130.40, 82.83, 75.19, 63.79, 39.24, 26.47, 24.98, 20.55, 14.43; **HRMS** (ESI) calcd for $C_{17}H_{29}BO_5$ $[M+Na]^+$: 347.2066, found 347.2075.

3e: 1:1 mixture of cis:trans isomers **1H NMR** (300 MHz, $CDCl_3$) Major: δ 6.22 (d, $J = 11.8$ Hz, 1H), 5.17 (s, 1H), 4.99 (dt, $J = 11.9, 7.7$ Hz, 1H), 2.65 (d, $J = 7.7$ Hz, 2H), 1.96 (s, 3H), 1.26 (s, 12H), 0.92 (s, 9H), 0.13 (s, 6H); Minor: δ 6.30 (d, $J = 11.5$ Hz, 1H), 5.10 (d, $J = 1.3$ Hz, 1H), 5.00 (dt, $J = 12.0, 7.7$ Hz, 1H), 2.95 (d, $J = 7.7$ Hz, 2H), 1.83 (d, $J = 1.3$ Hz, 3H), 1.25 (s, 12 H), 0.92 (s, 9H), 0.13 (s, 6H); **^{13}C NMR** (75 MHz, $CDCl_3$) Major: δ 162.50, 141.47, 110.00, 82.77, 34.30, 25.91, 25.05, 18.47, -5.00; Minor: δ 162.32, 141.88, 108.64, 82.81, 40.09, 25.89, 25.05, 21.21, -5.00; **HRMS** (ESI) calcd for $C_{18}H_{35}BO_3Si$ $[M+Na]^+$: 338.2449, found 338.2459.

3f: 1:1 mixture of cis:trans isomers **1H NMR** (300 MHz, $CDCl_3$) Major: δ 7.61 (m, 2H), 7.30 (m, 2H), 6.83 (d, $J = 14$ Hz, 1H), 5.13 (s, 1H), 4.70 (dt, $J = 14, 7.3$ Hz, 1H), 3.09 (d, $J = 7.3$ Hz, 2H), 2.81 (s, 3H), 2.42 (s, 3H), 1.78 (s, 3H), 1.26 (s, 12H); Minor: δ 7.61 (m, 2H), 7.30 (m, 2H), 6.73 (d, $J = 14$ Hz, 1H), 5.08 (s, 1H), 4.68 (dt, $J = 14, 7.3$ Hz, 1H), 2.82 (s, 3H), 2.78 (d, $J = 7.3$ Hz, 2H), 2.42 (s, 3H), 1.94 (s, 3H), 1.27 (s, 12H); **^{13}C NMR** (75 MHz, $CDCl_3$) δ 161.36, 161.26, 143.80, 143.75, 134.82, 132.83, 129.88, 129.78, 129.23, 127.64, 127.21, 119.21, 109.89, 108.71, 82.91, 82.88, 53.23, 42.41, 36.92, 34.37, 32.38, 26.21, 25.02, 24.99, 21.68, 21.30; **HRMS** (ESI) calcd for $C_{20}H_{30}BNO_4S$ $[M+Na]^+$: 414.1886, found 414.1892.

3g: **1H NMR** (300 MHz, $CDCl_3$) δ 5.42 (m, 2H), 5.11 (s, 1H), 4.03 (t, $J = 6.8$ Hz, 2H), 3.06 (d, $J = 5.4$ Hz, 2H), 2.05 (t, $J = 7.3$ Hz, 2H), 2.02 (s, 3H), 2.01 (m, 2H), 1.67 (p, $J = 8.0$ Hz,

2H), 1.40 (m, 2H), 1.24 (s, 12H), 1.23 (m, 6H), 0.85 (t, $J = 6.7$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.24, 165.42, 130.08, 129.92, 82.75, 64.16, 38.89, 38.48, 31.92, 29.22, 29.00, 28.58, 27.73, 24.98, 22.73, 21.11, 14.24; **HRMS** (ESI) calcd for $\text{C}_{22}\text{H}_{39}\text{BO}_4$ $[\text{M}+\text{Na}]^+$: 401.2839, found 401.2822.

3h: 11:1 mixture of cis:trans isomers ^1H NMR (300 MHz, CDCl_3) δ 5.45 (m, 2H), 5.42 (s, 1H), 4.04 (t, $J = 6.8$ Hz, 2H), 3.85 (s, 2H), 3.30 (s, 3H), 3.07 (d, $J = 5.5$ Hz, 2H), 2.06 (m, 2H), 2.04 (s, 3H), 1.67 (p, $J = 6.8$ Hz, 2H), 1.26 (s, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.29, 159.63, 130.48, 129.22, 82.99, 76.30, 64.13, 58.30, 35.32, 28.95, 28.51, 24.97, 21.14; **HRMS** (ESI) calcd for $\text{C}_{18}\text{H}_{31}\text{BO}_5$ $[\text{M}+\text{Na}]^+$: 361.2162, found 361.2158.

3i: ^1H NMR (300 MHz, CDCl_3) δ 5.91 (dt, $J = 18.5, 6.5$ Hz, 1H), 5.62 (dt, $J = 18.5, 1.2$ Hz, 1H), 5.40 (t, $J = 1.2$ Hz, 1H), 3.78 (d, $J = 1.2$ Hz, 2H), 3.24 (s, 3H), 3.13 (dd, $J = 6.5, 1.2$ Hz, 2H), 1.19 (s, 12H), -0.04 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3) δ 158.93, 144.60, 131.32, 83.02, 76.48, 58.34, 39.49, 24.99, -1.05; **HRMS** (ESI) calcd for $\text{C}_{16}\text{H}_{31}\text{BO}_3\text{Si}$ $[\text{M}+\text{Na}]^+$: 303.1572, found 303.1573.

3j: ^1H NMR (300 MHz, CDCl_3) δ 5.43 (m, 3H), 4.20 (q, $J = 6.4$ Hz, 1H), 4.04 (t, $J = 6.8$ Hz, 2H), 3.29 (d, $J = 14.0$ Hz, 1H), 2.84 (d, $J = 14.0$ Hz, 1H), 2.04 (m, 2H), 2.03 (s, 3H), 1.66 (p, $J = 8.0$ Hz, 2H), 1.25 (s, 12H), 1.19 (d, $J = 6.4$ Hz, 3H), 0.87 (s, 9H), 0.00 (s, 3H), -0.01 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.30, 166.95, 130.71, 129.69, 82.83, 72.08, 66.03, 64.20, 34.98, 29.00, 28.59, 26.11, 25.08, 24.96, 23.99, 21.26, 18.44, 15.46, -4.51, -4.72; **HRMS** (ESI) calcd for $\text{C}_{24}\text{H}_{45}\text{BO}_5\text{Si}$ $[\text{M}+\text{Na}]^+$: 475.3027, found 475.3024.

3k: ^1H NMR (300 MHz, CDCl_3) δ 5.81 (dt, $J = 16.3, 6.2$ Hz, 1H), 5.63 (dt, $J = 16.3, 6.5$ Hz, 1H), 5.47 (s, 1H), 4.54 (d, $J = 6.4$ Hz, 2H), 4.17 (q, $J = 7.1$ Hz, 2H), 4.19 (q, $J = 6.2$ Hz, 1H), 3.33 (dd, $J = 5.9$ Hz, 1H), 2.98 (dd, $J = 14.0, 7.4$ Hz, 1H), 1.29 (t, $J = 7.2$ Hz, 3H), 1.25 (s, 12H), 1.18 (d, $J = 6.2$ Hz, 3H), 0.87 (s, 9H), 0.01 (s, 3H), -0.01 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 165.81, 155.25, 136.58, 124.09, 82.96, 72.70, 68.53, 64.04, 34.34, 26.09, 25.07, 24.98, 23.97, 18.41, 14.47, -4.53, -4.71; **HRMS** (ESI) calcd for $\text{C}_{23}\text{H}_{43}\text{BO}_6\text{Si}$ $[\text{M}+\text{Na}]^+$: 477.2820, found 477.2832.

Procedure for Suzuki-Miyaura coupling: A flame-dried 25 mL round-bottom flask was charged with a solution of vinyl boronate **3a** (55 mg, 0.18 mmol), *trans*-2-iodoacrylate (45 mg, 0.20 mmol) and $\text{Pd}(\text{PPh}_3)_4$ (20 mg, 0.018 mmol) in THF (5 mL) under a nitrogen atmosphere. Thallium ethoxide (100 mg, 0.4 mmol) was then added slowly over 10 min.

The reaction stirred at room temperature for 12 hours. The brown suspension was then partitioned between ether and water, and the aqueous phase was extracted twice with ether. The combined organic phases were dried over MgSO_4 , filtered and concentrated in vacuo. The residue was purified by silica gel chromatography (Hex:EtOAc 8:1) yielding **10** (30 mg, 60%) as a colorless oil.

Procedure for Heck reaction: To a solution of **3a** (30 mg, 0.1 mmol) and ethyl acrylate (12 mg, 0.12 mmol) in dry DMF was added $\text{Pd}(\text{OAc})_2$ (1 mg, 0.005 mmol) and neocuproine (2 mg). The reaction was stirred at room temperature under O_2 atmosphere (balloon pressure) for 3 hours. After this time, water was added and the product extracted with ether. The combined ether extracts were dried over MgSO_4 , filtered and concentrated in vacuo. The residue was purified by silica gel chromatography (Hex:EtOAc 8:1).

7: $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 7.56 (dd, $J = 15.0, 11.5$ Hz, 1H), 6.01 (d, $J = 11.5$ Hz, 1H), 5.79 (d, $J = 15.0$ Hz, 1H), 5.42 (m, 2H), 4.20 (q, $J = 7.1$ Hz, 2H), 4.05 (t, $J = 6.6$ Hz, 2H), 2.96 (d, $J = 6.2$ Hz, 2H), 2.07 (m, 2H), 2.05 (s, 3H), 1.84 (s, 3H), 1.69 (p, $J = 7.0$ Hz, 2H), 1.30 (t, $J = 7.1$ Hz, 3H); $^{13}\text{C NMR}$ (75 MHz, CDCl_3) δ 171.36, 167.81, 148.22, 140.52, 131.44, 127.50, 124.31, 119.50, 64.08, 60.35, 36.14, 28.96, 24.43, 21.18, 14.54; **HRMS** (ESI) calcd for $\text{C}_{16}\text{H}_{24}\text{O}_4[\text{M}+\text{Na}]^+$: 303.1572, found 303.1573.

94.223

24.837

4.737

NOE data for **3a**

NOE for **3g**

NOE for **3k**

NOE for **10**

For comparison, we performed the Alder ene reaction with a silylated alkyne to verify that the stereochemical outcome is opposite that of the vinylboronates synthesized here.