Supporting Information

Synthesis of Heparin Oligosaccharides

Jinq-Chyi Lee,[‡] Xin-An Lu,[†] Suvarn S. Kulkarni,[†] Yuh-Sheng Wen,[†] and Shang-Cheng Hung^{*,†}

*Institute of Chemistry, Academia Sinica, Taipei 115, Taiwan, and Department of Chemistry,
National Tsing Hua University, Hsinchu 300, Taiwan*

§ The deposition number of Compound **15β** at the Cambridge Crystallographic Data Centre is CCDC 217878.

§ Experimental Section

General Procedures. Solvents were purified and dried from a safe purification system by passing through activated aluminium oxide under argon pressure.¹ Flash column chromatography was carried out on Silica Gel 60 (230-400 mesh, E. Merck).² TLC was performed on pre-coated glass plates of Silica Gel 60 F254 (0.25 mm, E. Merck); detection was executed by spraying with a solution of Ce(NH₄)₂(NO₃)₆, (NH₄)₆Mo₇O₂₄, and H₂SO₄ in water or ninhydrin and acetic acid solution in *n*-butanol and subsequent heating on a hot plate. Melting points were determined with a

1. Pangborn, A. B.; Giardello, A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518.
2. Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.

Büchi B-540 apparatus and are uncorrected. Optical rotations were measured with a Jasco DIP-370 polarimeter at $\sim 25^\circ\text{C}$. ^1H and ^{13}C NMR spectra were recorded with Bruker AMX400 and 500 MHz instruments. Chemical shifts are in ppm from Me_4Si , generated from the CDCl_3 lock signal at δ 7.24. IR spectra were taken with a Perkin-Elmer Paragon 1000 FT-IR spectrometer. Elemental analyses were measured with a Perkin-Elmer 2400CHN instrument. Mass spectra were obtained with a FAB JMS-700 double focusing mass spectrometer (JEOL, Tokyo, Japan), MALDI Voyager DE-PRO (Applied Biosystem Houston, USA) and ESI Finnigan LCQ mass spectrometer (Thermo Finnigan, San Jose, CA, United States) in negative mode. Gel-filtration chromatography (Sephadex® G-25 fine) was used in order to achieve purification of the final products.

Compound 9. To a solution of **8** (193 mg, 0.62 mmol) in CH_2Cl_2 (2.0 mL) was added pyridine (0.5 mL, 6.18 mmol) at 0°C under nitrogen. Benzoyl chloride (87 μL , 0.75 mmol) was added subsequently to the mixture in a dropwise manner. After stirring at 0°C for 2 h, methanesulfonyl chloride (73 μL , 0.94 mmol) was added, the reaction was warmed up to room temperature gradually and kept stirring overnight. The reaction was quenched by addition of H_2O (5 mL). The aqueous layer was extracted with EtOAc (3 x 10 mL). The combined organic layers were sequentially washed with 1 N $\text{HCl}_{(\text{aq})}$, saturated $\text{NaHCO}_{3(\text{aq})}$ and brine, dried over MgSO_4 , filtered, and concentrated *in vacuo*. The residue was purified by column chromatography ($\text{EtOAc}/\text{Hex} = 1/4$) to afford **9** (248 mg, 81%) as a white solid. $[\alpha]_D^{26}$ (*c* 0.65, CHCl_3) -14.52 ; mp $94\text{--}95^\circ\text{C}$; IR (CHCl_3) ν 2986, 1716, 1447, 1267, 1075, 921, 786, 709 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.06 (dd, $J = 7.5, 1.4$ Hz, 2H, Bz-H), 7.56 (t, $J = 7.5$ Hz, 1H, Bz-H), 7.45–7.26 (m, 7H, Ar-H), 5.91 (d, $J = 3.6$ Hz, 1H, H-1), 5.40 (ddd, $J = 8.0, 6.2, 2.0$ Hz, 1H, H-5), 4.91 (dd, $J = 12.8, 2.0$ Hz, 1H, H-6a), 4.70 (d, $J = 11.0$ Hz, 1H, CH_2Ph), 4.60 (d, $J = 3.6$ Hz, 1H, H-2), 4.59 (d, $J = 11.0$ Hz, 1H, CH_2Ph), 4.49 (dd, $J = 12.8, 6.2$ Hz, 1H, H-6b), 4.46 (dd, $J = 8.0, 3.1$ Hz, 1H, H-4), 4.13 (d, $J = 3.1$ Hz, 1H, H-3); ^{13}C NMR (100 MHz, CDCl_3) δ 166.0 (C), 137.1 (C), 133.2 (CH), 129.7 (CH), 129.6 (C), 128.5 (CH), 128.1 (CH), 128.0 (CH), 112.3 (C), 105.4 (CH), 81.5 (CH), 81.1 (CH), 78.3 (CH), 75.3 (CH), 72.4 (CH_2), 64.0 (CH_2), 39.1 (CH_3), 26.8 (CH_3), 26.3 (CH_3); HRMS (FAB, MH^+) calcd for $\text{C}_{24}\text{H}_{29}\text{O}_9\text{S}$ 493.1532, found 493.1543

Compound 10. To a solution of **9** (1.20 g, 2.44 mmol) in a mixed solvent ($\text{CH}_2\text{Cl}_2/t\text{BuOH} = 2/1$, 18 mL) was added potassium *t*-butoxide (0.60 g, 5.36 mmol) at 0°C under nitrogen. After stirring for 16 h, the reaction was neutralized with 0.6 N $\text{H}_2\text{SO}_{4(\text{aq})}$ (ca. 4.5 mL) and the flask was equipped with a simple distillation head to evaporate CH_2Cl_2 and *t*BuOH under reduced pressure.

0.6 N $\text{H}_2\text{SO}_{4(\text{aq})}$ (5 mL) and diglyme (10 mL) were added to the resulting solution and the mixture was kept stirring at 160 °C for another 16 h. After cooling to room temperature, the reaction was neutralized with 3 N $\text{NaOH}_{(\text{aq})}$ (2 mL), and the solvent was removed on rotary evaporator under vacuum. Water (10 mL) was added to the residue, and the mixture was extracted with EtOAc (3 x 10 mL). The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and concentrated *in vacuo*. The residue was purified by flash column chromatography (EtOAc/Hex = 1/1) to give a white solid, which was further recrystallized via vapor diffusion method to provide **10** (0.32 g, 52%) as colorless crystallines. mp 158-159 °C; ^1H NMR (400 MHz, CDCl_3) δ 7.36-7.28 (m, 5H, Ph-H), 5.27 (d, J = 1.8 Hz, 1H, H-1), 4.93 (d, J = 11.7 Hz, 1H, CH_2Ph), 4.72 (d, J = 11.7 Hz, 1H, CH_2Ph), 4.41 (t, J = 4.9 Hz, 1H, H-5), 4.01 (d, J = 7.8 Hz, 1H, H-6a), 3.85 (ddd, J = 7.8, 4.9, 3.3 Hz, 1H, H-4), 3.71 (dd, J = 7.8, 4.9 Hz, 1H, H-6b), 3.64 (dt, J = 7.8, 1.8 Hz, 1H, H-2), 3.37 (t, J = 7.8 Hz, 1H, H-3), 2.09 (d, J = 3.3 Hz, 1H, OH-4), 1.89 (d, J = 7.8 Hz, 1H, OH-2); ^{13}C NMR (75 MHz, CDCl_3) δ 138.4 (C), 128.7 (CH), 128.0 (CH), 127.9 (CH), 101.8 (CH), 84.2 (CH), 75.4 (CH), 75.0 (CH), 74.5 (CH_2), 71.1 (CH), 65.1 (CH_2); HRMS (FAB, M^+) calcd for $\text{C}_{13}\text{H}_{16}\text{O}_5$ 252.0997, found 252.0992. Anal. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_5$: C, 61.90; H, 6.39. Found: C, 61.65; H, 6.35.

Compounds 15 α & 15 β . A mixture of **14** (0.51 g, 0.95 mmol) and freshly dried 4 Å molecular sieves (1 g) in dichloromethane (5 mL) was stirred at room temperature for 30 min under nitrogen. The reaction flask was cooled to -78 °C, anhydrous potassium carbonate (0.26 g, 1.88 mmol) and trichloroacetonitrile (0.95 mL, 9.47 mmol) were sequentially added to the solution, and the mixture was gradually warmed up to room temperature. After stirring for 16 h, the resulting solution was filtered through celite, and the solid was washed with dichloromethane. The filtrate was concentrated *in vacuo* to afford the crude trichloroacetimidate (0.64 g, 98%, α/β = 1/1.9 determined by the ^1H NMR spectrum), which was directly used without further purification for the ensuing reaction.

A solution of this crude trichloroacetimidate (0.62 g, 0.91 mmol) and **11** (0.49 g, 1.37 mmol) in dichloromethane (11 mL) was added to a reaction flask containing freshly dried 4 Å molecular sieves (2 g) under nitrogen. The mixture was stirred at room temperature for 1 h, and the solution was cooled to -78 °C. Trimethylsilyl trifluoromethanesulfonate (25 μL , 0.14 mmol) was added to the reaction flask, the mixture was gradually warmed up to room temperature, and the resulting solution was kept stirring for 6 h. Triethylamine (50 μL) was added to quench the reaction, and the whole mixture was filtered through celite followed by wash with dichloromethane. The filtrate was concentrated *in vacuo* to furnish a residue, which was purified by flash column chromatography (EtOAc/Hex = 1/4) to yield **15 α** (0.49 g, 61%) and **15 β** (89 mg, 11%). The β -isomer was further recrystallized through vapor diffusion method to get colorless crystallines for X-ray single crystal analysis. **15 α** : ^1H NMR (400 MHz, CDCl_3) δ 8.04 (d, J = 8.4 Hz, 2H, Bz-H), 7.95 (d, J = 8.4 Hz, 2H, Bz-H), 7.81-7.73 (m, 3H, Ar-H), 7.69 (s, 1H, Ar-H), 7.60-7.52 (m, 2H, Ar-H), 7.46-7.32 (m, 13H, Ar-H), 7.25-7.17 (m, 4H, Ar-H), 5.53 (d, J = 1.6 Hz, 1H, H-1), 5.31 (d, J = 3.7 Hz, 1H, H-1'),

5.06 (dd, $J = 7.7, 1.6$ Hz, 1H, H-2), 5.04 (d, $J = 10.9$ Hz, 1H, CH₂Ar), 4.99 (d, $J = 10.6$ Hz, 1H, CH₂Ar), 4.95 (d, $J = 10.6$ Hz, 1H, CH₂Ar), 4.88 (d, $J = 10.9$ Hz, 1H, CH₂Ar), 4.82 (d, $J = 10.9$ Hz, 1H, CH₂Ar), 4.76 (d, $J = 10.9$ Hz, 1H, CH₂Ar), 4.65-4.60 (m, 2H, H-5, H-6a'), 4.47 (dd, $J = 12.0, 5.1$ Hz, 1H, H-6b'), 4.19 (d, $J = 7.8$ Hz, 1H, H-6a), 4.11 (t, $J = 7.7$ Hz, 1H, H-3), 4.09-4.03 (m, 2H, H-4, H-3'), 3.96 (ddd, $J = 9.6, 5.1, 1.7$ Hz, 1H, H-5'), 3.78 (dd, $J = 7.8, 5.2$ Hz, 1H, H-6b), 3.68 (t, $J = 9.6$ Hz, 1H, H-4'), 3.45 (dd, $J = 10.4, 3.7$ Hz, 1H, H-2'); ¹³C NMR (100 MHz, CDCl₃) δ 166.1 (C), 165.7 (C), 137.8 (C), 137.5 (C), 134.5 (C), 133.4 (CH), 133.3 (CH), 133.2 (C), 133.1 (C), 129.8 (CH), 129.51 (CH), 129.46 (C), 129.4 (C), 128.6 (CH), 128.54 (CH), 128.50 (CH), 128.4 (CH), 128.3 (CH), 128.1 (CH), 127.9 (CH), 127.8 (CH), 127.7 (CH), 127.1 (CH), 126.3 (CH), 126.2 (CH), 125.8 (CH), 99.5 (CH), 99.3 (CH), 80.1 (CH), 79.4 (CH), 78.6 (CH), 78.2 (CH), 75.7 (CH₂), 75.6 (CH₂), 74.9 (CH₂), 74.3 (CH), 70.1 (CH), 65.7 (CH₂), 63.5 (CH), 63.2 (CH₂); HRMS (MALDI, MNa⁺) calcd for C₅₁H₄₇N₃O₁₁Na 900.3108, found 900.3110. **15b**: mp 163-165 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.00 (dd, $J = 8.4, 1.3$ Hz, 2H, Bz-H), 7.91 (dd, $J = 8.4, 1.3$ Hz, 2H, Bz-H), 7.77-7.70 (m, 3H, Ar-H), 7.66 (s, 1H, Ar-H), 7.56 (t, $J = 7.4$ Hz, 1H, Ar-H), 7.49 (t, $J = 7.4$ Hz, 1H, Ar-H), 7.45-7.30 (m, 12H, Ar-H), 7.13-7.11 (m, 2H, Ar-H), 7.07-7.00 (m, 3H, Ar-H), 5.55 (d, $J = 1.7$ Hz, 1H, H-1), 5.02 (dd, $J = 8.2, 1.7$ Hz, 1H, H-2), 5.00 (d, $J = 10.7$ Hz, 1H, CH₂Ar), 4.96 (d, $J = 10.7$ Hz, 1H, CH₂Ar), 4.88 (d, $J = 10.7$ Hz, 1H, CH₂Ar), 4.83 (d, $J = 11.6$ Hz, 1H, CH₂Ar), 4.80 (d, $J = 10.7$ Hz, 1H, CH₂Ar), 4.69 (d, $J = 11.6$ Hz, 1H, CH₂Ar), 4.63 (t, $J = 4.9$ Hz, 1H, H-6a), 4.60 (dd, $J = 12.0, 2.1$ Hz, 1H, H-6a'), 4.49 (d, $J = 7.8$ Hz, 1H, H-1'), 4.50-4.45 (m, 1H, H-6b'), 4.26-4.20 (m, 2H, H-4, H-6b), 3.97 (t, $J = 8.2$ Hz, 1H, H-3), 3.83 (dd, $J = 7.4, 4.9$ Hz, 1H, H-5), 3.77 (dd, $J = 9.6, 8.6$ Hz, 1H, H-4'), 3.62 (ddd, $J = 9.6, 3.8, 2.1$ Hz, 1H, H-5'), 3.57-3.49 (m, 2H, H-2', H-3'); ¹³C NMR (100 MHz, CDCl₃) δ 165.9 (C), 165.7 (C), 138.1 (C), 137.6 (C), 134.6 (C), 133.2 (CH), 133.1 (CH), 129.9 (CH), 129.5 (CH), 129.4 (C), 128.6 (CH), 128.4 (CH), 128.3 (CH), 128.1 (CH), 128.0 (CH), 127.9 (CH), 127.7 (CH), 127.6 (CH), 127.4 (CH), 127.1 (CH), 126.2 (CH), 126.1 (CH), 125.8 (CH), 101.0 (CH), 99.2 (CH), 83.1 (CH), 78.8 (CH), 77.3 (CH), 77.0 (CH), 76.3 (CH), 75.8 (CH₂), 75.2 (CH₂), 74.1 (CH₂), 73.5 (CH), 73.0 (CH), 66.8 (CH), 65.5 (CH₂), 62.8 (CH₂); HRMS (FAB, MH⁺) calcd for C₅₁H₄₈N₃O₁₁ 878.3289, found 878.3287.

Compound 23. The same procedure as described before for **15a** was followed. It took 3 h to complete the reaction and compound **23** was afforded in 84% yield after flash column chromatography (EtOAc/Hex = 1/3). ¹H NMR (500 MHz, CDCl₃) δ 8.04-8.03 (m, 2H, Bz-H), 8.00-7.98 (m, 2H, Bz-H), 7.90-7.88 (m, 2H, Bz-H), 7.77-7.75 (m, 2H, Ar-H), 7.71 (dd, $J = 6.1, 3.4$ Hz, 1H, Ar-H), 7.62 (s, 1H, Ar-H), 7.51-7.47 (m, 2H, Ar-H), 7.45-7.41 (m, 2H, Ar-H), 7.38-7.18 (m, 23H, Ar-H), 5.30 (d, $J = 3.5$ Hz, 1H, H-1'), 5.17 (dd, $J = 4.1, 3.5$ Hz, 1H, H-2'), 4.99 (d, $J = 10.2$ Hz, 1H, CH₂Ar), 4.92 (d, $J = 10.9$ Hz, 1H, CH₂Ar), 4.80 (d, $J = 11.4$ Hz, 1H, CH₂Ar), 4.76 (d, $J = 3.6$ Hz, 1H, H-1), 4.74-4.71 (m, 4H, H-1'', 3 x CH₂Ar), 4.65 (dd, $J = 12.0, 1.5$ Hz, 1H, H-6a), 4.56-4.47 (m, 5H, H-5', H-6b, H-6a'', H-6b'', CH₂Ar), 4.36 (dd, $J = 11.5, 7.7$ Hz, 1H, H-6a'), 4.30

(d, $J = 10.5$ Hz, 1H, CH₂Ar), 4.11 (dd, $J = 11.5, 4.9$ Hz, 1H, H-6 b'), 4.07 (t, $J = 4.1$ Hz, 1H, H-3'), 3.98-3.93 (m, 4H, H-3, H-4, H-5, H-5''), 3.75 (t, $J = 10.1$ Hz, 1H, H-3''), 3.73 (t, $J = 4.3$ Hz, 1H, H-4'), 3.64 (t, $J = 10.1$ Hz, 1H, H-4''), 3.41 (s, 3H, OCH₃), 3.41-3.39 (m, 1H, H-2), 3.31 (dd, $J = 10.1, 3.7$ Hz, 1H, H-2''), 2.58-2.55 (m, 2H, 2 x OCOCH₂), 2.46-2.42 (m, 2H, 2 x CH₂COCH₃), 2.04 (s, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃) δ 206.3 (C), 172.2 (C), 166.0 (C), 165.9 (C), 165.4 (C), 137.9 (C), 137.4 (C), 137.2 (C), 134.7 (C), 133.2 (CH), 133.10 (CH), 133.05 (CH), 129.74 (CH), 129.68 (C), 129.7 (CH), 129.6 (C), 129.5 (CH), 128.52 (CH), 128.45 (CH), 128.4 (CH), 128.3 (CH), 128.2 (CH), 128.1 (CH), 128.0 (CH), 127.9 (CH), 127.7 (CH), 127.6 (CH), 127.1 (CH), 126.2 (CH), 126.1 (CH), 125.9 (CH), 98.6 (CH), 98.4 (CH), 98.0 (CH), 80.6 (CH), 79.0 (CH), 77.5 (CH), 75.8 (CH), 75.3 (CH₂), 74.6 (CH), 73.7 (CH), 73.2 (CH₂), 70.3 (CH), 70.0 (CH), 69.1 (CH), 67.2 (CH), 63.81 (CH), 63.75 (CH), 62.9 (CH₂), 62.7 (CH₂), 62.5 (CH₂), 55.3 (CH₃), 37.7 (CH₂), 29.7 (CH₃), 27.7 (CH₂); HRMS (FAB, MH⁺) calcd for C₇₇H₇₇N₆O₁₉ 1389.5244, found 1389.5264. Anal. Calcd for C₇₇H₇₆N₆O₁₉: C, 66.56; H, 5.51; N, 6.05. Found: C, 66.26; H, 5.62; N, 5.73.

Compound 24. To a stirred solution of **23** (0.36 g, 0.26 mmol) in a mixed solvent (CH₂Cl₂/H₂O = 18/1, 19 mL) was added DDQ (0.18 g, 0.79 mmol) in three equal portions at half-hour intervals at room temperature. After stirring for 4 h, the reaction was quenched by addition of saturated NaHCO_{3(aq)} (20 mL), and the mixture was extracted with CH₂Cl₂ (2 x 20 mL). The organic layers were consecutively washed with saturated NaHCO_{3(aq)} (2 x 20 mL) and brine, dried over anhydrous MgSO₄, filtered, and concentrated *in vacuo*. The residue was purified by flash column chromatography (EtOAc/ Hex = 1/1.8) to give the 4-OH compound (0.26 g, 80%). ¹H NMR (500 MHz, CDCl₃) δ 8.05-8.03 (m, 2H, Bz-H), 7.99 (d, $J = 8.0$ Hz, 4H, Bz-H), 7.54-7.50 (m, 2H, Ar-H), 7.47-7.33 (m, 9H, Ar-H), 7.32-7.17 (m, 13H, Ar-H), 5.27 (d, $J = 3.6$ Hz, 1H, H-1'), 5.19 (dd, $J = 4.2, 3.6$ Hz, 1H, H-2'), 4.99 (d, $J = 10.3$ Hz, 1H, CH₂Ph), 4.81 (d, $J = 3.7$ Hz, 1H, H-1''), 4.78 (d, $J = 11.5$ Hz, 2H, 2 x CH₂Ph), 4.76 (d, $J = 3.6$ Hz, 1H, H-1), 4.73 (s, 1H, CH₂Ph), 4.72 (d, $J = 11.5$ Hz, 1H, CH₂Ph), 4.65-4.61 (m, 2H, H-6a, CH₂Ph), 4.51-4.46 (m, 2H, H-5', H-6a''), 4.41 (d, $J = 10.7$ Hz, 1H, H-6b), 4.37 (dd, $J = 12.5, 2.0$ Hz, 1H, H-6b''), 4.33 (dd, $J = 11.6, 7.6$ Hz, 1H, H-6a'), 4.14 (dd, $J = 11.6, 5.0$ Hz, 1H, H-6b'), 4.07 (t, $J = 4.2$ Hz, 1H, H-3'), 3.97-3.92 (m, 3H, H-3, H-4, H-5), 3.84 (d, $J = 9.8$ Hz, 1H, H-5''), 3.79 (t, $J = 4.2$ Hz, 1H, H-4'), 3.62 (dd, $J = 10.2, 9.8$ Hz, 1H, H-3''), 3.48 (td, $J = 9.8, 4.4$ Hz, 1H, H-4''), 3.41-3.38 (m, 1H, H-2), 3.40 (s, 3H, OCH₃), 3.21 (dd, $J = 10.2, 3.7$ Hz, 1H, H-2''), 3.16 (d, $J = 4.4$ Hz, 1H, OH-4''), 2.66-2.61 (m, 2H, 2 x OCOCH₂), 2.49-2.35 (m, 2H, 2 x CH₂COCH₃), 2.11 (s, 3H, CH₃); ¹³C NMR (125 MHz, CDCl₃) δ 207.1 (C), 172.6 (C), 167.4 (C), 166.3 (C), 165.7 (C), 138.2 (C), 138.0 (C), 137.5 (C), 133.63 (CH), 133.57 (CH), 133.4 (CH), 130.0 (CH), 129.9 (CH), 129.64 (C), 129.61 (C), 128.99 (CH), 128.97 (CH), 128.9 (CH), 128.8 (CH), 128.7 (CH), 128.7 (CH), 128.6 (CH), 128.59 (CH), 128.51 (CH), 128.46 (CH), 128.4 (CH), 128.3 (CH), 128.2 (CH), 127.9 (CH), 98.7 (CH), 98.5 (CH), 98.2

(CH), 79.9 (CH), 79.2 (CH), 76.1 (CH), 75.52 (CH₂), 75.50 (CH₂), 74.2 (CH), 74.1 (CH), 73.6 (CH₂), 71.7 (CH), 71.0 (CH₂), 70.6 (CH), 69.4 (CH), 67.8 (CH), 64.0 (CH), 63.4 (CH₂), 63.3 (CH), 63.1 (CH₂), 62.6 (CH₂), 55.6 (CH₃), 38.1 (CH₂), 30.0 (CH₃), 28.0 (CH₂); HRMS [FAB, (M-H)⁺] calcd for C₆₆H₆₇N₆O₁₉ 1247.4460, found 1247.4465.

A mixture of this 4-alcohol (0.16 g, 0.13 mmol), the imidate **22** (0.15 g, 0.13 mmol), and freshly dried 4Å molecular sieves (1 g) in dichloromethane (6 mL) was stirred at room temperature for 1 h under nitrogen. The reaction flask was cooled to -40 °C, and trimethylsilyl trifluoromethanesulfonate (5 µL, 0.06 mmol) was added to the reaction mixture. After stirring for 1.5 h, more **22** (0.30 g, 0.26 mmol) in dichloromethane (1 mL) and trimethylsilyl trifluoromethanesulfonate (5 µL, 0.06 mmol) were consecutively added to the solution at the same temperature, and the mixture was stirred for an additional 3 h. The flask was gradually warmed up to room temperature and triethylamine (20 µL) was added to quench the reaction. The mixture was filtered through celite, the solid was washed with dichloromethane, and the filtrate was concentrated *in vacuo*. The residue was purified by flash column chromatography (EtOAc/Hex = 1/2) to provide **24** (0.21 g, 74%). ¹H NMR (400 MHz, CDCl₃) δ 8.08-8.01 (m, 8H, Bz-H), 7.91 (dd, *J* = 8.3, 1.1 Hz, 2H, Bz-H), 7.79-7.77 (m, 2H, Ar-H), 7.73 (dd, *J* = 6.0, 3.3 Hz, 1H, Ar-H), 7.63 (s, 1H, Ar-H), 7.55-7.49 (m, 3H, Ar-H), 7.47-7.42 (m, 3H, Ar-H), 7.40-7.18 (m, 37H, Ar-H), 5.32 (d, *J* = 3.7 Hz, 1H, H-1'), 5.25 (d, *J* = 3.0 Hz, 1H, H-1''), 5.20 (t, *J* = 3.7 Hz, 1H, H-2'), 5.17 (t, *J* = 3.0 Hz, 1H, H-2''), 5.02 (d, *J* = 10.3 Hz, 1H, CH₂Ar), 4.92 (d, *J* = 10.8 Hz, 1H, CH₂Ar), 4.84 (d, *J* = 11.5 Hz, 1H, CH₂Ar), 4.80-4.72 (m, 8H, H-1, H-1''), 6 x CH₂Ar), 4.69-4.64 (m, 2H), 4.67 (d, *J* = 3.3 Hz, 1H, H-1'), 4.53-4.39 (m, 8H), 4.31 (dd, *J* = 11.5, 7.8 Hz, 1H, H-6a'), 4.23 (d, *J* = 10.4 Hz, 1H, CH₂Ar), 4.19 (d, *J* = 10.3 Hz, 1H, CH₂Ar), 4.12 (dd, *J* = 11.1, 4.4 Hz, H-6b'), 4.09-3.99 (m, 3H, H-3', H-3'', H-6b'), 3.98-3.90 (m, 6H, H-3, H-4, H-4'', H-5, H-5'', H-5'''), 3.78 (t, *J* = 4.0 Hz, 1H, H-4'), 3.74-3.63 (m, 4H, H-3'', H-3''', H-4', H-4''), 3.44-3.41 (m, 1H, H-2), 3.43 (s, 3H, OCH₃), 3.33-3.28 (m, 2H, H-2'', H-2'''), 2.63 (t, *J* = 6.5 Hz, 2H), 2.56-2.50 (m, 2H), 2.48-2.44 (m, 2H), 2.41-2.37 (m, 2H), 2.14 (s, 3H, CH₃), 2.03 (s, 3H, CH₃); ¹³C NMR (100 MHz, CDCl₃) δ 206.33 (C), 206.26 (C), 172.2 (C), 172.1 (C), 166.0 (C), 165.90 (C), 165.87 (C), 165.4 (C), 137.9 (C), 137.7 (C), 137.4 (C), 137.2 (C), 134.7 (C), 133.3 (CH), 133.2 (CH), 133.12 (CH), 133.09 (CH), 133.0 (CH), 129.73 (CH), 129.65 (CH), 129.6 (CH), 129.5 (C), 129.48 (CH), 129.4 (C), 128.5 (CH), 128.49 (CH), 128.45 (CH), 128.4 (CH), 128.3 (CH), 128.24 (CH), 128.21 (CH), 128.1 (CH), 128.0 (CH), 127.92 (CH), 127.85 (CH), 127.7 (CH), 127.64 (CH), 127.58 (CH), 127.1 (CH), 126.14 (CH), 126.06 (CH), 125.9 (CH), 98.6 (CH), 98.4 (CH), 98.1 (CH), 97.9 (CH), 97.8 (CH), 80.6 (CH), 79.0 (CH), 78.8 (CH), 77.3 (CH), 76.0 (CH), 75.3 (CH₂), 75.2 (CH₂), 75.0 (CH₂), 74.9 (CH), 74.6 (CH), 74.2 (CH), 74.0 (CH), 73.3 (CH₂), 73.2 (CH), 73.1 (CH₂), 70.34 (CH), 70.28 (CH), 69.8 (CH), 69.1 (CH), 67.5 (CH), 66.8 (CH), 63.8 (CH), 62.8 (CH₂), 62.62 (CH₂), 62.56 (CH₂), 62.4 (CH₂), 62.2 (CH₂), 55.3 (CH₃), 37.8 (CH₂), 37.6 (CH₂), 29.71 (CH₃), 29.65 (CH₃), 27.7 (CH₂), 27.6 (CH₂); LRMS (MALDI, MNa⁺) calcd for C₁₂₂H₁₂₁N₉O₃₂Na 2246.80, found 2247.00.

Compound 27. $\text{H}_2\text{NNH}_2/\text{AcOH}$ (33 mg, 0.36 mmol) was added to a solution of compound **23** (100 mg, 72 μmol) in a mixed solvent (ethanol/toluene = 2/1, 7.5 mL) at room temperature under nitrogen. After stirring for 2 h, the solvent was removed under reduced pressure, the residue was diluted with H_2O (5 mL), and the mixture was extracted by EtOAc (5 mL x 3). The combined organic layers were sequentially washed with saturated $\text{NaHCO}_3(\text{aq})$ and brine, dried over anhydrous MgSO_4 , filtered, and concentrated *in vacuo* to furnish a residue, which was purified by flash column chromatography (EtOAc/Hex = 1/3) to afford the primary alcohol (90 mg, 97%). ^1H NMR (400 MHz, CDCl_3) δ 8.05 (dd, J = 7.8, 1.4 Hz, 2H, Bz-H), 7.99 (dd, J = 8.4, 1.3 Hz, 2H, Bz-H), 7.90 (dd, J = 8.4, 1.3 Hz, 2H, Bz-H), 7.77-7.74 (m, 2H, Ar-H), 7.71-7.67 (m, 1H, Ar-H), 7.61 (s, 1H, Ar-H), 7.52-7.48 (m, 2H, Ar-H), 7.44-7.40 (m, 2H, Ar-H), 7.39-7.25 (m, 21H, Ar-H), 7.23-7.20 (m, 2H, Ar-H), 5.192 (s, 1H, H-1'), 5.188 (s, 1H, H-2'), 4.91 (d, J = 11.0 Hz, 1H, CH_2Ar), 4.86 (d, J = 11.2 Hz, 1H, CH_2Ar), 4.81 (d, J = 10.7 Hz, 1H, CH_2Ar), 4.80 (d, J = 3.6 Hz, 1H, H-1), 4.78 (d, J = 3.6 Hz, 1H, H-1''), 4.77 (d, J = 10.7 Hz, 1H, CH_2Ar), 4.67-4.58 (m, 3H, H-6a, 2 x CH_2Ar), 4.52 (d, J = 10.4 Hz, 1H, CH_2Ar), 4.52-4.42 (m, 3H, H-6b, H-6a'', H-6b''), 4.35 (d, J = 10.4 Hz, 1H, CH_2Ar), 4.23 (td, J = 7.0, 2.6 Hz, 1H, H-5'), 4.08 (t, J = 4.0 Hz, 1H, H-3'), 4.04-3.96 (m, 3H, H-4, H-5, H-5''), 3.91 (dd, J = 9.9, 9.0 Hz, 1H, H-3), 3.75 (dd, J = 9.9, 9.2 Hz, 1H, H-3''), 3.66-3.65 (m, 1H, H-4'), 3.61 (t, J = 9.2 Hz, 1H, H-4''), 3.59-3.52 (m, 1H, H-6a'), 3.48 (dd, J = 9.9, 3.6 Hz, 1H, H-2), 3.44 (s, 3H, CH_3), 3.36-3.32 (m, 1H, H-6b'), 3.34 (dd, J = 9.9, 3.6 Hz, 1H, H-2''); ^{13}C NMR (125 MHz, CDCl_3) δ 166.1 (C), 166.0 (C), 165.7 (C), 137.7 (C), 137.5 (C), 137.4 (C), 134.7 (C), 133.2 (CH), 133.1 (CH), 129.8 (CH), 129.7 (CH), 129.52 (CH), 128.51 (CH), 128.43 (CH), 128.40 (CH), 128.35 (CH), 128.0 (CH), 127.9 (CH), 127.8 (CH), 127.7 (CH), 127.0 (CH), 126.1 (CH), 126.0 (CH), 125.9 (CH), 98.4 (CH), 97.907 (CH), 97.904 (CH), 80.7 (CH), 79.1 (CH), 77.6 (CH), 75.3 (CH_2), 75.2 (CH_2), 74.4 (CH), 73.6 (CH), 73.5 (CH), 72.9 (CH_2), 70.1 (CH), 69.4 (CH), 69.3 (CH), 68.0 (CH), 64.0 (CH), 63.8 (CH), 63.0 (CH_2), 61.2 (CH_2), 55.4 (CH_3); HRMS (MALDI, MNa^+) calcd for $\text{C}_{72}\text{H}_{70}\text{N}_6\text{O}_{17}\text{Na}$ 1313.4694, found 1313.4660.

To a solution of this primary alcohol (44 mg, 34 μmol) in CH_2Cl_2 (1.7 mL) was consecutively added H_2O (1.7 mL), 1 M $\text{KBr}(\text{aq})$ (34 μL , 34 μmol), TEMPO (5 mg, 34 μmol), 0.5 M $\text{NaHCO}_3(\text{aq})$ (1.7 mL), and $\text{Bu}_4\text{N}^+\text{Cl}^-$ (19 mg, 68 μmol) at room temperature. The reaction flask was immersed in an ice-bath, and NaOCl (0.22 mL, 3.6 mmol) was added to the mixture which was simultaneously calibrated with 0.5 N $\text{NaOH}(\text{aq})$ through micro-syringe to maintain at pH=10. The resulting solution was gradually warmed up to room temperature, and 0.5 N $\text{NaOH}(\text{aq})$ was added to keep at the same pH value. After stirring for 3 h, the mixture was extracted with CH_2Cl_2 (5 mL x 3), and the combined organic layers were acidified with 1 N $\text{HCl}(\text{aq})$, dried over MgSO_4 , filtered, and concentrated *in vacuo*. The residue was purified by flash column chromatography (EtOAc/Hex = 1/2) to give **27** (38 mg, 87%). ^1H NMR (400 MHz, CDCl_3) δ 8.04 (dd, J = 6.8, 1.6 Hz, 2H, Bz-H), 8.00 (dd, J = 7.1, 1.4 Hz, 2H, Bz-H), 7.88 (dd, J = 8.3, 1.2 Hz, 2H, Bz-H), 7.82-7.79 (m, 2H, Ar-H), 7.75-7.73 (m, 1H, Ar-H), 7.66 (s, 1H, Ar-H), 7.52-7.46 (m, 4H, Ar-H), 7.42-7.38 (m, 8H, Ar-H), 7.36-7.26 (m, 13H, Ar-H), 7.20-7.18 (m, 2H, Ar-H), 5.50 (d, J = 2.0 Hz, 1H, H-1'), 5.16 (t, J = 2.0 Hz, 1H, H-2'), 4.99 (d, J = 2.4 Hz, 1H, H-5'), 4.96 (d, J = 10.5 Hz, 1H, CH_2Ar), 4.94 (d, J = 11.0 Hz, 1H, CH_2Ar), 4.90 (d, J = 11.6 Hz, 1H, CH_2Ar), 4.83 (d, J = 10.5 Hz, 1H, CH_2Ar), 4.81 (d, J = 3.4 Hz, 1H, H-1), 4.80 (d, J = 11.6 Hz, 1H, CH_2Ar), 4.749 (d, J = 3.6 Hz, 1H, H-1''), 4.752 (d, J = 11.0 Hz, 1H, CH_2Ar), 4.72-4.67 (m, 2H, H-6a, H-6a''), 4.49 (dd, J = 12.1, 3.5 Hz, 1H, H-6b), 4.39 (dd, J = 12.4, 2.5 Hz, 1H, H-6b''), 4.33 (d, J = 10.5 Hz, 1H, CH_2Ar), 4.18 (d, J = 10.5 Hz, 1H, CH_2Ar), 4.14-4.03 (m, 3H, H-3', H-4', H-5''), 3.98-3.92 (m, 3H, H-3, H-4, H-5), 3.68-3.60 (m, 2H, H-3'', H-4''), 3.46-3.44 (m, 1H, H-2), 3.44 (s, 3H, OCH_3), 3.23 (dd, J = 9.9, 3.6 Hz, 1H, H-2''); ^{13}C

NMR (100 MHz, CDCl₃) δ 169.6 (C), 166.4 (C), 166.0 (C), 165.4 (C), 137.6 (C), 137.5 (C), 137.1 (C), 134.9 (C), 133.3 (CH), 133.13 (CH), 133.10 (CH), 133.05 (C), 129.9 (CH), 129.7 (CH), 129.6 (CH), 129.3 (C), 128.6 (CH), 128.43 (CH), 128.36 (CH), 128.3 (CH), 128.2 (CH), 128.1 (CH), 128.0 (CH), 127.9 (CH), 127.8 (CH), 127.7 (CH), 127.0 (CH), 126.2 (CH), 126.1 (CH), 126.0 (CH), 99.0 (CH), 98.5 (CH), 98.4 (CH), 80.3 (CH), 79.0 (CH), 77.6 (CH), 76.2 (CH), 75.2 (CH₂), 75.1 (CH₂), 75.0 (CH₂), 74.4 (CH), 72.9 (CH), 72.7 (CH₂), 70.1 (CH), 69.0 (CH), 68.8 (CH), 68.4 (CH), 63.8 (CH), 63.7 (CH), 62.9 (CH₂), 62.5 (CH₂), 55.4 (CH₃); HRMS (MALDI, MNa⁺) calcd for C₇₂H₆₈N₆O₁₈Na 1327.4488, found 1327.2898.

Compound 28. The same procedure as described before for the trisaccharide **27** was followed. LRMS (MALDI, MNa⁺) calcd for C₁₁₂H₁₀₅N₉O₃₀Na 2078.69, found 2078.47.

Compound 29. The same procedure as described before for the trisaccharide **27** was followed. LRMS (MALDI, MNa⁺) calcd for C₁₅₂H₁₄₂N₁₂O₄₂Na 2829.92, found 2830.86.

Compound 30. The same procedure as described before for trisaccharide **27** was followed. LRMS (MALDI, MNa⁺) calcd for C₁₉₂H₁₇₉N₁₅O₅₄Na 3581.16, found 3581.96.

Compound 2. (1) Debenzoylation. Sodium methoxide (6.0 mg, 0.11 mmol) was added to a solution of **27** (36 mg, 28 μ mol) in MeOH (0.6 mL) at room temperature under nitrogen. The mixture was kept stirring overnight, and the reaction solution was neutralized by Amberlite-120 acidic resin. The resulting mixture was filtered through paper, and the filtrate was concentrated *in vacuo* to yield the crude triol which was directly used for next step. ¹H NMR (500 MHz, CD₃OD) δ 8.00-7.94 (m, 3H, Ar-H), 7.88 (s, 1H, Ar-H), 7.66-7.56 (m, 5H, Ar-H), 7.53-7.50 (m, 2H, Ar-H), 7.47-7.37 (m, 11H, Ar-H), 5.49 (d, *J* = 3.0 Hz, 1H, H-1'), 5.32 (d, *J* = 3.8 Hz, 1H, H-1), 5.08-5.03 (m, 3H, 3 x CH₂Ar), 5.00-4.96 (m, 4H, H-5', 3 x CH₂Ar), 4.92-4.90 (m, 2H, H-1'', CH₂Ar), 4.67 (d, *J* = 11.1 Hz, 1H, CH₂Ar), 4.38 (t, *J* = 4.2 Hz, 1H, H-4'), 4.18 (t, *J* = 4.2 Hz, 1H, H-3'), 4.13-4.07 (m, 3H), 4.04-4.02 (m, 3H), 4.00-3.94 (m, 3H), 3.85-3.79 (m, 2H), 3.78 (dd, *J* = 10.3, 3.8 Hz, 1H, H-2), 3.56 (s, 3H, OCH₃), 3.45 (dd, *J* = 10.3, 3.6 Hz, 1H, H-2''); ¹³C NMR (125 MHz, CD₃OD) δ 175.0 (C), 139.5 (C), 139.4 (C), 137.1 (C), 134.7 (C), 134.4 (C), 129.40 (CH), 129.36 (CH), 129.3 (CH), 129.20 (CH), 129.18 (CH), 129.0 (CH), 128.88 (CH), 128.85 (CH), 128.7 (CH), 128.5 (CH), 127.5 (CH), 127.2 (CH), 127.0 (CH), 102.1 (CH), 100.0 (CH), 97.6 (CH), 82.0 (CH), 79.7 (CH), 79.4 (CH), 76.4 (CH₂), 76.367 (CH), 76.364 (CH), 75.96 (CH₂), 75.95 (CH₂), 74.9 (CH), 73.8 (CH₂), 73.5 (CH), 73.1 (CH), 70.2 (CH), 65.4 (CH), 64.7 (CH), 61.8 (CH₂), 61.7 (CH₂), 55.5 (CH₃); HRMS (MALDI, MNa⁺) calcd for C₅₁H₅₆N₆O₁₅Na 1015.3701, found 1015.5527.

(2) *O*-Sulfonation. A solution of this crude triol and sulfur trioxide-triethylamine complex (0.22 g, 1.2 mmol) in DMF (0.6 mL) was stirred at 50 °C under nitrogen overnight. The reaction flask was cooled down to room temperature, a solution of NaHCO₃ (0.42 g) in H₂O (5 mL) was added to the mixture, and the mixture was kept stirring for another 16 h. The solvent was coevaporated with ethanol under reduced pressure, and a mixed solvent of CH₂Cl₂/MeOH (1/1, 10

mL) was added to the residue. The mixture was filtered through paper, and the filtrate was concentrated *in vacuo* to give a syrup which was dissolved in a mixed solvent CH₂Cl₂/MeOH (4/1, 10 mL). Repeating of the filtration and concentration steps led to the crude tri-*O*-sulfonated derivative, which was used for next reaction without further purification. LRMS (ESI negative mode) calcd for [M-2H+Na]⁻ 1253.22, found 1253.28.

(3) Hydrogenolysis. A solution of this crude compound in a mixed solvent 1:9 H₂O/MeOH (1/9, 4 mL) was hydrogenated in the presence of 10% Pd/C (70 mg) under 50 psi pressure at room temperature. After 2 d, the mixture was filtered through celite, and the filtrate was concentrated *in vacuo*. The same procedure was repeated again until no signals of aryl groups could be detected by ¹H NMR spectrum.

(4) *N*-sulfonation. The above amino-alcohol was dissolved in water (3 mL), and the solution was adjusted to pH=9.5 through addition of 2 N NaOH_(aq). Sulfur trioxide-pyridine complex (0.17 g, 1.0 mmol) was added in five equal portions at half-hour intervals at room temperature, and the pH value was maintained at 9.5 via calibration of 2 N NaOH_(aq). After stirring for 3 h, the reaction mixture was concentrated *in vacuo*, and the residue was purified by column chromatography on Sephadex G-25 using 0.2 N NaCl_(aq) as an eluent. The crude product portion was lyophilized followed by desalting through a Sephadex G-25 column eluted with water to give compound **2** (10 mg, 36%) as a white solid. ¹H NMR (500 MHz, D₂O) δ 5.47 (d, *J* = 3.3 Hz, 1H, H-1''), 5.30 (d, *J* = 2.2 Hz, 1H, H-1'), 5.08 (d, *J* = 3.5 Hz, 1H, H-1), 4.43-4.37 (m, 5H), 4.28-4.26 (m, 3H), 4.16 (t, *J* = 3.1 Hz, 1H), 4.05-4.03 (m, 2H), 3.88-3.68 (m, 4H), 3.63 (t, *J* = 10.0 Hz, 1H, H-3''), 3.49 (s, 3H, OCH₃), 3.46-3.42 (m, 1H), 3.35 (dd, *J* = 10.3, 3.5 Hz, 1H, H-2), 3.33 (dd, *J* = 10.0, 3.3 Hz, 1H, H-2''); ¹³C NMR (125 MHz, D₂O) δ 99.1 (CH), 98.2 (CH), 96.6 (CH), 76.7 (CH), 75.6 (CH), 75.5 (CH), 70.9 (CH), 69.9 (CH), 69.8 (CH), 69.2 (CH), 69.1 (CH), 68.7 (CH), 68.6 (CH), 68.4 (CH), 66.7 (CH₂), 66.5 (CH₂), 57.7 (CH), 57.6 (CH), 55.4 (CH₃); LRMS (ESI negative mode) calcd for [M-2H+Na]⁻ 950.95, found 951.17; calcd for [M-3H+2Na]⁻ 972.94, found 973.25.

Compound 3. The same protocol as described before for the trisaccharide **2** was followed to obtain the pentasaccharide **3** in 21% overall yield. (1) Debenzoylation. HRMS (MALDI, MNa⁺) calcd for C₇₇H₈₅N₉O₂₅Na 1558.5533, found 1558.5508. (2) *O*-Sulfonation. LRMS (ESI negative mode) calcd for [M-5H+4Na]⁻ 2022.27, found 2021.00. (3) Hydrogenolysis and *N*-Sulfonation. ¹H NMR (500 MHz, D₂O) δ 5.47-5.44 (m, 1H), 5.29-5.25 (m, 2H, H-1', H-1''), 5.07 (d, *J* = 3.5 Hz, 1H, H-1), 4.92-4.79 (m, 3H), 4.46-4.22 (m, 12H), 4.15-3.97 (m, 6H), 3.85-3.60 (m, 7H), 3.48 (s, 3H, OCH₃), 3.45-3.41 (m, 1H), 3.34 (dd, *J* = 10.4, 3.3 Hz, 1H), 3.31 (dd, *J* = 10.3, 3.6 Hz, 1H); ¹³C NMR (125 MHz, D₂O) δ 99.2 (CH), 99.1 (CH), 98.2 (CH), 96.7 (CH), 96.5 (CH), 76.8 (CH), 75.9 (CH), 75.8 (CH), 75.5 (CH), 75.4 (CH), 70.9 (CH), 69.9 (CH), 69.8 (CH), 69.5 (CH), 69.2 (CH), 69.11 (CH), 69.05 (CH), 68.9 (CH), 68.4 (CH), 66.8 (CH₂), 66.7 (CH₂), 66.4 (CH₂), 57.82 (CH), 57.75 (CH), 57.6 (CH), 55.4 (CH₃); LRMS (ESI negative mode) calcd for [M-6H+5Na]⁻ 1615.85, found 1616.00; calcd for [M-7H+6Na]⁻ 1637.84, found 1636.00; calcd for [M-9H+8Na]⁻ 1681.80, found 1681.00.

Compound 4. The same protocol as described before for the trisaccharide **2** was followed to obtain the heptasaccharide **4** in 8% overall yield. (1) Debenzoylation. LRMS (MALDI, MNa⁺) calcd for C₁₀₃H₁₁₄N₁₂O₃₅Na 2101.74, found 2101.92. (2) *O*-Sulfonation. LRMS (ESI negative mode) calcd for [M-8H+7Na]⁻ 2791.32, found 2792.36. (3) Hydrogenolysis and *N*-Sulfonation. LRMS (ESI negative mode) calcd for [M-3H+2Na]⁻ 2126.88, found 2127.00; calcd for [M-9H+8Na]⁻ 2258.77, found 2259.00.

Compound 5. The same protocol as described before for the trisaccharide **2** was followed to obtain the nonasaccharide **5** in 7% overall yield. (1) Debenzoylation. LRMS (MALDI, MNa⁺)

calcd for $C_{129}H_{143}N_{15}O_{45}Na$ 2644.93, found 2644.44. (2) *O*-Sulfonation. LRMS (ESI negative mode) calcd for $[M-8H+7Na]^-$ 3494.41, found 3494.00; calcd for $[M-10H+9Na]^-$ 3538.38, found 3538.00. (3) Hydrogenolysis and *N*-Sulfonation. 1H NMR (500 MHz, D_2O) δ 5.50-5.46 (m, 2H), 5.31-5.28 (m, 3H), 5.13-5.08 (m, 2H), 5.00 (s, 1H), 4.91-4.89 (m, 2H), 4.87-4.80 (m, 3H), 4.48-4.28 (m, 20H), 4.22-4.03 (m, 13H), 3.83-3.62 (m, 15H), 3.50 (s, 3H, OCH_3), 3.37-3.32 (m, 4H); ^{13}C NMR (125 MHz, D_2O) δ 99.24 (CH), 99.09 (CH), 98.98 (CH), 98.93 (CH), 98.25 (CH), 96.80 (CH), 96.72 (CH), 96.67 (CH), 76.12 (CH), 76.07 (CH), 75.81 (CH), 75.68 (CH), 75.58 (CH), 75.47 (CH), 75.37 (CH), 75.32 (CH), 75.24 (CH), 69.96 (CH), 69.85 (CH), 69.74 (CH), 69.62 (CH), 69.51 (CH_2), 69.43 (CH), 69.32 (CH), 69.27 (CH), 69.17 (CH), 69.05 (CH), 68.91 (CH), 68.65 (CH), 68.52 (CH), 66.90 (CH_2), 66.56 (CH_2), 66.46 (CH_2), 66.40 (CH_2), 62.61 (CH_2), 57.91 (CH), 57.84 (CH), 57.75 (CH), 55.51 (CH_3); LRMS (ESI negative mode) calcd for $[M-H]^-$ 2659.89, found 2660.00; $[M-4H+3Na]^-$ 2725.83, found 2726.00; calcd for $[M-8H+7Na]^-$ 2813.76, found 2816.00. LRMS (ESI negative mode) calcd for $[M-H]^-$ 2659.89, found 2660.00; $[M-4H+3Na]^-$ 2725.83, found 2726.00; calcd for $[M-8H+7Na]^-$ 2813.76, found 2816.00.

1.55936
1.90100
1.90281
1.92309
1.92508
2.10273
2.11094
3.37565
3.39508
3.41504
3.63601
3.65561
3.65861
3.66296
3.71156
3.72448
3.73105
3.74379
3.86394
3.87310
3.88432
4.02530
4.04476
4.42153
4.43330
4.44471
4.72787
4.75723
4.94393
4.97327
5.29461
5.29902
7.26170
7.26376
7.30513
7.31722
7.32719
7.33868
7.37301
7.38376

Current Data Parameters
NAME sc0913
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date 990913
Time 13.56
PULPROG zg
SOLVENT CDCl3
AQ 1.7039560 sec
FIDRES 0.293438 Hz
DQ 104.0 usec
RG 1024
NUCLEUS 1H
HL1 3 dB
D1 1.0000000 sec
P1 6.0 usec
DE 148.8 usec
SF01 400.1319596 MHz
SWH 4807.69 Hz
TD 16384
NS 16
DS 2

F2 - Processing parameters
SI 16384
SF 400.1300086 MHz
WDW EM
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 23.90 cm
FIP 7.758 ppm
F1 3104.38 Hz
F2 6.885 ppm
F2 2755.05 Hz
PPMCH 0.03653 ppm/
HZDM 14.61619 Hz/c

Current Data Parameters
NAME sc0430
EXPNO 1
PROCNO 1

F2 - Acquisition Parameter
Date 990430
Time 12.21
PULPROG zg
SOLVENT CDCl3
AQ 1.7039560 se
FIDRES 0.293438 Hz
AQ 104.0 us
RG 512
NUCLEUS 1H
HL1 3 dB
D1 1.0000000 se
P1 6.0 us
DE 130.0 us
SF01 400.1320110 MHz
SMH 4807.69 Hz
TD 16384
NS 16
DS 0

F2 - Processing parameters
SI 32768
SF 400.1300099 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 0.80

1D NMR plot parameters
CX 23.90 cm
F1P 10.000 pp
F1 4001.30 Hz
F2P -0.500 pp
F2 -200.07 Hz
PPMCH 0.43933 pp
HZCH 175.78934 Hz

Current Data Parameters
NAME LJC-132-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20000320
Time 17:56
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT DMSO
NS 32
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 512
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SF01 400.1319596 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300106 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters
NAME LJC-134-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20000320
Time 17:48
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 32
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 512
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SFO1 400.131956 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300172 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters
NAME LJC-131-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 970131
Time 20.02
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 32
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 16384
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SFO1 400.1319596 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300114 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters	
NAME	LJC-140-P
EXPNO	5
PROCNO	1

P2 - Acquisition	
Title.....	960306
Date.....	2-08
INSTRUM ..	spect
PROBHD ..	5 mm Multist
PULPRG ..	#3
TD.....	32768
SOLVENT ..	CDCl ₃
NS.....	4
DS.....	2
SWH.....	6410.256 Hz
FIDRES ..	0.195625 Hz
AQ.....	2.5559540 sec
RG.....	64
DW.....	78.000 usec
DE.....	111.43 usec
TE.....	300.0 K
H1L.....	3 dB
D1.....	1.000000000 sec
P1.....	6.00 usec
SFO1.....	400.1319596 MHz
NUCLEUS ..	¹ H

Current Data Parameters
NAME LJC-138-P-2
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20021006
Time 1.54
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl₃
NS 40
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 512
DE 78.000 usec
TE 111.43 usec
HL1 300.0 K
D1 3 dB
P1 1.00000000 sec
SFO1 400.1319596 MHz
NUCLEUS ¹H

F2 - Processing parameters
SI 32768
SF 400.1300174 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters
NAME LJC-NH3-10H
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20021028
Time 23.58
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 41
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 2048
DE 78.000 usec
TE 111.43 usec
HL1 300.0 K
D1 3.00 usec
P1 1.00000000 sec
SFO1 400.131956 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300172 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters		F2 - Acquisition Parameters		F2 - Processing parameters	
NAME	LJC-685-P	Date	20020928	SI	32768
EXPNO	1	Time	15.25	SF	400.1300052 MHz
PROCNO	1	INSTRUM	5 mm spect	WDW	no
		PROBHD	5 mm PULPROG	SSB	0
		TD	32768	LB	0.00 Hz
		SOLVENT	CDCl3	GB	0
		NS	33	PC	1.00
		DS	2		
		SWH	6410.256 Hz		
		FIDRES	0.195626 Hz		
		AQ	2.5559540 sec		
		RG	512		
		DW	78.000 usec		
		DE	111.43 usec		
		TE	300.0 K		
		HL1	3 dB		
		P1	1.00000000 sec		
		F1	6.00 usec		
		SF01	400.1319596 MHz		
		NUCLEUS	1H

Current Data Parameters
NAME LJC-754b
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20021004
Time 3.45
INSTRUM spect
PROBHD 5 mm Maltinu
PULPROG zg
TD 32768
SOLVENT CDCl₃
NS 43
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 64
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SFO1 400.131956 MHz
NUCLEUS ¹H

F2 - Processing parameters
SI 32768
SF 400.1300172 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters		F2 - Acquisition Parameters		F2 - Processing parameters	
NAME	LJC-807-P	Date	20021112	SI	32768
EXPNO	1	Time	0.35	SF	400.1300172 MHz
PROCNO	1	INSTRUM	5 mm spect	WDW	no
		PROBHD	1H	SSB	0
		PULPROG	zg	LB	0.00 Hz
		TD	32768	GB	0
		SOLVENT	CDCl3	PC	1.00
		NS	33		
		DS	2		
		SWH	6410.256 Hz		
		FIDRES	0.195626 Hz		
		AQ	2.5559540 sec		
		RG	128		
		DW	78.000 usec		
		DE	111.43 usec		
		TE	300.0 K		
		HL1	3 dB		
		D1	1.00000000 sec		
		P1	6.00 usec		
		SF01	400.1319596 MHz		
		NUCLEUS	1H

Current Data Parameters	
NAME	LJC-166-P
EXPNO	1
PROCNO	1

P2 - Acquisition Parameters	
Date_	970329
Time	10:32
INSTRUM	5 mm Multispec
PROBHD	zgpg30
PULPROG	zgpg30
TD	32768
SOLVENT	CDCl3
DS	32
NS	2
SWH	6410.256 Hz
FIDRES	0.195625 Hz
AQ	2.5559540 sec
RG	128
DW	78.000 usec
DE	111.43 usec
TE	300.0 K
HL1	3 dB
D1	1.00000000 sec
P1	6.00 usec
SFO1	400.1319596 MHz
NUC1	1H

F2 - Processing parameters	
SI	32768
SF	400.1300176 MHz
WDW	no
SSB	0
LB	0.00 Hz
GB	0
PC	1.00

Current Data Parameters
NAME LJC-241-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 970516
Time 4:44
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl₃
NS 8
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 128
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SFO1 400.1319596 MHz
NUCLEUS ¹H

F2 - Processing parameters
SI 32768
SF 400.1300052 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

7.3925
7.3858
7.3738
7.3645
7.3577
7.3372
7.3181
7.3015
7.2828
7.2649
7.2611
7.2405
7.2316
7.2286
7.2243
7.2189
7.2140
7.2065
5.3116
5.1849
5.1760
5.1673
5.0219
4.9962
4.9448
4.9175
4.8266
4.7982
4.7762
4.7670
4.7626
4.7590
4.7503
4.7416
4.7348
4.7138
4.6778
4.6481
4.5804
4.5753
4.5495
4.5452
4.5342
4.5071
4.4972
4.4814
4.4743
4.4663
4.4539
4.3839
4.3653
4.3550
4.3366
4.3246
4.2984
4.1304
4.1185
4.1015
4.0955
4.0853
4.0747
3.9948
3.9873
3.9553
3.9424
3.9356
3.9114
3.7812
3.7567
3.7336
3.7290
3.7188
3.6656
3.6413
3.6186
3.4298
3.4163

Current Data Parameters
NAME LJC-244-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 970526
Time 12.15
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 8
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 64
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SFO1 400.1319596 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300172 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters
NAME LJC-467-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20010725
Time 18.21
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
DS 32
NS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 128
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SFO1 400.1319596 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300174 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

2.182	4.309	2.129	10.058	14.089	1.000	1.038	1.085	6.954	1.084	1.159	1.248	1.288	3.379	1.072	1.083	3.431	1.075	1.040	1.137	4.404	0.988	0.967	3.205
-------	-------	-------	--------	--------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

Current Data Parameters
NAME schung010402
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20020104
Time 20.24
INSTRUM spect
PROBHD 5 mm txi2 1h/v1
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 8
DS 2
SWH 5995.204 Hz
FIDRES 0.182959 Hz
AQ 2.7329011 sec
RG 57
DE 83.400 usec
TE 15.50 usec
D1 300.0 K
1.00000000 sec
P1 5.00 usec
SFO1 500.1325156 MHz
NUC1 1H
PL1 -2.00 dB

F2 - Processing parameters
SI 16384
SF 500.1300275 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

The chemical structure shows a complex molecule. On the left, a 2-NAPO group is attached to a BnO group. This is connected via an oxygen atom to a sugar derivative. The sugar derivative has a BnO group and a HO group. The N3 group is also present, connected to the sugar derivative.

7.2027
 7.1890
 7.1851
 7.1526
 7.1453
 7.1335
 7.1290
 6.2460
 6.2167
 6.2111
 5.2664
 5.2296
 5.2240
 5.2175
 5.2118
 5.1069
 4.9153
 4.9107
 4.8965
 4.8880
 4.8832
 4.8676
 4.8443
 4.8157
 4.7831
 4.7718
 4.7546
 4.7415
 4.7137
 4.6591
 4.6500
 4.6124
 4.6072
 4.5991
 4.5818
 4.5767
 4.5307
 4.5215
 4.5131
 4.5000
 4.4911
 4.4823
 4.4787
 4.4638
 4.4507
 4.4410
 4.4283
 4.4086
 4.3808
 4.3619
 4.3418
 4.3346
 4.3194
 4.3078
 4.2912
 4.2796
 4.2678
 4.2627
 4.2490
 4.1913
 4.1840
 4.1672
 4.1158
 4.1120
 4.1081
 4.0870
 4.0716
 4.0660
 4.0429
 4.0169
 3.7532
 3.7281
 3.7221
 3.7057
 3.6984
 3.6756
 3.6510
 3.6369
 3.6255

Current Data Parameters
 NAME LJC-608-P
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20020425
 Time 16.38
 INSTRUM spect
 PROBHD 5 mm Multinu
 PULPROG zg
 TD 32768
 SOLVENT CDCl3
 NS 13
 DS 2
 SWH 6410.256 Hz
 FIDRES 0.195625 Hz
 AQ 2.5559540 sec
 RG 64
 DW 78.000 usec
 DE 111.43 usec
 TE 300.0 K
 HL1 3 dB
 P1 1.00000000 sec
 D1 6.00 usec
 SF01 400.1319596 MHz
 NUCLEUS 1H
 F2 - Processing parameters
 SI 32768
 SF 400.1300168 MHz
 WDW no
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

7.2400
 7.2391
 7.1408
 7.1328
 7.1237
 7.1201
 5.2752
 5.2618
 5.2569
 5.2375
 5.2325
 5.0685
 5.0596
 5.0550
 5.0498
 5.0445
 4.8897
 4.8676
 4.8622
 4.8448
 4.8391
 4.8159
 4.7777
 4.7591
 4.7488
 4.7300
 4.7195
 4.6922
 4.6267
 4.6175
 4.5930
 4.5877
 4.5623
 4.5573
 4.5500
 4.5405
 4.5222
 4.5078
 4.5027
 4.4985
 4.4930
 4.4771
 4.4721
 4.4674
 4.4428
 4.4230
 4.4148
 4.4020
 4.3956
 4.3170
 4.3079
 4.2986
 4.2858
 4.2732
 4.2673
 4.2508
 4.2329
 4.2262
 4.2070
 4.0512
 4.0356
 4.0220
 4.0140
 3.9880
 3.9736
 3.9476
 3.9036
 3.6570
 3.6496
 3.6244
 3.6128
 3.6070
 3.6034
 3.5882
 3.5831
 3.5688
 3.5601
 3.5465

Current Data Parameters
 NAME LJC-61
 EXPNO
 PROCNO
 F2 - Acquisition Parameters
 Date_ 20020201
 Time 16:51
 INSTRUM spect
 PROBHD 5 mm Mult
 PULPROG zgpg30
 TD 32768
 SOLVENT CDCl3
 NS 1024
 DS 4
 SWH 6410.3 Hz
 FIDRES 0.195 Hz
 AQ 2.5555 sec
 RG 327.68
 DE 78.50 mm
 TE 300.2 K
 HL1 5 mm
 NUC1 1H
 P1 1.00000 sec
 SFO1 400.1315 MHz
 NUC2 13C
 F2 - Processing parameters
 SI 32768
 SF 400.1300 MHz
 WDW EM
 SSB 0
 LB 0.3 Hz
 GB 0
 PC 1

7.3172
7.3128
7.3050
7.3001
7.2959
7.2844
7.2800
7.2708
7.2699
7.2673
7.2566
7.1723
7.1647
7.1534
7.1485
6.5060
6.4991
6.4556
5.3472
5.3472
4.9713
4.9573
4.9441
4.9280
4.9001
4.8783
4.7717
4.7617
4.7426
4.7342
4.7270
4.7177
4.6655
4.6520
4.6390
4.6248
4.6135
4.5880
4.5828
4.5611
4.5514
4.5425
4.5207
4.5113
4.5029
4.4561
4.4377
4.4279
4.4092
4.3677
4.3450
4.3188
4.3131
4.2998
4.2842
4.2325
4.1241
4.1186
4.0997
4.0744
4.0484
3.7742
3.7316
3.7090
3.6845
3.6622
3.6570
3.6333
3.6104
3.4999
3.4197
3.3991
3.3940
3.3901
3.3852
3.3743
3.3653
2.8874

Current Data Parameters
NAME LJC-598-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20020418
Time 15:33
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl₃
NS 19
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 512
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
P1 1.00000000 sec
D1 6.00 usec
SFO1 400.131956 MHz
NUCLEUS ¹H

F2 - Processing parameters
SI 32768
SF 400.1300052 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters
NAME schung050802
EXPNO 21
PROCNO 1

F2 - Acquisition Parameters
Date_ 20020508
Time 21:34
INSTRUM spect
PROBHD 5 mm tx12 lh/l
PULPROG zg
TD 16384
SOLVENT CDCl3
NS 8
DS 2
SWH 5518.764 Hz
FIDRES 0.336839 Hz
AQ 1.484404 sec
RG 101.6
DE 90.600 usec
TE 15.50 usec
D1 280.0 K
1.00000000 sec
P1 3.00 usec
SFO1 500.1322463 MHz
NUC1 1H
PC1 -2.00 dB

F2 - Processing parameters
SI 16384
SF 500.1300236 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

7.1834 3.4732
7.1730 3.4855
7.1664 3.4972
7.1486 3.5516
5.2858 3.5759
5.2768 3.5986
5.2228 3.6349
5.2144 3.6561
5.1825 3.6812
5.1726 3.6976
5.1664 3.7075
5.1577 3.7175
5.1513 3.7407
4.9970 3.7510
4.9713 3.7614
4.9085 3.7790
4.8085 3.7857
4.7799 3.8100
4.7577 3.8729
4.7577 3.8792
4.7577 3.8980
4.7577 3.9174
4.7577 3.9251
4.7577 3.9334
4.7577 3.9586
4.7577 4.0244
4.7577 4.0367
4.7577 4.0498
4.7577 4.0621
4.7577 4.0796
4.7577 4.0954
4.7577 4.1079
4.7577 4.1121
4.7577 4.1590
4.7577 4.1850
4.7577 4.2345
4.7577 4.2530
4.7577 4.2629
4.7577 4.2817
4.7577 4.3313
4.7577 4.3472
4.7577 4.3589
4.7577 4.3745
4.7577 4.3940
4.7577 4.4115
4.7577 4.4209
4.7577 4.4359
4.7577 4.4680
4.7577 4.4750
4.7577 4.5025
4.7577 4.5104
4.7577 4.5621
4.7577 4.5890
4.7577 4.6076
4.7577 4.6395
4.7577 4.6974
4.7577 4.7049
4.7577 4.7103
4.7577 4.7157
4.7577 4.7266
4.7577 4.7384
4.7577 4.7478
4.7577 4.7577
4.7577 4.7744
4.7577 4.7799
4.7577 4.8085
4.7577 4.9713
4.7577 4.9970
4.7577 5.1577
4.7577 5.1664
4.7577 5.1726
4.7577 5.1825
4.7577 5.2144
4.7577 5.2228
4.7577 5.2768
4.7577 5.2858
4.7577 7.1486
4.7577 7.1664
4.7577 7.1730
4.7577 7.1834

Current Data Parameters
NAME LJC-735-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20020917
Time 3:54
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl₃
NS 32
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 128
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SFO1 400.131956 MHz
NUCLEUS ¹H

F2 - Processing parameters
SI 32768
SF 400.1300174 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

```
Current Data Parameters
NAME      schung11402
EXPNO     1
PROCNO    1
```

F2 - Acquisition Parameters	
Date_	20021114
Time	21:04
INSTRUM	spect
PROBHD	5 mm tx1z lvy1
PULPROG	zgpg30
TD	16384
SOLVENT	CDCl3
NS	8
DS	2
SWH	5518.7664 Hz
FIDRES	0.336839 Hz
AQ	1.4644404 sec
RG	32
DW	90.600 usec
DE	15.50 usec
TE	280.0 K
TD1	1.00000000 sec
PD1	4.00 usec
SP1	500.1322463 MHz
NUC1	¹ H
PL1	-2.00 dB

F2 - Processing parameters	
SI	16384
SF	500.1300236 MHz
WDW	no
SSB	0
LB	0.00 Hz
GB	0
PC	1.00

5.2947
 5.2876
 5.2323
 5.2258
 5.2174
 5.2109
 5.1917
 5.1834
 5.1752
 5.1675
 5.1601
 5.1528
 5.0597
 5.0051
 4.9845
 4.8101
 4.7943
 4.7871
 4.7708
 4.7627
 4.7534
 4.7412
 4.7328
 4.7239
 4.7184
 4.7027
 4.6816
 4.6743
 4.6668
 4.6348
 4.6202
 4.6112
 4.5869
 4.5655
 4.5111
 4.4875
 4.4518
 4.4475
 4.4284
 4.4214
 4.4031
 4.3874
 4.3800
 4.3556
 4.3452
 4.3341
 4.3237
 4.3039
 4.2963
 4.2812
 4.2614
 4.2464
 4.2383
 4.2239
 4.2078
 4.1879
 4.1672
 4.1161
 4.1012
 4.0940
 4.0808
 4.0531
 4.0446
 4.0373
 4.0220
 4.0139
 3.9978
 3.9846
 3.9751
 3.9550
 3.9487
 3.9359
 3.9290
 3.9182
 3.9083
 3.8807
 3.8592
 3.8396
 3.8095

Current Data Parameters
 NAME schung010903
 EXPNO 1
 PROCNO 1
 F2 - Acquisition Parameters
 Date_ 20030109
 Time 17.19
 INSTRUM spect
 PROBD 5 mm tx12 lh/l
 PULPROG zg
 TD 16384
 SOLVENT CDCl3
 NS 16
 DS 2
 SWH 5518.764 Hz
 FIDRES 0.336839 Hz
 AQ 1.4844404 sec
 RG 64
 DW 90.600 usec
 DE 15.50 usec
 TE 280.0 K
 D1 1.00000000 sec
 P1 4.00 usec
 SFO1 500.1322743 MHz
 NUC1 1H
 PL1 -2.00 dB
 F2 - Processing parameters
 SI 16384
 SF 500.1300141 MHz
 MDW no
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

Current Data Parameters
NAME schung010903
EXPNO 11
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030109
Time 19:29
INSTRUM spect
PROBHD 5 mm txi2 1h/1
PULPROG zg
TD 16384
SOLVENT CDCl3
NS 16
DS 2
SWH 5518.764 Hz
FIDRES 0.336839 Hz
AQ 1.484404 sec
RG 64
DW 90.600 usec
DE 15.50 usec
TE 280.0 K
D1 1.00000000 sec
P1 4.00 usec
SFO1 500.1322743 MHz
NUC1 1H
PL1 -2.00 dB

F2 - Processing parameters
SI 16384
SF 500.1300141 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters
NAME LTC-268-NOE
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20010621
Time 20:38
INSTRUM spect
PROBHD 5 mm txi2 1h/v1
PULPROG zg
TD 16384
SOLVENT CDCl3
NS 8
DS 2
SWH 5518.764 Hz
FIDRES 0.336839 Hz
AQ 1.484404 sec
RG 64
DW 90.600 usec
DE 15.50 usec
TE 280.0 K
D1 2.0000000 sec
P1 3.00 usec
SFO1 500.132506 MHz
NUC1 1H
PL1 -2.00 dB

F2 - Processing parameters
SI 16384
SF 500.1300233 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Current Data Parameters
NAME LJC-803-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20021107
Time 17:17
INSTRUM spect
PROBHD 5 mm txi2 1h/v1
PULPROG zg
TD 16384
SOLVENT CDCl3
NS 16
DS 2
SWH 5518.764 Hz
FIDRES 0.336839 Hz
AQ 1.484404 sec
RG 64
DW 90.600 usec
DE 15.50 usec
TE 280.0 K
D1 1.0000000 sec
P1 4.00 usec
SFO1 500.1322463 MHz
NUC1 1H
PL1 -2.00 dB

F2 - Processing parameters
SI 16384
SF 500.1300236 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

```
Current Data Parameters
NAME      schung012803
EXPNO     1
PROCNO    1
```

F2 - Acquisition Parameters	
Date_	20030128
Time_	22:41
INSTRUM	ac241
PROBHD	5 mm QNP 1H/1
PULPROG	zg30
TD	16384
SOLVENT	CDCl3
NS	1
DS	2
SWH	518.766 Hz
FIDRES	0.336839 Hz
AQ	1.4844357 sec
RG	64
DW	90.600 usec
DE	15.50 usec
TE	280.0 K
D1	1.00000000 sec
F1	4.00 usec
SFO1	500.1324798 MHz
NUC1	1H
PL1	-2.00 dB

F2 - Processing parameters	
SI	16384
SF	500.1300236 MHz
WDW	no
SSB	0
LB	0.00 Hz
GB	0
PC	1.00

Current Data Parameters
NAME LJC-442-P
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20010803
Time 17.02
INSTRUM spect
PROBHD 5 mm Multinu
PULPROG zg
TD 32768
SOLVENT CDCl3
NS 59
DS 2
SWH 6410.256 Hz
FIDRES 0.195625 Hz
AQ 2.5559540 sec
RG 256
DW 78.000 usec
DE 111.43 usec
TE 300.0 K
HL1 3 dB
D1 1.00000000 sec
P1 6.00 usec
SF01 400.1319596 MHz
NUCLEUS 1H

F2 - Processing parameters
SI 32768
SF 400.1300176 MHz
WDW no
SSB 0
LB 0.00 Hz
GB 0
PC 1.00

Voyager Spec #1=>BC=>NR(2.00)=>DI=>MC[BP = 1570.7, 84309]

Voyager Spec #1=>BC=>NR(2.00)[BP = 875.4, 2939]

Mass (m/z)

<<0530_schung_LJC-843-P_DHB_0001>> Voyager Spec #1=>BC=>NR(2.00)[BP = 875.4, 2939]

Mass (m/z)

Voyager Spec #1=>BC=>NR(2.00)[BP = 2645.4, 2615]

Mass (m/z)

<<0530_schung_LJC-877-P_DHB_0001>> Voyager Spec #1=>BC=>NR(2.00)[BP = 2645.4, 2615]

Mass (m/z)

0625_LJC-667 #53-62 RT: 1.11-1.34 AV: 10 NL: 7.62E6

T: - c Full ms [50.00-2000.00]

LJC-664

1 RT: 0.00 P: - NL: 2.62E5
T: - c Full ms [50.00-2000.00]

LJC-880-P

Current Data Parameters
 NAME schung052203
 EXPNO 11
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20030522
 Time 22.01
 INSTRUM spect
 PROBHD 5 mm b2 1h/1
 PULPROG zgpgpr
 TD 16384
 SOLVENT D2O
 NS 16
 DS 4
 SWH 4990.020 Hz
 FIDRES 0.304567 Hz
 AQ 1.6417269 sec
 RG 406.4
 DW 100.200 usec
 DE 15.50 usec
 TE 280.0 K
 D12 0.00002000 sec
 PL9 65.00 dB
 D1 1.00000000 sec
 SFO1 500.1323196 MHz
 NUC1 ¹H
 PL1 -2.00 dB
 D13 0.00000300 sec
 P1 11.60 usec

F2 - Processing parameters
 SI 16384
 SF 500.1239184 MHz
 WDW no
 SSB 0
 LB 0.00 Hz
 GB 0
 PC 1.00

0829_LJC_489-23' #45-48 RT: 0.98-1.03 AV: 3 NL: 9.34E3

T: -c Full ms [800.00-2000.00]