Synthesis of Jenamidines A₁/A₂

Barry B. Snider* and Jeremy R. Duvall

Department of Chemistry, MS 015, Brandeis University, Waltham, Massachusetts 02454-9110.

Supporting Material

Experimental ProceduresS2-S8

Copies of ¹H and ¹³C NMR Spectra.....S9-S35

***N*-Cyano-*N'*-methoxy-*N'*-methylprolinamide (12).** *N*-Boc Weinreb amide **11** (316 mg, 1.2 mmol) was stirred in 4 mL of 1:1 TFA:CH₂Cl₂ solution for 18 h and concentrated. A solution of the resulting oil in 4 mL of EtOH was added to a solution of sodium bicarbonate (500 mg, 6.0 mmol) and cyanogen bromide (140 mg, 1.32 mmol) in 9 mL of EtOH. The mixture was stirred for 2 h at 25 °C, treated with 2 mL of H₂O, stirred for 15 min, and extracted with CH₂Cl₂. The combined organic extracts were dried over MgSO₄ and concentrated. Flash chromatography on silica gel (98:2 CH₂Cl₂:MeOH) gave 185 mg (84%) of nitrile **12**: ¹H NMR (CDCl₃) 4.57 (dd, 1, *J* = 3.7, 8.5), 3.74 (s, 3), 3.62-3.70 (m, 1), 3.45-3.55 (m, 1), 3.23 (s, 3), 2.20-2.28 (m, 1), 1.90-2.10 (m, 3).

***t*-Butyl 2-(1-Carbamoyl-pyrrolidine)-3-hydroxypropenoate (16), *t*-Butyl (2Z)-(2,5,6,7-Tetrahydro-3-oxo-3*H*-pyrrolo[1,2-*c*]imidazol-1-ylidene)acetate (15), and *t*-Butyl 1-(2,5,6,7-Tetrahydro-3-oxo-3*H*-pyrrolo[1,2-*c*]imidazole)acetate (14).** *t*-Butyl acetate (0.18 mL, 1.3 mmol) was added to a freshly prepared solution of LDA (1.3 mmol in 4 mL THF) at -78 °C under N₂ and the mixture was stirred for 15 min. Weinreb amide nitrile **12** (215 mg, 1.17 mmol) was added in 2 mL of dry THF. The mixture was stirred for 2 h at -78 °C, quenched with 5% HCl solution, and extracted with CH₂Cl₂. The combined organic extracts were dried over MgSO₄ and concentrated. Flash chromatography on silica gel (98:2 CH₂Cl₂:MeOH) gave 90 mg (30%) of **16**, followed by 50 mg (21%) of **15**, and 75 mg (32%) of **14**. A solution of urea **15** in CDCl₃ containing several drops of TFA completely converted **15** to **14** over 2 h.

Data for **14**: ¹H NMR (CDCl₃) 9.30 (br s, 1, OH), 3.66 (t, 2, *J* = 6.7), 3.27 (s, 2), 2.66 (t, 2, *J* = 7.3), 2.40 (tt, 2, *J* = 6.7, 7.3), 1.45 (s, 9); ¹³C NMR (CDCl₃) 169.0, 151.1, 124.6, 105.4, 81.5, 41.8, 32.2, 28.3, 27.9 (3 C), 22.0.

Data for **15**: ¹H NMR (CDCl₃) 8.70 (br s, 1, NH), 5.19 (s, 1), 4.87 (dd, 1, *J* = 7.3, 7.3), 3.59 (ddd, 1, *J* = 8.0, 8.0, 11.0), 3.24-3.34 (m, 1), 2.58-2.66 (m, 1), 2.00-2.10 (m, 2), 1.40-1.52 (m, 1), 1.46 (s, 9); ¹³C NMR (CDCl₃) 166.5, 161.8, 155.1, 91.8, 79.5, 64.6, 44.8, 30.1, 28.3 (3 C), 26.4; HRMS (DCI/NH₃) calcd for C₁₂H₁₉N₂O₃ (MH⁺) 239.1396, found 239.1402.

Data for **16**: ¹H NMR (CDCl₃) 8.95 (br s, 1, NH), 4.87 (s, 1), 4.33 (dd, 1, *J* = 7.9, 7.9), 3.59 (ddd, 1, *J* = 7.9, 7.9, 11.0), 3.15-3.22 (m, 1), 1.95-2.20 (m, 3), 1.55-1.65 (m, 1), 1.48 (s, 9).

***t*-Butyl 3-Amino-5,6,7,7a-tetrahydro-1-oxo-1*H*-pyrrolizine-2-carboxylate (**19**).**

t-Butyl acetate (1.5 mL, 11.1 mmol) was added dropwise over a period of 10 min to a freshly prepared solution of LDA (10 mmol in 27 mL THF) at -45 °C under N₂. The solution was stirred for 15 min and treated with methyl (*S*)-1-cyano-2-pyrrolidinecarboxylate (**17**)⁸ (680 mg, 4.4 mmol) in 5 mL of dry THF. The mixture was stirred for 1 h at -45 °C, treated dropwise with LHMDS (5 mL, 1.0 M in THF) and allowed to warm to 25 °C. The mixture was stirred for 2 h at 25 °C, quenched with H₂O, and extracted with CH₂Cl₂. The combined organic extracts were dried over MgSO₄ and concentrated. Flash chromatography on silica gel (95:5 CH₂Cl₂:MeOH) gave 250 mg (24%) of byproduct **21** followed by 280 mg (27%) of pure **19**.

Data for **21**: ¹H NMR (CDCl₃) 9.95 (br s, 1, NH), 4.34 (s, 1), 3.98 (dd, 1, *J* = 8.6, 8.6), 3.31-3.38 (m, 1), 3.21 (ddd, 1, *J* = 7.9, 7.9, 10.4), 1.98-2.26 (m, 3), 1.64-1.75 (m, 1), 1.47 (s, 9); ¹³C NMR (CDCl₃) 173.4, 169.4, 160.9, 79.1, 72.6, 63.6, 49.9, 28.4 (3 C), 27.3, 27.1. A 1D NOESY experiment showed an NOE from the alkene hydrogen (δ 4.34) to the *t*-butyl hydrogens (δ 1.47) and the methylene group adjacent to the ring fusion nitrogen (δ 3.21 and δ 3.31-3.38) establishing that the stereochemistry is *E* as drawn.

Data for **19**: mp 200-203 °C; [α]_D²² -66.9 (*c* 1.2, MeOH); ¹H NMR (CDCl₃) 8.00 (br, 1, NH), 5.42 (br, 1, NH), 3.87 (dd, 1, *J* = 6.7, 9.8), 3.34-3.42 (m, 1), 3.20-3.28 (m, 1), 2.06-2.26 (m, 3), 1.54-1.60 (m, 1), 1.55 (s, 9); ¹³C NMR (CDCl₃) 193.1, 174.0, 164.9, 90.5, 79.9, 69.7, 46.6, 28.5 (3 C), 27.6, 26.7; IR (KBr) 3475, 3077, 1714, 1638; HRMS (DEI) calcd for C₁₂H₁₈N₂O₃ (M⁺) 238.1317, found 238.1315.

3-(2-Methyl-2*E*-butenoylamino)-5,6,7,7a-tetrahydro-1*H*-pyrrolizin-1-one (24**).** 2-Methyl-2*E*-butenoyl chloride (27 mg, 0.23 mmol) was treated with **19** (25 mg, 0.10 mmol) and NaH (10 mg, 0.25 mmol) as described below for the preparation of **32d** to give 15 mg (69%) of **24** after flash chromatography on silica gel (98:2 CH₂Cl₂/MeOH): [α]_D²² 1.3 (*c* 0.7, MeOH); ¹H NMR (CD₃OD) 6.60 (q, 1, *J* = 6.7), 5.65 (s, 1), 3.95 (dd, 1, *J* = 8.2, 8.2), 3.42-3.50 (m, 1), 3.22

(ddd, 1, $J = 7.3, 7.3, 11.0$), 2.06-2.22 (m, 3), 1.88 (s, 3), 1.87 (d, 3, $J = 6.7$), 1.50-1.60 (m, 1); ^{13}C NMR (CD_3OD) 204.5, 173.7, 169.9, 136.3, 133.2, (93.6, 93.5), 70.7, 28.8, (27.5, 27.4), 14.4, 12.3 (one C is obscured by the CD_3OD peak between 49-50).

***t*-Butyl (4*S*)-4-[[*(1,1*-Dimethylethyl)dimethylsilyl]oxy]-2-methylpent-2*E*-enoate (**27**).**

A solution of (1-*t*-butoxycarbonylethylidene)triphenylphosphorane (**25**)¹² (1.9 g, 4.7 mmol) in 18 mL of dry CH_2Cl_2 was added to 2-[[*(1,1*-dimethylethyl)dimethylsilyl]oxy]-propanal (**26**)¹³ (730 mg, 3.8 mmol) in 8 mL of dry CH_2Cl_2 . The mixture was stirred at 25 °C for 2 h and quenched with 25 mL of H_2O . The layers were separated and the organic layer was dried over MgSO_4 and concentrated. Flash chromatography on silica gel (98:2 hexanes:EtOAc) yielded 770 mg (67%) of **27**: ^1H NMR (CDCl_3) 6.57 (br d, 1, $J = 8.5$), 4.59 (dq, 1, $J = 8.5, 6.7$), 1.78 (d, 3, $J = 1.2$), 1.49 (s, 9), 1.22 (d, 3, $J = 6.7$), 0.88 (s, 9), 0.05 (s, 3), 0.04 (s, 3); ^{13}C NMR (CDCl_3) 167.3, 144.7, 126.8, 80.1, 66.0, 28.0 (3 C), 25.8 (3C), 23.4, 18.1, 12.5, -4.7, -4.8; IR (neat) 1710; HRMS (DCI/ NH_3) calcd for $\text{C}_{16}\text{H}_{36}\text{NO}_3\text{Si}$ (MNH_4^+) 318.2464, found 318.2455.

***t*-Butyl (5*S*)-4-Hydroxy-2-methylpent-2*E*-enoate (**28**).** Silyl ether **27** (300 mg, 1.0 mmol) in 2 mL of THF was added to 18 mL of a 1.4 M solution of pyridine•HF (5 mL pyridine•(HF)_x, 20 mL THF, 20 mL pyridine). The mixture was stirred for 4 h in an Eppendorf tube, slowly quenched with saturated sodium bicarbonate solution, and extracted with EtOAc. The combined organic extracts were dried over MgSO_4 and concentrated to yield 185 mg (99%) of pure **28**: ^1H NMR (CDCl_3) 6.58 (br d, 1, $J = 8.3$), 4.67 (dq, 1, $J = 8.3, 6.1$), 1.83 (d, 3, $J = 1.5$), 1.49 (s, 9), 1.31 (d, 3, $J = 6.1$); ^{13}C NMR (CDCl_3) 167.2, 143.0, 129.0, 80.5, 64.9, 28.0 (3 C), 22.6, 12.6; IR (neat) 3421, 1707; HRMS (DCI/ NH_3) calcd for $\text{C}_{10}\text{H}_{22}\text{NO}_3$ (MNH_4^+) 204.1600, found 204.1606.

***t*-Butyl (5*S*)-4-Methoxyacetoxy-2-methylpent-2*E*-enoate (**29d**).** Methoxyacetyl chloride (0.15 mL, 1.6 mmol) was added slowly to a solution of alcohol **28** (185 mg, 0.99 mmol), pyridine (0.17 mL, 2.12 mmol), and DMAP (12 mg, 0.1 mmol) in 15 mL of dry THF under N_2 resulting in a white precipitate. The mixture was stirred for 2 h at 25 °C, filtered, diluted with EtOAc, washed successively with 2 M HCl, water, saturated sodium bicarbonate solution, and

water, dried over MgSO₄, and concentrated to give 255 mg (99%) of pure **29d**: ¹H NMR (CDCl₃) 6.51 (br d, 1, *J* = 8.9), 5.73 (dq, 1, *J* = 8.9, 6.1), 4.03 (s, 2), 3.45 (s, 3), 1.88 (d, 3, *J* = 1.5), 1.49 (s, 9), 1.36 (d, 3, *J* = 6.1); ¹³C (CDCl₃) 169.4, 166.6, 137.6, 131.3, 80.7, 69.8, 68.2, 59.3, 27.9 (3 C), 19.7, 12.8; IR (neat) 1756, 1709; HRMS (DCI/NH₃) calcd for C₁₃H₂₆NO₅ (MNH₄⁺) 276.1811, found 276.1809.

(S)-4-Methoxyacetoxy-2-methylpent-2E-enoic acid (30d). A solution of ester **29d** (255 mg, 0.98 mmol) in 5 mL of 9:1 CH₂Cl₂:TFA was stirred for 12 h at 25 °C. The mixture was diluted with H₂O and extracted with CH₂Cl₂. The combined organic extracts were dried over MgSO₄ and concentrated to give 194 mg (98%) of pure **30d**: ¹H NMR (CDCl₃) 11.0 (br s, 1, OH), 6.74 (d, 1, *J* = 8.8), 5.74 (dq, 1, *J* = 8.8, 6.4), 4.05 (s, 2), 3.45 (d, 3, *J* = 1.9), 1.93 (s, 3), 1.38 (dd, 3, *J* = 1.9, 6.4); ¹³C NMR (CDCl₃) 172.8, 169.5, 141.3, 129.0, 69.7, 68.1, 59.3, 19.4, 12.4; IR (neat) 3187, 1753, 1696; HRMS (DCI/NH₃) calcd for C₉H₁₈NO₅ (MNH₄⁺) 220.1185, found 220.1189.

(S)-4-Methoxyacetoxy-2-methylpent-2E-enoyl chloride (31d). A solution of acid **30d** (194 mg, 0.96 mmol) in 8 mL of oxalyl chloride was stirred for 4 h at 25 °C under N₂. Excess oxalyl chloride was removed under reduced pressure to give acid chloride **31d**, which was used immediately: ¹H NMR (CDCl₃) 6.98 (d, 1, *J* = 8.6), 5.73 (dq, 1, *J* = 8.6, 6.7), 4.06 (s, 2), 3.46 (s, 3), 2.00 (3, s), 1.42 (d, 3, *J* = 6.7).

Jenamidines A₁/A₂ 4'-Acetate (32a). (S)-4-Acetoxy-2-methylpent-2E-enoyl chloride (**31a**, 80 mg, 0.44 mmol) (prepared as described for **31d** using AcCl instead of methoxyacetyl chloride) was treated with **19** (36 mg, 0.15 mmol) and NaH (14 mg, 0.33 mmol) as described below for the preparation of **32d** to give 37 mg (84%) of **32a** as a 1:1 mixture of diastereomers upon flash chromatography (95:5 CH₂Cl₂:MeOH): [α]_D²² -27.5 (*c* 0.8, MeOH); ¹H NMR (CD₃OD) 6.28-6.38 (m, 1), 5.59-5.68 (m, 1), 5.62 (s, 1), 3.94 (dd, 1, *J* = 8.5, 7.9), 3.39-3.46 (m, 1), 3.16-3.24 (m, 1), 2.06-2.20 (m, 3), 2.03 (s, 3 × 0.5), 2.02 (s, 3 × 0.5), 1.95 (s, 3 × 0.5), 1.94 (s, 3 × 0.5), 1.48-1.57 (m, 1), 1.35 (d, 3 × 0.5, *J* = 6.7), 1.34 (d, 3 × 0.5, *J* = 6.7); ¹³C NMR (CD₃OD) 204.6, 173.5, 172.0, (169.5, 169.4), (138.74, 139.69), 133.7, 93.9, 70.8, (68.9, 68.8),

28.8, 27.5, 21.0, 19.7, 13.2, one C is obscured by the CD₃OD peak between 49-50; HRMS (MeOH/NBA) calcd for C₁₅H₂₁N₂O₄ (MH⁺) 293.1501, found 293.1504.

Jenamidines A₁/A₂ (3), Jenamidines A₁/A₂ 4'-Methoxyacetate (32d), and 3-Amino-5,6,7,7a-tetrahydro-1H-pyrrolizin-1-one (33). Vinylogous amide **19** (106 mg, 0.44 mmol) and NaH (60% dispersion in mineral oil, 45 mg, 1.1 mmol) were stirred in 10 mL of dry THF under N₂ for 10 min. Acid chloride **31d** (0.096 mmol) in 3 mL of dry THF was added and the resulting solution was stirred for 2 h, quenched with brine, and extracted with CH₂Cl₂. The combined organic extracts were dried over MgSO₄ and concentrated to give 196 mg of a mixture of mono and bis acylated products. A solution of the mixture in 10 mL of 9:1 CH₂Cl₂:TFA was stirred for 15 h, neutralized with saturated sodium bicarbonate solution, saturated with NaCl and extracted with EtOAc. The combined organic extracts were dried over MgSO₄ and concentrated to afford jenamidines A₁/A₂ 4'-methoxyacetate (**32d**) as the major product and no jenamidines A₁/A₂ (**3**), as expected. Flash chromatography on silica gel (97:3 CH₂Cl₂:MeOH to 90:10 CH₂Cl₂:MeOH) gave 55 mg (39%) of **32d**, followed by 20 mg (18%) of **3** and a complex mixture of products resulting from the decomposition of **32d**.

A solution of a second batch of crude **32d** (prepared as before from 96 mg (0.40 mmol) of **19**, 42 mg (1.0 mmol) of NaH, and 0.96 mmol of **31d** followed by TFA hydrolysis) was stirred in 10 mL of MeOH containing 10 drops of H₂O and 200 mg of Na₂CO₃ at 0 °C for 24 h. The mixture was filtered through silica gel and concentrated to give a 4:1:1 mixture of **3**, **32d**, and **33**. Flash chromatography on silica gel (95:5 CH₂Cl₂:MeOH) gave 14 mg (11% from **19**) of **32d** and 45 mg (45% from **19**) of **3**, while **33** was recovered along with a complex mixture of products upon flushing the silica gel with MeOH.

Data for **32d** as a 1:1 mixture of diastereomers: ¹H NMR (CD₃OD) 6.345 (br d, 1 × 0.5, *J* = 8.0), 6.340 (br d, 1 × 0.5, *J* = 8.0), 5.72-5.80 (m, 1), 5.64 (s, 1), 4.07 (s, 2 × 0.5), 4.06 (s, 2 × 0.5), 3.95 (dd, 1, *J* = 8.8, 8.8), 3.40-3.47 (m, 1), 3.42 (s, 3 × 0.5), 3.41 (s, 3 × 0.5), 3.20-3.26 (m, 1), 2.10-2.25 (m, 3), 1.994 (s, 3 × 0.5), 1.991 (s, 3 × 0.5), 1.50-1.60 (m, 1), 1.41 (d, 3 × 0.5, *J* = 6.1), 1.40 (d, 3 × 0.5, *J* = 6.1); ¹³C NMR (CD₃OD) 204.7, 173.4, (171.39, 171.37), (169.4,

169.3), (138.04, 138.00), (134.20, 134.18), 93.9, 70.7, 70.5, (69.4, 69.3), (59.54, 59.52), 49.3, 28.8, 27.5, 19.8, (13.29, 13.26); IR (neat) 1744, 1702, 1636, 1563, 1506; HRMS (DCI/NH₃) calcd for C₁₆H₂₃N₂O₅ (MH⁺) 323.1607, found 323.1609.

Data for jenamidines A₁/A₂ (**3**) as a 1:1 mixture of diastereomers: $[\alpha]_D^{22}$ 4.2 (*c* 0.6, MeOH), lit.¹ $[\alpha]_D^{22}$ 6.8 (*c* 0.7, MeOH); UV $\lambda_{\max}^{\text{MeOH}}$ nm (log ϵ) 240 (4.29), 282 (3.84), 326 (4.02); ¹H NMR (CD₃OD) 6.37 (d, 1, *J* = 7.9), 5.67 (s, 1 × 0.5), 5.63 (s, 1 × 0.5), 4.67 (dq, 1, *J* = 6.5, 7.9), 3.92-3.98 (m, 1), 3.42-3.49 (m, 1), 3.18-3.26 (m, 1), 2.08-2.24 (m, 3), 1.92 (s, 3), 1.50-1.60 (m, 1), 1.31 (d, 3 × 0.5, *J* = 6.5), 1.29 (d, 3 × 0.5, *J* = 6.5); ¹³C NMR (CD₃OD) (204.69, 204.57), (173.61, 173.48), (169.84, 169.82), (143.67, 143.65), (131.62, 131.60), (93.82, 93.80, 93.76, 93.75), (70.77, 70.67), (65.31, 65.25), 49.29, 28.80, (27.51, 27.47, 27.40, 27.37), (22.68, 22.64), 12.92; IR (neat) 2973, 1699, 1634, 1557, 1504, 1404, 1244, 1139, 1059; HRMS (DCI/NH₃) calcd for C₁₃H₁₉N₂O₃ (MH⁺) 251.1396, found 251.1394. The ¹H and ¹³C NMR spectrum are identical to those of the natural product. H-7a is approximately 50% exchanged with deuterium. C-2 (93.7-93.8) and C-7 (27.3-27.5) absorb as four peaks since the carbon for each diastereomer absorbs separately for the protio and deuterio isomers due to the isotope shift.¹⁵ As was noted for the natural product, H-2 slowly exchanges with CD₃OD over the course of several hours.

Pure **32d** (30 mg, 0.093 mmol) was stirred in 2 mL of MeOH containing 5 drops of H₂O and 50 mg of Na₂CO₃ at 0 °C for 6 h. The mixture was filtered through silica gel and concentrated. Flash chromatography on silica gel (90:10 CH₂Cl₂:MeOH) gave 6 mg (20%) of recovered **32d**, followed by 13 mg (56%, 70% based on recovered **32d**) of **3** and 2.5 mg (18%) of **33**.

Data for **33**: ¹H NMR (CD₃OD) 4.50 (s, 1), 3.90 (dd, 1, *J* = 6.7, 9.1), 3.28-3.35 (m, 1), 3.12-3.20 (m, 1), 2.05-2.25 (m, 3), 1.40-1.48 (m, 1); ¹³C NMR (CD₃OD) 197.5, 178.5, 72.1, 47.7, 29.3, 28.2 (C-2 was not observed because H-2 exchanged with D); HRMS (DEI) calcd for C₇H₁₀N₂O (M⁺) 138.0793, found 138.0789.

Preparation of 19 from Cbz-Proline *N*-Hydroxysuccinimide Ester (37). *t*-Butyl cyanoacetate (0.68 mL, 4.8 mmol) was added dropwise to a solution of NaH (160 mg, 4.0 mmol, 60% dispersion in mineral oil) in 8 mL of dry benzene at RT. The resulting slurry was stirred for 1 h and treated with **37** (554 mg, 1.6 mmol) in 2 mL of dry benzene. The mixture was stirred for an additional 3 h at 25 °C, quenched with H₂O, and extracted with Et₂O. The aqueous layer was acidified with 2 M HCl resulting in the formation of a white precipitate, which dissolved upon addition of CH₂Cl₂. The layers were separated and the aqueous layer was extracted with CH₂Cl₂. The combined CH₂Cl₂ extracts were dried over MgSO₄ and concentrated to give **38**.

A solution of crude **38** in 25 mL of MeOH was stirred with 10% Pd/C (170 mg, 0.16 mmol) under 1 atm of H₂ at 25 °C for 2 h. The mixture was filtered through Celite and concentrated to give crude **39**, which cyclized on standing for 24 h at 25 °C to give crude **19**. Flash chromatography on silica gel (95:5 EtOAc/MeOH) gave 190 mg (50%) of **19** with spectral data identical to those of a sample obtained from **17**: $[\alpha]_D^{22}$ -62.2 (*c* 1.1, MeOH).