Supporting Information

Synthesis of N-confused Phlorins via Addition/Cyclization Pathway

Xiaofang Li, ¹ Piotr J. Chmielewski, ² Junfeng Xiang, ³ Jialiang Xu, ¹ Li Jiang, ¹ Yuliang Li*, ¹ Huibiao Liu, ¹ Daoben Zhu¹

1 CAS Key Laboratory of Organic Solids, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100080, P.R. China

2 Department of Chemistry, University of Wrocław, 14 F. Joliot-Curie Street, 50 383 Wrocław, Poland

3 Center for Physiochemical Analysis & Measurement, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100080, P. R. China

E-mail: ylli@iccas.ac.cn; zhudb@iccas.ac.cn

29 Pages

Table of contents

General Experimental Methods	S2-4
Selected data for 4b-g	S4-6
Selected data for 6b-d	S6-8
NMR spectra of 4a-g	S8-18
NMR spectra of 6a-d	S18-29
UV-Vis spectrum of 6a	S29

General Experimental Methods. Instrumentation. Absorption spectra were recorded on a Hitachi U-3010 spectrometer. Mass spectra were recorded on a Bruker Biflex □ MALDI-TOF spectrometer and APEX II FT-ICRMS spectrometer. NMR spectra were recorded on a Bruker Avance 600 and Bruker Avance 400 spectrometer. The 1D and 2D experiments (COSY, NOESY, HMQC, and HMBC) were performed by means of standard experimental procedures of the Bruker library. The peaks were referenced to the residual CHCl₃ resonances in ¹H and ¹³C NMR (7.26 ppm and 77.2 ppm, respectively).

Preparation of the precursor: 5, 10, 15, 20-Tetrakis (*p*-tolyl)-2-aza-21-carbaporphyrin **1a**, 5, 10, 15, 20-Tetraphenyl-2-aza-21-carbaporphyrin **1b**: These compounds were prepared according to Lindsey's procedure.¹ 5, 10, 15, 20-Tetraphenyl-2-aza-21-nitro-21-carbaporphyrin **1c** was synthesized according to the Ishikawa's procedure.² Preparation of 5,10,15,20-tetrakis (3,4,5-trimethoxyphenyl)-2-Aza-21-carbaporphyrin (**1d**): To a solution of pyrrole (0.52 mL, 7.5 mmol) and 3,4,5-trimethoxybenzaldehyde (1.47 g, 7.5 mmol) in CH₂Cl₂ (750 mL) was added methanesulfonic acid (0.34 mL, 5.2 mmol). The mixture was stirred for 30 min after which 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ)(1.50 g, 6.6 mmol) was added. After 1 min, triethylamine (1.5 mL) was added. The crude reaction mixture was chromatographed with silica gel (200 mL) under vacuum and eluted with ethyl acetate. All eluent was collected as a single fraction, concentrated under vacuum and then absorbed onto 7.5 g of activity III basic alumina. The absorbed sample was added to the top of a column with 150 g activity III basic alumina in 100:0.2 CH₂Cl₂/CH₃OH. The polarity

1 Geier III, G. R.; Haynes, D. M.; Lindsey, J. S. *Org. Lett.* **1999**, *1*, 1455-1458.

2 Ishikawa, Y.; Yoshida, I.; Akaiwa, K.; Koguchi, E.; Sasaki, T.; Furuta, H. *Chem. Lett.* **1997**, 453-454.

of the eluent was increased from 100:0.2 to 100:1 CH₂Cl₂/CH₃OH, the **1d** was eluted with 100:1 CH₂Cl₂/CH₃OH. The solvent was removed under vacuum and the residue was triturated with CH₂Cl₂/hexanes to yield the product 450 mg (25%). ¹H NMR (400 MHz, CDCl₃, 298K) δ: -4.95 (s, 1H), -2.25 (br, 2H), 3.99 (s, 12H), 4.09 (s, 6H), 4.13 (s, 6H), 4.14 (s, 3H), 4.17 (s, 6H), 4.18 (s, 3H), 7.41 (s, 2H), 7.42 (s, 2H), 7.58 (s, 2H), 7.60 (s, 2H), 8.67-8.71 (m, 3H), 8.73 (d, *J* = 4.4 Hz, 1H), 8.87 (s, 1H), 8.99 (d, *J* = 4.4 Hz, 1H), 9.08 (d, *J* = 4.4 Hz, 1H). ¹³C NMR (100 MHz, CDCl₃, 298 K) δ: 56.2, 56.3, 56.4, 56.5, 56.6, 56.7, 61.1, 61.3, 99.5, 112.6, 112.7, 112.8, 112.9, 114.8, 115.0, 117.4, 118.9, 125.4, 125.7, 126.5, 127.8, 128.4, 128.6, 134.3, 134.7, 135.2, 135.5, 135.7, 137.3, 137.4, 137.9, 138.0, 138.6, 138.8, 140.0, 140.5, 149.4, 151.7, 152.1, 152.4, 155.7, 156.5, 157.4, 157.5. UV-vis (CHCl₃) λ_{max}/nm (logε): 446 (5.25), 544 (3.98), 588 (4.08), 730 (4.06). MS (MALDI-TOF): *m/z* 974.3 (974.4 for C₅₆H₅₄N₄O₁₂). N-phenylbenzenecarbohydrazonoyl chloride **2a**,³ N-phenyl-C-ethoxy-carbonylformohydrazidoyl chloride **2b**,⁴ N-(p-CH₃)-phenyl-C-ethoxycarbonylformohydrazidoyl bromide **2c**⁵ and N-(p-CH₃)-phenyl-C-acetylformohydrazidoyl bromide **2d**⁵ were prepared according to literatures.

General method for synthesis of 4a-g.

A solution of NCP **1** (0.1 mmol), hydrazonoyl halide **2** (0.2 mmol) and triethylamine (30 mg, 0.3 mmol) in dry benzene were stirred and refluxed under a nitrogen atmosphere for 5-26 h.

³ Wolkoff, P. *Can. J. Chem.* **1975**, 53, 1333-1336.

⁴ Shawali, A. S.; Albar, H. A. *Can. J. Chem.* **1986**, 64, 871-875.

⁵ Tewarl, R. S.; Parthar, P. *J. Chem. Eng. Data* **1981**, 26, 418-420.

The reaction progress was monitored by TLC (CH₂Cl₂ as developing solvent). After reaction completion (full consumption of **1**), the reaction mixture was cooled to the room temperature, and solvent was evaporated under vacuum. The residue was chromatographed on a silica gel column with CH₂Cl₂ as eluent. The fastest moving green band was collected and solvent was removed to give the N-confused phlorin **4**.

Selected data for 4b: ¹H NMR (400MHz,CDCl₃, 298K) δ: 4.58 (d, *J*=0.8 Hz, 1H), 6.51-6.52 (m, 2H), 6.55(d, *J*=4.8 Hz, 1H), 6.73(d, *J*=5.6 Hz, 1H), 6.78 (d, *J*=0.8 Hz, 1H), 6.85(d, *J*=4.8 Hz, 1H), 6.88(d, *J*=4.8 Hz, 1H), 7.07(d, *J*=5.6 Hz, 1H), 7.13 (m, 2H), 7.09-7.18(m, 3H) 7.28-7.34(m, 9H), 7.36-7.50(m, 12H), 7.59-7.62(m, 3H), 8.22(s, 1H), 8.69(s, 1H); ¹³C NMR (100MHz, CDCl₃, 298 K) δ: 66.2, 98.8, 107.7, 113.0, 116.3, 120.2, 122.0, 126.2, 127.3, 127.7, 127.9, 128.2, 128.3, 128.4, 128.7, 129.1, 131.2, 132.2, 134.6, 140.3, 140.5, 142.2, 146.1, 146.7, 152.0, 167.1. UV-vis (CHCl₃) λ_{max}/nm (logε):396 (4.50), 693 (4.30). HRMS (SIMS) calc. for C₅₇H₄₀N₆ [M⁺]: 808.3314, Found: 808.3300.

Selected data for 4c: ¹H NMR (600MHz,CDCl₃, 298K) δ: 1.39(t, *J*=7.2Hz, 3H), 4.30-4.33 (m, 1H), 4.42-4.50 (m, 1H), 4.51 (s, 1H), 6.52(m, 1H), 6.55(d, *J*=4.8 Hz, 1H), 6.61(m, 1H), 6.74(d, *J*=5.4Hz, 1H), 6.83(d, *J*=5.4Hz, 1H), 6.94(m, 1H), 7.08(d, *J*=4.8Hz, 1H), 7.11-7.14 (m, 5H), 7.19-7.20 (m, 2H), 7.30-7.33(m, 3H), 7.34-7.42(m, 8H), 7.46(d, *J*=6.6Hz, 2H), 7.54(d, *J*=7.8Hz, 4H), 7.82(s, 1H), 8.13(s, 1H), 8.57(s, 1H); ¹³C NMR (150MHz, CDCl₃, 298 K) δ: 14.3, 62.1, 67.2, 98.8, 107.5, 112.4, 118.9, 120.9, 122.7,123.5, 126.8, 127.3, 127.5, 127.7, 127.8, 127.9, 128.0, 128.1, 128.2, 128.3, 129.3, 131.4, 132.3, 135.0, 135.2, 141.9, 144.8, 146.9, 152.3, 160.3, 167.2. UV-vis (CHCl₃) λ_{max}/nm (logε):397 (4.57), 696 (4.37).

HRMS (SIMS) calc. for $C_{54}H_{40}N_6O_2$ [M^+]: 804.3213, Found: 804.3243.

Selected data for 4d: 1H NMR (600MHz, $CDCl_3$, 298K) δ : 1.35(t, $J=7.2$ Hz, 3H), 2.21(s, 3H), 4.26-4.32 (m, 1H), 4.36-4.42 (m, 1H), 4.45 (s, 1H), 6.52(m, 1H), 6.55(d, $J=4.8$ Hz, 1H), 6.59(m, 1H), 6.74(d, $J=5.4$ Hz, 1H), 6.83(d, $J=5.4$ Hz, 1H), 6.91-6.93(m, 2H), 7.08(d, $J=4.8$ Hz, 1H), 7.08-7.24 (m, 8H), 7.40-7.46(m, 11H), 7.54(d, $J=7.8$ Hz, 4H), 7.81(s, 1H), 8.11(s, 1H), 8.53(s, 1H); ^{13}C NMR (150MHz, $CDCl_3$, 298 K) δ : 14.3, 20.7, 29.7, 62.0, 67.2, 98.6, 107.5, 112.4, 118.8, 120.9, 122.6, 122.9, 126.8, 127.3, 127.5, 127.8, 127.9, 128.2, 128.3, 128.5, 129.3, 131.4, 131.7, 132.4, 133.2, 135.0, 135.2, 141.9, 142.6, 146.9, 152.2, 160.4, 167.1. UV-vis ($CHCl_3$) λ_{max}/nm (log ϵ): 394 (4.63), 693 (4.41). HRMS (SIMS) calc. for $C_{55}H_{42}N_6O_2$ [M^+]: 818.3369, Found: 818.3336.

Selected data for 4e: 1H NMR (400MHz, $CDCl_3$, 298K) δ : 6.51-6.55(m, 2H), 6.68-6.69(m, 3H), 6.86-6.87(m, 1H), 6.92-6.95(m, 1H), 7.05(d, $J=5.6$ Hz, 1H), 7.10-7.14 (m, 5H), 7.21-7.23 (m, 4H), 7.28-7.30 (m, 1H), 7.33-7.44(m, 15H), 7.47-7.49(m, 2H), 7.55(d, $J=7.6$ Hz, 2H), 7.86(br, 1H), 9.44(s, 1H); ^{13}C NMR (100MHz, $CDCl_3$, 298 K) δ : 66.9, 109.3, 111.2, 113.8, 114.1, 114.3, 118.1, 121.6, 123.8, 124.0, 126.3, 126.9, 127.4, 127.7, 127.8, 128.6, 128.8, 129.3, 129.9, 130.0, 130.9, 131.6, 132.3, 135.2, 135.3, 135.6, 136.6, 139.9, 140.0, 140.4, 145.3, 145.5, 148.4, 151.1, 166.2. UV-vis ($CHCl_3$) λ_{max}/nm (log ϵ): 390 (4.57), 664 (4.31). HRMS (SIMS) calc. for $C_{57}H_{39}N_7O_2$ [M^+]: 853.3165, Found: 853.3179.

Selected data for 4f: 1H NMR (600MHz, $CDCl_3$, 298K) δ : 2.23(s, 3H), 2.50(s, 3H), 6.51-6.52(m, 1H), 6.53(d, $J=4.8$ Hz, 1H), 6.62-6.63(m, 1H), 6.68(d, $J=5.4$ Hz, 1H), 6.83(d, $J=4.8$ Hz, 1H), 6.96(d, $J=8.4$ Hz, 2H), 7.07(d, $J=5.4$ Hz, 1H), 7.09-7.10(m, 3H), 7.22-7.24 (m,

3H), 7.28-7.29 (m, 2H), 7.32-7.35(m,3H), 7.36-7.42(m, 7H), 7.43-7.45(m, 2H), 7.58(d, $J=7.8\text{Hz}$, 2H), 7.81(br, 1H), 7.97(s, 1H), 9.42(s, 1H); ^{13}C NMR (150MHz, CDCl_3 , 298 K) δ : 20.9, 25.6, 68.1, 109.0, 112.9, 117.3, 120.7, 124.3, 126.9, 127.4, 127.8, 128.2, 128.3, 128.5, 128.6, 129.1, 129.6, 129.8, 130.0, 131.1, 132.3, 135.3, 135.9, 140.0, 144.7, 151.9, 166.8, 190.9. UV-vis (CHCl_3) $\lambda_{\text{max}}/\text{nm}$ ($\log\epsilon$):387 (4.72), 666 (4.42). HRMS (SIMS) calc. for $\text{C}_{54}\text{H}_{39}\text{N}_7\text{O}_3$ [M^+]: 833.3114, Found: 833.3141.

Selected data for 4g : ^1H NMR (600MHz, CDCl_3 , 298K) δ : 1.37(t, $J=7.2\text{Hz}$, 3H), 4.30-4.33 (m, 1H), 4.41-4.44 (m, 1H), 6.52-6.54(m, 2H), 6.68-6.70(m, 2H), 6.84(d, $J=4.2\text{Hz}$, 1H), 7.01-7.02(m, 1H), 7.06-7.09 (m, 4H), 7.12-7.15 (m, 2H), 7.23-7.24(m,2H), 7.33-7.46(m, 15H), 7.59(d, $J=7.2\text{Hz}$, 2H), 7.83(br, 1H), 7.85(s, 1H), 9.49(s, 1H); ^{13}C NMR (150MHz, CDCl_3 , 298 K) δ : 14.2, 62.4, 67.9, 109.1, 113.2, 115.4, 116.7, 120.9, 124.5, 125.2, 126.9, 127.4, 127.7, 127.8, 127.9, 128.2, 128.4, 129.0, 129.6, 131.0, 132.3, 135.3, 135.9, 136.0, 140.3, 144.3, 148.0, 151.8, 159.7, 166.8. UV-vis (CHCl_3) $\lambda_{\text{max}}/\text{nm}$ ($\log\epsilon$):389 (4.53), 666 (4.30). HRMS (SIMS) calc. for $\text{C}_{54}\text{H}_{39}\text{N}_7\text{O}_4$ [M^+]: 849.3064, Found: 849.3054.

General method for synthesis of 6a-d.

A sample of 5,10,15,20-tetraaryl-2-aza-21-carbaporphyrin **1** (0.1mmol) was placed in a flame-dried 25 mL two-neck round-bottom flask and dissolved in 10 mL of dry benzene under nitrogen. The dialkyl acetylenedicarboxylate **5** (0.2mmol) was added to the solution, which was stirred and refluxed for 3-10 h(for **1d**, 3 h; for **1a** and **1b**, 10 h). Lack of the **1** in the reaction mixture (TLC-controlled) indicated the reaction completion. The mixture was cooled to the room temperature, and solvent was evaporated under vacuum. The residue was

chromatographed on a silica gel column with CH₂Cl₂/ethyl acetate (90:10 v/v) as an eluent (in the case of compounds **6c** and **6d**, the polarity of the eluent was increased from CH₂Cl₂/hexanes (2:1 v/v) to CH₂Cl₂ to elute the product). The fastest moving brown band was collected and solvent was removed to give the N-confused phlorin **6** (**6b**, 73mg, yield 64%; **6c**, 48mg, yield 62%; **6d**, 56mg, yield 67%).

Selected data for 6b : ¹H NMR (400 MHz, CDCl₃, 298K) δ: 1.16 (t, *J* = 7.2 Hz, 3H), 1.37 (t, *J* = 7.2 Hz, 3H), 3.79 (s, 6H), 3.82 (s, 12H), 3.84 (s, 3H), 3.86 (s, 6H), 3.89 (s, 3H), 3.92 (s, 6H), 4.17 (q, *J* = 7.2 Hz, 2H), 4.41 (q, *J* = 7.2 Hz, 2H), 6.24 (d, *J* = 4.8 Hz, 1H), 6.28 (d, *J* = 4.8 Hz, 1H), 6.41 (d, *J* = 5.6 Hz, 1H), 6.47 (d, *J* = 4.0 Hz, 1H), 6.58 (d, *J* = 4.8 Hz, 1H), 6.55 (s, 2H), 6.56 (s, 2H), 6.64 (s, 2H), 6.65 (s, 2H), 6.71 (d, *J* = 5.6 Hz, 1H), 7.07 (s, 2H), 10.65 (br, 1H), 10.82 (br, 1H). ¹³C NMR (150 MHz, CDCl₃, 298 K) δ: 13.9, 14.2, 52.9, 56.2, 56.3, 56.4, 58.2, 60.0, 60.8, 60.9, 61.4, 62.9, 101.7, 105.7, 108.6, 109.0, 113.1, 118.6, 123.1, 127.6, 127.7, 128.7, 128.8, 130.5, 133.2, 135.2, 135.4, 135.6, 137.0, 137.6, 137.8, 137.9, 138.3, 138.8, 142.5, 152.2, 152.7, 152.8, 152.9, 153.1, 159.8, 162.3, 168.9. UV-vis (CHCl₃) λ_{max}/nm (logε): 412 (4.67), 760 (4.17). HRMS (SIMS) calc. for C₆₄H₆₄N₄O₁₆ [M⁺]: 1144.4317, Found: 1144.4303

Selected data for 6c : ¹H NMR (400 MHz, CDCl₃, 298K) δ: 1.10 (t, *J* = 7.2 Hz, 3H), 1.35 (t, *J* = 7.2 Hz, 3H), 4.08-4.15 (m, 2H), 4.40 (q, *J* = 7.2 Hz, 2H), 6.12(d, *J* = 4.8 Hz, 1H), 6.15(d, *J* = 3.6 Hz, 1H), 6.27 (d, *J* = 5.6 Hz, 1H), 6.31 (d, *J* = 3.6 Hz, 1H), 6.42 (d, *J* = 4.8 Hz, 1H), 6.59 (d, *J* = 5.6 Hz, 1H), 6.94 (s, 1H), 7.16 (s, 1H), 7.27-7.44 (m, 20H), 10.81 (br, 1H), 11.17 (br, 1H). ¹³C NMR (150 MHz, CDCl₃, 298 K) δ: 13.8, 13.9, 57.9, 61.2, 62.8,

101.8, 112.9, 117.7, 123.4, 126.6, 126.8, 127.3, 127.5, 127.6, 127.7, 127.8, 127.9, 128.0, 128.2, 128.5, 128.7, 130.5, 131.0, 131.2, 131.4, 131.9, 133.4, 135.7, 136.3, 138.8, 139.8, 140.0, 140.7, 142.8, 148.5, 152.8, 160.0, 162.2, 168.9. UV-vis (CHCl₃) λ_{max} /nm (log ϵ): 406 (4.80), 756 (4.26). HRMS (SIMS) calc. for C₅₂H₄₀N₄O₄ [M⁺]: 784.3050, Found: 784.3022.

Selected data for 6d : ¹H NMR (400 MHz, CDCl₃, 298K) δ : 1.13 (t, J = 7.2 Hz, 3H), 1.35 (t, J = 7.2 Hz, 3H), 2.33 (s, 3H), 2.37 (s, 3H), 2.38 (s, 3H), 2.40 (s, 3H), 4.10-4.16 (m, 2H), 4.38 (q, J = 7.2 Hz, 2H), 6.12(d, J = 4.8 Hz, 1H), 6.15(d, J = 3.6 Hz, 1H), 6.27 (d, J = 5.6 Hz, 1H), 6.31 (d, J = 3.6 Hz, 1H), 6.42 (d, J = 4.8 Hz, 1H), 6.59 (d, J = 5.6 Hz, 1H), 6.94 (s, 1H), 7.16 (s, 1H), 7.27-7.44 (m, 20H), 10.81 (br, 1H), 11.17 (br, 1H). ¹³C NMR (150 MHz, CDCl₃, 298 K) δ : 13.9, 14.0, 21.2, 21.3, 21.4, 21.5, 57.7, 61.2, 62.8, 101.8, 112.7, 117.7, 123.3, 126.5, 127.5, 128.0, 128.6, 128.7, 128.9, 129.2, 129.5, 131.0, 131.4, 133.5, 135.7, 136.3, 137.2, 137.6, 138.8, 140.1, 142.5, 148.5, 152.9, 160.1, 162.3, 169.2. UV-vis (CHCl₃) λ_{max} /nm (log ϵ): 406 (4.80), 756 (4.26). HRMS (SIMS) calc. for C₅₆H₄₈N₄O₄ [M⁺]: 840.3676, Found: 840.3649.

Figure S1. ¹H NMR spectrum of **4a**, 223 K, CDCl₃.

Figure S2. ^{13}C NMR spectrum of **4a**, 298 K, CDCl_3 .

Figure S3. COSY spectrum of **4a** (CDCl_3 , 223 K).

Figure S4. NOESY spectrum (CDCl₃, 223 K) and correlation scheme for **4a**.

Figure S5. HMQC spectrum of **4a** (CDCl_3 , 298 K)

Figure S6. HMBC spectrum of **4a** (CDCl₃, 298 K)

Figure S7. ^1H NMR spectrum of **4b** (298 K, CDCl_3).

Figure S8. ^{13}C NMR spectrum of **4b** (298 K, CDCl_3).

Figure S9. ^1H NMR spectrum of **4c** (298 K, CDCl_3).

Figure S10. ^{13}C NMR spectrum of **4c** (298 K, CDCl_3)

Figure S11. ^1H NMR spectrum of **4d** (298 K, CDCl_3).

Figure S12. ^{13}C NMR spectrum of **4d** (298 K, CDCl_3)

Figure S13. ^1H NMR spectrum of **4e** (298 K, CDCl_3).

Figure S14. ^{13}C NMR spectrum of **4e** (298 K, CDCl_3)

Figure S15. ^1H NMR spectrum of **4f** (298 K, CDCl_3).

Figure S16. ^{13}C NMR spectrum of **4f** (298 K, CDCl_3).

Figure S17. ^1H NMR spectrum of **4g** (298 K, CDCl_3).

Figure S18. ^{13}C NMR spectrum of **4g** (298 K, CDCl_3).

Figure S19. ^1H NMR spectrum of **6a** (223 K, CDCl_3).

Figure S20. ^1H NMR spectrum of **6a** (298 K, CDCl_3).

Figure S21. ^{13}C NMR spectrum of **6a** (298 K, CDCl_3).

Figure S22. NOESY spectrum (223 K, CDCl₃) and correlation scheme of **6a**.

Figure S23. COSY spectrum of **6a** (223 K, CDCl_3).

Figure S24. HMQC spectrum of **6a** (298 K, CDCl_3).

Figure S25. HMBC spectrum of **6a** (298 K, CDCl₃).

Figure S26. ^1H NMR spectrum of **6b** (298 K, CDCl_3)

Figure S27. ^{13}C NMR spectrum of **6b** (298 K, CDCl_3).

Figure S28. ^1H NMR spectrum of **6c** (298 K, CDCl_3).

Figure S29. ^1H NMR spectrum of **6c** (223 K, CDCl_3).

Figure S30. ^{13}C NMR spectrum of **6c** (298 K, CDCl_3).

Figure S31. ^1H NMR spectrum of **6d** (298 K, CDCl_3).

Figure S32. ^{13}C NMR spectrum of **6d** (298 K, CDCl_3).

Figure S33. UV-Vis spectra of **6a** (solid line) and **1d** (dotted line) in CHCl_3 ($8.5 \times 10^{-6}\text{M}$).