Synthesis of Novel 11-Desmethyl Analogues of Laulimalide by Nozaki-Kishi Coupling

Ian Paterson,^{*,†} Hermann Bergmann,[†] Dirk Menche,[†] and Albrecht Berkessel[‡][†]University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW, UK and [‡]Institut für Organische Chemie der Universität zu Köln, Greinstrasse 4, D-50939 Köln, Germany

ip100@cam.ac.uk

Revised Supporting Information (20 pages): In the original version of the Supporting Information published March 19, 2004, the spectral data and copies of NMR spectra of compound **2** were incorrect. The corrected NMR data (S9–S10) and copies of spectra (S11–S20) are included here, together with detailed NMR comparisons with laulimalide.

(3Z,9E)-(1R,7S,11S,17R)-11-(tert-Butyl-dimethyl-silanyloxy)-7-(tert-butyl-diphenyl-silanyloxymethyl)-6,21-dioxo-bicyclo[15.3.1]henicosa-3,9,19-triene-5,13-dione

R_f = 0.25 (hexane/EtOAc = 80/20); **IR** $\tilde{\nu}$ (cm⁻¹, thin film) = 2930, 1719, 1113, 703; **¹H-NMR** (500 MHz, CDCl₃): δ = 0.04 (s, 6H), 0.88 (s, 9H), 1.03 (s, 9H), 1.30-1.40 (m, 1H), 1.45-1.55 (m, 1H), 1.57-1.64 (m, 2H), 1.83-2.00 (m, 2H), 2.12 (dd, *J* = 16.5 Hz, *J* = 2.8 Hz, 1H), 2.18-2.25 (m, 1H), 2.25-2.40 (m, 3H), 2.52-2.68 (m, 2H), 3.60-3.70 (m, 3H), 3.86 (dt, *J* = 16.4 Hz, *J* = 11.0 Hz, 1H), 4.36 (br d, *J* = 11.3 Hz, 1H), 4.55-4.60 (m, 1H), 5.05-5.11 (m, 1H), 5.52-5.65 (m, 2H), 5.71 (d, *J* = 11.1 Hz, 1H), 5.80-5.85 (m, 1H), 5.90 (d, *J* = 11.5 Hz, 1H), 6.35 (dt, *J* = 11.0 Hz, *J* = 3.4 Hz, 1H), 7.32-7.44 (m, 6H), 7.64 (d, *J* = 7.1 Hz, 4H); **¹³C-NMR** (100 MHz, CDCl₃): δ = -5.0, -4.6, 18.1, 19.2, 20.0, 25.8, 26.7, 31.1, 32.8, 32.9, 34.9, 43.5, 51.6, 65.5, 65.8, 68.7, 73.0, 73.2, 121.2, 125.1, 125.6, 127.6, 129.0, 129.6, 133.2, 133.2, 135.5, 135.5, 135.6, 148.3, 165.8, 209.2; **[α]_D²⁰** = -120 (*c* = 0.98, CH₂Cl₂); **m/z** (+CI, NH₃): 734 ([M+NH₄]⁺, 2), 717 ([M+H]⁺, 1), 586 (23), 274 (19), 196 (28), 91 (100); **HRMS** (+ES) calculated for C₄₂H₆₀O₆Si₂NH₄ [M+NH₄]⁺ 734.4272, found 734.4273.

(3Z,9E)-(1R,7S,11S,17R)-11-(tert-Butyl-dimethyl-silanyloxy)-7-(tert-butyl-diphenyl-silanyloxymethyl)-13-methylene-6,21-dioxo-bicyclo[15.3.1]henicosa-3,9,19-trien-5-one

R_f = 0.46 (hexane/EtOAc = 80/20); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 2931, 1722, 1167, 1113, 702; **$^1\text{H-NMR}$** (400 MHz, CDCl_3): δ = 0.00 (s, 3H), 0.02 (s, 3H), 0.87 (s, 9H), 1.03 (s, 9H), 1.35-1.55 (m, 4H), 1.85-2.02 (m, 4H), 2.10-2.25 (m, 3H), 2.30-2.40 (m, 2H), 3.62-3.75 (m, 3H), 3.91 (dt, J = 16.0 Hz, J = 11.0 Hz, 1H), 4.14-4.22 (m, 1H), 4.35 (br d, J = 10.3 Hz, 1H), 4.73 (d, J = 17.6 Hz, 2H), 5.03-5.12 (m, 1H), 5.45-5.60 (m, 2H), 5.72 (br d, J = 10.3 Hz, 1H), 5.80-5.86 (m, 1H), 5.90 (d, J = 11.6 Hz, 1H), 6.35 (dt, J = 11.0 Hz, J = 3.6 Hz, 1H), 7.35-7.47 (m, 6H), 7.60-7.70 (m, 4H); **$^{13}\text{C-NMR}$** (100 MHz, CDCl_3): δ = -4.8, -4.4, -0.5, 1.1, 7.1, 18.3, 19.3, 23.2, 26.0, 26.7, 31.2, 33.1, 35.2, 35.9, 44.9, 65.5, 66.7, 72.4, 73.0, 73.5, 121.3, 122.6, 125.2, 125.3, 127.7, 129.1, 129.7, 133.4, 133.4, 135.6, 136.2, 148.5, 165.7; $[\alpha]_D^{20}$ = -99.8 (c = 0.51, CH_2Cl_2); **m/z** (+CI, NH_3): 732 ($[\text{M}+\text{NH}_4]^+$, 25), 715 ($[\text{M}+\text{H}]^+$, 3), 583 (100), 465 (18), 327 (30), 274 (30), 159 (48), 132 (90), 91 (55); **HRMS** (+ES) calculated for $\text{C}_{43}\text{H}_{62}\text{O}_5\text{Si}_2\text{NH}_4$ $[\text{M}+\text{NH}_4]^+$ 732.4480, found 732.4480.

(3Z,9E)-(1R,7S,11S,17R)-11-(tert-Butyl-dimethyl-silanyloxy)-7-hydroxymethyl-13-methylene-6,21-dioxo-bicyclo[15.3.1]henicosa-3,9,19-trien-5-one

R_f = 0.20 (hexane/EtOAc = 70/30); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 2931, 1718, 1168, 1075, 831; **$^1\text{H-NMR}$** (500MHz, CDCl_3): δ = 0.01 (s, 3H), 0.02 (s, 3H), 0.87 (s, 9H), 1.30-1.50 (m, 4H), 1.86-2.00 (m, 4H), 2.12 (dd, J = 14.0 Hz, J = 6.2 Hz, 1H), 2.17-2.40 (m, 4H), 3.60-3.70 (m, 2H), 3.72 (dd, J = 11.9 Hz, J = 3.7 Hz, 1H), 3.85 (dt, J = 16.2 Hz, J = 10.4 Hz, 1H), 4.15-4.23

(m, 1H), 4.35 (br d, $J = 11.0$ Hz, 1H), 4.73 (d, $J = 13.4$ Hz, 2H), 4.94-5.04 (m, 1H), 5.48-5.58 (m, 2H), 5.70-5.76 (m, 1H), 5.80-5.86 (m, 1H), 5.90-5.96 (m, 1H), 6.39 (dt, $J = 11.0$ Hz, $J = 4.1$ Hz, 1H); $^{13}\text{C-NMR}$ (62.5MHz, CDCl_3): $\delta = -4.8, -4.4, 18.2, 23.2, 25.8, 31.1, 33.1, 34.3, 35.1, 35.7, 44.8, 65.3, 66.6, 72.3, 72.9, 74.7, 112.2, 120.8, 124.7, 125.3, 128.9, 136.5, 146.0, 149.3, 166.4$; $[\alpha]_{\text{D}}^{20} = -118$ ($c = 0.60$, CH_2Cl_2); m/z (+CI, NH_3): 494 ($[\text{M}+\text{NH}_4]^+$, 5), 477 ($[\text{M}+\text{H}]^+$, 2), 394 (8), 345 (71), 242 (24), 159 (32), 132 (65), 90 (100); **HRMS** (+CI, NH_3) calculated for $\text{C}_{27}\text{H}_{44}\text{O}_5\text{SiNH}_4$ $[\text{M}+\text{NH}_4]^+$ 494.3302, found 494.3301.

(3Z,9E)-(1R,7S,11S,17R)-11-(tert-Butyl-dimethyl-silanyloxy)-13-methylene-5-oxo-6,21-dioxo-bicyclo[15.3.1]henicosa-3,9,19-triene-7-carbaldehyde

$R_f = 0.29$ (hexane/EtOAc = 70/30); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 2928, 1726, 1713, 1162, 821; $^1\text{H-NMR}$ (500MHz, CDCl_3): $\delta = 0.01$ (s, 3H), 0.03 (s, 3H), 0.88 (s, 9H), 1.30-1.50 (m, 4H), 1.81-1.98 (m, 4H), 2.13 (dd, $J = 13.9$ Hz, $J = 6.9$ Hz, 1H), 2.20-2.40 (m, 3H), 2.57-2.66 (m, 1H), 3.55-3.65 (m, 1H), 3.83 (dt, $J = 16.0$ Hz, $J = 11.3$ Hz, 1H), 4.23 (q, $J = 6.4$ Hz, 1H), 4.32-4.44 (m, 1H), 4.73 (d, $J = 13.0$ Hz, 2H), 5.09 (dd, $J = 11.2$ Hz, $J = 2.7$ Hz, 1H), 5.42-5.61 (m, 2H), 5.68-5.75 (m, 1H), 5.80-5.88 (m, 1H), 5.98-6.05 (m, 1H), 6.47 (dt, $J = 11.1$ Hz, $J = 4.1$ Hz, 1H), 9.54 (s, 1H); $^{13}\text{C-NMR}$ (62.5MHz, CDCl_3): $\delta = -4.8, -4.5, 18.2, 23.4, 25.9, 31.0, 31.1, 33.5, 35.2, 35.7, 45.2, 66.8, 72.1, 72.8, 77.9, 112.1, 120.0, 123.4, 125.4, 128.8, 137.3, 146.0, 150.3, 165.6, 198.2$; $[\alpha]_{\text{D}}^{20} = -132$ ($c = 0.08$, CH_2Cl_2); m/z (+CI, NH_3): 492 ($[\text{M}+\text{NH}_4]^+$, 15), 475 ($[\text{M}+\text{H}]^+$, 4), 343 (100), 185 (18), 132 (18), 92 (22); **HRMS** (+CI, NH_3) calculated for $\text{C}_{27}\text{H}_{42}\text{O}_5\text{SiNH}_4$ $[\text{M}+\text{NH}_4]^+$ 492.3145, found 492.3137.

(S)-5-((E)-2-Iodo-vinyl)-1-methyl-cyclohexene

R_f = 0.6 (hexane/Et₂O = 95/5); **IR** $\tilde{\nu}$ (cm⁻¹, thin film) = 3368, 2953, 1725; **¹H-NMR** (250 MHz, CDCl₃): δ = 1.70 (s, 3H), 1.90-1.94 (br d, J = 9.3 Hz, 1H), 2.04-2.06 (m, 1H), 3.97-4.01 (m, 1H), 4.14-4.21 (m, 2H), 5.41 (br s, 1H), 6.41 (dd, J = 14.5 Hz, J = 1.3 Hz, 1H), 6.63 (dd, J = 14.6 Hz, J = 5.4 Hz, 1H); **¹³C-NMR** (62.9 MHz, CDCl₃): δ = 22.9, 34.9, 65.5, 75.4, 77.8, 119.7, 130.9, 145.9; $[\alpha]_D^{20}$ = -80 (c = 0.26, CHCl₃); **m/z** (+EI): 123 ([M-I]⁺, 100); **HRMS** (+EI) calculated for C₈H₁₁O [M-I]⁺ 123.0810, found 123.0810.

(3Z,9E)-(1R,7S,11S,17R)-11-(tert-Butyl-dimethyl-silanyloxy)-7-[(E)-(S)-1-hydroxy-3-((S)-3-methyl-cyclohex-3-enyl)-allyl]-13-methylene-6,21-dioxo-bicyclo[15.3.1]henicosa-3,9,19-trien-5-one

(diastereomeric purity of the compound was determined to be 9:1 by chiral HPLC)

R_f = 0.24 (hexane/EtOAc = 70/30); **IR** $\tilde{\nu}$ (cm⁻¹, thin film) = 3430, 2926, 1722, 1462, 1078; **¹H-NMR** (500 MHz, CDCl₃): δ = 0.02 (s, 3H), 0.01 (s, 3H), 0.87 (s, 9H), 0.92 (m, 1H), 0.98 (m, 1H), 1.26 (s, 3H), 1.40 (m, 3H), 1.55 (m, 1H), 1.70 (br s, 3H), 1.92 (m, 3H), 2.01 (m, 1H), 2.16 (m, 1H), 2.18 (m, 1H), 2.34 (m, 1H), 3.66 (br s, 1H), 3.86-3.83 (m, 1H), 4.06-4.02 (m, 1H), 4.18-4.15 (m, 5H), 4.34 (d, J = 9.9 Hz, 1H), 4.71 (s, 1H), 4.75 (s, 1H), 4.95 (dd, J = 5.8, J = 5.8 Hz, s, 1H), 5.42 (s, 1H), 5.55-5.46 (m, 1H), 5.77-5.70 (m, 2H), 5.86-5.83 (m, 2H), 5.90 (d, J = 11.4 Hz, 1H), 6.40 (dt, J = 11.1, J = 3.8 Hz, 1H); **¹³C-NMR** (125.8 MHz, CDCl₃): δ = -4.8, -4.4, 18.2, 22.9, 23.3, 25.9, 31.2, 33.0, 33.4, 35.2, 35.7, 36.0, 44.7, 65.6, 66.6, 72.3, 72.9, 73.3, 73.8, 75.3, 112.3, 119.7, 120.7, 124.7, 125.3, 128.9, 129.2, 131.3, 133.5, 136.7, 146.2, 149.4, 165.9; $[\alpha]_D^{22}$ = -215 (c = 0.02, CH₂Cl₂); **m/z** (+EI, Na): 621 ([M+Na]⁺, 100), 598 (29); **HRMS** (+EI) calculated for C₃₅H₅₄O₆SiNa [M+Na]⁺ 621.3587, found 621.3564.

(3Z,9E)-(1R,7S,11S,17R)-11-(tert-Butyl-dimethyl-silanyloxy)-7-[(E)-3-((S)-3-methyl-cyclohex-3-enyl)-acryloyl]-13-methylene-6,21-dioxa-bicyclo[15.3.1]henicosa-3,9,19-trien-5-one

R_f = 0.62 (hexane/EtOAc = 70/30); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 2927, 1726, 1638, 1263, 835; **$^1\text{H-NMR}$** (400MHz, CDCl_3): δ = -0.02 (s, 3H), -0.01 (s, 3H), 0.86 (s, 9H), 1.22 (s, 3H), 1.38 (m, 3H), 1.55 (s, 1H), 1.70 (br s, 2H), 1.92 (br s, 3H), 2.01 (m, 2H), 2.13 (m, 1H), 2.25 (m, 1H), 2.36 (m, 1H), 2.50 (m, 1H), 3.58 (br s, 1H), 3.82 (m, 1H), 4.40-4.10 (m, 5H), 4.73 (d, J = 6.7 Hz, 2H), 5.18 (d, J = 9.1 Hz, 1H), 5.42 (s, 1H), 5.57 (br s, 2H), 5.68 (d, J = 9.0 Hz, 1H), 5.80 (m, 1H), 6.02 (d, J = 10.7 Hz, 1H), 6.38 (m, 1H), 6.55 (d, J = 15.1 Hz, 1H), 7.02 (dd, J = 15.5, J = 3.7 Hz, 1H); **$^{13}\text{C-NMR}$** (125.8MHz, CDCl_3): δ = -4.8, -4.5, 18.2, 22.8, 23.4, 25.9, 29.7, 31.1, 33.0, 33.3, 34.9, 35.3, 35.5, 45.2, 65.8, 66.8, 72.2, 72.4, 72.8, 77.8, 112.0, 119.7, 120.4, 122.5, 124.2, 125.2, 128.9, 131.1, 137.1, 146.0, 147.6, 149.6, 165.5, 195.5; $[\alpha]_D^{22}$ = -172 (c = 0.05, CH_2Cl_2); **m/z** (+EI, Na): 619 ($[\text{M}+\text{Na}]^+$, 35), 576 (42), 465 (100); **HRMS** (+EI) calculated for $\text{C}_{35}\text{H}_{52}\text{O}_6\text{SiNa}$ $[\text{M}+\text{Na}]^+$ 619.3431, found 619.3422.

(3Z,9E)-(1R,7S,11S,17R)-11-Hydroxy-7-[(E)-3-((S)-1-hydroxy-3-((S)-3-methyl-cyclohex-3-enyl)-allyl)-13-methylene-6,21-dioxo-bicyclo[15.3.1]henicosa-3,9,19-trien-5-one

R_f = 0.22 ($\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$ = 2/1); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 3260, 2925, 1723, 1463, 1098; **$^1\text{H-NMR}$** (500 MHz, CDCl_3): δ = 1.30-1.69 (m, 5H), 1.69 (s, 3H), 1.86-2.14 (m, 6H), 2.17-2.28 (m, 2H), 2.33-2.36 (m, 2H), 3.63-3.70 (m, 1H), 3.70-3.82 (m, 1H), 4.04-4.10 (m, 1H), 4.15-4.25 (m, 4H), 4.36 (br d, 1H), 4.80 (s, 1H), 4.86 (s, 1H), 4.97-5.04 (m, 1H), 5.44 (br s, 1H), 5.60-5.67 (m, 2H), 5.69-5.77 (m, 2H), 5.84-5.95 (m, 2H), 5.94 (br d, J = 11.6 Hz, 1H), 6.43

(ddd, $J = 11.6, 10, 3.6$ Hz, 1H); $[\alpha]_D^{20} = -108$ ($c = 0.2$, CH_2Cl_2); **m/z** (+EI, Na): 507 ($[\text{M}+\text{Na}]^+$, 58), 351 (52), 242 (100); **HRMS** (+EI) calculated for $\text{C}_{29}\text{H}_{40}\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 507.2801, found 507.2783.

*C*₁₁-*desmethyl*-Laulimalide

R_f = 0.18 ($\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O} = 2/1$); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 3276, 2924, 1720, 1647, 1262, 1090; **¹H-NMR** (700 MHz, $[\text{D}_4]-\text{MeOH}$): $\delta =$ see tabulated data on page S9; **¹³C-NMR** (125.8 MHz, $[\text{D}_4]-\text{MeOH}$): $\delta =$ see tabulated data on page S10; $[\alpha]_D^{20} = -176$ ($c = 0.2$, CH_2Cl_2); **m/z** (+EI, Na): 523 ($[\text{M}+\text{Na}]^+$, 46), 413 (100); **HRMS** (+EI) calculated for $\text{C}_{29}\text{H}_{40}\text{O}_7\text{Na}$ $[\text{M}+\text{Na}]^+$ 523.2672, found 523.2692.

(3Z,9E)-(1R,7S,11S,17R)-11-(*tert*-Butyl-dimethyl-silanyloxy)-7-[(E)-(S)-1-methoxy-3-((S)-3-methyl-cyclohex-3-enyl)-allyl]-13-methylene-6,21-dioxabicyclo[15.3.1]henicosa-3,9,19-trien-5-one

R_f = 0.7 (hexane/EtOAc = 70/30); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 2920, 2856, 1723, 1164, 1095; **¹H-NMR** (500 MHz, CDCl_3): $\delta = -0.04$ (s, 3H), -0.02 (s, 3H), 0.85 (s, 9H), 0.90-1.30 (m, 5H), 1.69 (s, 3H), 1.85-2.35 (m, 9H), 3.32 (s, 3H), 3.63-3.75 (m, 2H), 3.91-4.21 (m, 5H), 4.25-4.37 (m, 1H), 4.68 (br s, 1H), 4.73 (br s, 1H), 4.92 (m, 1H), 5.40-5.58 (m, 5H), 5.66-5.74 (m, 1H), 5.77-5.84 (m, 2H), 5.91 (dd, $J = 11.4$ Hz, $J = 1.4$ Hz, 1H), 6.36 (ddd, $J = 11.4$ Hz, $J = 10.8$ Hz, $J = 3.7$ Hz, 1H); **¹³C-NMR** (125.8 MHz, CDCl_3): $\delta = -4.8, -4.6, 18.2, 22.9, 25.9, 29.7, 31.1, 31.9, 33.2, 35.1, 35.7, 35.9, 44.7, 57.2, 65.6, 66.6, 72.3, 72.9, 73.3, 73.6, 51.4, 53.4, 57.0, 112.2, 119.7, 120.7, 125.0, 125.2, 126.4, 128.9, 130.9, 135.0, 136.4, 146.1, 148.9,$

165.6; $[\alpha]_D^{20} = -99$ ($c = 0.09$, CH_2Cl_2); m/z (+EI, Na): 635 ($[\text{M}+\text{Na}]^+$, 100); **HRMS** (+EI) calculated for $\text{C}_{36}\text{H}_{56}\text{O}_6\text{SiNa}$ $[\text{M}+\text{Na}]^+$ 635.3744, found 635.3730.

(3Z,9E)-(1R,7S,11S,17R)-11-Hydroxy-7-[(E)-(S)-1-methoxy-3-((S)-3-methyl-cyclohex-3-enyl)-allyl]-13-methylene-6,21-dioxabicyclo[15.3.1]henicosa-3,9,19-trien-5-one

$R_f = 0.3$ (hexane/EtOAc = 60/40); **IR** $\tilde{\nu}$ (cm^{-1} , thin film) = 2927, 1726, 1265, 1098; **$^1\text{H-NMR}$** (500 MHz, CDCl_3): $\delta = 1.25\text{--}1.48$ (m, 5H), 1.69 (s, 3H), 1.85–2.33 (m, 10H), 3.29 (s, 3H), 3.62–3.69 (m, 1H), 3.72–3.80 (m, 1H), 4.02–4.18 (m, 5H), 4.29–4.34 (m, 1H), 4.78 (br s, 1H), 4.86 (br s, 1H), 5.08 (dd, $J = 8.4$ Hz, $J = 5.2$ Hz, 5.40 (br s, 1H), 5.48–5.62 (m, 4H), 5.68–5.72 (m, 1H), 5.78–5.84 (m, 2H), 5.89 (dd, $J = 10.2$ Hz, $J = 1.2$ Hz, 1H), 6.38 (ddd, $J = 13.3$ Hz, $J = 9.1$ Hz, $J = 5.2$ Hz, 1H); **$^{13}\text{C-NMR}$** (125.8 MHz, CDCl_3): $\delta = 22.9, 23.1, 31.4, 32.6, 33.3, 35.7, 36.3, 43.2, 60.4, 65.6, 66.8, 70.5, 72.9, 73.3, 73.4, 82.5, 112.6, 119.7, 120.8, 125.1, 126.4, 126.4, 128.9, 131.2, 135.3, 135.5, 146.0, 149.1, 165.6$; $[\alpha]_D^{20} = -143$ ($c = 0.04$, CH_2Cl_2); m/z (+EI, Na): 522 ($[\text{M}+\text{Na}]^+$, 100); **HRMS** (+EI) calculated for $\text{C}_{30}\text{H}_{42}\text{O}_6\text{Na}$ $[\text{M}+\text{Na}]^+$ 521.2879, found 521.2862.

C₁₁-desmethyl-Laulimalide-O₂₀-methylether

R_f = 0.28 (hexane/EtOAc = 60/40); **IR** $\tilde{\nu}$ (cm⁻¹, thin film) = 3485, 2924, 1721, 1261, 1097, 807; **¹H-NMR** (500 MHz, CDCl₃): δ = 1.22-1.48 (m, 5H), 1.62 (s, 3H), 1.82-2.20 (m, 10H), 2.90 (br s, 1H), 3.14 (m, 1H), 3.29 (s, 3H), 3.74 (m, 1H), 3.86 (m, 1H), 3.92-4.18 (m, 4H), 4.26 (br d, J = 7.4 Hz, 1H), 4.38 (br d, J = 9.2 Hz, 1H), 4.79 (br s, 1H), 4.89 (br s, 1H), 5.18 (m, 1H), 5.38 (br s, 1H), 5.52-5.58 (m, 1H), 5.71 (br d, J = 10.2 Hz, 1H), 5.78-5.85 (m, 2H), 5.89 (br d, J = 12.0 Hz, 1H), 6.38 (ddd, J = 12.0 Hz, J = 9.8 Hz, J = 3.2 Hz, 1H); **¹³C-NMR** (125.8 MHz, CDCl₃): δ = 25.3, 25.5, 31.4, 31.7, 33.3, 33.8, 36.9, 37.2, 39.4, 52.1, 57.2, 60.4, 61.6, 64.9, 65.3, 66.0, 69.6, 71.9, 71.9, 106.7, 114.8, 115.3, 120.1, 120.4, 123.2, 125.5, 129.4, 139.2, 143.4, 161.2; **[α]_D²⁰** = -112 (c = 0.02, CH₂Cl₂); **m/z** (+EI, Na): 576 (100), 537 ([M+Na]⁺, 70); **HRMS** (+EI) calculated for C₃₀H₄₂O₇Na [M+Na]⁺ 537.2828, found 537.2829.

¹H NMR data comparison of (-)-laulimalide and 11-desmethyl-laulimalide recorded on a Bruker av700 in [D₄]-MeOH.

H	δ [ppm], mult., J [Hz] laulimalide	δ [ppm], mult., J [Hz] 11-desmethyl-laulimalide
2	5.98, ddd, 11.3/ 2.4/ 1.2	5.94, ddd, 11.5/ 2.6/ 1.2
3	6.43, ddd, 11.3/ 10.1/ 4.0	6.42, ddd, 11.5/ 10.4/ 3.4
4	3.74, ddd, 16.8/ 10.1/ 1.2 2.23, ddd, 16.8/ 4.0/ 2.9	3.82, ddd, 16.9/ 10.4/ 1.9 2.22, ddd, 16.9/ 4.4/ 2.5
5	4.23, brs	4.34, ddd, 10.4/ 2.5/ 2.2
6	5.74, ddd, 10.3/ 3.8/ 1.2	5.75, ddd, 10.4/ 3.8/ 1.3
7	5.82, m	5.85, m
8	2.07, m 1.90, m	1.95, ddd, 17.5/ 4.0/ 3.8 1.88, ddd, 17.5/ 10.0/ 2.5
9	3.81, dddd, 8.2/ 8.1/ 4.1/ 4.0	3.69, dddd, 8.6/ 8.0/ 4.0/ 3.0
10	1.51, ddd, 14.3/ 4.8/ 3.6 1.25, ddd, 14.3/ 4.9/ 3.6	1.45, m 1.38, m
11	1.73, m	1.45, m 1.45, m
12	2.28, dd, 12.6/ 5.2 1.88, m	2.06, ddd, 14.3/ 8.9/ 5.7 1.96, ddd, 14.3/ 9.3/ 5.6
14	2.07, dd, 15.5/ 3.1 2.00, dd, 15.5/ 8.5	2.12, m 1.92, dd, 15.3/ 8.9
15	3.96, ddd, 8.9/ 3.1/ 2.2	4.02, ddd, 8.3/ 3.2/ 2.2
16	2.82, dd, 2.5/ 2.2	2.87, dd, 2.2/ 2.2
17	3.00, ddd, 8.2/ 2.5/ 2.2	2.97, ddd, 7.9/ 3.6/ 2.2
18	2.10, ddd, 14.4/ 2.2/ 1.2, 1.54, ddd, 14.4/ 11.2/ 8.2	2.14, ddd, 14.3/ 3.6/ 1.9 1.58, ddd, 14.3/ 11.3/ 7.9
19	5.14, ddd, 11.2/ 4.6/ 1.2	5.15, ddd, 11.3/ 4.4/ 1.8
20	4.17, dd, 4.9/ 4.6	4.17, dd, 4.8/ 4.4
21	5.70, ddd, 15.6/ 4.9/ 1.2	5.70, ddd, 15.6/ 5.5/ 1.3
22	5.79, ddd, 15.6/ 5.5/ 1.2	5.80, 15.6/ 5.4/ 1.2
23	4.02, ddd, 9.0/ 5.5/ 3.6	4.02, m
24	1.98, m 1.90, m	2.01 m 1.92, m
26	5.41, dd, 2.2/ 1.3	5.42, dd, 1.7/ 1.3
27	4.12, m	4.14, m
28	1.69, s	1.69, s
29	4.81, brs 4.77, brs	4.82, brs 4.78, brs
30	0.83, d, 6.5	

¹³C NMR data of 11-desmethyl-laulimalide (recorded on a Bruker DRX500 in [D₄]-MeOH)

C	δ [ppm]
1	165.7
2	120.2
3	149.5
4	32.6
5	73.1
6	128.6
7	124.7
8	30.5
9	67.1
10	35.1
11	24.3
12	37.6
13	146.3
14	37.0
15	66.4
16	61.1
17	51.9
18	32.9
19	72.2
20	72.1
21	129.3
22	132.3
23	73.5
24	35.2
25	131.1
26	119.2
27	65.1
28	21.6
29	110.0

Laulimalide **11-Desmethyl-laulimalide**

Laulimalide **11-Desmethyl-laulimalide**

Laulimalide **11-Desmethyl-laulimalide**