Supporting information for

Synthesis of Novel polycyclic Indolyldiamines

Sabrina Cutri,[§] Anna Diez,^{*,#} Martine Bonin,^{*,§} Laurent Micouin[§] and Henri-Philippe Husson[§]

Laboratoire de Chimie Thérapeutique, UMR 8638 associée au CNRS et à l'Université René Descartes, Faculté des Sciences Pharmaceutiques et Biologiques, 4 av de l'Observatoire, 75270 Paris cedex 06 (France) and Parc Científic de Barcelona c/Josep Samitier, 1-5 08028- Barcelona (Spain).

martine.bonin@univ-paris5.fr

General methods

Silica gel (230-400 mesh) was used for product purification by flash column chromatography. Solvents were purified and dried by standard techniques, and distilled prior to use. All organic extracts were dried over MgSO₄ or Na₂SO₄. IR spectra were recorded on a *Perkin-Elmer* FTIR 1600 spectrometer as thin film. ¹H and ¹³C NMR spectra were recorded on a *Bruker* AC-300 or AC-400 spectrometer with TMS as internal standard and CDCl₃ as solvent. HRMS were determined on a *Kratos* MS 80 RF or *Micromass* Q ToF 1 spectrometer.

2-[7-(1-Methyl-1H-indol-2-yl)-6,8-diazabicyclo[3.2.1]oct-6-en-8-yl]-2-phenylethanol **6**

To a stirred solution of TMEDA (2 eq., 44 mmol, 6.62 mL) was added *n*-BuLi (2 eq., 44 mmol, 1.6 M, 27.5 mL) under argon at -20 °C. After 5 min. stirring, N-methylindole (3 eq., 66 mmol, 8.42 mL) was dropwise added. Stirring was maintained for 1 h while proceeding to two ultrasonic irradiations (2 x 1 min.). A solution of (-)-2-cyano-6-phenyloxazolopiperidine **1** (22 mmol, 5 g) in anhydrous ether (200 mL) was then added under argon. Stirring was maintained for 5 h until the temperature of the reaction mixture was warmed gradually to room temperature. The reaction mixture was cautiously diluted with a saturated, aqueous solution of NH₄Cl, extracted with CH₂Cl₂ (2 x 150 mL), and the combined organic layers were dried (MgSO₄). The solvents were evaporated to give an oily residue which was purified by chromatography (CH₂Cl₂/MeOH 98/2 then 95/5) to provide imine **6** (yellow oil, 7.27 g, 92 %). [α]_D²⁰ -197 (c 1.0, MeOH); IR (KBr) $\nu_{\max}$ 3350, 3027, 2990, 2855, 1648, 1610, 1468, 1087, 1059 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.56 (d, *J* = 8.0 Hz, CH), 7.20-7.40 (7 H, m, 7CH), 7.10 (1 H, td, *J*₁ = 7.8 Hz, *J*₂ = 1.2 Hz, CH), 6.56 (1 H, s, CH), 5.54 (1 H, t, *J* = 2.4 Hz, CH), 4.09 (3 H, s, CH₃), 3.98 (1 H, t, *J* = 3.0 Hz, CH), 3.94 (1 H, dd, *J*₁ = 11.1 Hz, *J*₂ = 5.4 Hz, CH₂), 3.81 (1 H, dd, *J*₁ = 11.1 Hz, *J*₂ = 5.4 Hz, CH₂), 3.50 (1 H, t, *J* = 5.4 Hz, CH), 1.20-1.90 (7 H, m, 3CH₂, OH); ¹³C NMR (CDCl₃, 75 MHz): δ 165.3, 140.2, 131.0, 128.7, 128.5, 127.9, 126.8, 121.7, 124.4, 120.2, 110.0, 108.8, 87.0, 66.1, 67.6, 64.3, 32.3, 16.9, 25.5, 26.1; MS (CI) *m/z* 360 (MH⁺); HRMS (CI) calcd for (MH)⁺, 360.2076, found for (MH)⁺: 360.2074.

2-(1-Methyl-1H-indol-2-yl)-azepan-3-ylamine **7**

LiAlH₄ (8 eq., 71.4 mmol, 2.72 g) was carefully added at 0 °C to a solution of imine **6** (8.92 mmol, 3.20 g) in dry *i*Pr₂O (160 mL) under argon. The reaction mixture was stirred for 24 h at reflux, allowed to cool and treated at 0 °C successively with aqueous NaOH (1M, 5.4 mL) and H₂O (8.2 mL). After filtration through celite, the residue was washed several times with Et₂O and the organic layer was concentrated in vacuo. The crude (3.24 g) was then dissolved in methanol (120 mL) and stirred under hydrogen in presence of aqueous HCl (12N, pH 2) and 10% Pd/C (647 mg) for 48 h at room temperature. After filtration through celite and concentration *in vacuo*, the residue was dissolved in ether and extracted three times with an aqueous 2N HCl solution. Aqueous layers were made basic (NaOH 6N) and then extracted with AcOEt. The organic layers were dried (Na₂SO₄) and concentrated to provide an oily residue. The residue was purified by chromatography (AcOEt/MeOH/NH₄OH 175/20/5) to provide azepane **7** (yellow oil, 1.76 g, 81%). [α]_D²⁰ -3 (c 0.7, CHCl₃); IR (KBr) ν_{max} 3360, 2928, 1468, 1315, 1135 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.56 (1 H, d, *J* = 7.3 Hz, CH), 7.30 (1 H, d, *J* = 8.2 Hz, CH), 7.20 (1 H, td, *J*₁ = 7.7 Hz, *J*₂ = 0.7 Hz, CH), 7.08 (1 H, td, *J*₁ = 7.2 Hz, *J*₂ = 0.6 Hz, CH), 6.44 (1 H, s, CH), 3.80 (3 H, s, CH₃), 3.60 (1 H, d, *J* = 9.6 Hz, CH), 3.39 (1 H, m, CH), 3.15 (1 H, dt, *J*₁ = 13.7 Hz, *J*₂ = 4.9 Hz, CH₂), 2.84 (1 H, m, CH₂), 1.60-2.10 (9 H, m, 3CH₂, NH, NH₂); ¹³C NMR (CDCl₃, 75 MHz): δ 142.3, 137.8, 127.6, 121.5, 120.5, 119.6, 109.2, 98.9, 65.0, 56.8, 48.6, 35.5, 30.7, 30.2, 22.4; MS (CI) *m/z* 244 (MH⁺); HRMS (CI) calcd for (MH)⁺, 244.1814, found for (MH)⁺: 244.1816.

[2-(tert-Butyl-dimethyl-silanyloxy)-1-phenyl-ethyl]-[7-methyl-2-(1-methyl-1H-indol-2-yl)-azepan-3-yl]-amine **9**

NaH (60% suspension in mineral oil, 2 eq., 2.03 mmol, 81 mg) was added at 0 °C under argon to a solution of imine **6** (1.02 mmol, 365 mg) in THF (15 mL). After 30 min. at room temperature, *tert*-butyldimethylsilyl chloride (2 eq., 2.03 mmol, 307 mg) was added. The resulting mixture was stirred for an additional 16 h at 20 °C. The reaction mixture was cautiously diluted with a saturated, aqueous NH₄Cl solution and extracted with CH₂Cl₂ (2 x). The combined organic layers were dried (MgSO₄) and concentrated in vacuo. The oily residue (600 mg) was then dissolved in Et₂O (20 mL). LiAlH₄ (5 eq., 5.10 mmol, 194 mg) was carefully added at 0 °C under argon. The reaction mixture was stirred for 1h30 at 0 °C, and treated successively with aqueous NaOH (1M, 389 μ L) and H₂O (583 μ L). After filtration through celite the residue was washed several times with Et₂O and the organic layer was concentrated in vacuo providing the aminal **8** as colourless oil (550 mg). This aminal was dissolved in Et₂O (24 mL) under argon. An ethereal solution of methylmagnesium bromide (3

M/Et₂O, 3.5 eq., 3.57 mmol, 1.19 mL) was carefully added at room temperature. The reaction mixture was stirred for 3 h at reflux. The reaction mixture was cautiously treated with a saturated, aqueous solution of NH₄Cl, extracted with CH₂Cl₂ (2 x 40 mL). The combined organic layers were dried (MgSO₄). The solvents were evaporated to give an oily residue. The residue was purified by chromatography (AcOEt/CH₃CN/NH₄OH 195/4/1) to provide azepane **9** (yellow oil, 176 mg, 35 %). [α]_D²⁰ -26 (c 1.4, CH₂Cl₂); IR (KBr) $\nu_{\max}$ 2952, 2926, 2855, 1470, 1314, 1255, 1091 cm⁻¹; ¹H NMR (CDCl₃, 400 MHz): δ 7.64 (1 H, d, J = 7.7 Hz, CH), 7.22-7.44 (7 H, m, 7CH), 7.15 (1 H, t, J = 7.6 Hz, CH), 6.56 (1 H, s, CH), 4.18 (1 H, d, J = 7.6 Hz, CH), 3.86 (3 H, s, CH₃), 3.70 (1 H, dd, J_1 = 8.0 Hz, J_2 = 4.2 Hz, CH), 3.58 (1 H, dd, J_1 = 10.0 Hz, J_2 = 4.2 Hz, CH₂), 3.52 (1 H, t, J = 9.0 Hz, CH₂), 3.26 (1 H, td, J_1 = 7.6 Hz, J_2 = 2.1 Hz, CH), 2.90 (1 H, m, CH₂), 2.48 (2 H, bs, 2NH), 1.18-1.94 (6 H, m, 3CH₂), 0.95 (3 H, d, J = 6.4 Hz, CH₃), 0.74 (9 H, s, 3CH₃), -0.10 (3 H, s, CH₃), -0.13 (3 H, s, CH₃); ¹³C NMR (CDCl₃, 75 MHz): δ 142.1, 141.5, 137.9, 128.3, 127.9, 127.6, 127.4, 121.2, 120.6, 119.3, 109.1, 99.9, 68.0, 64.0, 62.2, 56.8, 48.3, 38.5, 33.9, 30.1, 25.8, 24.7, 24.5, 18.2, 1.9, 1.6; MS (CI) m/z 492.3 (MH⁺); HRMS (CI) calcd for (MH)⁺, 492.3410, found for (MH)⁺: 492.3411.

Tetracycline **10**

Aminal **8** (550 mg) was dissolved in ethyl acetate (19.6 mL) at room temperature. CH₃CN (0.4 mL) and silica (3 g) were added. The reaction mixture was stirred for 4 h at room temperature and filtered. The organic layer was concentrated in vacuo give an oily residue. The residue was purified by chromatography (AcOEt/CH₃CN 95/5) to provide tetracycline **10** (yellow oil, 102 mg, 21 % from **6**). [α]_D²⁰ +61 (c 0.7, CH₂Cl₂); IR (KBr) $\nu_{\max}$ 2928, 2855, 1470, 1372, 1254, 1190, 1096, 1007 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.40 (1 H, dd, J_1 = 7.8 Hz, J_2 = 1.6 Hz, CH), 7.20-7.32 (6 H, m, 6CH), 7.14 (1 H, td, J_1 = 7.5 Hz, J_2 = 1.2 Hz, CH), 6.98 (1 H, td, J_1 = 7.1 Hz, J_2 = 0.9 Hz, CH), 4.32 (1 H, t, J = 4.8 Hz, CH), 4.06 (1 H, bs, CH), 3.83 (3 H, s, CH₃), 3.80-3.87 (3 H, m, CH, CH₂), 3.57 (1 H, d, J = 5.4 Hz, CH), 1.89 (2 H, bs, NH₂), 1.13-2.20 (6 H, m, 3CH₂), 0.82 (9 H, s, 3CH₃), -0.12 (3 H, s, CH₃), -0.14 (3 H, s, CH₃); ¹³C NMR (CDCl₃, 75 MHz): δ 142.6, 138.3, 137.4, 128.4, 127.2, 124.4, 121.0, 118.6, 118.4, 112.4, 109.0, 67.8, 64.9, 59.3, 50.0, 48.6, 29.7, 26.0, 24.6, 22.2, 18.3, 16.8, 1.5; MS (CI) m/z 476.3 (MH⁺); HRMS (CI) calcd for (MH)⁺, 476.3097, found for (MH)⁺: 476.3098.

Tetracycline **11**

To a stirred solution of diisopropylamine (2 eq., 1.75 mmol, 247 μ L) in THF (1 mL) was added *n*-BuLi (1.1 eq., 0.965 mmol, 2.5 M, 386 μ L) under argon at -70 $^{\circ}$ C. After being stirred for 30 min. at -70 $^{\circ}$ C, the reaction mixture was warmed gradually to 0 $^{\circ}$ C. A solution of N-phenylsulfonylindole (2.5 eq., 2.19 mmol, 563 mg) in THF (1 mL) was added dropwise. Stirring was maintained for 30 min. before cooling the reaction mixture to -70 $^{\circ}$ C. A solution of (-)-2-cyano-6-phenyloxazolopiperidine **1** (0.877 mmol, 200 mg) in THF (1 mL) was then added under argon. Stirring was maintained for 3 h until the temperature of the reaction mixture was warmed gradually to room temperature. The reaction mixture was cautiously treated with a saturated, aqueous solution of NH_4Cl , extracted with CH_2Cl_2 (2 x 50 mL), and the combined organic layers were dried (MgSO_4). The solvents were evaporated to give an oily residue. The residue was purified by chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$ 98/2) to provide imine **10** (yellow oil, 121 mg, 40 %). $[\alpha]_{\text{D}}^{20}$ -37 (c 0.8, CH_2Cl_2); IR (KBr) ν_{max} 3266, 3061, 2948, 2872, 1607, 1454, 1420, 1349, 1333, 1310, 1265, 1082, 1053 cm^{-1} ; ^1H NMR (CDCl_3 , 200 MHz): δ 7.54 (1 H, d, $J = 8.1$ Hz, CH), 7.15-7.31 (7 H, m, 7CH), 7.09 (1 H, m, CH), 6.55 (1 H, s, CH), 5.52 (1 H, bs, CH), 4.45 (1 H, bs, NH), 3.98 (2 H, m, CH_2), 3.89 (1 H, bs, CH), 3.43 (1 H, t, $J = 5.1$ Hz, CH), 1.10-1.94 (7 H, m, 3 CH_2 , OH); ^{13}C NMR (CDCl_3 , 50 MHz): δ 165.3, 140.3, 137.7, 129.6, 128.6, 128.4, 127.7, 124.8, 121.6, 120.3, 112.0, 108.5, 85.8, 66.7, 66.6, 65.4, 25.9, 25.4, 16.7; MS (EI) m/z 345 (M^+); HRMS (CI) calcd for (M) $^+$, 345.1841, found for (M) $^+$: 345.1843.

Tetracycline **12**

LiAlH_4 (4 eq., 1.16 mmol, 44 mg) was carefully added at -20 $^{\circ}$ C to a solution of imine **11** (0.290 mmol, 100 mg) in anhydrous Et_2O (5 mL) under argon. The reaction mixture was stirred for 3 h and the temperature reaction was gradually warmed to room temperature. The reaction mixture was treated successively with aqueous NaOH (1M, 88 μ L) and H_2O (132 μ L). After filtration through celite the residue was washed several times with Et_2O and the organic layer was concentrated in vacuo to give an oily residue. The residue was purified by chromatography ($\text{AcOEt}/\text{MeOH}/\text{NH}_4\text{OH}$ 179/20/1) to provide tetracycline **12** (yellow oil, 38 mg, 38 %). $[\alpha]_{\text{D}}^{20}$ -4 (c 0.8, CH_2Cl_2); IR (KBr) ν_{max} 3395, 2932, 1639, 1453, 1341, 1265, 1237, 1050 cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz): δ 9.00 (1 H, bs, NH), 7.19-7.42 (7 H, m, 7CH), 7.13 (1 H, td, $J_1 = 7.4$ Hz, $J_2 = 1.1$ Hz, CH), 7.04 (1 H, td, $J_1 = 7.4$ Hz, $J_2 = 1.0$ Hz, CH), 4.34 (1 H, bs, CH), 4.07 (1 H, t, $J = 4.9$ Hz, CH), 3.92 (1 H, dd, $J_1 = 11.1$ Hz, $J_2 = 4.9$ Hz, CH_2), 3.66 (2 H, m, CH, CH_2), 3.23 (1 H, bd, $J = 4.0$ Hz, CH), 2.30 (2 H, bs, NH_2), 1.12-2.02 (7 H, m, 3 CH_2 , OH) ppm; ^{13}C NMR (CDCl_3 , 75 MHz): δ 141.5, 136.4, 136.2, 128.8, 128.5, 127.7, 125.3, 121.7, 119.3, 118.2, 112.2, 111.3, 64.9, 64.7, 59.6, 49.7, 48.4, 28.0, 26.2, 16.6 ppm, MS (CI) m/z 348 (MH^+); HRMS (CI) calcd for (MH) $^+$, 348.2076, found for (MH) $^+$: 348.2075.

Tetracycline **14**

To a solution of azepane **7** (2.14 mmol, 519 mg) in THF (10 mL), distilled Et₃N (1.5 eq., 3.21 mmol, 638 μ L) and N-carbethoxyphthalimide (1.5 eq., 3.21 mmol, 701 mg) were added at room temperature under nitrogen. The reaction mixture was stirred for 24 h at reflux, cautiously diluted with H₂O, extracted with CH₂Cl₂ (2 x 100 mL), and the combined organic layers were dried (MgSO₄). The solvents were evaporated to give an oily residue. The crude was diluted in CH₂Cl₂ (16 mL). Glyoxal (40% in H₂O, 3 eq., 6.43 mmol, 736 μ L) and acetic acid (4 eq., 8.58 mmol, 488 μ L) were added at room temperature. Stirring was maintained for 5 h. The reaction mixture was cautiously diluted with a saturated, aqueous solution of NaHCO₃, extracted with CH₂Cl₂ (2 x 10 mL), and the combined organic layers were dried (MgSO₄). The solvents were evaporated to give an oily residue. The residue was purified by chromatography (CH₂Cl₂ then CH₂Cl₂/MeOH 99/1 and finally CH₂Cl₂/MeOH 96/4) to provide ketone **14** (yellow oil, 574 mg, 65 %). [α]_D²⁰ +470 (c 0.9, CH₂Cl₂); IR (KBr) ν_{max} 2927, 2861, 1771, 1713, 1652, 1614, 1480, 1464, 1422, 1391, 1357, 1329, 1132, 1090, 1063 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 8.23 (1 H, d, J = 7.5 Hz, CH), 7.90 (1 H, m, CH), 7.70 (3 H, m, 3CH), 7.26 (1 H, td, J_1 = 7.4 Hz, J_2 = 1.0 Hz, CH), 7.18 (1 H, td, J_1 = 7.6 Hz, J_2 = 1.2 Hz, CH), 7.03 (1 H, d, J = 7.6 Hz, CH), 4.90 (1 H, d, J = 10.0 Hz, CH), 4.67 (1 H, td, J_1 = 9.6 Hz, J_2 = 3.7 Hz, CH), 4.01 (1 H, d, J = 18.0 Hz, CH₂), 3.56 (1 H, d, J = 18.0 Hz, CH₂), 3.40 (2 H, dd, J_1 = 6.0 Hz, J_2 = 3.0 Hz, CH₂), 3.35 (3 H, s, CH₃), 2.70 (1 H, dt, J_1 = 14.6 Hz, J_2 = 10.0 Hz, CH₂), 1.70-2.30 (5 H, m, CH₂); ¹³C NMR (CDCl₃, 300 MHz): δ 190.6, 153.3, 137.7, 134.5, 124.6, 124.0, 123.4, 123.2, 122.1, 110.8, 109.4, 58.1, 55.9, 55.0, 53.7, 32.5, 31.1, 26.3, 24.1; MS (CI) m/z 414 (MH⁺); HRMS (CI) calcd for (MH)⁺, 414.1818, found for (MH)⁺: 414.1815.

N-[1-(2-Chloro-acetyl)-2-(1-methyl-1H-indol-2-yl)-azepan-3-yl]-2-nitro-benzenesulfonamide **15**

To a solution of diamine **7** (4.12 mmol, 1 g) in THF (13 mL), NaHCO₃ (3 eq., 12.3 mmol, 1.03 g) was added at 0 °C under nitrogen. A solution of nosyl chloride (1.5 eq., 6.17 mmol, 1.37 g) in THF (13 mL) was slowly added dropwise (1 mL/h) at 0 °C. Stirring was maintained for 20 h, keeping the reaction mixture at 0 °C. The reaction mixture was cautiously diluted with H₂O and extracted with CH₂Cl₂ (2 x 50 mL). The combined organic layers were dried (MgSO₄) and concentrated in vacuo. The residue was purified by chromatography (AcOEt/CH₃CN 95/5) to provide the protected azepane (yellow oil, 883 mg, 50%). This product was dissolved in CH₃CN (23 mL) at room temperature. K₂CO₃ (4 eq.,

8.24 mmol, 1.14 g) was added and then chloroacetyl chloride (4 eq., 8.24 mmol, 659 μL) was added dropwise. Stirring was maintained for 24 h at room temperature and the reaction mixture was concentrated *in vacuo*. The residue was dissolved in CH_2Cl_2 and H_2O and extracted with CH_2Cl_2 (2 x 50 mL). The combined organic layers were dried (MgSO_4) and concentrated *in vacuo* to give an oily residue. The residue was purified by chromatography (AcOEt/cyclohexane 1/1) to provide chloroacetamide **15** (yellow oil, 729 mg, 70 %). $[\alpha]_{\text{D}}^{20}$ -240 (c 1.1, CH_2Cl_2); IR (KBr) ν_{max} 3322, 2927, 1637, 1538, 1442, 1341, 1163 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 7.63 (1 H, d, $J = 7.7$ Hz, CH), 7.48 (1 H, d, $J = 7.9$ Hz, CH), 7.42 (1 H, d, $J = 7.8$ Hz, CH), 7.24 (1 H, d, $J = 7.9$ Hz, CH), 7.18 (1 H, t, $J = 7.5$ Hz, CH), 7.09 (2 H, m, 2CH), 6.98 (1 H, t, $J = 7.7$ Hz, CH), 6.43 (1 H, s, CH), 6.06 (1 H, d, $J = 8.4$ Hz, SO_2NH), 5.73 (1 H, d, $J = 6.9$ Hz, CH), 4.33 (1 H, dd, $J_1 = 8.4$ Hz, $J_2 = 6.9$ Hz, CH), 4.10 (2 H, s, CH_2), 3.50 (1 H, bd, $J = 13.5$ Hz, CH_2), 3.39 (3 H, s, CH_3), 3.14 (1 H, td, $J_1 = 13.5$ Hz, $J_2 = 2.2$ Hz, CH_2), 2.24 (1 H, dd, $J_1 = 13.6$ Hz, $J_2 = 7.2$ Hz, CH_2), 2.03 (1 H, m, CH_2), 1.50-1.90 (4 H, m, CH_2); ^{13}C NMR (CDCl_3 , 100 MHz): δ 167.8, 148.9, 137.5, 136.5, 134.2, 132.7, 132.0, 130.2, 126.7, 124.5, 122.7, 120.0, 119.1, 109.7, 102.2, 60.0, 55.0, 43.8, 40.9, 35.1, 29.8, 25.3; MS (CI) m/z 505.1 (MH^+); HRMS (CI) calcd for (MH) $^+$, 505.1312, found for (MH) $^+$: 505.1315.

10-(1-Methyl-1H-indol-2-yl)-7-(2-nitrobenzenesulfonyl)-1,7-diazabicyclo[4.3.1]decan-9-one **16**

To a solution of chloroacetamide **15** (0.129 mmol, 65 mg) in CH_3CN (2 mL), K_2CO_3 (2 eq., 0.258 mmol, 36 mg) and KI (2 eq., 0.258 mmol, 43 mg) were added at room temperature under nitrogen. The reaction mixture was stirred for 24 h at reflux and concentrated *in vacuo*. The residue was diluted in CH_2Cl_2 and H_2O and extracted with CH_2Cl_2 (2 x 10 mL). The combined organic layers were dried (MgSO_4) and concentrated *in vacuo*. The residue was purified by chromatography (AcOEt/cyclohexane 7/3) to provide piperazinone **16** (yellow oil, 30 mg, 50 %). $[\alpha]_{\text{D}}^{20}$ +285 (c 0.6, CH_2Cl_2); IR (KBr) ν_{max} 2944, 1679, 1543, 1470, 1418, 1372, 1166, 1127, 1067 cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz): δ 8.06 (1 H, dd, $J_1 = 4.2$ Hz, $J_2 = 2.6$ Hz, CH), 7.71 (2 H, td, $J_1 = 4.2$ Hz, $J_2 = 1.1$ Hz, 2CH), 7.61 (2 H, m, 2CH), 7.30 (2 H, m, 2CH), 7.16 (1 H, td, $J_1 = 7.8$ Hz, $J_2 = 1.1$ Hz, CH), 6.70 (1 H, s, CH), 5.00 (1 H, s, CH), 4.92 (1 H, s, CH), 4.32 (2 H, m, CH), 3.97 (1 H, d, $J = 14.2$ Hz, CH_2), 3.63 (3 H, s, CH_3), 2.52 (1 H, dd, $J_1 = 13.7$ Hz, $J_2 = 8.4$ Hz, CH_2), 2.35 (2 H, m, CH_2), 1.72 (2 H, m, CH_2), 1.36 (2 H, m, CH_2); ^{13}C NMR (CDCl_3 , 100 MHz): δ 169.9, 148.8, 138.0, 134.8, 132.3, 132.0, 131.6, 131.1, 127.2, 124.8, 123.3, 121.5, 120.6, 109.6, 103.3, 56.5, 56.2, 48.5, 41.2, 35.6, 30.4, 27.3, 20.4; MS (CI) m/z 469 (MH^+); HRMS (CI) calcd for (MH) $^+$, 469.1546, found for (MH) $^+$: 469.1547.

10

15

8a 322 carbon
C13PPT1 C0C13 u npt1 1