Supporting Information

The Synthesis of β -Substituted α -Amino Acids via Lewis Acid Promoted Radical Conjugate Additions to α,β -Unsaturated α -Nitro Esters and Amides

G. S. C. Srikanth and Steven L. Castle*

*Department of Chemistry and Biochemistry
Brigham Young University, Provo, Utah, 84602*

General Experimental Details

Methylene chloride and ether were dried by passage through a Glass Contour solvent drying system containing cylinders of activated alumina.¹ Flash chromatography was carried out using 60–230 mesh silica gel. ¹H NMR spectra were obtained on Varian 200 MHz or 500 MHz spectrometers with chloroform (7.27 ppm) as internal reference. Signals are reported as follows: s (singlet), d (doublet), t (triplet), q (quartet), dd (doublet of doublets), dt (doublet of triplets), br s (broad singlet), m (multiplet). Coupling constants are reported in hertz (Hz). ¹³C NMR spectra were obtained on a Varian spectrometer operating at 125 MHz with chloroform (77.23 ppm) as internal reference. Infrared spectra were obtained on a Nicolet Avatar 360 FT-IR Spectrometer. Mass spectral data (EI, FAB) were obtained from the Brigham Young University mass spectrometry facility.

Synthesis of α,β -unsaturated α -nitro esters 1a–d. Compounds **1b** and **1c** were prepared using the Knoevenagel condensation conditions reported by Montgomery for the synthesis of **1a** and **1d**.²

¹ Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometallics* **1996**, *15*, 1518.

² Fornicola, R. S.; Oblinger, E.; Montgomery, J. J. *Org. Chem.* **1998**, *63*, 3528.

Methyl 2-Nitro-3-phenylacrylate (1b) (2:1 mixture of diastereomers): Pale yellow oil, 76% yield; ^1H NMR (CDCl_3 , 500 MHz) δ 8.11 and 7.56 (2s, 1H), 7.52–7.47 (m, 5H), 3.97 and 3.93 (2s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 161.9 and 159.9, 137.2, 133.5, 132.7 and 132.5 (2C), 130.6 and 130.0, 129.6 (2C), 129.0, 53.9 and 53.8; IR (film) ν_{max} 3016, 2892, 1732, 1643, 1601, 1229 cm^{-1} ; HRMS (EI) m/z 207.0534 (M^+ , $\text{C}_{10}\text{H}_9\text{NO}_4$ requires 207.0531).

Methyl 3-(4-fluorophenyl)-2-nitroacrylate (1c) (2:1 mixture of diastereomers): White solid, 66% yield; ^1H NMR (CDCl_3 , 500 MHz) δ 8.06 and 7.51 (2s, 1H), 7.45–7.42 (m, 2H), 7.15–7.10 (m, 2H), 3.97 and 3.92 (2s, 3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 165.0 (d, $J = 254$ Hz), 161.8 and 159.8, 141.6, 139.7, 136.0 and 135.9, 132.6 and 132.5, 132.3 and 132.0, 125.3, 117.0 (d, $J = 22$ Hz) and 116.9 (d, $J = 16$ Hz), 53.9 and 53.7; IR (film) ν_{max} 3023, 2959, 1732, 1644, 1603, 1229 cm^{-1} ; HRMS (EI) m/z 225.0439 (M^+ , $\text{C}_{10}\text{H}_8\text{FNO}_4$ requires 225.0438).

Scheme S1. Synthesis of α,β -unsaturated α -nitro amides **4a–c**.

***N*-Benzyl-2-iodoacetamide (S1).** To a solution of *N*-Benzyl-2-chloroacetamide (2.1 g, 11.4 mmol) in acetone (20 mL) was added NaI (2.05 g, 13.6 mmol). The resulting solution was stirred at rt for 6–7 h. The mixture was concentrated in vacuo, and the residue was dissolved in CHCl_3 (20 mL) and washed with H_2O (20 mL). The aqueous layer was extracted with CHCl_3 (10 mL), and the combined organic layers were dried

(Na₂SO₄) and concentrated in vacuo. The remaining solid was washed with Et₂O (20 mL) and dried under high vacuum to give **S1** (2.7 g, 9.6 mmol, 85%) as a white solid: ¹H NMR (CDCl₃, 500 MHz) δ 7.37–7.26 (m, 5H), 6.44 (br s, 1H), 4.46 (m, 2H), 3.72 (m, 2H); ¹³C NMR (CDCl₃, 125 MHz) δ 167.1, 137.7, 129.0 (2C), 127.9 (3C), 44.6, –0.36; IR (film) ν_{max} 2971, 1730, 1620, 1340 cm^{–1}; HRMS (FAB) *m/z* 275.9885 (MH⁺, C₉H₁₀INOH requires 275.9885).

***N*-Benzyl-2-nitroacetamide (S2).** To a solution of **S1** (1.0 g, 3.6 mmol) in THF–Et₂O (1:1, 20 mL) was added AgNO₂ (0.83 g, 5.4 mmol). The resultant mixture was stirred at rt for 16–18 h and filtered through a pad of Celite. The solution was concentrated in vacuo, and the residue was purified by chromatography (SiO₂, 5 × 18 cm, CH₂Cl₂ elution) to give **S2** (0.60 g, 3.0 mmol, 84%) as a white solid: ¹H NMR (CDCl₃, 500 MHz) δ 7.37–7.26 (m, 5H), 6.87 (br s, 1H), 5.09 (s, 2H), 4.47 (d, *J* = 5.5 Hz, 2H); ¹³C NMR (CDCl₃, 125 MHz) δ 160.2, 136.8, 129.1 (2C), 128.2, 128.0 (2C), 77.9, 44.3; IR (film) ν_{max} 2975, 1733, 1628, 1360 cm^{–1}; HRMS (EI) *m/z* 195.0760 (MH⁺, C₉H₁₀N₂O₃H requires 195.0769).

General procedure for Knoevenagel condensation of S2 with aldehydes. To a stirred solution of **S2** (0.60 g, 3.1 mmol) and aldehyde (4.6 mmol) in anhydrous THF (25 mL) at –10 °C was added TiCl₄ (1.0 M solution in CH₂Cl₂, 5.2 mL, 5.2 mmol). The resultant yellow mixture was stirred for 10 min and treated with *N*-methyl morpholine (1.3 mL, 1.2 g, 12.4 mmol) dropwise over 20 min. The mixture was then allowed to warm to rt and stirred for an additional 18 h. The reaction was quenched with H₂O (20 mL), and the organic layer was separated. The aqueous layer was extracted with THF (2 × 15 mL),

and the combined organic extracts were washed with brine (15 mL), dried (Na₂SO₄), and concentrated in vacuo. Flash chromatography (5 × 15 cm, 5–20% EtOAc in hexanes gradient elution) afforded α,β -unsaturated methyl nitroacetamides **4a–c** as single isomers of undetermined olefin stereochemistry.

***N*-Benzyl-3-(4-methoxyphenyl)-2-nitroacrylamide (4a):** Yellow solid, 0.68 g, 2.0 mmol, 66% yield; ¹H NMR (CDCl₃, 500 MHz) δ 8.02 (s, 1H), 7.47 (d, *J* = 9.0 Hz, 2H), 7.36–7.25 (m, 5H), 6.83 (d, *J* = 9.0 Hz, 2H), 6.33 (br s, 1H), 4.62 (d, *J* = 6.0 Hz, 2H), 3.86 (s, 3H); ¹³C NMR (CDCl₃, 125 MHz) δ 163.3, 161.0, 142.2, 136.8, 136.7 (2C), 133.4 (2C), 129.0 (2C), 128.4 (2C), 121.6, 115.0 (2C), 55.7, 44.5; IR (film) ν_{max} 3287, 2933, 1660, 1603, 1516, 1028 cm⁻¹; HRMS (FAB) *m/z* 335.1001 (MNa⁺, C₁₇H₁₆N₂O₄Na requires 335.1007).

***N*-Benzyl-2-nitro-3-phenylacrylamide (4b):** Pale yellow solid, 0.55 g, 1.8 mmol, 58% yield; ¹H NMR (CDCl₃, 500 MHz) δ 8.01 (s, 1H), 7.40–7.21 (m, 10 H), 6.32 (br s, 1H), 4.66 (d, *J* = 6.0 Hz, 2H); ¹³C NMR (CDCl₃, 125 MHz) δ 160.5, 144.4, 136.7, 136.4, 132.4, 130.9 (2C), 129.6, 129.5, 129.3, 129.0 (2C), 128.3 (2C), 128.1, 44.5; IR (film) ν_{max} 3280, 2946, 1658, 1601, 1516, 1078 cm⁻¹; HRMS (FAB) *m/z* 305.0901 (MNa⁺, C₁₆H₁₄N₂O₃Na requires 305.0902).

***N*-Benzyl-3-(4-fluorophenyl)-2-nitroacrylamide (4c):** Pale yellow solid, 0.50 g, 1.5 mmol, 49% yield; ¹H NMR (CDCl₃, 500 MHz) δ 8.01 (s, 1H), 7.52–7.50 (m, 2H), 7.36–7.26 (m, 5H), 7.04–7.01 (m, 2H), 6.38 (br s, 1H), 4.60 (d, *J* = 6.0 Hz, 2H); ¹³C NMR (CDCl₃, 125 MHz) δ 165.1 (d, *J* = 255 Hz), 160.3, 144.1, 136.6, 135.4 (2C), 133.4, 133.3, 129.1 (2C), 128.4, 128.2, 125.4, 116.8 (d, *J* = 22 Hz, 2C), 44.5; IR (film) ν_{max}

3023, 2959, 1732, 1644, 1603, 1229 cm^{-1} ; HRMS (FAB) m/z 323.0796 (MNa^+ , $\text{C}_{16}\text{H}_{13}\text{FN}_2\text{O}_3\text{Na}$ requires 323.0808).

General procedure for radical conjugate additions to 1a–c. To a solution of **1** (0.12 mmol) in CH_2Cl_2 – Et_2O (4:1, 1 mL) was added a Lewis acid (0.14 mmol). The resultant mixture was stirred at rt for 30 min. The clear solution was cooled to $-78\text{ }^\circ\text{C}$ and treated with Bu_3SnH (77 μL , 81 mg, 0.25 mmol), *i*-PrI (71 μL , 106 mg, 0.63 mmol) and Et_3B (1.0 M solution in hexanes, 0.66 mL, 0.63 mmol). O_2 was added via syringe (2 mL) every 30 min, and the reaction was stirred at $-78\text{ }^\circ\text{C}$ for 3h. The mixture was diluted with CH_2Cl_2 (10 mL) followed by 0.25 N HCl (5 mL), and the layers were separated. The organic layer was dried (Na_2SO_4) and concentrated in vacuo. Flash chromatography (SiO_2 , $2 \times 12\text{ cm}$, 0–0.75% EtOAc in hexanes gradient elution) afforded the adducts as colorless oils (**2a–c**) or the reduced compounds (**3a**, **3c**) along with tin impurities. Analytically pure samples were prepared by subjecting the compound to further chromatography (same conditions).

Methyl 3-(4-Methoxyphenyl)-4-methyl-2-nitropentanoate (2a) (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.11–7.01 (m, 2H), 6.86–6.82 (m, 2H), 5.60 and 5.56 (2d, $J = 10.5$ and 11 Hz, 1H), 3.85 and 3.79 (2s, 3H), 3.78 and 3.52 (2s, 3H), 3.60–3.55 (m, 1H), 2.10–2.02 (m, 1H), 0.92–0.84 (m, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 164.8 and 164.0, 159.2, 132.5, 130.8, 130.4, 127.0 and 126.8, 113.9, 91.2 and 90.8, 55.4, 53.8 and 53.4, 51.8, 29.5 and 29.4, 21.7 and 21.2, 18.1 and 18.0; IR (film) ν_{max} 3240, 1656, 1545 cm^{-1} ; HRMS (EI) m/z 281.1256 (M^+ , $\text{C}_{14}\text{H}_{19}\text{NO}_5$ requires 281.1263).

Methyl 2-Deuterio-3-(4-methoxyphenyl)-4-methyl-2-nitropentanoate (*d*-2a):

^1H NMR (CDCl_3 , 500 MHz) was identical to **2a** except for the absence of the two doublets at δ 5.60 and 5.56. Also, the multiplet at δ 3.60–3.55 appeared as two doublets centered at δ 3.60 and 3.56 ($J = 6.5$ Hz).

Methyl 4-Methyl-2-nitro-3-phenylpentanoate (2b) (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.43–7.14 (m, 5H), 5.66 and 5.62 (2d, $J = 11$ and 10.5 Hz, 1H), 3.93 and 3.86 (2s, 3H), 3.68–3.59 (m, 1H), 2.06–2.00 (m, 1H), 0.86 and 0.79 (2d, $J = 7.0$ Hz, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 164.7 and 163.9, 135.2 and 135.1, 132.7 and 132.5, 130.6 and 130.0, 129.7 and 129.6, 129.3 and 129.0, 128.4 and 127.9, 91.0 and 90.8, 53.8 and 53.6, 52.5 and 52.4, 29.6 and 29.4, 21.7 and 21.1, 18.1 and 18.0; IR (film) ν_{max} 3242, 1651, 1555 cm^{-1} ; HRMS (FAB) m/z 274.1054 (MNa^+ , $\text{C}_{13}\text{H}_{17}\text{NO}_4\text{Na}$ requires 274.1055).

Methyl 2-Deuterio-4-methyl-2-nitro-3-phenylpentanoate (*d*-2b): ^1H NMR (CDCl_3 , 500 MHz) was identical to **2b** except for the absence of the two doublets at δ 5.66 and 5.62. Also the multiplet at δ 3.68–3.59 appeared as two doublets centered at δ 3.66 and 3.61 ($J = 8.5$ Hz).

Methyl 3-(4-Fluorophenyl)-4-methyl-2-nitropentanoate (2c) (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.49–7.00 (m, 4H), 5.62 and 5.59 (2d, $J = 11.0$ and 10.5 Hz, 1H), 3.89 and 3.88 (2s, 3H), 3.63 and 3.58 (two dd, $J = 9.0$, 15.5 Hz and 5.0, 12.0 Hz, 1H), 2.06–2.02 (m, 1H), 0.90 and 0.86 (2d, $J = 11.0$ and 11.5 Hz, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 165.7 (d, $J = 272$ Hz), 164.3 and 160.8, 132.5 and 132.4, 131.3 and 131.2, 131.0 and 130.9, 117.0 and 115.5 (2d, $J = 37$ and 35 Hz, 2C), 90.9 and

90.7, 53.9 and 53.8, 53.5 and 51.8, 29.6 and 29.4, 21.7 and 21.1, 18.1 and 17.9; IR (film) ν_{max} 3267, 1655, 1551 cm^{-1} .

Methyl 2-Deuterio-3-(4-fluorophenyl)-4-methyl-2-nitropentanoate (*d*-2c): ^1H NMR (CDCl_3 , 500 MHz) was identical to **2c** except for the absence of the two doublets at δ 5.62 and 5.59. Also the two doublet of doublets at δ 3.63 and 3.58 appeared as two doublets ($J = 9.0$ and 9.5 Hz).

Methyl 3-(4-Fluorophenyl)-2-nitropropionate (3c): ^1H NMR (CDCl_3 , 500 MHz) δ 7.20–7.17 (m, 2H), 7.03–7.05 (m, 2H), 5.31 (dd, $J = 5.0, 9.0$ Hz, 1H), 3.84 (s, 3H), 3.53 (dd, $J = 6.0, 14.5$ Hz, 1H), 3.44 (dd, $J = 5.5, 14.0$ Hz, 1H). This compound was obtained as a mixture with **2c** and hence was not fully characterized.

General procedure for radical conjugate additions to 4a–c: To a solution of **4** (0.080 mmol) in CH_2Cl_2 –THF (1:1, 1.5 mL) was added a Lewis acid (0.088 mmol). The resultant mixture was stirred at rt for 30 min. The clear solution was cooled to -78 °C and treated with Bu_3SnH (50 μL , 52 mg, 0.17 mmol), *i*-PrI (47 μL , 68 mg, 0.40 mmol), and Et_3B (1.0 M solution in hexanes, 0.41 mL, 0.40 mmol). O_2 was added via syringe (2 mL) every 30 min, and the reaction was stirred at -78 °C for 1–4 h. The mixture was diluted with CH_2Cl_2 (15 mL) followed by 1 N HCl (5 mL), and the layers were separated. The organic layer was dried (Na_2SO_4) and concentrated in vacuo. Flash chromatography (SiO_2 , 2×12 cm, 2–8% EtOAc in hexanes gradient elution) afforded mixtures of **5** and **6** as white solids. Analytically pure samples of **5a–c** were prepared by washing the solid with cold hexanes. Reduction products **6a–c** were contaminated with **5a–c** or unreacted starting material and thus are not fully characterized.

***N*-Benzyl-3-(4-methoxyphenyl)-4-methyl-2-nitropentanamide (5a)** (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.36–7.30 (m, 5H), 7.03–7.01 (m, 2H), 6.87 and 6.40 (2br s, 1H), 6.83–6.78 (m, 2H), 5.59 and 5.57 (2d, J = 10.0 and 10.5 Hz, 1H), 4.45 (dd, J = 5.5, 15 Hz, 1H), 4.27–4.23 (m, 1H), 3.81 and 3.77 (2s, 3H), 3.50 and 3.44 (2dd, J = 5.0, 11 Hz and 5.0, 10.0 Hz, 1H), 2.04–2.03 and 1.99–1.95 (2m, 1H), 0.94–0.83 (m, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 162.9 and 162.2, 159.4, 137.0 and 136.9, 130.9 and 130.3, 129.1 and 128.8 (2C), 128.1, 127.9 and 127.85 (2C), 127.80 (2C), 126.8 and 126.2, 114.0 and 113.9, 93.1 and 92.9, 55.4, 54.0 and 53.3, 44.2 and 44.0, 29.5 and 28.9, 21.8 and 21.3, 18.2 and 17.6; IR (film) ν_{max} 3259, 1659, 1552 cm^{-1} ; HRMS (FAB) m/z 379.1647 (MNa^+ , $\text{C}_{20}\text{H}_{24}\text{N}_2\text{O}_4\text{Na}$ requires 379.1634).

***N*-Benzyl-4-Methyl-2-Nitro-3-phenylpentanamide (5b)** (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.36–7.14 (m, 9H), 6.89 and 6.41 (2br s, 1H), 6.85 and 6.80 (2d, J = 7.0 Hz, 1H), 5.64 and 5.62 (2d, J = 7.5 and 10.5 Hz, 1H), 4.51–4.33 (m, 1H), 4.24–4.15 (m, 1H), 3.57–3.51 (m, 1H), 2.12–2.10 and 2.10–1.98 (2m, 1H), 0.98–0.85 (m, 6H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 162.8 and 162.1, 137.0 and 136.8, 134.9 and 134.5, 129.9 and 129.3, 129.1 and 128.8 (2C), 128.7, 128.6 and 128.5 (2C), 128.18 and 128.10, 127.9 and 127.7, 128.0 and 127.8 (2C), 93.0 and 92.8, 54.4 and 54.0, 44.3 and 44.0, 29.5 and 28.9, 21.8 and 21.3, 18.1 and 17.7; IR (film) ν_{max} 3221, 1650, 1564 cm^{-1} ; HRMS (FAB) m/z 327.1704 (MH^+ , $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_3\text{H}$ requires 327.1709).

***N*-Benzyl-3-(4-fluorophenyl)-4-methyl-2-nitropentanamide (5c)** (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.38–6.84 (m, 9H), 6.53 (br s, 1H), 5.62 and 5.58 (2d, J = 9.0 and 10.5 Hz, 1H), 4.86–4.82 (m, 1H), 4.24 (dd, J = 3.0, 9.0 Hz, 1H), 3.58–3.56 (m, 1H), 2.10–1.99 (m, 1H), 0.92 (d, J = 6.5 Hz, 3H), 0.79 (d, J = 7.0 Hz,

3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 163.1 (d, $J = 283$ Hz), 162.7 and 160.9, 136.9 and 136.8, 131.5 and 131.4, 130.9 and 130.8, 129.1 and 128.9 (2C), 128.2 (2C), 128.0 and 127.8 (2C), 115.6 and 115.4 (2d, $J = 26$ and 27 Hz, 2C), 92.8 and 92.6, 53.7 and 53.3, 44.3 and 44.0, 28.1 and 27.1, 21.7 and 21.3, 17.78 and 17.73; IR (film) ν_{max} 3261, 1650, 1561 cm^{-1} ; HRMS (FAB) m/z 367.1432 (MNa^+ , $\text{C}_{19}\text{H}_{21}\text{FN}_2\text{O}_3\text{Na}$ requires 367.1434).

***N*-Benzyl-3-(4-methoxyphenyl)-2-nitropropionamide (6a):** ^1H NMR (CDCl_3 , 500 MHz) δ 7.41–7.39, (m, 5H), 7.21–7.19 (m, 2H), 6.90–6.88 (m, 2H), 6.81 (br s, 1H), 5.28 (dd, $J = 8.5, 11.5$ Hz, 1H), 4.55–4.52 (m, 2H), 3.79 (s, 3H), 3.52 (m, 2H).

***N*-Benzyl-3-phenyl-2-nitropropionamide (6b):** ^1H NMR (CDCl_3 , 500 MHz) δ 7.37–7.22 (m, 10H), 6.73 (br s, 1H), 5.27 (dd, $J = 8.0, 10.0$ Hz, 1H), 4.56–4.54 (m, 2H), 3.58–3.54 (m, 2H).

***N*-Benzyl-3-(4-fluorophenyl)-2-nitropropionamide (6c):** ^1H NMR (CDCl_3 , 500 MHz) δ 7.38–7.29 (m, 5H), 6.84–6.78 (m, 4H), 6.50 (br s, 1H), 5.22 (dd, $J = 9.0, 12$ Hz, 1H), 4.57–4.54 (m, 2H), 3.57 (dd, $J = 9.5, 14.5$ Hz, 1H), 3.55–3.52 (m, 1H).

General procedure for nitro reduction and *N*-Cbz protection: A solution of **2** (0.12 mmol) or **5** (0.080 mmol) in degassed MeOH (7 mL) was treated with 10% Pd/C (20 mg). The resulting mixture was stirred at rt under H_2 (1 atm) for 12 hr, then filtered through a pad of Celite and concentrated in vacuo. The crude amine was dissolved in dioxane– H_2O (10:1, 1 mL), treated with NaHCO_3 (20 mg, 0.20 mmol), and stirred at rt for 10 min. Benzyl chloroformate (15 μL , 23 mg, 0.12 mmol) was added to the mixture, and it was stirred for 1 h then concentrated in vacuo. The residue was taken up in CH_2Cl_2 (10 mL) and H_2O (5 mL), the layers were separated, and the aqueous layer was extracted

with CH₂Cl₂ (2 × 5 mL). The combined organic layers were dried (Na₂SO₄) and concentrated in vacuo. Flash chromatography (SiO₂, 2 × 12 cm, 5–15% EtOAc in hexanes gradient elution) afforded **7** or **8** as colorless oils.

Methyl 2-Benzyloxycarbonylamino-3-(4-methoxyphenyl)-4-methylpentanoate (7a) (1:1 mixture of diastereomers): ¹H NMR (CDCl₃, 500 MHz) δ 7.38–7.34 (m, 5H), 6.98 and 6.94 (2d, *J* = 9.0 and 8.5 Hz, 2H), 6.84–6.81 (m, 2H), 5.11 (s, 2H), 5.00 (br s, 1H), 4.92–4.90 and 4.85–4.84 (2m, 1H), 3.78 (s, 3H), 3.65 and 3.63 (2s, 3H), 2.84–2.79 and 2.62–2.59 (2m, 1H), 2.26–2.23 and 2.08–2.03 (2m, 1H), 1.22–1.19 and 0.79–0.72 (m, 6H); ¹³C NMR (CDCl₃, 125 MHz) δ 172.7 and 171.7, 158.9, 155.8, 141.1, 136.5 and 136.4, 130.5, 130.1, 129.7 (2C), 128.8 and 128.7, 128.4 and 128.3 (2C), 127.9 and 127.2, 114.2 and 113.9, 67.3 and 67.1, 65.6, 56.2 and 56.0, 55.3 and 54.5, 52.3 and 52.1, 29.9 and 29.0, 28.5, 21.5 and 21.1; IR (film) ν_{max} 3346, 2959, 1726, 1602, 1222, 1048 cm⁻¹; HRMS (FAB) *m/z* 386.1975 (MH⁺, C₂₂H₂₇NO₅H requires 386.1967).

Methyl 2-Benzyloxycarbonylamino-2-deuterio-3-(4-methoxyphenyl)-4-methylpentanoate (d-7a): ¹H NMR (CDCl₃, 500 MHz) was identical to **7a** except for the absence of the two signals at δ 4.92–4.90 and 4.85–4.84. Also, the multiplets at δ 2.84–2.79 and 2.62–2.59 appeared as two doublets centered at δ 2.82 and 2.61 (*J* = 10.0 and 10.5 Hz). HRMS (FAB) *m/z* 387.2018 (M⁺, C₂₂H₂₆DNO₅H requires 387.2012).

Methyl 2-Benzyloxycarbonylamino-4-methyl-3-phenylpentanoate (7b) (1:1 mixture of diastereomers): ¹H NMR (CDCl₃, 500 MHz) δ 7.42–7.36 (m, 8H), 7.07 and 7.02 (2d, *J* = 8.0 Hz, 2H), 5.12 (s, 2H), 5.03 (br s, 1H), 4.99–4.97 and 4.92–4.89 (2m, 1H), 3.64 and 3.63 (2s, 3H), 2.88 and 2.66 (2dd, *J* = 3.5, 9.5 Hz and 6.0, 9.0 Hz, 1H),

2.31–2.28 and 2.15–2.10 (2m, 1H), 1.18 and 0.97, (2d, $J = 5.0$ and 6.0 Hz, 3H), 0.79 and 0.75 (2d, $J = 6.5$ Hz, 3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 172.6 and 171.6, 156.5 and 155.7, 138.7 and 138.6, 136.5 and 136.4 (2C), 129.2 (3C), 128.8 and 128.7 (2C), 128.5 and 128.4 (2C), 128.3, 127.5 and 127.4, 67.3 and 67.2, 57.1 and 56.0, 55.9 and 55.4, 52.3 and 52.1, 28.9 and 28.3, 21.5 and 21.4, 21.2; IR (film) ν_{max} 3337, 2941, 1718, 1603, 1034 cm^{-1} ; HRMS (FAB) m/z 378.1691 (MNa^+ , $\text{C}_{21}\text{H}_{25}\text{NO}_4\text{Na}$ requires 378.1701).

Methyl 2-Benzyloxycarbonylamino-2-deuterio-4-methyl-3-phenylpentanoate (*d*-7b): ^1H NMR (CDCl_3 , 500 MHz) was identical to **7b** except for the absence of the two signals at δ 4.99–4.97 and 4.92–4.89. Also the two doublet of doublets at δ 2.88 and 2.66 appeared as doublets ($J = 10.5$ Hz). HRMS (FAB) m/z 379.1744 (MNa^+ , $\text{C}_{21}\text{H}_{24}\text{DNO}_4\text{Na}$ requires 379.1744).

Methyl 2-Benzyloxycarbonylamino-3-(4-fluorophenyl)-4-methylpentanoate (7c) (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.41–7.38 (m, 5H), 7.31–7.26 (m, 4H), 5.12 (s, 2H), 4.96–4.94 (br s, 1H, obscured), 4.96–4.94 and 4.88–4.85 (2m, 1H), 3.77 and 3.65 (2s, 3H), 2.87–2.81 and 2.69–2.67 (2m, 1H), 2.27–2.24 and 2.09–2.05 (2m, 1H), 1.18 and 1.14 (2d, $J = 6.5$ Hz, 3H), 0.74 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 172.5 and 171.6, 162.2 (d, $J = 244$ Hz), 156.5 and 155.7, 136.4 and 134.4, 130.6 and 130.2 (2C), 128.9, 128.6 (3C), 128.4, 128.2, 115.7 and 115.4 (2d, $J = 21$ Hz, 2C), 67.4 and 67.2, 56.2 and 56.1, 55.8 and 52.5, 52.3 and 52.1, 28.7 and 28.6, 21.5, 21.4 and 21.0; IR (film) ν_{max} 3340, 2949, 1728, 1603, 1048 cm^{-1} ; HRMS (FAB) m/z 396.1593 (MNa^+ , $\text{C}_{21}\text{H}_{24}\text{FNO}_4\text{Na}$ requires 396.1587).

Methyl 2-Benzyloxycarbonylamino-2-deuterio-3-(4-fluorophenyl)-4-methylpentanoate (*d*-7c): ^1H NMR (CDCl_3 , 500 MHz) was identical to **7c** except for

the absence of the two multiplets at δ 4.96–4.94 and 4.88–4.85 (the amide resonance which was obscured by the multiplet at δ 4.96–4.94 became visible as a broad singlet). Also, the two multiplets at δ 2.87–2.81 and 2.69–2.67 appeared as two doublets centered at δ 2.85 and 2.67 (J = 10.5 and 9.0 Hz). HRMS (FAB) m/z 375.1826 (MH^+ , $C_{21}H_{23}DFNO_4H$ requires 375.1831).

***N*-Benzyl 2-Benzyloxycarbonylamino-3-(4-methoxyphenyl)-4-methylpentanamide (8a)** (1:1 mixture of diastereomers): 1H NMR ($CDCl_3$, 200 MHz) δ 7.35–7.21 (m, 10H), 7.05 (d, J = 8.8 Hz, 2H), 6.78 (d, J = 8.0 Hz, 2H), 5.85 (br s, 1H), 5.25 (br s, 1H), 5.12 and 5.09 (2s, 2H), 4.64–4.62 (m, 1H), 4.16–4.13 (m, 1H), 4.11–4.09 (m, 1H), 3.87 (s, 3H), 3.14–3.12 and 2.82–2.79 (2m, 1H), 2.33–2.28 and 2.09–2.05 (2m, 1H), 1.05 and 0.91 (2d, J = 6.0 Hz, 3H), 0.80 (d, J = 6.2 Hz, 3H); ^{13}C NMR ($CDCl_3$, 125 MHz) δ 170.6, 159.4, 156.3, 138.4 and 137.5, 136.3 (2C), 129.8, 128.79 and 128.72 (2C), 128.5 (2C), 128.4, 128.36, 128.32 (3C), 127.8, 127.6, 127.4 and 127.3 (2C), 67.4, 57.2, 55.7 and 55.6, 54.8 and 53.4, 43.5, 31.1 and 28.5, 21.7, 20.7; IR (film) ν_{max} 3281, 2930, 1649, 1520 cm^{-1} ; HRMS (FAB) m/z 483.2252 (MNa^+ , $C_{28}H_{32}N_2O_4Na$ requires 483.2260).

***N*-Benzyl 2-Benzyloxycarbonylamino-2-deuterio-3-(4-methoxyphenyl)-4-methylpentanamide (*d*-8a)**: 1H NMR ($CDCl_3$, 500 MHz) was identical to **8a** except for the absence of the signal at δ 4.64–4.62. Also, the two multiplets at δ 3.14–3.12 and 2.82–2.79 appeared as two doublets centered at δ 3.11 and 2.80 (J = 8.5 and 9.0 Hz). HRMS (FAB) m/z 484.2308 (MNa^+ , $C_{28}H_{31}DN_2O_4Na$ requires 484.2322).

***N*-Benzyl 2-Benzyloxycarbonylamino-4-methyl-3-phenylpentanamide (8b)** (1:1 mixture of diastereomers): 1H NMR ($CDCl_3$, 500 MHz) δ 7.35–7.31 (m, 15H), 6.21

and 5.93 (2br s, 1H), 5.09 and 5.08 (2s, 2H), 4.96 and 4.75 (2 br s, 1H), 4.66 and 4.64 (2d, $J = 9.0$ and 9.5 Hz, 1H), 4.29–4.27 (m, 1H), 4.11–4.08 (m, 1H), 3.20–3.17 and 2.90–2.87 (2m, 1H), 2.35–2.31 and 2.06–2.03 (2m, 1H), 1.09 and 0.92 (2d, $J = 5.5$ and 6.0 Hz, 3H), 0.81 and 0.77 (2d, $J = 6.5$ Hz, 3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 170.9, 156.3, 138.4, 137.6 and 136.3, 129.8 and 129.3, 128.79, 128.72, 128.5, 128.4(2C), 128.36, 128.3(2C), 127.8, 127.7(3C), 127.6, 127.4, 127.3, 67.4, 57.2, 54.8 and 53.6, 43.5, 31.1 and 28.4, 21.8, 20.5; IR (film) ν_{max} 3280, 2933, 1645, 1526, 1251 cm^{-1} ; HRMS (FAB) m/z 453.2141 (MNa^+ , $\text{C}_{27}\text{H}_{30}\text{N}_2\text{O}_3\text{Na}$ requires 453.2155).

***N*-Benzyl 2-Benzylloxycarbonylamino-2-deuterio-4-methyl-3-phenylpentanamide (*d*-8b):** ^1H NMR (CDCl_3 , 500 MHz) was identical to **8b** except for the absence of the two doublets at δ 4.66 and 4.64. Also, the two multiplets at δ 3.20–3.17 and 2.90–2.87 appeared as two doublets centered at δ 3.19 and δ 2.88 ($J = 9.0$ Hz). HRMS (FAB) m/z 454.2184 (MNa^+ , $\text{C}_{27}\text{H}_{29}\text{DN}_2\text{O}_3\text{Na}$ requires 454.2193).

***N*-Benzyl 2-Benzylloxycarbonylamino-3-(4-fluorophenyl)-4-methylpentanamide (**8c**)** (1:1 mixture of diastereomers): ^1H NMR (CDCl_3 , 500 MHz) δ 7.38–7.33 (m, 10H), 7.10–7.09 (m, 2H), 6.96–6.93 (m, 2H), 6.83 (br s, 1H), 5.12 and 5.08 (2s, 2H), 4.89 and 4.71 (2br s, 1H), 4.57 and 4.56 (2d, $J = 9$ Hz, 1H), 4.38–4.35 (m, 1H), 4.10–4.07 (m, 1H), 3.19–3.16 and 2.89–2.86 (2m, 1H), 2.20–2.16 and 2.05–2.02 (2m, 1H), 0.90 (d, $J = 6.5$ Hz, 3H), 0.79 (d, $J = 6.0$ Hz, 3H); ^{13}C NMR (CDCl_3 , 125 MHz) δ 170.6 and 170.4, 164.3 (d, $J = 297$ Hz), 156.3, 137.5 (2C), 136.2, 134.1 (3C), 131.2 (2C), 129.8 and 128.6 (3C), 128.2 (2C), 127.7 (2C), 115.4 (d, $J=20$ Hz, 2C), 67.58 and 67.52, 57.2, 43.6, 43.5, 29.9, 28.4, 21.8; IR (film) ν_{max} 3288, 2946, 1644, 1526, 1254, 1039 cm^{-1} ; HRMS (FAB) m/z 471.2076 (MNa^+ , $\text{C}_{27}\text{H}_{29}\text{FN}_2\text{O}_3\text{Na}$ requires 471.2050).

***N*-Benzyl 2-Benzoyloxycarbonylamino-2-deuterio-3-(4-fluorophenyl)-4-methylpentamide (*d*-8c):** ^1H NMR (CDCl_3 , 500 MHz) was identical to **8c** except the two doublets at δ 4.57 and 4.56 integrated as 0.11H. Also, the multiplet at δ 3.19–3.16 appeared as a doublet centered at δ 3.17 ($J = 9.5$ Hz), and the multiplet at δ 2.89–2.86 appeared as a broad singlet at δ 2.88. HRMS (FAB) m/z 472.2114 (MNa^+ , $\text{C}_{27}\text{H}_{28}\text{DFN}_2\text{O}_3\text{Na}$ requires 472.2123).