Supporting Information for:

Synthesis of a pentasaccharide fragment of varianose, a cell wall polysaccharide from *Penicillium varians*

Yu Bai and Todd L. Lowary*

Alberta Ingenuity Centre for Carbohydrate Science and Department of Chemistry, The

University of Alberta, Gunning-Lemieux Chemistry Centre,

Edmonton, AB T6G 2G2, Canada.

Email: tlowary@ualberta.ca

Table of Contents

General experimental methods	S3
Figure S1. Residue designations in 4 used in assignments of NMR spectra.	S4
Additional experimental details and data for new compounds	S4–S7
References for Supporting Information	S7
¹ H NMR spectrum of 4	S8
¹³ C NMR spectrum (APT) of 4	S9
¹ H NMR spectrum of 5	S10
¹³ C NMR spectrum (APT) of 5	S11
¹ H NMR spectrum of 6	S12
¹³ C NMR spectrum (APT) of 6	S13
¹ H NMR spectrum of 10	S14
¹³ C NMR spectrum (APT) of 10	S15

¹ H NMR spectrum of 12	S16
¹³ C NMR spectrum (APT) of 12	S17
¹ H NMR spectrum of 13	S18
¹³ C NMR spectrum (APT) of 13	S19
¹ H NMR spectrum of 14	S20
¹³ C NMR spectrum (APT) of 14	S21
¹ H NMR spectrum of 15	S22
¹³ C NMR spectrum (APT) of 15	S23
¹ H NMR spectrum of 16	S24
¹³ C NMR spectrum (APT) of 16	S25
¹ H NMR spectrum of 17	S26
¹³ C NMR spectrum (APT) of 17	S27
¹ H NMR spectrum of 18	S28
¹³ C NMR spectrum (APT) of 18	S29
¹ H NMR spectrum of 19	S30
¹³ C NMR spectrum (APT) of 19	S31
¹ H NMR spectrum of 20	S32
¹³ C NMR spectrum (APT) of 20	S33
¹ H NMR spectrum of 21	S34
¹³ C NMR spectrum (APT) of 21	S35
¹ H NMR spectrum of 22	S36
¹³ C NMR spectrum (APT) of 22	S37
¹ H NMR spectrum of 23	S38
¹³ C NMR spectrum (APT) of 23	S39
¹ H NMR spectrum of 24	S40
¹³ C NMR spectrum (APT) of 24	S41

General Methods. All reagents used were purchased from commercial sources and were used without further purification unless noted. Solvents used in reactions were purified by successive passage through columns of alumina and copper under nitrogen. Unless stated otherwise, all reactions were carried out under a positive pressure of argon and were monitored by TLC on silica gel G-25 UV₂₅₄ (0.25 mm). Spots were detected under UV light and/or by charring with 10% H₂SO₄ in ethanol, or in acidified ethanolic anisaldehyde. Solvents were evaporated under reduced pressure and below 50 °C (water bath). Column chromatography was performed on Silica Gel 60 (40–60 mM). The ratio between silica gel and crude product ranged from 100:1 to 20:1 (w/w). ¹H NMR spectra were recorded at 500 or 600 MHz, and chemical shifts are referenced to either TMS (0.0, CDCl₃) or HOD (4.78, D₂O). ¹³C NMR APT spectra¹ were recorded at 125 MHz, and ¹³C chemical shifts are referenced to CDCl₃ (77.23, CDCl₃) or external acetone (31.07, D₂O). ¹H NMR data are reported as though they are first order, and the peak assignments were made on the basis of 2D-NMR (¹H–¹H COSY and HMQC) experiments. The monosaccharide residues in the pentasaccharide are labeled A–E as shown in Figure S1 and these letters are maintained in the assignment of NMR spectra of the intermediates synthesized en route to the target, i.e., disaccharide **12** contains only residues A and B. ESI-MS spectra were recorded on samples suspended in THF or CH₃OH and added NaCl. Optical rotations were measured at 22 ± 2 °C at the sodium D line (589 nm) and are in units of deg·mL(dm·g)⁻¹.

Figure S1. Residue designations in **4** used in assignments of NMR spectra.

Methyl 2,3,5-tri-*O*-benzyl- β -D-galactofuranoside (5**).** To a solution of compound **10** (0.89 g, 1.3 mmol) in CH₃OH (5 mL) and CH₂Cl₂ (20 mL) was added *p*-TsOH until the pH of the solution was 3–4. The reaction mixture was stirred for 12 h at rt and then Et₃N was added to neutralize the solution, which was then concentrated to a residue that was purified by chromatography (3:1, hexanes:EtOAc) to provide **5** (0.50 g, 85%). *R*_f 0.25 (2:1, hexanes:EtOAc); [α]_D –47.4 (*c* 0.2, CH₂Cl₂); ¹H NMR (500 MHz, CDCl₃, δ _H) 7.40–7.25 (m, 15 H, Ar), 4.97 (s, 1 H, H-1), 4.70 (d, 1 H, *J* = 11.7 Hz, PhCH₂), 4.65 (d, 1 H, *J* = 11.7 Hz, PhCH₂), 4.57 (d, 1 H, *J* = 11.8 Hz, PhCH₂), 4.54 (d, 1 H, *J* = 11.8 Hz, PhCH₂), 4.49 (d, 1 H, *J* = 11.8 Hz, PhCH₂), 4.44 (d, 1 H, *J* = 11.8 Hz, PhCH₂), 4.22 (dd, 1 H, *J* = 6.8, 5.1 Hz, H-4), 4.04 (dd, 1 H, *J* = 6.8, 2.8 Hz, H-3), 4.01–3.99 (m, 1 H, H-2), 3.79–3.65 (m, 3 H, H-6 $\times$ 2, H-5), 3.39 (s, 3 H, OCH₃), 2.24 (br s, 1 H, OH); ¹³C NMR (125 MHz, CDCl₃, δ _C) 138.3 (Ar), 137.5 (Ar), 137.4 (Ar), 128.5 (Ar), 128.4 (Ar), 128.1 (Ar), 128.03 (Ar), 128.0 (Ar), 127.9 (Ar), 127.8 (Ar), 107.1 (C-1), 87.8 (C-2), 83.1 (C-3), 82.2 (C-4), 78.3 (C-5), 72.8 (PhCH₂), 72.2 (PhCH₂), 72.0 (PhCH₂), 62.2

(C-6), 54.8 (OCH₃). HRMS (ESI) calcd. for (M + Na) C₂₈H₃₂O₆Na: 487.2091, found: 487.2093.

***p*-Tolyl 2,3-di-*O*-acetyl-5,6-*O*-isopropylidene- α/β -D-galactofuranoside (6).** To a solution of **11** (3:7 α/β , 1.32 g, 4.6 mmol) and 2,2-dimethoxypropane (4.5 mL, 37 mmol) in acetone (30 mL) was added *p*-TsOH (9 mg) at rt. The solution was stirred for 4 h and then neutralized with Et₃N. The mixture was concentrated and the resulting oil was dissolved in pyridine (7 mL), cooled to 0 °C and then acetic anhydride (1.3 mL, 14 mmol) was added. The reaction mixture was stirred for 6 h, diluted with CH₂Cl₂ (50 mL), and then washed with a satd aq NaHCO₃ soln (3 $\times$ 30 mL). The organic layer was dried (Na₂SO₄), concentrated and the resulting residue was purified by chromatography (4:1, hexanes:EtOAc) to yield **6** (1.58 g, 3:7 α/β , 84% over two steps) as a colorless oil. *R*_f 0.29 (3:1, hexanes:EtOAc); ¹H NMR (500 MHz, CDCl₃, δ _H) 7.43–7.38 (m, 2 H, Ar), 7.13–7.10 (m, 2 H, Ar), 5.48 (d, 0.27 H, *J* = 4.7 Hz, H-1 α), 5.45 (d, 0.73 H, *J* = 2.7 Hz, H-1 β), 5.42 (dd, 0.27 H, *J* = 4.7, 2.9 Hz, H-2 α), 5.23–5.21 (m, 1 H, H-2 β , H-3 α), 5.17 (dd, 0.73 H, *J* = 5.3, 2.4 Hz, H-3 β), 4.46 (ddd, 0.27 H, *J* = 6.7, 6.6, 6.2 Hz, H-5 α), 4.37 (ddd, 0.73 H, *J* = 6.8, 6.2, 4.8 Hz, H-5 β), 4.29 (dd, 0.73 H, *J* = 5.3, 4.8 Hz, H-4 β), 4.06 (dd, 0.27 H, *J* = 8.2, 6.7 Hz, H-6 α), 4.04 (dd, 0.73 H, *J* = 8.4, 6.8 Hz, H-6 β), 3.92 (dd, 0.27 H, *J* = 6.2, 4.4 Hz, H-4 α), 3.89 (dd, 0.73 H, *J* = 8.4, 6.2 Hz, H-6 β), 3.88 (dd, 0.27 H, *J* = 8.2, 6.6 Hz, H-6 α), 2.32 (s, 3 H, tolyl CH₃), 2.15 (s, 0.81 H, acyl CH₃ α), 2.11 (s, 2.19 H, acyl CH₃ β), 2.08 (s, 2.19 H, acyl CH₃ β), 2.07 (s, 0.81 H, acyl CH₃ α), 1.46 (s, 0.81 H, isopropylidene CH₃ α), 1.40 (s, 2.19 H, isopropylidene CH₃ β), 1.38 (s, 0.81 H, isopropylidene CH₃ α), 1.36 (s, 2.19 H, isopropylidene CH₃ β); ¹³C NMR (125 MHz, CDCl₃, δ _C) 170.0 (C=O β), 169.6 (C=O β), 169.55

(C=O α), 169.5 (C=O α), 138.1 (Ar β), 137.9 (Ar α), 132.9 (Ar β), 132.6 (Ar α), 130.0 (Ar α), 129.8 (Ar β), 129.7 (Ar α), 129.6 (Ar β), 109.9 (isopropylidene C α), 109.8 (isopropylidene C β), 91.0 (C-1 β), 89.7 (C-1 α), 83.4 (C-4 α), 81.7 (C-4 β), 81.3 (C-2 β), 77.6 (C-2 α), 77.2 (C-3 α/β), 75.6 (C-5 α), 75.57 (C-5 β), 65.5 (C-6 α), 65.4 (C-6 β), 26.4 (isopropylidene CH $_3\alpha$), 26.2 (isopropylidene CH $_3\beta$), 25.4 (isopropylidene CH $_3\beta$), 25.2 (isopropylidene CH $_3\alpha$), 21.1 (tolyl CH $_3\alpha/\beta$), 20.8 (acyl CH $_3\alpha/\beta$), 20.7 (acyl CH $_3\beta$), 20.6 (acyl CH $_3\alpha$). HRMS (ESI) calcd. for (M + Na) C $_{20}$ H $_{26}$ O $_7$ SNa: 433.1292, found: 433.1289.

Methyl 2,3,5-tri-*O*-benzyl-6-*O*-trityl- β -D-galactofuranoside (10). To a solution of **9** (2.5 g, 6.9 mmol) in CH $_3$ OH (10 mL) and CH $_2$ Cl $_2$ (10 mL) was added NaOCH $_3$ until the pH was $\sim$ 10. The mixture was stirred for 12 h, neutralized by Amberlite IR-120 H $^+$ resin, filtered, and the filtrate was concentrated. The resulting crude residue was dissolved in pyridine (10 mL), followed by the addition of chlorotriphenylmethane (2.3 g, 8.3 mmol) and DMAP (0.17 g, 1.4 mmol). The reaction mixture was stirred at 50 $^{\circ}$ C for 6 h at which point TLC showed one major spot at R_f 0.59 (EtOAc). DMF (30 mL) was added to the mixture, the solution was cooled to 0 $^{\circ}$ C, and then NaH (1.24 g, 31 mmol) and BnBr (3.0 mL, 24.8 mmol) were added. The resulting mixture was then stirred for 12 h at rt and then a few drops of CH $_3$ OH were added. The resulting solution was diluted with CH $_2$ Cl $_2$ (60 mL), and then washed with a satd aq NaHCO $_3$ soln (3 $\times$ 40 mL). The organic layer was subsequently dried (Na $_2$ SO $_4$), concentrated and the residue was purified by chromatography (8:1, hexanes: EtOAc) to give **10** (3.71 g, 76%) as a light yellow oil. R_f 0.29 (8:1, hexanes: EtOAc); $[\alpha]_D -39.0$ (c 1.0, CH $_2$ Cl $_2$); 1 H NMR (500 MHz, CDCl $_3$, δ_H) 7.50–7.21 (m, 30 H, Ar), 4.94 (s, 1 H, H-1), 4.74 (d, 1 H, J = 11.7 Hz, PhCH $_2$),

4.53 (d, 1 H, $J = 11.8$ Hz, PhCH_2), 4.52 (d, 1 H, $J = 11.7$ Hz, PhCH_2), 4.44 (d, 2 H, $J = 11.8$ Hz, PhCH_2), 4.29 (d, 1 H, $J = 11.8$ Hz, PhCH_2), 4.19 (dd, 1 H, $J = 7.2, 3.5$ Hz, H-4), 3.98–3.95 (m, 2 H, H-2, H-3), 3.74–3.70 (m, 1 H, H-5), 3.46 (dd, 1 H, $J = 9.9, 6.6$ Hz, H-6), 3.38 (dd, 1 H, $J = 9.9, 5.3$ Hz, H-6), 3.34 (s, 3 H, OCH_3); ^{13}C NMR (125 MHz, CDCl_3 , δ_{C}) 144.1 (Ar), 138.5 (Ar), 138.0 (Ar), 137.7 (Ar), 129.2 (Ar), 128.8 (Ar), 128.4 (Ar), 128.33 (Ar), 128.3 (Ar), 128.0 (Ar), 127.9 (Ar), 127.82 (Ar), 127.8 (Ar), 127.7 (Ar), 127.3 (Ar), 127.0 (Ar), 106.9 (C-1), 88.5 (C-2), 87.0 (Ph_3C), 83.0 (C-3), 81.2 (C-4), 77.0 (C-5), 73.4 (PhCH_2), 72.2 (PhCH_2), 71.8 (PhCH_2), 63.9 (C-6), 54.7 (OCH_3). HRMS (ESI) calcd. for ($\text{M} + \text{Na}$) $\text{C}_{47}\text{H}_{46}\text{O}_6\text{Na}$: 729.3187, found: 729.3184.

References

1. (a) Patt, S.; Shoolery, J. N. *J. Magn. Reson.* **1982**, *46*, 535–539; (b) Torres, A. M.; Nakashima, T. T.; McClung, R. E. D. *J. Magn. Reson. Ser. A* **1993**, *101*, 285–289.
2. Veeneman, G. H.; Notermans, S.; Hoogerhout, P.; van Boom, J. H. *Recl. Trav. Chim. Pays-Bas* **1989**, *108*, 344–350.

compound-4
600 MHz 1D in D2O (ref. to 0.1 % ext. acetone @ 2.225 ppm)

Pulse Sequence: zgpg30

compound-4
 125 MHzAPT in D2O (ref. to 1% acetone @ 31.07 ppm)
 C & CH2 opposite side of CH & CH3

compound-5
500 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

compound-5
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

Compound-6
500 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

Compound-6
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

compound-10
500 MHz 1D 1H CDCl3 (ref. to CDCl3 @ 7.26 ppm)

Pulse Sequence: s2pu1

compound-10
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apt

compound-12
600 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

Compound-12
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₃ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

compound-13
600 MHz 1D in CDCl3 (ref. to CDCl3 @ 7.26 ppm)

Pulse Sequence: zgpg30

compound-13
 100 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

compound-14
500 MHz 1D in CDCl3 (ref. to CDCl3 @ 7.26 ppm)

Pulse Sequence: zgpg30

compound-14
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

Compound-15
600 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

compound-15
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

compound-16
500 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

Pulse Sequence: apt

compound-17
500 MHz LD in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

Pulse Sequence: apt

Pulse Sequence: s2pul

compound-18
 125 MHz APT in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apt

compound-19
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

compound-20
500 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

Compound-20
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

compound-21
500 MHz 1D in CDCl3 (ref. to CDCl3 @ 7.26 ppm)

Pulse Sequence: zgpg30

Pulse Sequence: apt

Compound-22
500 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

compound-22
 125 MHz APT in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apt

compound-23
600 MHz 1D in CDCl₃ (ref. to CDCl₃ @ 5.32 ppm)

Pulse Sequence: zgpg30

Compound-23
 125 MHz APF in CDCl₃ (ref. to CDCl₃ @ 53.8 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apc

compound-24
600 MHz ID in CDCl₃ (ref. to CDCl₃ @ 7.26 ppm)

Pulse Sequence: zgpg30

compound-24
 125 MHzAPT in CDCl₃ (ref. to CDCl₃ @ 77.0 ppm)
 C & CH₂ same, CH & CH₃ opposite side of solvent signal

Pulse Sequence: apt