Supporting Information

Synthesis of an Isostere of an *O*-Linked Glycopeptide

Lisa J. Whalen and Randall L. Halcomb*

*Department of Chemistry and Biochemistry,
University of Colorado, Boulder, Colorado 80309-0215*

Table of Contents:

<u>Page</u>	<u>Supporting Information</u>
S2-S29	Experimental details and full characterization of all new compounds.
S30-S62	¹ H and ¹³ C NMR spectra for compounds 7, 8, 9a, 9b, 11, 12, 21, 2, 16, 17, 18, 19, 20, 22, and 23.

General Experimental Section. Commercially available reagents were purchased from Aldrich Chemical Co. and used as supplied unless stated otherwise. Solvents were purified and dried by the standard procedures and distilled prior to use. Silica gel for flash chromatography (230-450 mesh) was not neutral unless otherwise specified; neutral silica gel (60-200 mesh) was used in specific cases noted in the preparation of individual compounds. The same preparative thin-layer chromatography plates (Merck PLC, regular phase, Silica 60 F254, 0.5 mm thickness) were used each time. Proton NMR spectra (^1H NMR) were recorded at 500 MHz or 400 MHz. Chemical shifts are expressed in parts per million (δ) and are referenced to residual protium in the NMR solvent: CD_2HOD , δ 3.31; CHCl_3 , δ 7.24. Carbon NMR (^{13}C NMR) spectra were recorded at 100 MHz. Chemical shifts (δ ppm) are referenced to the carbon signal for the solvent: CD_3OD , 49.15; CDCl_3 , 77.23. Infrared spectra were recorded (16-32 scans, 2-5 mg) on a Nicolet Avatar 360 FT-IR with thin films on KBr plates. Optical rotation measurements were taken on a Jasco P-1030 Polarimeter using a 3.5 mm by 100 mm cell and an aperture of 3 mm.

Benzyl-protected ketone. To 80 mL CH_2Cl_2 was added 1.1 mL (16 mmol) anhydrous DMSO. After cooling the reaction solution to $-78\text{ }^\circ\text{C}$, 1.0 mL (11.9 mmol) oxalyl chloride was added slowly dropwise. After 1 h, a solution of **6** (2.6 g, 4.0 mmol) in 18 mL CH_2Cl_2 at $-78\text{ }^\circ\text{C}$ was transferred to the reaction by cannula with 1 x 2 mL rinse with CH_2Cl_2 . After 1.5 h, 4.5 mL Et_3N (32 mmol) was added to the reaction dropwise and stirred for 15 min at $-78\text{ }^\circ\text{C}$. The reaction was then warmed to $0\text{ }^\circ\text{C}$ and 2 mL glacial

AcOH added. The reaction was partitioned between CH₂Cl₂ and sat. aq. NaHCO₃, the layers were separated, and the aqueous layer washed with 200 mL CH₂Cl₂. The combined organic layers were washed once with 200 mL sat. aq. NaCl, dried over Na₂SO₄, and concentrated. The syrup was purified by flash chromatography on 250 mL silica, eluting with 9:1 hexanes:ethyl acetate, to yield 2.2 g ketone (84%). $[\alpha]_D^{27} = -1.4$ (c 0.83, CHCl₃). IR (cm⁻¹): 3088, 3063, 3031, 2928, 1734. MS (EI): m/z 654.4 (M⁺). ¹H NMR (500 MHz, CDCl₃): δ 7.33-7.25 (m, 20H), 4.66-4.33 (m, 10H), 4.27 (d, $J = 4.4$ Hz, 1H), 4.11 (app t, $J = 4.6$ Hz, 1H), 3.88-3.83 (m, 2H), 3.74 (dd, $J = 4.9, 9.7$ Hz, 1H), 0.91 (s, 9H), 0.06 (s, 3H), 0.05 (s, 3H). ¹³C NMR (CDCl₃): δ 207.7, 138.6, 138.1, 137.6, 137.4, 128.6-127.7, 82.6, 80.3, 80.1, 74.7, 74.5, 73.4, 73.2, 72.8, 62.9, 26.1, 18.4, -5.2.

Vinyl bromide 7.

The above ketone (0.46 g, 0.71 mmol) was dissolved in 5 mL THF and cooled to -78 °C. The ylide was generated by suspending (bromomethyl)triphenylphosphonium bromide (0.62 g, 1.4 mmol) in 13 mL THF, cooling to 0 °C, and adding a solution of potassium *tert*-butoxide (1.4 mL, 1.0 M in 2-methyl-2-propanol) dropwise. After stirring the ylide solution for 5 min at 0 °C, it was transferred by cannula to the ketone solution. Two more portions of ylide were generated identically and added to the ketone solution 15 min and 25 min after the first addition. After stirring 2 h at -78 °C, the reaction was poured through a silica plug that was then eluted with 500 mL Et₂O. The filtrate was then concentrated and purified by flash chromatography, eluting with 1:1 hexanes:CH₂Cl₂

followed by 7:3 CH₂Cl₂:hexanes, yielding 0.17 g (33%, 6:1 *Z:E* ratio) of a clear syrup. The isomers were separated by HPLC (Rainin Dynamax-100 Å Si R0080120C5, eluting with 7:3 CH₂Cl₂:hexanes at 4 mL/min) to provide the pure (*Z*)-vinyl bromide for characterization. $[\alpha]_D^{27} = +8.2$ (*c* 1.4, CHCl₃). IR (cm⁻¹): 3088, 3064, 3031, 2952, 1454, 1100. MS (ESI): *m/z* 731/733 ([M + H]⁺), 753/755 ([M + Na]⁺), 769/771 ([M + K]⁺), 863/865 ([M + Cs]⁺). ¹H NMR (500 MHz, CDCl₃): δ 7.30-7.21 (m, 20H), 6.70 (s, 1H), 4.90 (d, *J* = 8.3 Hz, 1H), 4.65-4.40 (m, 7H), 4.14-4.05 (m, 3H), 3.88 (dd, *J* = 2.4, 8.1 Hz, 1H), 3.76 (ddd, *J* = 2.6, 6.2, 8.9 Hz, 1H), 3.65 (dd, *J* = 6.5, 10.5 Hz, 1H), 3.57 (dd, *J* = 6.3, 10.3 Hz, 1H), 0.85 (s, 9H), -0.025 (s, 3H), -0.034 (s, 3H). ¹³C NMR (CDCl₃): δ 139.5, 139.2, 138.7, 128.6-127.6, 110.1, 79.4, 78.4, 77.1, 74.2, 73.8, 73.1, 70.7, 63.1, 26.1, 18.4.

Deprotected vinyl bromide.

To a solution of vinyl bromide **7** (1.2 g, 1.6 mmol) in 8 mL THF was added 3.2 mL TBAF (3.2 mmol, 1.0 M solution in THF). After 1 h, the reaction was partitioned between ethyl acetate and water. The layers were separated and the aqueous layer washed once with 100 mL ethyl acetate. The combined organic layers were dried over Na₂SO₄ then concentrated to give a clear syrup. The syrup was purified by flash chromatography on 50 mL silica, eluting with 9:1 hexanes:ethyl acetate followed by 8:2 hexanes:ethyl acetate, to yield 0.8 g alcohol (88%). $[\alpha]_D^{27} = +15.7$ (*c* 1.5, CHCl₃). IR (cm⁻¹): 3447, 3030, 2861. MS (ESI): *m/z* 641/643 ([M + Na]⁺). ¹H NMR (500 MHz, CDCl₃): δ 7.31-7.21 (m, 20H), 6.74 (m, 1H), 4.86 (d, *J* = 7.6 Hz, 1H), 4.67 (d, *J* = 11.3

Hz, 1H), 4.58-4.53 (m, 3H), 4.50-4.44 (m, 3H), 4.21 (d, $J = 11.3$ Hz, 1H), 4.13 (dd, $J = 1.6, 13.3$ Hz, 2H), 3.87 (dd, $J = 4.2, 7.7$ Hz, 1H), 3.73-3.70 (m, 1H), 3.67-3.64 (m, 2H), 1.76 (t, $J = 6$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 139.3, 138.6, 138.3, 137.9, 137.8, 128.6-127.8, 110.4, 79.5, 79.2, 77.6, 74.2, 73.3, 73.2, 70.9, 69.9, 62.2.

Aldehyde 8.

To a solution of 0.22 mL oxalyl chloride (2.6 mmol) in 30 mL CH_2Cl_2 at -78 $^\circ\text{C}$ was added 0.5 mL DMSO (7.0 mmol). After 45 min, a solution of the above alcohol (0.80 g, 1.3 mmol) in 8 mL CH_2Cl_2 at -78 $^\circ\text{C}$ was transferred through cannula to the DMSO solution with 1 x 2 mL CH_2Cl_2 rinse. After stirring 1 h at -78 $^\circ\text{C}$, 0.89 mL Et_3N (6.4 mmol) was added. After 25 min at -78 $^\circ\text{C}$, the reaction solution was warmed to 0 $^\circ\text{C}$ for 20 min, then 2 mL glacial AcOH was added. The reaction was partitioned between CH_2Cl_2 and sat. aq. NaHCO_3 . The layers were separated and the aqueous layer was washed once with 100 mL CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 and concentrated to give a clear syrup. The syrup was purified by flash chromatography on 100 mL neutral silica, eluting with 1:1 CH_2Cl_2 :hexanes then 9:1 hexanes:ethyl acetate, to yield 0.85 g aldehyde (98%). $[\alpha]_{\text{D}}^{28} = +43.0$ (c 0.89, CHCl_3). IR (cm^{-1}): 3087, 3063, 3030, 2920, 2861, 1729. MS (ESI): m/z 669/671 ($[\text{M} + \text{Na}]^+$), 685/687 ($[\text{M} + \text{K}]^+$), 779/781 ($[\text{M} + \text{Cs}]^+$). ^1H NMR (400 MHz, CDCl_3): δ 9.55 (d, $J = 1.5$ Hz, 1H), 7.31-7.15 (m, 20H), 6.73 (br s, 1H), 4.87 (d, $J = 8.7$ Hz, 1H), 4.60 (d, $J = 11.8$ Hz, 1H), 4.51-4.35 (m, 6H), 4.13-3.98 (m, 5H). ^{13}C NMR (CDCl_3): δ 202.9, 138.6,

137.9, 137.7, 137.5, 137.4, 128.7-127.8, 111.2, 84.3, 79.2, 76.5, 73.9, 73.7, 73.1, 70.9, 69.6.

Allylic alcohols **9a** and **9b**.

CrCl₂ (2.4 g, 19 mmol) and NiCl₂ (0.25 g, 1.9 mmol) were weighed out in a glove box and then dissolved in DMF (21 mL, degassed by 3 freeze-pump-thaw cycles). After stirring for 30 min at rt, a solution of **8** (0.47 g, 0.77 mmol) in DMF (19 mL, degassed by three 7-minute freeze-pump-thaw cycles) was cannulated over to the CrCl₂/NiCl₂ solution. The reaction was then heated to 40 °C for 18 h. After cooling to 0 °C, 1 M sodium serinate (pH 8) was added and the reaction was warmed to rt and stirred for 20 min, then partitioned between Et₂O and water. After separation of layers, the aqueous layer was washed with two 200 mL portions of Et₂O. The combined organic layers were washed once with 200 mL water, dried over Na₂SO₄, and concentrated to give a yellow syrup. The syrup was purified by flash chromatography on 100 mL silica, eluting with 7:3 hexanes:ethyl acetate, to yield 0.27 g alcohols (58%, 6:1 **9a**:**9b** ratio). The isomers were separated by HPLC (Rainin Dynamax-100 Å Si R0080120C5, eluting with 19:1 CH₂Cl₂:ethyl acetate at 4 mL/min) for characterization. **Allylic alcohol 9a**. R_t 120 min. [α]_D²⁷ = + 6.1 (*c* 0.97, CHCl₃). IR (cm⁻¹): 3422, 3087, 3063, 3029, 2861. MS (ESI): *m/z* 543 ([M + Li]⁺), 559 ([M + Na]⁺), 575 ([M + K]⁺). ¹H NMR (400 MHz, CDCl₃): δ 7.39-7.29 (m, 20H), 5.76 (d, *J* = 2.5 Hz, 1H), 4.84 (AB, *J* = 11.6 Hz, Δ*v* = 65.8 Hz, 2H), 4.80 (AB, *J* = 12.1 Hz, Δ*v* = 6.3 Hz, 2H), 4.76 (AB, *J* = 11.3 Hz, Δ*v* = 129.9 Hz, 2H),

4.41 (AB, $J = 11.8$ Hz, $\Delta\nu = 12.3$ Hz, 2H), 4.27 (d, $J = 3.4$ Hz, 1H), 4.14 (m, 1H), 3.98 (AB, $J = 12.0$ Hz, $\Delta\nu = 79.1$ Hz, 2H), 3.96 (dd, $J = 6.5, 9.2$ Hz, 1H), 3.64 (dd, $J = 3.5, 9.2$ Hz, 1H), 2.34 (d, $J = 6.1$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 138.84, 138.77, 138.5, 138.2, 138.5, 128.6-127.8, 80.7, 79.7, 74.53, 74.49, 73.5, 73.4, 72.6, 71.3, 70.7. **Allylic alcohol 9b**. R_t 110 min. $[\alpha]_D^{28} = +75.6$ (c 0.85, CHCl_3). IR (cm^{-1}): 3446, 3087, 3062, 3029, 2861. MS (ESI): m/z 537 ($[\text{M} + \text{H}]^+$), 554 ($[\text{M} + \text{NH}_4]^+$), 559 ($[\text{M} + \text{Na}]^+$), 575 ($[\text{M} + \text{K}]^+$). ^1H NMR (400 MHz, CDCl_3): δ 7.36-7.25 (m, 20H), 5.80 (d, $J = 4.0$ Hz, 1H), 4.75 (AB, $J = 12$ Hz, $\Delta\nu = 12.3$ Hz, 2H) 4.70 (AB, $J = 11.6$ Hz, $\Delta\nu = 31$ Hz, 2H), 4.68 (AB, $J = 11.3$ Hz, $\Delta\nu = 119.4$ Hz, 2H), 4.40 (AB, $J = 11.9$ Hz, $\Delta\nu = 22.9$ Hz, 2H), 4.34 (m, 1H), 4.22 (d, $J = 3.4$ Hz, 1H), 4.03 (dd, $J = 4.3, 8.8$ Hz, 1H), 3.98 (AB, $J = 12.5$ Hz, $\Delta\nu = 91.2$ Hz, 2H) 3.91 (dd, $J = 3.4, 8.9$ Hz, 1H), 2.68 (d, $J = 4.3$ Hz, 1H). ^{13}C NMR (CDCl_3): δ 138.83, 138.8, 138.4, 138.3, 137.4, 128.6-127.7, 126.5, 76.8, 74.5, 73.8, 73.64, 73.61, 72.3, 70.7, 65.9.

Benzyl-protected enone 10.

To **9a/9b** (0.058 g, 0.11 mmol) dissolved in CH_2Cl_2 (1.0 mL) were added NaHCO_3 (0.046 g, 0.55 mmol) and Dess-Martin periodinane (0.092 g, 0.22 mmol) at rt. After 40 min. sat. aq. $\text{Na}_2\text{S}_2\text{O}_3$ was added and the reaction stirred vigorously until it turned clear. The reaction was partitioned between CH_2Cl_2 and sat. aq. NaHCO_3 , the layers were separated, and the aqueous layer was washed once with 20 mL CH_2Cl_2 . The combined

organic layers were dried over Na₂SO₄ and concentrated to give a clear glass. Flash chromatography over 25 mL silica, eluting with 8:2 hexanes:ethyl acetate, provided 0.051 g enone **10** (86%). $[\alpha]_D^{29} = + 66.8$ (*c* 1.3, CHCl₃). IR (cm⁻¹): 3030, 2866, 1687. ¹H NMR (500 MHz, CDCl₃): δ 7.41-7.22 (m, 20H), 6.11 (t, *J* = 1.6 Hz, 1H), 4.83 (AB, *J* = 11.5 Hz, $\Delta\nu$ = 108.2 Hz, 2H), 4.78 (AB, *J* = 12 Hz, $\Delta\nu$ = 66.1 Hz, 2H), 4.72 (AB, *J* = 11.3 Hz, $\Delta\nu$ = 132.0 Hz, 2H), 4.45 (AB, *J* = 11.9 Hz, $\Delta\nu$ = 15.3 Hz, 2H), 4.43 (d, *J* = 9 Hz, 1H), 4.32 (d, *J* = 3.1 Hz, 1H), 4.06 (ABX, (*J*_{AB} = 16.0 Hz, *J*_{AX} = 1.43 Hz, $\Delta\nu$ = 12.1 Hz, 2H), 3.91 (dd, *J* = 3.2, 8.8 Hz, 1H). ¹³C NMR (CDCl₃): δ 196.9, 156.1, 138.32, 138.07, 137.9, 137.6, 128.67-127.92, 124.93, 80.26, 79.28, 74.5, 74.2, 73.8, 73.7, 73.1, 69.6.

Acetate-protected enone **11**.

To **10** (0.010 g, 0.019 mmol) dissolved in CH₂Cl₂ (1 mL) at -78 °C was added BCl₃ (0.1 mL, 1.0 M soln in CH₂Cl₂) dropwise. The solution was gradually warmed to 0 °C over 1.5 h, then re-cooled to -78 °C. Pyridine (0.1 mL) and CH₃OH (0.2 mL) were then added and the reaction was warmed to rt. The reaction was concentrated and 1 mL pyridine then 0.5 mL acetic anhydride were added to the crude residue. After 1 h, the reaction was concentrated and co-evaporated once with 2 mL toluene. The crude residue was partitioned between ethyl acetate and water; after separation of layers the aqueous layer was washed with 1 x 20 mL ethyl acetate. The combined organic layers were dried over Na₂SO₄ and concentrated to give a brown oil. Purification of the oil using preparative thin-layer chromatography in 1:1 hexanes:ethyl acetate yielded 0.0049 g **11** (75%, two

steps). $[\alpha]_D^{29} = +119.1$ (c 0.45, CHCl_3). IR (cm^{-1}): 2930, 1753, 1703. ^1H NMR (500 MHz, CDCl_3): δ 6.25 (t, $J = 1.76$ Hz, 1H), 5.85 (d, $J = 3.7$ Hz, 1H), 5.72 (d, $J = 11.4$ Hz, 1H), 5.41 (dd, $J = 3.7, 11.2$ Hz, 1H), 4.72 (ABX, $J_{AB} = 16.7$ Hz, $J_{AX} = 1.7$ Hz, $\Delta\nu = 12.1$ Hz, 2H), 2.17 (s, 3H), 2.15 (s, 3H), 2.12 (s, 3H), 2.02 (s, 3H). ^{13}C NMR (CDCl_3): δ 190.2, 170.25, 170.18, 170.04, 169.8, 150.6, 127.5, 71.5, 68.8, 65.1, 62.7, 20.8, 20.75.

Allylic alcohol **12**.

Enone **11** (0.012 g, 0.034 mmol) was dissolved in CH_3OH (0.5 mL) at rt, then cooled to -7°C (ice/ $\text{NaCl}/\text{H}_2\text{O}$ bath). $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (0.014 g, 0.037 mmol) was added followed by NaBH_4 (0.0026 g, 0.069 mmol). After 1.5 h, water was added at -5°C and the reaction was partitioned between ethyl acetate and water. The layers were separated and the aqueous layer was washed with 1 x 20 mL ethyl acetate. The combined organic layers were dried over Na_2SO_4 and then concentrated to give a clear oil. The oil was purified by preparative thin-layer chromatography, eluting with 9:1 ethyl acetate:hexanes, yielding 0.0095 g **12** (81%). $[\alpha]_D^{29} = +66.8$ (c 0.26, CHCl_3). IR (cm^{-1}): 3475, 2941, 1751.

^1H NMR (CDCl_3): δ 5.94 (ddd, $J = 1.20, 2.55$ Hz, 1H), 5.67 (d, $J = 3.87$ Hz, 1H), 5.21 (dd, $J = 7.3, 11.2$ Hz, 1H), 5.10 (dd, $J = 3.9, 11.1$ Hz, 1H), 4.54 (dd, $J = 1.71, 13.6$ Hz, 1H), 4.48 (dd, $J = 1.25, 13.5$ Hz, 1H), 4.31 (m, 1H), 2.86 (d, $J = 5.2$ Hz, 1H), 2.11 (s, 3H), 2.10 (s, 3H), 2.06 (s, 3H), 1.99 (s, 3H). ^{13}C NMR (CDCl_3): δ 172.08, 170.59, 170.48, 170.06, 131.7, 130.7, 73.7, 71.1, 68.5, 65.7, 63.7, 21.1, 20.9, 20.7.

PMB-protected thioglycoside.

p-Tolyl 1-thio- β -D-galactopyranoside **13** (2.0 g, 7 mmol) was dissolved in 35 mL DMF then cooled to 0 °C. NaH (1.7 g, 60% suspension, 42 mmol) was added slowly over 5 min. After 10 min, PMB-Br (7.8 mL, 56 mmol) was added and the reaction stirred for another 10 min, at which time the temperature was raised to rt for 2 h. The reaction was re-cooled to 0 °C and water was added slowly dropwise, then diluted with ethyl acetate and water. The layers were separated and the organic layer was washed twice with 100 mL water, once with 100 mL sat. aq. NaCl, dried over Na₂SO₄ and concentrated to give a clear syrup. Flash chromatography over 400 mL silica, eluting with 8:2 hexanes:ethyl acetate followed by 7:3 hexanes:ethyl acetate, yielded 5.26 g thiogalactoside (96%). $[\alpha]_D^{29} = + 11.9$ (*c* 2.3, CHCl₃). IR (cm⁻¹): 2998, 2908, 2833. ¹H NMR (400 MHz, CDCl₃): δ 7.43 (d, *J* = 8.2 Hz, 2H), 7.30 (d, *J* = 8.0 Hz, 2H), 7.25 (d, *J* = 9.0 Hz, 2H), 7.20 (d, *J* = 8.3 Hz, 2H), 7.17 (d, *J* = 8.6 Hz, 2H), 6.96 (d, *J* = 7.2 Hz, 2H), 6.85-6.82 (m, 8H), 4.67 (AB, *J* = 11.2 Hz, $\Delta\nu$ = 135 Hz, 2H), 4.66 (AB, *J* = 9.8 Hz, $\Delta\nu$ = 21 Hz, 2H), 4.62 (AB, *J* = 11.4 Hz, $\Delta\nu$ = 7.9 Hz, 2H), 4.54 (AB, *J* = 15 Hz, $\Delta\nu$ = 20 Hz, 2H), 4.50 (d, *J* = 11.1 Hz, 1H), 4.34 (AB, *J* = 11.4 Hz, $\Delta\nu$ = 22 Hz, 2H), 3.88 (d, *J* = 2.6 Hz, 1H), 3.83 (t, *J* = 9.5 Hz, 1H), 3.79 (s, 3H), 3.78 (s, 6H), 3.77 (s, 3H), 3.57-3.48 (m, 4H), 2.26 (s, 3H). ¹³C NMR (CDCl₃): δ 159.48, 159.45, 159.37, 159.24, 137.24, 132.23, 131.17, 130.85, 130.68, 130.62, 130.18, 129.79, 129.74, 129.71, 129.36, 113.99, 113.98, 113.92,

113.70, 88.34, 84.14, 77.49, 77.28, 75.43, 74.11, 73.38, 73.19, 72.54, 68.70, 55.48, 55.45, 21.29.

PMB-protected pyranose 14.

The thioglycoside prepared above (5.3 g, 6.9 mmol) was dissolved in 150 mL acetone at rt. Water (15 mL, 820 mmol) was added and the reaction cooled to 0 °C. *N*-bromosuccinimide (2.9 g, 16.4 mmol) was added which produced a bright orange color. After 15 min, the color had disappeared; after 1h TLC showed completion so the reaction was concentrated. The residue was dissolved in ethyl acetate and washed with 200 mL water. The layers were separated and the aqueous layer was washed once with 100 mL ethyl acetate. The combined organic layers were dried over Na₂SO₄ and concentrated. Flash chromatography over 300 mL silica, eluting with CH₂Cl₂, 6:4 hexanes:ethyl acetate, and 1:1 hexanes:ethyl acetate afforded 3.74 g **14** (82%) as a clear syrup. $[\alpha]_D^{29} = +0.40$ (*c* 1.7, CHCl₃). IR (cm⁻¹): 3420, 2999, 2835. ¹H NMR (400 MHz, CDCl₃): δ 7.32-7.17 (m, 14H), 6.89-6.80 (m, 14H), 5.21 (t, *J* = 2.7 Hz, 1H), 4.85-4.82 (m, 2H), 4.74-4.69 (m, 3H), 4.67-4.59 (m, 4H), 4.53-4.49 (m, 2H), 4.42-4.28 (m, 4H), 4.12 (t, *J* = 6.1 Hz, 1H), 4.05 (d, *J* = 6.6 Hz, 1H), 3.97 (dd, *J* = 3.3, 10.3 Hz, 1H), 3.88-3.85 (m, 2H), 3.80-3.77 (m, 22H), 3.75-3.68 (m, 2H), 3.62-3.36 (m, 7H). ¹³C NMR (CDCl₃): δ 159.42, 159.37, 159.35, 159.29, 159.26, 159.22, 131.01, 130.99, 130.88, 130.80, 130.73, 130.61, 130.09, 130.05, 129.95, 129.78, 129.75, 129.30, 129.26, 113.89, 113.81, 113.72, 113.69, 97.92, 91.99, 82.07, 80.60, 78.62, 76.27, 74.84, 74.37, 74.24, 74.11, 73.66, 73.24, 73.16, 73.13, 72.74, 69.51, 68.94, 68.78, 55.38, 55.36.

PMB-protected diol **15**.

The mixture of pyranoses **14** (3.7 g, 5.6 mmol) was dissolved in 30 mL CH₃OH then cooled to 0 °C. NaBH₄ (0.64 g, 17 mmol) was added and the reaction stirred for 1h, at which time the reaction was warmed to rt and then stirred overnight. The reaction was concentrated and partitioned between ethyl acetate and water. The layers were separated and the organic layer washed once with 200 mL sat. aq. NaCl, then dried over Na₂SO₄ and concentrated to give 3.6 g (95%) diol **15** which was not further purified. $[\alpha]_D^{26} = -4.6$ (*c* 0.67, CHCl₃). IR (cm⁻¹): 3457, 2999, 2854. ¹H NMR (400 MHz, CDCl₃): δ 7.25-7.12 (m, 8H), 6.85-6.79 (m, 8H), 4.61 (AB, *J* = 11 Hz, Δ*v* = 26 Hz, 2H), 4.54 (AB, *J* = 11.5 Hz, Δ*v* = 19.3 Hz, 2H), 4.45 (AB, *J* = 10.9 Hz, Δ*v* = 61.4 Hz, 2H), 4.36 (AB, *J* = 11.7 Hz, Δ*v* = 20 Hz, 2H), 3.99-3.95 (m, 1H), 3.77-3.70 (m, 14 H), 3.74-3.70 (m, 1H), 3.67-3.61 (m, 2H), 3.48 (dd, *J* = 5.8, 9.2 Hz, 1H), 3.42 (dd, *J* = 6.8, 9.2 Hz, 1H), 3.30 (d, *J* = 4.5 Hz, 1H), 2.33 (m, 1H). ¹³C NMR (CDCl₃): δ 159.56, 159.54, 159.43, 130.33, 130.26, 130.18, 130.02, 129.81, 129.70, 114.07, 114.03, 113.96, 80.16, 78.79, 77.44, 77.16, 74.08, 73.54, 73.13, 72.13, 70.74, 70.05, 61.09, 55.46.

TBS-protected alcohol.

Diol **15** (3.6 g, 5.4 mmol) was dissolved in 22 mL DMF then cooled to 0 °C. Imidazole (0.92 g, 13.6 mmol) and TBSCl (0.89 g, 5.97 mmol) were added. After 5 min the reaction was warmed to rt. The reaction was partitioned between ethyl acetate and water. The layers were separated and the aqueous layer was washed once with 100 mL ethyl acetate. The combined organic layers were washed three times with 200 mL water and once with 200 mL sat. aq. NaCl, then dried over Na₂SO₄ and concentrated to give a clear syrup. Flash chromatography over 300 mL silica, eluting with CH₂Cl₂ then 7:3 hexanes:ethyl acetate, yielded 4.1 g alcohol (98%). $[\alpha]_D^{29} = -6.9$ (*c* 0.42, CHCl₃). IR (cm⁻¹): 3489, 2932, 2856. ¹H NMR (400 MHz, CDCl₃): δ 7.24-7.10 (m, 8H), 6.84-6.78 (m, 8H), 4.61 (AB, *J* = 11 Hz, Δ*v* = 23 Hz, 2H), 4.58 (AB, *J* = 11.5 Hz, Δ*v* = 51 Hz, 2H), 4.37 (AB, *J* = 11.4 Hz, Δ*v* = 30 Hz, 2H), 4.33 (AB, *J* = 10.9 Hz, Δ*v* = 41 Hz, 2H), 4.05-4.00 (m, 1H), 3.87-3.61 (m, 17 H), 3.47 (dd, *J* = 6.1, 9.2 Hz, 1H), 3.42 (dd, *J* = 6.6, 9.2 Hz, 1H), 3.17 (d, *J* = 4.9 Hz, 1H), 0.86 (s, 9H), 0.003 (s, 3H), -0.003 (s, 3H). ¹³C NMR (CDCl₃): δ 159.42, 159.38, 159.33, 130.99, 130.73, 130.63, 130.50, 130.10, 129.76, 129.72, 129.64, 113.93, 113.90, 113.87, 79.85, 79.18, 74.51, 73.07, 72.95, 72.67, 71.13, 70.17, 63.38, 55.46, 26.16, 18.48, -5.14.

Ketone 16.

To 110 mL CH₂Cl₂ was added 1.5 mL (22 mmol) anhydrous DMSO. After cooling the reaction solution to -78 °C, 1.4 mL (16 mmol) oxalyl chloride was added slowly dropwise. After 30 min, a solution of the alcohol prepared above (4.2 g, 5.4 mmol) in 22

mL CH₂Cl₂ at -78 °C was transferred to the reaction by cannula with 1 x 5 mL rinse with CH₂Cl₂. After 1 h, 6.0 mL Et₃N (43 mmol) was added to the reaction dropwise and stirred for 40 min at -78 °C. The reaction was then warmed to 0 °C and 4 mL glacial AcOH added. The reaction was partitioned between CH₂Cl₂ and sat. aq. NaHCO₃, the layers were separated, and the aqueous layer washed with 200 mL CH₂Cl₂. The combined organic layers were washed once with 200 mL water, once with 200 mL sat. aq. NaCl, dried over Na₂SO₄ and concentrated to yield a brown syrup. Flash chromatography over 300 mL silica, eluting with 8:2 hexanes:ethyl acetate then 7:3 hexanes:ethyl acetate provided 3.69 g **16** (88%). [α]_D²⁶ = -1.8 (*c* 0.275, CHCl₃). IR (cm⁻¹): 2999, 2856, 1732. ¹H NMR (400 MHz, CDCl₃): δ 7.21-7.12 (m, 8H), 6.84-6.79 (m, 8H), 4.51-4.20 (m, 10H), 4.16 (d, *J* = 4.8 Hz, 1H), 3.96 (t, *J* = 4.7 Hz, 1H), 3.77-3.69 (m, 14H), 3.65-3.62 (m, 1H), 0.87 (s, 9H), 0.009 (s, 3H), 0.003 (s, 3H). ¹³C NMR (CDCl₃): δ 208.1, 159.55, 159.44, 159.37, 159.30, 130.81, 130.30, 129.93, 129.89, 129.84, 129.77, 129.73, 129.56, 113.95, 113.89, 113.80, 113.78, 82.03, 79.79, 74.48, 74.10, 73.06, 72.90, 72.47, 63.07, 55.39, 55.37, 26.12, 18.44, -5.16, -5.18.

TBS-protected alkene.

Ketone **16** (1.4 g, 1.8 mmol) was dissolved in 18 mL THF at rt, then cooled to -78 °C. Methyltriphenylphosphonium bromide (1.1 g, 3.2 mmol) was suspended in THF at rt, then cooled to 0 °C and stirred vigorously. *n*-Butyllithium (0.87 mL of a 2.6 M solution in hexanes, 2.3 mmol) was added dropwise to the suspension, which turned yellow but

The alkene prepared above (1.4 g, 0.25 mmol) was dissolved in 10 mL THF at rt. TBAF (3.7 mL of a 1.0 M solution in THF, 3.7 mmol) was added dropwise. After 1 h the reaction was diluted with ethyl acetate and water. The layers were separated and the aqueous layer was washed once with 100 mL ethyl acetate. The combined organic layers were dried over Na₂SO₄ and concentrated to give a clear syrup. Flash chromatography over 100 mL silica, eluting with 6:4 hexanes:ethyl acetate then 1:1 hexanes:ethyl acetate afforded 1.1 g **17** (92%). $[\alpha]_D^{26} = +28.3$ (c 1.78, CHCl₃). IR (cm⁻¹): 3466, 2998, 2836. ¹H NMR (400 MHz, CDCl₃): δ 7.27-7.14 (m, 8H), 6.86-6.78 (m, 8H), 5.54 (d, $J = 1.4$ Hz, 1H), 5.42 (br s, 1H), 4.53-4.42 (m, 7H), 4.14-4.03 (m, 4H), 3.77 (s, 3H), 3.76 (s, 6H), 3.75 (s, 3H), 3.70-3.50 (m, 4H), 1.86 (dd, $J = 4.8, 7.1$ Hz, 1H). ¹³C NMR (CDCl₃): δ 159.41, 159.35, 144.08, 130.91, 130.55, 130.50, 130.28, 129.77, 129.68, 129.51, 116.07, 113.95, 113.93, 113.82, 80.76, 79.82, 79.26, 74.11, 72.86, 72.69, 70.44, 70.16, 62.27, 55.42.

PMB-protected aldehyde.

To 28 mL CH₂Cl₂ was added 0.5 mL (7.2 mmol) anhydrous DMSO. After cooling the reaction solution to -78 °C, 0.47 mL (5.4 mmol) oxalyl chloride was added slowly dropwise. After 30 min, a solution of alcohol **17** (1.1 g, 1.8 mmol) in 7 mL CH₂Cl₂ at -78 °C was transferred to the reaction by cannula with 1 x 5 mL rinse with CH₂Cl₂. After 1 h, 2.0 mL Et₃N (14 mmol) was added to the reaction dropwise and stirred for 45 min at -78 °C. The reaction was then warmed to 0 °C. After 15 min, the reaction was partitioned between CH₂Cl₂ and sat. aq. NaHCO₃, the layers were separated, and the

aqueous layer washed with 100 mL CH₂Cl₂. The combined organic layers were washed once with 50 mL sat. aq. NaCl, dried over Na₂SO₄ and concentrated to yield a syrup. Flash chromatography over 200 mL silica, eluting with 6:4 hexanes:ethyl acetate provided 1.15 g aldehyde (97%). $[\alpha]_D^{27} = +41.9$ (*c* 1.4, CHCl₃). IR (cm⁻¹): 2999, 2906, 2836, 1728. ¹H NMR (400 MHz, CDCl₃): δ 9.44 (d, *J* = 1.8 Hz, 1H), 7.27-7.23 (m, 2H), 7.21-7.18 (m, 2H), 7.10-7.04 (m, 4H), 6.87-6.75 (m, 8H), 5.54 (q, *J* = 1.5 Hz, 1H), 5.42 (br s, 1H), 4.47 (AB, *J* = 11.6 Hz, $\Delta\nu$ = 13.8 Hz, 2H), 4.46 (AB, *J* = 11.6 Hz, $\Delta\nu$ = 14.2 Hz, 2H), 4.35 (d, *J* = 10.7 Hz, 1H), 4.32 (AB, *J* = 10.7 Hz, $\Delta\nu$ = 23 Hz, 2H), 4.07-3.99 (m, 5H), 3.89 (dd, *J* = 3.2, 8.9 Hz, 1H), 3.77 (s, 3H), 3.76 (s, 3H), 3.75 (s, 3H), 3.74 (s, 3H). ¹³C NMR (CDCl₃): δ 203.21, 159.70, 159.46, 159.38, 159.35, 143.48, 130.51, 130.28, 130.16, 130.03, 129.89, 129.83, 129.52, 129.45, 116.78, 114.02, 113.95, 113.89, 113.76, 84.24, 80.51, 78.69, 74.05, 73.27, 72.69, 70.05, 69.84, 55.42, 55.40, 55.39.

Allylic alcohol 18.

The aldehyde prepared above (0.53 g, 0.80 mmol) was dissolved in 8 mL THF at rt, then cooled to -78 °C. Vinylmagnesium bromide (1.2 mL of a 1.0 M solution in THF, 1.2 mmol) was added slowly dropwise. After 1.5 h, the reaction was warmed to 0 °C for 40 min, then diluted with Et₂O and quenched with water at 0 °C. The reaction was poured into a separatory funnel and solid NaCl was used to break the resulting emulsion. The layers were separated and the aqueous layer was washed once with 100 mL Et₂O. The combined organic layers were washed once with 100 mL water, once with 100 mL sat.

aq. NaCl, then dried over Na₂SO₄ and concentrated to give a clear syrup. Flash chromatography over 200 mL silica, eluting with 7:3 hexanes:ethyl acetate, yielded 0.41 g **18** (75%) of a 4:1 mixture of diastereomers. $[\alpha]_D^{26} = +17.9$ (*c* 0.73, CHCl₃). IR (cm⁻¹): 3491, 2909, 2835. ¹H NMR (400 MHz, CDCl₃): (mixture of isomers) δ 7.27-7.23 (m, 10H), 7.19-7.10 (m, 24H), 6.86-6.77 (m, 32H), 5.89 (ddd, *J* = 5.0, 10.4, 17.3 Hz, 3H), 5.82 (ddd, *J* = 5.7, 10.5, 17.2 Hz, 1H), 5.55 (q, *J* = 1.6 Hz, 3H), 5.52 (q, *J* = 1.6 Hz, 1H), 5.42 (br s, 3H), 5.40 (br s, 1H), 5.35 (dt, *J* = 1.7, 17.2 Hz, 3H), 5.24 (dt, *J* = 1.5, 17.2 Hz, 1H), 5.18 (dt, *J* = 1.7, 10.6 Hz, 3H), 5.12 (dt, (*J* = 1.6, 10.5 Hz, 1H), 4.58-4.35 (m, 32H), 4.20-4.00 (m, 18H), 3.77-3.69 (m, 54H), 3.63 (dd, *J* = 3.0, 5.4 Hz, 3H), 3.60-3.58 (m, 1H), 2.70 (d, *J* = 6.1 Hz, 3H), 2.69 (d, *J* = 5.3 Hz, 1H). ¹³C NMR (CDCl₃): (mixture of isomers) δ 159.43, 159.34, 144.09, 138.61, 138.23, 130.86, 130.68, 130.62, 130.45, 130.40, 130.34, 129.96, 129.79, 129.76, 129.68, 129.49, 129.45, 116.32, 116.12, 113.95, 113.90, 113.88, 113.85, 113.79, 81.83, 80.91, 80.35, 80.21, 79.98, 79.52, 74.83, 74.11, 73.65, 73.07, 72.80, 72.73, 72.62, 72.21, 70.34, 69.92, 55.45.

Allylic alcohols **19**.

The mixture of alcohols **18** (0.33 g, 0.483 mmol) was dissolved in 16 mL toluene at rt. Grubbs' second generation catalyst (0.041 g, 10 mol%) was added and a reflux condenser was added to the flask. The dark maroon reaction was heated to reflux for 2 h, at which time TLC showed no sm. The reaction was cooled to rt and concentrated. Flash chromatography over 150 mL silica, eluting with 65:35 hexanes:ethyl acetate then 1:1

hexanes:ethyl acetate provided 0.24 g of the mixture of alcohols **19** (76%). $[\alpha]_D^{26} = +35.6$ (*c* 1.1, CHCl₃). IR (cm⁻¹): 3488, 2999, 2907, 2836. ¹H NMR (400 MHz, CDCl₃): (mixture of isomers) δ 7.30-7.16 (m, 44H), 6.87-6.80 (m, 41H), 5.76 (d, *J* = 4.3 Hz, 3.81H), 5.71 (d, *J* = 2.7 Hz, 1H), 4.82-4.55 (m, 27H), 4.49-4.43 (m, 6H), 4.38-4.26 (m, 15H), 4.19 (d, *J* = 3.5 Hz, 1H), 4.17 (d, *J* = 3.3 Hz, 4H), 4.08-3.99 (m, 6H), 3.96 (dd, *J* = 4.4, 8.6 Hz, 4H), 3.89-3.57 (m, 77H), 2.68 (d, *J* = 4.6 Hz, 4H), 2.20 (d, *J* = 6.5 Hz, 1H). ¹³C NMR (CDCl₃): δ 159.55, 159.38, 159.34, 137.47, 135.69, 131.05, 131.02, 130.73, 130.52, 130.46, 130.39, 130.01, 129.94, 129.84, 129.79, 129.69, 129.60, 129.52, 128.09, 126.52, 114.09, 114.04, 113.99, 113.94, 113.90, 113.86, 80.20, 79.29, 76.45, 76.34, 74.04, 73.95, 73.35, 73.21, 73.18, 73.07, 72.24, 71.92, 70.44, 65.86, 55.46.

PMB-protected enone **20**.

The mixture of cyclohexenols **19** (0.24 g, 0.36 mmol) was dissolved in 3 mL CH₂Cl₂ at rt. NaHCO₃ (0.15 g, 1.8 mmol) and Dess-Martin periodinane (0.31 g, 0.73 mmol) were added and the reaction was stirred for 1 h. Sat. aq. Na₂S₂O₃ was added and the reaction was stirred until clear, then partitioned between CH₂Cl₂ and sat. aq. NaHCO₃. The layers were separated and the aqueous layer was washed once with 20 mL CH₂Cl₂. The combined organic layers were washed once with 20 mL sat. aq. NaCl, then dried over Na₂SO₄ and concentrated to give a clear syrup. Flash chromatography over 100 mL silica, eluting with 75:25 hexanes:ethyl acetate, 7:3 hexanes:ethyl acetate, and 6:4 hexanes:ethyl acetate yielded 0.2 g enone **20** (80%). $[\alpha]_D^{29} = +65.0$ (*c* 1.2, CHCl₃). IR

(cm⁻¹): 2999, 2934, 2836, 1685. ¹H NMR (400 MHz, CDCl₃): δ 7.32 (d, *J* = 8.5 Hz, 2H), 7.23 (d, *J* = 8.5 Hz, 2H), 7.18 (d, *J* = 8.5 Hz, 2H), 7.14 (d, *J* = 8.5 Hz, 2H), 6.87-6.81 (m, 8H), 6.06 (br s, 1H), 4.74 (AB, *J* = 10.9 Hz, Δ*ν* = 103 Hz, 2H), 4.68 (AB, *J* = 10.9 Hz, Δ*ν* = 65 Hz, 2H), 4.62 (AB, *J* = 10.7 Hz, Δ*ν* = 130 Hz, 2H), 4.37 (AB, *J* = 11.6 Hz, Δ*ν* = 20 Hz, 2H), 4.36 (d, *J* = 8.6 Hz, 1H), 4.25 (d, *J* = 2.8 Hz, 1H), 4.00 (AB, *J* = 16.1 Hz, Δ*ν* = 8.9 Hz, 2H), 3.84 (dd, *J* = 3.3, 8.8 Hz, 1H), 3.79 (s, 12 H). ¹³C NMR (CDCl₃): δ 197.20, 159.63, 159.58, 159.45, 156.42, 130.51, 130.28, 130.21, 130.13, 130.05, 129.71, 129.65, 129.61, 124.87, 114.06, 113.99, 113.91, 79.91, 78.90, 74.12, 73.89, 73.39, 73.30, 72.75, 69.32, 55.49.

Allylic alcohol **12**.

Enone **11** (0.012 g, 0.034 mmol) was dissolved in CH₃OH (0.5 mL) at rt, then cooled to -7°C (ice/NaCl/H₂O bath). CeCl₃·7H₂O (0.014 g, 0.037 mmol) was added followed by NaBH₄ (0.0026 g, 0.069 mmol). After 1.5 h, water was added at -5°C and the reaction was partitioned between ethyl acetate and water. The layers were separated and the aqueous layer was washed once with 20 mL ethyl acetate. The combined organic layers were dried over Na₂SO₄ and then concentrated to give a clear oil. The oil was purified by preparative thin-layer chromatography, eluting with 9:1 ethyl acetate:hexanes, yielding 0.0095 g **12** (81%). [α]_D²⁹ = + 66.8 (*c* 0.26, CHCl₃). IR (cm⁻¹): 3475, 2941, 1751. ¹H NMR (500 MHz, CDCl₃): δ 5.94 (ddd, *J* = 1.20, 2.55 Hz, 1H), 5.67 (d, *J* = 3.87 Hz, 1H), 5.21 (dd, *J* = 7.3, 11.2 Hz, 1H), 5.10 (dd, *J* = 3.9, 11.1 Hz, 1H), 4.54 (dd, *J* = 1.71, 13.6 Hz, 1H), 4.48 (dd, *J* = 1.25, 13.5 Hz, 1H), 4.31 (m, 1H), 2.86 (d, *J* = 5.2 Hz, 1H),

2.11 (s, 3H), 2.10 (s, 3H), 2.06 (s, 3H), 1.99 (s, 3H). ^{13}C NMR (CDCl_3): δ 172.08, 170.59, 170.48, 170.06, 131.7, 130.7, 73.7, 71.1, 68.5, 65.7, 63.7, 21.1, 20.9, 20.7.

Unsaturated thioether **21**.

Alcohol **12** (0.021 g, 0.061 mmol) was dissolved in 1 mL CH_2Cl_2 at rt. Pyridine (0.099 mL, 1.2 mmol) and methanesulfonyl chloride (0.094 mL, 1.2 mmol) were added. After 1 h water was added and the reaction was partitioned between CH_2Cl_2 and water. The layers were separated and the aqueous layer was washed once with 20 mL CH_2Cl_2 . The combined organic layers were washed once with 15 mL sat. aq. NaCl, dried over Na_2SO_4 and concentrated to give the crude mesylate, which was dissolved in 0.3 mL DMF.

N-Boc-cysteine methyl ester (0.1 g, 0.43 mmol) was dissolved in 0.3 mL DMF at rt. Cs_2CO_3 (0.140 g, 0.43 mmol) and water (0.1 mL) were added and the solution was stirred vigorously for 20 min. The crude mesylate was then added dropwise by cannula to the amino acid solution, rinsing with 0.2 mL DMF. After 2.5 h, the reaction was partitioned between ethyl acetate and water, washing the aqueous layer once with 20 mL ethyl acetate. The organic layers were dried over Na_2SO_4 and concentrated. Flash chromatography over 25 mL silica, eluting with 8:2 hexanes:ethyl acetate, 6:4 hexanes:ethylacetate, and 1:1 hexanes:ethyl acetate yielded 0.026 g **21** (76%). $[\alpha]_{\text{D}}^{29} = +76.0$ (*c* 0.18, CHCl_3). IR (cm^{-1}): 3360, 2924, 2846, 1748, 1709. ^1H NMR (500 MHz, CDCl_3): δ 5.97 (d, *J* = 5.0 Hz, 1H), 5.68 (s, 1H), 5.45 (m, 2H), 5.30 (d, *J* = 7.2 Hz, 1H),

4.52 (m, 1H), 4.46 (AB, $J = 13.3$ Hz, $\Delta\nu = 49.7$ Hz, 2H), 3.97 (m, 1H), 3.75 (s, 3H), 3.11 (dd, $J = 4.3, 13.6$ Hz, 1H), 2.90 (dd, $J = 6.3, 13.7$ Hz, 1H), 2.11 (s, 3H), 2.06 (s, 3H), 2.05 (s, 3H), 1.99 (s, 3H), 1.43 (s, 9H). ^{13}C NMR (CDCl_3): δ 171.29, 170.65, 170.41, 170.36, 169.98, 155.35, 131.06, 128.96, 80.55, 68.51, 67.44, 66.18, 63.87, 53.29, 52.94, 44.3, 35.50, 28.45, 20.98, 20.86, 20.82.

Saturated α -galactosylserine isostere **2**.

To **21** was added 0.3 mL benzene, 0.1 mL absolute ethanol, and 0.009 g Wilkinson's catalyst. The reaction was then pressurized to 750 psi H_2 for 14 h. ^1H NMR showed about 50% completion, so 0.008 g Wilkinson's catalyst, 0.3 mL benzene, and 0.1 mL absolute ethanol were added to the crude residue and it was re-pressurized to 750 psi H_2 for another 20 h. At this time the reaction had reached completion, so it was concentrated and purified twice by preparative thin-layer chromatography, eluting with 1:1 hexanes:ethyl acetate, to yield 0.0013 g **2** (18%). $[\alpha]_{\text{D}}^{29} = +18.4$ (c 0.18, CHCl_3). IR (cm^{-1}): 3360, 2923, 1745, 1700. ^1H NMR (500 MHz, CDCl_3): δ 5.47 (br s, 1H), 5.32 (d, $J = 6.4$ Hz, 1H), 5.24 (m, 2H), 4.50 (m, 1H), 3.95 (dd, $J = 10.1$ Hz, 1H), 3.84 (dd, $J = 6.2, 11.1$ Hz, 1H), 3.74 (s, 3H), 3.53 (m, 1H), 3.04 (d, $J = 12.7$ Hz, 1H), 2.95 (dd, $J = 5.0, 13.0$ Hz, 1H), 2.46 (m, 1H), 2.1 (s, 3H), 2.06 (s, 3H), 2.01 (s, 3H), 1.95 (s, 3H), 1.95 (ddd, $J = 3.6, 13.3, 14.1$ Hz, 1H), 1.77 (dt, $J = 3.6, 14.1$ Hz, 1H), 1.42 (s, 9H). ^{13}C NMR (CDCl_3): δ 171.21, 171.03, 170.44, 170.27, 170.21, 80.38, 71.21, 70.05, 68.56, 63.07,

53.28, 52.87, 45.26, 35.14, 33.89, 28.45, 28.07, 21.05, 20.97, 20.96, 20.89. (missing carbamate carbonyl)

Scheme S1. Synthesis of *N*-Boc-Gly-Cys(SH)-Phe-OEt.

***N*-(9-fluorenylmethoxycarbonyl)-*S*-trityl-L-cysteine-L-phenylalanine ethyl ester.**

N-(9-Fluorenylmethoxycarbonyl)-*S*-trityl-L-cysteine (0.2 g, 0.34 mmol), L-phenylalanine ethyl ester hydrochloride (0.12 g, 0.51 mmol), PyBOP (0.265 g, 0.51 mmol) and HOBT (0.078 g, 0.51 mmol) were dissolved in 2.5 mL DMF at rt. CH₂Cl₂ (2.5 mL) and 2,4,6-collidine (0.16 mL, 1.2 mmol) were added and the reaction was stirred for 5 h. The reaction was concentrated then partitioned between ethyl acetate and water. The layers were separated and the aqueous layer washed once with 50 mL ethyl acetate. The combined organic layers were washed with 50 mL sat. aq. NaCl then dried over Na₂SO₄ and concentrated to give a clear syrup. Flash chromatography over 100 mL silica, eluting with 8:2 hexanes:ethyl acetate then 75:25 hexanes:ethyl acetate provided 0.21 g dipeptide (80%). $[\alpha]_D^{26} = + 3.4$ (*c* 0.73, CHCl₃). IR (cm⁻¹): 3302, 3060, 1736, 1664. ¹H NMR (400 MHz, CDCl₃): δ 7.74 (d, *J* = 7.1 Hz, 1H), 7.72 (d, *J* = 7.6 Hz, 1H), 7.55 (d, *J* = 7.3

Hz, 2H), 7.40-7.35 (m, 8H), 7.28-7.23 (m, 7H), 7.19-7.14 (m, 6H), 7.03-7.02 (m, 2H), 6.30 (d, $J = 7.0$ Hz, 1H), 4.88 (m, 1H), 4.72 (q, $J = 5.7$ Hz, 1H), 4.39-4.36 (m, 1H), 4.30-4.27 (m, 1H), 4.17 (t, $J = 5.8$ Hz, 1H), 4.10 (q, $J = 6.4$ Hz, 2H), 3.67 (m, 1H), 3.07 (dd, $J = 6.0, 14.4$ Hz, 1H), 3.02 (dd, $J = 5.3, 13.6$ Hz, 1H), 2.65 (dd, $J = 7.9, 13.6$ Hz, 1H), 2.58 (dd, $J = 5.3, 13.2$ Hz, 1H), 1.2 (t, $J = 6.7$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 170.97, 169.82, 144.48, 143.86, 141.45, 135.80, 129.75, 129.54, 128.63, 128.28, 127.94, 127.28, 127.23, 127.11, 125.29, 125.24, 120.18, 67.54, 67.23, 61.70, 54.06, 53.44, 47.22, 37.95, 33.84, 14.25.

S-trityl-L-cysteine-L-phenylalanine ethyl ester.

The dipeptide made above (0.19 g, 0.25 mmol) was dissolved in 2 mL DMF at rt. Piperidine (0.4 mL) was added and the reaction stirred for 30 min. The reaction was diluted with ethyl acetate and water. The layers were separated and the aqueous layer was washed once with 50 mL ethyl acetate. The combined organic layers were washed once with 100 mL water, once with 100 mL sat. aq. NaCl, and dried over Na_2SO_4 and concentrated. Flash chromatography over 100 mL silica, eluting with 1:1 hexanes:ethyl acetate then 6:4 ethyl acetate:hexanes, provided 0.12 g (93%) as a clear syrup. $[\alpha]_{\text{D}}^{28} = +41.4$ (c 1.4, CHCl_3). IR (cm^{-1}): 3368, 3059, 2927, 1739, 1675. ^1H NMR (400 MHz, CDCl_3): δ 7.49 (d, $J = 8.6$ Hz, 1H), 7.42 (d, $J = 7.6$ Hz, 6H), 7.29-7.16 (m, 12H), 7.05-7.04 (m, 2H), 4.74 (q, $J = 6.3$ Hz, 1H), 4.11 (q, $J = 6.8$ Hz, 1H), 3.07 (dd, $J = 5.7, 13.6$ Hz, 1H), 3.00 (dd, $J = 6.5, 13.9$ Hz, 1H), 2.94 (m, 1H), 2.64 (dd, $J = 3.4, 12.7$ Hz, 1H),

2.44 (dd, $J = 8.9, 12.6$ Hz, 1H), 1.19 (m, 2H), 1.19 (t, $J = 7.1$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 172.76, 171.49, 144.73, 136.08, 129.75, 129.50, 128.54, 128.13, 127.10, 126.96, 67.12, 61.52, 53.99, 52.88, 38.18, 37.38, 14.26.

***N*-(*tert*-Butoxycarbonyl)-glycine-*S*-trityl-*L*-cysteine-*L*-phenylalanine ethyl ester.**

The amino dipeptide prepared above (0.37 g, 0.69 mmol), *N*-Boc-glycine (0.18 g, 1.0 mmol), PyBOP (0.53 g, 1.0 mmol), and HOBt (0.16 g, 1.0 mmol) were dissolved in 3 mL CH_2Cl_2 at rt. DMF (3 mL) and 2,4,6-collidine (0.29 mL, 2.2 mmol) were added and the reaction stirred for 7 h. The reaction was concentrated then partitioned between ethyl acetate and water. The layers were separated and the aqueous layer was washed once with 100 mL ethyl acetate. The combined organic layers were washed once with 100 mL water, once with 100 mL sat. aq. NaCl, then dried over Na_2SO_4 and concentrated to give a white solid. Flash chromatography over 100 mL silica, eluting with 1:1 hexanes:ethyl acetate, provided 0.44 g tripeptide (91%). $[\alpha]_{\text{D}}^{26} = +16.9$ (c 1.6, CHCl_3). IR (cm^{-1}): 3282, 3060, 2978, 1747, 1703, 1653. mp: 188-190 °C. ^1H NMR (400 MHz, CDCl_3): δ 7.39-7.37 (m, 6H), 7.27-7.17 (m, 12H), 7.07-7.05 (m, 2H), 6.44 (d, $J = 7.6$ Hz, 1H), 6.13 (d, $J = 7.0$ Hz, 1H), 4.97 (br s, 1H), 4.71 (q, $J = 6.6$ Hz, 1H), 4.10 (q, $J = 7.2$ Hz, 2H), 3.91 (m, 1H), 3.61 (d, $J = 5.7$ Hz, 2H), 3.08 (dd, $J = 5.8, 13.8$ Hz, 1H), 3.00 (dd, $J = 6.2, 13.5$ Hz, 1H), 2.75 (dd, $J = 7.3, 13.1$ Hz, 1H), 2.48 (dd, $J = 5.2, 13.2$ Hz, 1H), 1.40 (s, 9H), 1.2 (t, $J = 6.6$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 170.99, 169.52, 169.34, 155.97, 144.46, 135.99, 129.69, 129.51, 128.59, 128.22, 127.14, 127.04, 80.37, 67.35, 61.61, 53.48, 52.11, 44.19, 37.85, 33.27, 28.44, 14.22.

***N*-tert-Butoxycarbonyl-glycine-L-cysteine-L-phenylalanine ethyl ester.**

The tripeptide above (0.44 g, 0.63 mmol) was dissolved in 63 mL CH₂Cl₂ at rt. Triisopropylsilane (3.1 mL, 5 vol. %) and trifluoroacetic acid (1.2 mL, 2 vol. %) were added to the reaction, turning it a pale yellow color. After 25 min the color had disappeared and TLC showed no sm. The reaction was partitioned between CH₂Cl₂ and sat. aq. NaHCO₃. The layers were separated and the organic layer was washed once with 50 mL sat. aq. NaHCO₃ and once with 100 mL sat. aq. NaCl, then dried over Na₂SO₄ and concentrated. Flash chromatography over 100 mL silica, eluting with 1:1 hexanes:ethyl acetate then 7:3 ethyl acetate:hexanes, yielded 0.25 g free thiol (89%). $[\alpha]_D^{28} = +1.6$ (*c* 1.2, CHCl₃). IR (cm⁻¹): 3299, 2979, 2575, 1737, 1651. mp: 39-40 °C. ¹H NMR (400 MHz, CDCl₃): δ 7.28-7.19 (m, 3H), 7.13-7.11 (m, 2H), 7.00 (d, *J* = 8.2 Hz, 1H), 6.89 (d, *J* = 7.3 Hz, 1H), 5.18 (br s, 1H), 4.79 (ddd, *J* = 5.9, 7.5, 13.4 Hz, 1H), 4.61 (ddd, *J* = 4.5, 6.4, 8.1 Hz, 1H), 4.15 (q, *J* = 7.1 Hz, 2H), 3.77 (dd, *J* = 5.5, 16.8 Hz, 1H), 3.72 (dd, *J* = 5.9, 17.4 Hz, 1H), 3.16 (dd, *J* = 5.6, 13.9 Hz, 1H), 3.03 (dd, *J* = 7.2, 14.0 Hz, 1H), 3.02-2.97 (m, 1H), 2.64 (ddd, *J* = 6.4, 10.5, 14.0 Hz, 1H), 1.53 (t, *J* = 8.4 Hz, 1H), 1.44 (s, 9H), 1.22 (t, *J* = 7.2 Hz, 3H). ¹³C NMR (CDCl₃): δ 171.29, 169.78, 169.37, 156.31, 136.06, 129.46, 128.78, 127.33, 80.72, 61.86, 54.03, 53.66, 44.59, 37.86, 28.51, 26.80, 14.28.

Unsaturated thioether **22**.

Alcohol **12** (0.010 g, 0.029 mmol) was dissolved in 0.5 mL CH₂Cl₂ at rt. Pyridine (0.047 mL, 0.58 mmol) and methanesulfonyl chloride (0.045 mL, 0.58 mmol) were added. After 1 h water was added and the reaction was partitioned between CH₂Cl₂ and water. The layers were separated and the aqueous layer was washed once with 20 mL CH₂Cl₂. The organic layers were dried over Na₂SO₄ and concentrated to give the crude mesylate, which was then dissolved in 0.5 mL DMF.

N-Boc-Gly-Cys(SH)-Phe-OEt (0.066 g, 0.14 mmol) was dissolved in 0.5 mL DMF at rt. Cs₂CO₃ (0.047 g, 0.14 mmol) and water (0.15 mL) were added and the solution was stirred vigorously for 15 min. The crude mesylate was then added dropwise by cannula to the tripeptide solution, rinsing with 0.3 mL DMF. After 6 h, the reaction was partitioned between ethyl acetate and water, washing the aqueous layer once with 20 mL ethyl acetate. The combined organic layers were washed once with water, dried over Na₂SO₄ and concentrated. Flash chromatography over 75 mL silica, eluting with 1:1 hexanes:ethyl acetate, 6:4 ethyl acetate:hexanes, and 7:3 ethyl acetate:hexanes yielded 0.019 g **22** (84%). $[\alpha]_D^{29} = +63.2$ (*c* 0.67, CHCl₃). IR (cm⁻¹): 3304, 2978, 1748, 1653. ¹H NMR (500 MHz, CDCl₃): δ 7.28-7.11 (m, 5H), 6.91 (d, *J* = 7.9 Hz, 1H), 6.79 (d, *J* = 7.7 Hz, 1H), 5.96 (d, *J* = 4.8 Hz, 1H), 5.69 (br s, 1H), 5.41 (m, 2H), 5.12 (br s, 1H), 4.75 (dd, *J* = 7.1 Hz, 1H), 4.50 (m, 1H), 4.47 (AB, *J* = 13.5 Hz, $\Delta\nu$ = 27 Hz, 2H), 4.14 (q, *J* =

7.2 Hz, 2H), 4.02 (m, 1H), 3.77 (d, $J = 7.2$ Hz, 2H), 3.15 (dd, $J = 6.1, 14.4$ Hz, 1H), 3.05 (dd, $J = 7.3, 13.9$ Hz, 1H), 2.86 (m, 2H), 2.06 (s, 3H), 2.04 (s, 3H), 2.03 (s, 3H), 2.01 (s, 3H), 1.43 (s, 9H), 1.21 (t, $J = 7.0$ Hz, 3H). ^{13}C NMR (CDCl_3): δ 171.18, 170.84, 170.60, 170.43, 170.12, 169.69, 169.61, 156.17, 135.99, 131.05, 129.55, 129.45, 128.78, 127.55, 80.68, 67.73, 66.17, 64.84, 63.97, 61.91, 53.94, 52.51, 44.80, 44.54, 37.91, 34.87, 28.52, 21.12, 21.05, 20.97, 20.91, 14.33.

Glycopeptide isostere **23**.

To **22** was added 0.6 mL benzene, 0.2 mL absolute ethanol, and 0.012 g Wilkinson's catalyst. The reaction was then pressurized to 750 psi H_2 for 16 h, then concentrated. ^1H NMR showed trace reduction, so 0.005 g Wilkinson's catalyst, 0.6 mL benzene, and 0.2 mL absolute ethanol were added to the crude residue and it was re-pressurized to 500 psi H_2 for 24 h. After concentration of the reaction, flash chromatography over 25 mL silica, eluting with 1:1 hexanes:ethyl acetate, 6:4 ethyl acetate:hexanes, 7:3 ethyl acetate:hexanes, and 8:2 ethyl acetate:hexanes yielded a product still contaminated with catalyst. Two rounds of preparative thin-layer chromatography in 9:1 ethyl acetate:hexanes yielded 0.0029 g clean product (28%). $[\alpha]_{\text{D}}^{28} = +7.2$ (c 0.15, CHCl_3). IR (cm^{-1}): 3308, 2923, 1745, 1657. ^1H NMR (400 MHz, CDCl_3): δ 7.26-7.23 (m, 3H), 7.11 (m, 2H), 6.92 (d, $J = 7.2$ Hz, 1H), 6.81 (d, $J = 7.8$ Hz, 1H), 5.46 (br s, 1H), 5.31 (dd, $J = 3.9, 10.2$ Hz, 1H), 5.21 (dd, $J = 2.8, 10.7$ Hz, 1H), 5.09 (br s, 1H), 4.76 (q, $J = 7.0$ Hz, 1H), 4.43 (q, $J = 7.2$ Hz, 1H), 4.16 (q, $J = 7.2$ Hz, 1H), 3.96-3.83 (m, 2H), 3.76 (d, J

= 5.8 Hz, 2H), 3.59 (q, J = 3.1 Hz, 1H), 3.17 (dd, J = 5.1, 14.1 Hz, 1H), 3.05 (dd, J = 6.8, 13.8 Hz, 1H), 2.87 (dd, J = 5.9, 13.8 Hz, 1H), 2.81 (dd, J = 7.6, 14.0 Hz, 1H), 2.45-2.34 (m, 1H), 2.07 (s, 3H), 2.00 (s, 3H), 1.97 (s, 3H), 1.93 (ddd, J = 6.6, 9.9, 13.5 Hz, 1H), 1.94 (s, 3H), 1.78 (dt, J = 3.4, 13.5 Hz, 1H), 1.44 (s, 9H), 1.22 (t, J = 6.8 Hz, 3H).
 ^{13}C NMR (CDCl_3): δ 171.23, 171.21, 170.49, 170.37, 170.32, 169.73, 169.56, 156.31, 136.05, 129.50, 128.86, 127.34, 80.76, 71.03, 70.45, 68.59, 62.98, 61.86, 53.90, 52.80, 45.91, 44.45, 37.78, 34.73, 34.42, 28.51, 28.47, 21.07, 21.02, 20.99, 20.98, 14.35.

Relax. delay 0.500 sec
Pulse 33.0 degrees
Acq. time 3.000 sec
W dth 6505.1 Hz
16 repetitions

Relax. delay 0.500 sec
 Pulse 33.0 degrees
 Acq. time 3.000 sec
 Width 6000.6 Hz
 16 repetitions

OBSERVE H1, 400.1547354
 DATA PROCESSING
 Line broadening 0.2 Hz
 FT size 131072
 Total time 1 minute

Solvent: CDCl₃
 Ambient temperature

11

Relax. delay 1.000 sec
Pulse 27.8 degrees
Acq. time 3.000 sec
Width 7200.1 Hz
32 repetitions

Solvent: CDCl₃
Ambient temperature

STANDARD 1H OBSERVE

Pulse Sequence: gHMBC

Solvent: CDCl₃

Ambient temperature

INNOVA-500 "phantom

Relax. delay 1.000 sec

Acq. time 0.146 sec

Wdth 3502.6 Hz

2D Wdth 24147.3 Hz

8 repetitions

400 increments

OBSERVE H1, 400.1547262 MHz

DATA PROCESSING

Sine bell 0.073 sec

F1 DATA PROCESSING

Sine bell 0.027 sec

FT size 1024 x 8192

Total time 1 hr, 6 min, 35 sec

STANDARD 1H OBSERVE

Pulse Sequence: gHSQC

Solvent: CDCl₃

Ambient temperature

1 NOVA-500 "phantom

Relax. delay 1.000 sec

Acq. time 0.146 sec

Width 3502.6 Hz

2D Width 17105.0 Hz

4 repetitions

2 x 256 increments

OBSERVE H1, 400.1547362 MHz

DECOUPLE C13, 100.6265087 MHz

Power 40 dB

on during acquisition

off during delay

W0_3mmID_5784 modulated

DATA PROCESSING

Sq. sine bell 0.146 sec

Shifted by -0.146 sec

F1 DATA PROCESSING

Sq. sine bell 0.059 sec

Shifted by -0.059 sec

FT size 1024 x 4096

Total time 42 min, 21 sec

CH₂ Carbons from gHSQC projection

CH₃, CH Carbons from gHSQC projection

Relax. delay 0.500 sec
Pulse 33.4 degrees
Acq. time 3.000 sec
Width 6000.6 Hz
16 repetitions

OBSERVE H1, 400.1547356
DATA PROCESSING
Line broadening 0.1 Hz
FT size 131072
Total time 1 minute

Solvent: CDCl₃
Ambient temperature

Solvent: CDCl₃
Ambient temperature

DATA PROCESSING
Line broadening 0.1 Hz
FT size 65536
Total time 4 minutes

7 6 OBSERVE H1, 500.3674489

Relax. delay 1.000 sec
Pulse 26.5 degrees
Acq. time 3.000 sec
Width 7200.1 Hz
64 repetitions

UCB Varian Inova-500
Standard Proton Parameters
NALORAC 4N Probe

Pulse Sequence: gHSQC

Solvent: CDCl₃
Ambient temperature

INOVA-500 "phantom

Relax. delay 1.000 sec

Acq. time 0.248 sec

W dth 4130.7 Hz

2D W dth 21390.4 Hz

3 repetitions

2 x 256 increments

OBSERVE H1, 500.3674484 MHz

DECOUPLE C13, 125.8268952 MHz

Power 42 dB

on during acquisition

off during delay

GARP-1 modulated

DATA PROCESSING

Gauss apodization 0.115 sec

F1 DATA PROCESSING

Gauss apodization 0.022 sec

FT size 2048 x 4096

Total time 34 min, 33 sec

LJW X-075C, gHMBC (J nxh=9), 400 NMR, 3mm D Probe

Pulse Sequence: gHMBC

Solvent: CDCl₃

Ambient temperature

INOVA-500 "phantom

Relax. delay 1.000 sec

Acq. time 0.151 sec

Wdth 3397.7 Hz

2D Wdth 24147.3 Hz

12 repetitions

400 increments

OBSERVE H1, 400.1547351 MHz

DATA PROCESSING

Sine bell 0.075 sec

F1 DATA PROCESSING

Sine bell 0.025 sec

FT size 1024 x 4096

Total time 1 hr, 39 min, 20 sec

Pulse Sequence: gHSQC

Solvent: CDCl₃

Ambient temperature

INOVA-500 "phantom

CH₂ Carbons from gHSQC projection

Relax. delay 1.000 sec

Acq. time 0.151 sec

Width 3397.7 Hz

2D Width 17105.0 Hz

8 repetitions

2 x 180 increments

OBSERVE HI, 400.1547251 MHz

DECOUPLE C13, 100.6265087 MHz

Power 40 dB

on during acquisition

off during delay

W40_3mm_ID_5784 modulated

DATA PROCESSING

Gauss apodization 0.070 sec

F1 DATA PROCESSING

Gauss apodization 0.019 sec

FT size 1024 x 4096

Total time 59 min, 25 sec

CH₃, CH Carbons from gHSQC projection