Contents of the Supporting Information

General information	S2
^1H NMR and ^{13}C NMR spectra of 3a-3i	S4

SUPPORTING INFORMATION

Synthesis of *N*-Diphenylphosphinoylimines Using the Kresze Reaction

Caroline Lauzon, Jean-Nicolas Desrosiers, and André B. Charette*

*Département de chimie, Université de Montréal, P.O. Box 6128, Station Downtown,
Montréal, Québec, Canada H3C 3J7. E-Mail: andre.charette@umontreal.ca*

General information. All the reactions were run under argon atmosphere with rigid exclusion of moisture from reagents and glassware using standard techniques for manipulating air-sensitive compounds.¹ All glassware was stored in the oven and/or flame-dried under an argon atmosphere prior to use. Dichloromethane, tetrahydrofuran and toluene were obtained by filtration through drying columns. Analytical thin-layer chromatography (TLC) was performed on precoated, glass-backed silica gel. Visualization of the developed chromatogram was performed by UV absorbance and aqueous potassium permanganate. Flash column chromatography was performed using 230-400 mesh silica of the indicated system according to standard technique.

Melting points were obtained on a melting point apparatus and are uncorrected. NMR spectra (¹H, ¹³C, ³¹P) were recorded on AV 300, AMX 300, AV 400 or ARX 400 spectrometers. Chemical shifts for ¹H NMR spectra are recorded in ppm with the solvent resonance as the internal standard (CDCl₃, δ 7.27 ppm or C₆D₆, δ 7.15 ppm). Data are reported as follows: chemical shift, multiplicity (s, singlet; d, doublet; m, multiplet), coupling constant in Hz, and integration. Chemical shifts for ¹³C NMR spectra are recorded in ppm from tetramethylsilane by using the central peak of deuteriochloroform (77.00 ppm) or deuterated benzene (128.62 ppm) as the internal standard. All spectra were obtained with complete proton decoupling. Infrared spectra were taken on a FTIR and are reported in reciprocal centimeters (cm⁻¹).

Reagents. Thionyl chloride was distilled over a small amount of quinoline. 2-furylaldehyde was distilled prior to use. Unless otherwise stated, commercial reagents were used without purification. *P,P*-diphenylphosphinic amide was prepared according to the procedure developed by Stec.²

-
- (1) Shriver, D. F.; Drezdson, M. A. *The manipulation of air-sensitive compounds*; 2nd Edition ed.; Wiley: New York, 1986.
- (2) Krzyzanowska, B.; Stec, W. J. *Synthesis* **1978**, 521-524.

***N*-diphenylphosphinoylimines **3a** to **3h**.** The *N*-diphenylphosphinoylimines **3a**,^{2,3} **3b**,^{3,4} **3c**,⁵ **3d**,⁶ **3e**,⁷ **3f**,^{3,4} **3g**³ and **3h**⁸ were prepared by the general procedure and were identical in all respects to authentic materials. (¹H NMR, ¹³C NMR and ³¹P NMR).

***N*-(mesitylmethylene)-*P,P*-diphenylphosphinic amide (**3i**).** The general procedure was followed. White solid, Mp 156.0–157.0 °C; R_f 0.43 (100% EtOAc); ¹H NMR (400 MHz, CDCl₃) δ 9.77 (d, *J* = 34.9 Hz, 1H), 7.94–8.00 (m, 4H), 7.43–7.54 (m, 6H), 6.95 (s, 2H), 2.66 (s, 6H), 2.34 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 173.7 (d, *J*_{C–P} 8.1 Hz), 143.4, 142.1, 133.9 (d, *J*_{C–P} 127.0 Hz), 132.0 (d, *J*_{C–P} 2.8 Hz), 131.9 (d, *J*_{C–P} 9.2 Hz), 131.0, 128.9 (d, *J*_{C–P} 12.4 Hz), 22.3, 21.9; ³¹P NMR (121 MHz, CDCl₃) δ 25.3; IR (Neat) 3071, 2961, 2916, 1659, 1596, 1442, 1434, 1200, 1122, 1105, 827, 727, 694 cm^{–1}; Elemental Analysis calcd for C₂₂H₂₂NOP: C, 76.06; H, 6.38; N, 4.03; O, 4.61; P, 8.92 found: C, 75.93; H, 6.21; N, 4.01.

-
- (3) Boyd, D. R.; Malone, J. F.; McGuckin, R. M.; Jennings, W. B.; Rutherford, M.; Saket, B. M. *J. Chem. Soc., Perkin. Trans. 2* **1988**, 1145-1150.
 - (4) Jennings, W. B.; Lovely, C. J. *Tetrahedron Lett.* **1991**, 47, 5561-5568.
 - (5) Shi, M.; Wang, C.-J. *Adv. Synth. Catal.* **2003**, 345, 971-973.
 - (6) Wipf, P.; Kendall, C.; Stephenson C. R. J. *J. Am. Chem. Soc.* **2003**, 125, 761-768.
 - (7) Pinho, P.; Andersson, P. G. *Tetrahedron* **2001**, 57, 1615-1618.
 - (8) Suzuki, T.; Shibata, T.; Soai, K. *J. Chem. Soc., Perkin. Trans. 1* **1997**, 2757-2760.

^{13}C NMR (C_6D_6 , 100 MHz)

^{13}C NMR (C_6D_6 , 100 MHz)

^{13}C NMR (CDCl_3 , 100 MHz)

^1H NMR (CDCl_3 , 400 MHz)

^{13}C NMR (CDCl_3 , 100 MHz)

¹H NMR (CDCl₃, 400 MHz)

^1H NMR (CDCl_3 , 400 MHz)