Synthesis of *erythro*- α -amino β -hydroxy carboxylic acid esters by diastereoselective photocycloadditions of 5-methoxyoxazoles with aldehydes.

Axel G. Griesbeck*, Samir Bondock, and Johann Lex

Institute of Organic Chemistry, University of Cologne, Greinstr. 4

D-50939 Köln, Germany

Supporting Information

1)	Yields and spectroscopic data for of the oxazole substrates 2-6 :	2
2)	Yields and spectroscopic data for <i>exo</i> - 8b-8f :	2-3
3)	Yields and spectroscopic data for <i>erythro</i> - 9b-9f :	3-4
4)	Yields and spectroscopic data for Z-10d , Z-10e , and Z-10f :	4
5)	Yields and spectroscopic data for C1-substituted bicyclic oxetanes:	4-8
6)	Yields and spectroscopic data for C3-quaternary <i>erythro</i> (2S*,3S*)- α -acetamido- α -alkylated- β -hydroxy esters:	8-13
7)	¹³ C-NMR-spectra of bicyclic oxetanes:	13-23
8)	X-ray crystallographic data for compound 23 :	24-27

1) Yields and spectroscopic data for of the oxazole substrates **2-6**.

2,4-Dimethyl-5-methoxyoxazole (2) 60 %, b.p: 61-63 °C, 10 torr. $^1\text{H-NMR}$ δ 1.96 (s, 3H, CH₃), 2.27 (s, 3H, CH₃), 3.83 (s, 3H, OCH₃). $^{13}\text{C-NMR}$ δ 9.8 (q, CH₃), 14.1 (q, CH₃), 61.2 (q, OCH₃), 111.2 (s, C-4), 151.9 (s, C-5), 154.6 (s, C-2).

4-Ethyl-2-methyl-5-methoxyoxazole (3) 74 %, b.p: 64-67 °C, 10 torr. $^1\text{H-NMR}$ δ 1.07 (t, J = 7.5 Hz, 3H, CH₃), 2.22 (s, 3H, CH₃), 2.28 (q, J = 7.5 Hz, 2H, CH₂), 3.77 (s, 3H, OCH₃). $^{13}\text{C-NMR}$ δ 13.0 (q, CH₃), 14.1 (q, CH₃), 17.9 (t, CH₂), 61.2 (q, OCH₃), 117.1 (s, C-4), 151.8 (s, C-5), 154.0 (s, C-2).

2-Methyl-4-propyl-5-methoxyoxazole (4) 68 %, b.p: 75-78 °C, 10 torr. $^1\text{H-NMR}$ δ 0.83 (t, J = 7.5 Hz, 3H, CH₃), 1.52 (sextet, J = 7.5 Hz, 2H, CH₂), 2.21 (t, J = 7.5 Hz, 2H, CH₂), 2.22 (s, 3H, CH₃), 3.77 (s, 3H, OCH₃). $^{13}\text{C-NMR}$ δ 13.6 (q, CH₃), 14.1 (q, CH₃), 21.7 (t, CH₂), 26.5 (t, CH₂), 115.6 (s, C-4), 151.9 (s, C-5), 154.6 (s, C-2).

4-Isopropyl-2-methyl-5-methoxyoxazole (5) 65 %, b.p: 71-73 °C, 10 torr, $^1\text{H-NMR}$ δ 1.09 (d, J = 7.1 Hz, 6H, 2CH₃), 2.20 (s, 3H, CH₃), 2.65 (sept, J=7.1 Hz, 1H, CH), 3.75 (s, 3H, OCH₃). $^{13}\text{C-NMR}$ δ 14.1 (q, CH₃), 21.5 (q, CH₃), 24.7 (d, CH), 61.2 (q, OCH₃), 121.1 (s, C-4), 151.8 (s, C-5), 153.2 (s, C-2).

4-Isobutyl-2-methyl-5-methoxyoxazole (6) 70 %, b.p: 86-88 °C, 10 torr, $^1\text{H-NMR}$ δ 0.80 (d, J = 6.8 Hz, 6H, 2CH₃), 1.79 (sext, J = 6.6 Hz, 1H, CH), 2.08 (d, J = 6.8 Hz, 2H, CH₂), 2.19 (s, 3H, CH₃), 3.74 (s, 3H, OCH₃). $^{13}\text{C-NMR}$ δ 14.0 (q, CH₃), 22.1 (q, CH₃), 22.2 (q, CH₃), 27.6 (t, CH₂), 33.5 (d, CH), 60.9 (q, OCH₃), 114.8 (s, C-4), 151.7 (s, C-5), 154.9 (s, C-2).

2) Yields and spectroscopic data for *exo*-**8b-8f**.

***exo*-5-Methoxy-3-methyl-7-naphth-2-yl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-8b).** From 2-naphthaldehyde (0.78 g, 5 mmol) according to the general procedure. Distillation of the solvent under vacuum afforded 1.1 g of the oxetane **8b** as a pale yellow oil. 82 %, $^1\text{H-NMR}$ δ 2.00 (s, 3H, CH₃), 3.76 (s, 3H, OCH₃), 4.62 (d, J = 7.8 Hz, 1H, 1-H), 5.91 (d, J = 7.8 Hz, 1H, 7-H), 7.25-8.00 (m, 7H, H_{arom}). $^{13}\text{C-NMR}$ δ 15.5 (q, CH₃), 51.9 (q, OCH₃), 78.0 (d, C-1), 84.1 (d, C-7), 123.1 (s, C-5), 126.1 (d, CH_{arom}), 128.1 (d, CH_{arom}), 129.3 (d, CH_{arom}), 134.5 (s, Cq_{arom}), 137.1 (s, Cq_{arom}), 139.1 (s, Cq_{arom}), 168.0 (s, C-3). HRMS: (C₁₆H₁₅NO₃, M = 269.11 g/mol) Calcd: 269.1052 Found: 269.1058.

***exo*-5-Methoxy-3-methyl-7-phenethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-8c).** From 3-phenylpropionaldehyde (0.67 g, 5 mmol) according to the general procedure. Distillation of the solvent under vacuum afforded 0.98 g of the oxetane **8c** as a pale yellow oil. 79 %. $^1\text{H-NMR}$ δ 1.99 (s, 3H, CH₃), 2.08 (m, 2H, CH₂), 2.82 (m, 2H, CH₂), 3.69 (s, 3H, OCH₃), 4.87 (d, J = 5.2 Hz, 1H, 1-H), 4.91 (dd, J = 5.2, 3.4 Hz, 1H, 7-H), 7.20-7.28 (m, 5H, H_{arom}). $^{13}\text{C-NMR}$ δ 15.9 (q, CH₃), 27.9 (t, CH₂), 35.1 (t, CH₂), 52.2 (q, OCH₃), 73.3 (d, C-1), 73.6 (d, C-7), 124.5 (s, C-5), 125.7 (d, C_{arom}), 127.6 (d, C_{arom}), 135.5 (s, Cq_{arom}), 169.6 (s, C-3).

***exo*-7-Ethyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-8d).** From propion-aldehyde (0.29 g, 5 mmol) according to the general procedure. Distillation of the solvent under vacuum afforded 0.76 g of the oxetane **8d** as a pale yellow oil. 89 %. $^1\text{H-NMR}$ δ 0.98 (t, J = 7.5 Hz, 3H, CH₃), 1.25 (m, 2H, CH₂), 2.07 (s, 3H, CH₃), 3.83 (s, 3H, OCH₃), 5.04 (d, J = 7.4 Hz, 1H, 1-H), 5.24 (ddd, J = 11.6, 7.4, 4.7 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 8.7 (q, CH₃), 14.7 (q, CH₃), 24.7 (t, CH₂), 52.2 (q, OCH₃), 73.3 (d, C-1), 75.5 (d, C-7), 124.6 (s, C-5), 167.5 (s, C-3).

***exo*-7-Isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-8e).** From isobutyraldehyde (0.36 g, 5 mmol) according to the general procedure. Distillation of the solvent under vacuum afforded 0.78 g of the oxetane **8e** as a pale yellow oil. 84 %. $^1\text{H-NMR}$ δ 1.05 (d, J = 6.2 Hz, 6H, 2CH₃), 1.54 (m, 1H, CH), 1.95 (s, 3H, CH₃), 3.65 (s, 3H, OCH₃), 4.52 (d, J = 3.5 Hz, 1H, 1-H), 4.64 (dd, J=3.5, 7.9 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 15.7 (q,

CH₃), 18.8 (q, CH₃), 18.9 (q, CH₃), 31.1 (d, CH), 52.0 (q, OCH₃), 78.4 (d, C-1), 87.3 (d, C-7), 124.6 (s, C-5), 170.3 (s, C-3).

***exo*-7-Isobutyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-8f).**

From 3-methylbutyraldehyde (0.43 g, 5 mmol) according to the general procedure.

Distillation of the solvent under vacuum afforded 0.85 g of the oxetane **8f** as a pale yellow oil. 85 %. ¹H-NMR δ 0.82 (d, J = 6.6 Hz, 6H, 2CH₃), 1.62 (m, 1H, CH), 1.90 (m, 2H, CH₂), 1.97 (s, 3H, CH₃), 3.65 (s, 3H, OCH₃), 4.32 (d, J = 4.2 Hz, 1H, 1-H), 4.44 (dd, J = 4.2, 6.8 Hz, 1H, 7-H). ¹³C-NMR δ 13.4 (q, CH₃), 22.2 (q, CH₃), 24.2 (q, CH₃), 27.5 (d, CH), 37.2 (t, CH₂), 51.7 (q, OCH₃), 80.8 (d, C-1), 84.5 (d, C-7), 144.7 (s, C-5), 167.4 (s, C-3). HRMS: (C₁₀H₁₇NO₃, M = 199.12 g/mol) Calcd: 199.1275 Found: 199.1271.

3) Yields and spectroscopic data for *erythro*-9b-9f.

Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-3-naphth-2-yl-propionate (*erythro*-9b).

According to the general procedure, oxetane **8b** (0.54 g, 2 mmol) was hydrolyzed in 4h.

Preparative chromatography yielded 0.43 g of **9b** as a colorless oil. 75 %. ¹H-NMR δ 2.12 (s, 3H, CH₃CO), 3.85 (s, 3H, OCH₃), 4.47 (d, J = 9.6 Hz, 1H, CHN), 5.91 (d, J = 9.6 Hz, 1H, CHOH), 6.24 (bs, 1H, NH), 7.25-8.04 (m, 7H, H_{arom}). ¹³C-NMR δ 23.0 (q, CH₃), 51.9 (q, OCH₃), 76.1 (d, C-2), 83.1 (d, C-3), 126.7 (d, CH_{arom}), 128.5 (d, CH_{arom}), 129.1 (d, CH_{arom}), 134.5 (s, C_{qarom}), 135.7 (s, C_{qarom}), 139.1 (s, C_{qarom}), 169.3 (s, CON), 172.0 (s, COO). HRMS: (C₁₆H₁₇NO₄, M = 287.32 g/mol) Calcd: 287.1158 Found: 287.1163.

Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-5-phenylpentanoate (*erythro*-9c). According to the general procedure, oxetane **8c** (0.49 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.35 g of **9c** as a colorless oil. 65 %. ¹H-NMR δ 1.97 (s, 3H, CH₃CO), 2.43 (m, 2H, CH₂), 2.89 (m, 2H, CH₂), 3.74 (s, 3H, OCH₃), 4.84 (dd, J = 3.2, 8.2 Hz, 1H, CHN), 5.06 (dd, J = 3.2, 7.4 Hz, 1H, CHOH), 6.24 (bs, 1H, NH), 7.12-7.28 (m, 5H, H_{arom}). ¹³C-NMR δ 23.1 (q, CH₃), 30.6 (t, CH₂), 36.4 (t, CH₂), 52.6 (q, OCH₃), 55.0 (d, C-2), 73.8 (d, C-3), 126.2 (d, CH_{arom}), 128.5 (d, CH_{arom}), 129.3 (d, CH_{arom}), 140.6 (s, C_{qarom}), 169.8 (s, CON), 172.8 (s, COO). HRMS: (C₁₅H₂₁NO₄, M = 279.15 g/mol) Calcd: 279.1465 Found: 279.1458.

Methyl (2S*,3S*) 2-acetylamino-3-hydroxypentanoate (*erythro*-9d). According to the general procedure, oxetane **8d** (0.34 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.27 g of **9d** as a colorless oil. 72 %. IR: $\tilde{\nu}$ (cm⁻¹) = 3320, 3300, 2989, 1725, 1648, 1440, 1341, 1052, 967. ¹H-NMR δ 1.06 (t, J = 7.5 Hz, 3H, CH₃), 1.67 (m, 2H, CH₂), 2.03 (s, 3H, CH₃CO), 3.77 (s, 3H, OCH₃), 4.01 (dd, J = 8.2, 4.7 Hz, 1H, CHN), 4.88 (dd, J = 8.2, 3.4 Hz, 1H, CHOH). ¹³C-NMR δ 11.5 (q, CH₃), 23.1 (q, CH₃), 28.6 (t, CH₂), 52.6 (q, OCH₃), 56.5 (d, C-2), 65.3 (d, C-3), 169.1 (s, CON), 169.6 (s, COO). HRMS: (C₈H₁₅NO₄, M = 189.21 g/mol) Calcd: 189.1001 Found: 189.1008.

Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-4-methylpentanoate (*erythro*-9e). According to the general procedure, oxetane **8e** (0.37 g, 2 mmol) was hydrolyzed in 4h. Preparative chromatography yielded 0.32 g of **9e** as a colorless oil. 78 %. ¹H-NMR δ 0.93 (d, J = 6.6 Hz, 3H, CH₃), 0.98 (d, J = 6.8 Hz, 3H, CH₃), 1.67 (m, 1H, CH), 2.04 (s, 3H, CH₃CO), 3.76 (s, 3H, OCH₃), 4.45 (dd, J = 8.6, 3.3 Hz, 1H, CHN), 4.75 (dd, J = 7.4, 3.3 Hz, 1H, CHOH). ¹³C-NMR δ 18.5 (q, CH₃), 19.9 (q, CH₃), 20.5 (q, CH₃), 32.7 (d, CH), 53.1 (q, OCH₃), 54.3 (d, C-2), 69.8 (d, C-3), 169.3 (s, CON), 171.1 (s, COO). HRMS: (C₉H₁₇NO₄, M = 203.24 g/mol) Calcd: 203.1158 Found: 203.1151.

Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-5-methylhexanoate (*erythro*-9f). According to the general procedure, oxetane **8f** (0.4 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.32 g of **9f** as a colorless oil. 74 %. ¹H-NMR δ 0.98 (d, J = 6.6 Hz, 6H, 2CH₃), 1.25 (m, 1H, CH), 1.54 (m, 2H, CH₂), 2.04 (s, 3H, CH₃CO), 3.75 (s, 3H, OCH₃), 4.18 (ddd, J = 7.6, 3.2, 4.7 Hz, 1H, CHN), 4.87 (dd, J = 8.2, 3.2 Hz, 1H, CHOH). ¹³C-NMR δ 19.5 (q, CH₃), 21.3 (q, CH₃), 21.4 (q, CH₃), 28.3 (d, CH), 34.1 (t, CH₂), 52.7 (q, OCH₃), 56.8

(d, C-2), 61.8 (d, C-3), 169.3 (s, CON), 171.3 (s, COO). HRMS: ($C_{10}H_{19}NO_4$, $M = 217.13$ g/mol) Calcd: 217.1287 Found: 217.1281.

4) Yields and spectroscopic data for **Z-10d**, **Z-10e**, and **Z-10f**.

Methyl 2-acetylamino-pent-2-enoate (Z-10d): According to the general procedure, methyl 2-acetylamino-3-hydroxypentanoate **9d** (0.19 g, 1 mmol) was dehydrated in 24h. Preparative chromatography yielded 0.13 g of the product as a colorless oil which was solidified by treatment with petroleum ether as a colorless powder. 75 %, m.p: 55-56°C (ref. 58-60°C).

Methyl 2-acetylamino-4-methyl-pent-2-enoate (Z-10e). According to the general procedure, methyl 2-acetylamino-3-hydroxy-4-methylpentanoate **9e** (0.2 g, 1 mmol) was dehydrated in 24h. Preparative chromatography yielded 0.14 g of the product as a colorless oil which was solidified by treatment with petroleum ether as a colorless powder. 78 %, m.p: 71-72°C (ref. 73-75°C).

Methyl 2-acetylamino-5-methyl-hex-2-enoate (Z-10f). According to the general procedure, methyl 2-acetylamino-3-hydroxy-5-methylhexanoate **9f** (0.22 g, 1 mmol) was dehydrated in 24h. Preparative chromatography yielded 0.17 g of the product as a colorless oil which was solidified by treatment with petroleum ether as a colorless powder. 83 %, m.p: 66-68°C (ref. 70-71°C).

5) Yields and spectroscopic data for C1-substituted bicyclic oxetanes

exo-1-Ethyl-5-methoxy-3-methyl-7-phenyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-13a). From benzaldehyde (0.53 g, 5 mmol) and 4-ethyl-2-methyl-5-methoxyoxazole (**3**) (0.71 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.31 g of the oxetane **13a** as a pale yellow oil. 87 %, 1H -NMR δ 0.84 (t, $J = 7.2$ Hz, 3H, CH_3), 1.06 (q, $J = 7.2$ Hz, 2H, CH_2), 2.00 (s, 3H, CH_3), 3.89 (s, 3H, OCH_3), 5.79 (s, 1H, 7-H), 7.26-7.35 (m, 5H, H_{arom}). ^{13}C -NMR δ 8.5 (q, CH_3), 23.7 (q, CH_3), 24.9 (t, CH_2), 51.7 (q, OCH_3), 79.2 (s, C-1), 82.0 (d, C-7), 124.5 (s, C-5), 125.6 (d, CH_{arom}), 127.3 (d, CH_{arom}), 129.2 (d, CH_{arom}), 136.5 (s, Cq_{arom}), 164.9 (s, C-3). HRMS: ($C_{14}H_{17}NO_3$, $M = 247.12$ g/mol) Calcd: 247.1204, Found: 247.1197

exo-5-Methoxy-3-methyl-7-phenyl-1-propyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-15a). From benzaldehyde (0.53g, 5 mmol) and 2-methyl-4-propyl-5-methoxyoxazole (**4**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.14 g of the oxetane **15a** as a pale yellow oil. 87 %, IR: (film) $\tilde{\nu}$ (cm^{-1}) = 2992, 1628, 1598, 1440, 1348, 1042, 960. 1H -NMR δ 0.55 (t, $J = 7.1$ Hz, 3H, CH_3), 0.78 (sextet, $J = 7.1$ Hz, 2H, CH_2), 1.12 (t, $J = 7.1$ Hz, 2H, CH_2), 2.00 (s, 3H, CH_3), 3.55 (s, 3H, OCH_3), 5.17 (s, 1H, 7-H), 7.25-7.32 (m, 5H, H_{arom}). ^{13}C -NMR δ 14.1 (q, CH_3), 15.1 (q, CH_3), 16.4 (t, CH_2), 29.6 (t, CH_2), 51.3 (q, OCH_3), 78.9 (s, C-1), 89.9 (d, C-7), 124.6 (s, C-5), 126.4 (d, CH_{arom}), 128.2 (d, CH_{arom}), 137.1 (s, Cq_{arom}), 165.1 (s, C-3). HRMS: ($C_{15}H_{19}NO_3$, $M = 261.14$ g/mol) Calcd: 261.1360, Found: 261.1356.

1-Isopropyl-5-methoxy-3-methyl-7-phenyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (17a). From benzaldehyde (0.53 g, 5 mmol) and 4-isopropyl-2-methyl-5-methoxyoxazole (**5**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.1 g (85% total yield) of an inseparable mixture of *exo* and *endo* oxetanes **17a** as a pale yellow oil.

exo-diastereoisomer (exo-17a). 1H -NMR δ 0.53 (d, $J = 6.8$ Hz, 3H, CH_3), 0.76 (d, $J = 6.8$ Hz, 3H, CH_3), 1.89 (m, 1H, CH), 2.02 (s, 3H, CH_3), 3.61 (s, 3H, OCH_3), 5.26 (s, 1H, 7-H), 7.28-7.35 (m, 5H, H_{arom}). ^{13}C -NMR δ 14.5 (q, CH_3), 15.3 (q, CH_3), 16.3 (q, CH_3), 24.9 (d, CH), 50.5 (q, OCH_3), 82.2 (s, C-1), 90.2 (d, C-7), 124.3 (s, C-5), 127.4 (d, CH_{arom}), 127.8 (d, CH_{arom}), 128.2 (d, C_{arom}), 135.9 (s, Cq_{arom}), 164.8 (s, C-3). **endo-diastereoisomer (endo-17a)**. 1H -NMR δ 0.93 (d, $J = 6.8$ Hz, 3H, CH_3), 0.98 (d, $J = 6.9$ Hz, 3H, CH_3), 1.95 (m, 1H, CH), 1.67 (s, 3H, CH_3), 3.71 (s, 3H, OCH_3), 5.28 (s, 1H, 7-H), 7.30-7.35 (m, 5H, H_{arom}). ^{13}C -NMR δ 14.7 (q, CH_3), 15.4 (q, CH_3), 16.3 (q, CH_3), 24.7 (d, CH), 50.8 (q, OCH_3), 82.7 (s, C-1), 90.7 (d, C-7), 124.3 (s, C-5), 127.3 (d, CH_{arom}), 127.8 (d, CH_{arom}), 128.2 (d, CH_{arom}), 135.5 (s, Cq_{arom}), 165.1 (s, C-3).

exo-1-Isobutyl-5-methoxy-3-methyl-7-phenyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-19a). From benzaldehyde (0.53 g, 5 mmol) and 4-isobutyl-2-methyl-5-methoxyoxazole

(6) (0.84 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.16 g of the oxetane **19a** as a pale yellow oil. 84%, $^1\text{H-NMR}$ δ 0.58 (d, J = 6.8 Hz, 3H, CH_3), 0.72 (d, J = 6.6 Hz, 3H, CH_3), 0.90 (m, 1H, CH), 1.20 (dd, J = 5.3, 7.8 Hz, 2H, CH_2), 2.21 (s, 3H, CH_3), 3.67 (s, 3H, OCH_3), 5.26 (s, 1H, 7-H), 7.30-7.49 (m, 5H, H_{arom}). $^{13}\text{C-NMR}$ δ 15.1 (q, CH_3), 22.9 (q, CH_3), 23.7 (q, CH_3), 23.8 (d, CH), 35.6 (t, CH_2), 51.3 (q, OCH_3), 78.7 (s, C-1), 90.4 (d, 7-H), 124.8 (s, C-5), 126.8 (d, C_{arom}), 127.9 (d, C_{arom}), 128.8 (d, C_{arom}), 164.7 (s, C-3).

exo-5-Methoxy-1,3-dimethyl-7-naphth-2-yl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-11b). From 2-naphthaldehyde (0.64 g, 5 mmol) and 2,4-dimethyl-5-methoxyoxazole (**2**) (0.64 g, 5 mmol) in benzene (50 mL). Distillation of the solvent under vacuum afforded 1.3 g of the oxetane **11b** as a pale yellow oil. 92 %, $^1\text{H-NMR}$ δ 0.82 (s, 3H, CH_3), 2.00 (s, 3H, CH_3), 3.54 (s, 3H, OCH_3), 5.21 (s, 1H, 7-H), 7.42-7.81 (m, 7H, H_{arom}). $^{13}\text{C-NMR}$ δ 13.6 (q, CH_3), 14.9 (q, CH_3), 51.4 (q, OCH_3), 75.9 (s, C-1), 89.5 (d, C-7), 124.6 (s, C-5), 125.9 (d, CH_{arom}), 128.2 (d, CH_{arom}), 129.5 (d, CH_{arom}), 134.3 (s, C_{qarom}), 136.2 (s, C_{qarom}), 137.0 (s, C_{qarom}), 165.3 (s, C-3).

exo-5-Methoxy-1,3-dimethyl-7-phenylethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-11c). From 3-phenylpropionaldehyde (0.67 g, 5 mmol) and 2,4-dimethyl-5-methoxyoxazole (**2**) (0.64 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.27 g of the oxetane **11c** as a pale yellow oil. 87 %, $^1\text{H-NMR}$ δ 1.18 (s, 3H, CH_3), 1.95 (s, 3H, CH_3), 2.70 (m, 2H, CH_2), 2.92 (m, 2H, CH_2), 3.43 (s, 3H, OCH_3), 4.15 (dd, J = 4.3, 9.5 Hz, 1H, 7-H), 7.26 (m, 5H, H_{arom}). $^{13}\text{C-NMR}$ δ 12.1 (q, CH_3), 14.9 (q, CH_3), 27.9 (t, CH_2), 45.1 (t, CH_2), 51.3 (q, OCH_3), 73.4 (s, C-1), 87.9 (d, C-7), 124.5 (s, C-5), 128.0 (d, CH_{arom}), 128.8 (d, CH_{arom}), 129.5 (d, CH_{arom}), 140.7 (s, C_{qarom}), 164.8 (s, C-3). HRMS: ($\text{C}_{15}\text{H}_{19}\text{NO}_3$, M = 261.14 g/mol) Calcd: 261.1360, Found: 261.1358.

exo-1-Isopropyl-5-methoxy-3-methyl-7-phenylethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-17c). From 3-phenylpropionaldehyde (0.67 g, 5 mmol) and 4-isopropyl-2-methyl-5-methoxyoxazole (**5**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.27 g of the oxetane **17c** as a pale yellow oil. 87 %, $^1\text{H-NMR}$ δ 0.71 (d, J = 6.6 Hz, 3H, CH_3), 0.85 (d, J = 6.8 Hz, 3H, CH_3), 1.87 (m, 2H, CH_2), 1.95 (s, 3H, CH_3), 2.44 (m, 2H, CH_2), 2.75 (m, 1H, CH), 3.46 (s, 3H, OCH_3), 4.18 (dd, J = 11.2, 2.7 Hz, 1H, 7-H), 7.21-7.34 (m, 5H, H_{arom}). $^{13}\text{C-NMR}$ δ 14.0 (q, CH_3), 14.8 (q, CH_3), 16.6 (q, CH_3), 17.7 (d, CH), 30.6 (t, CH_2), 33.4 (t, CH_2), 50.8 (q, OCH_3), 79.9 (s, C-1), 88.3 (d, C-7), 124.5 (s, C-5), 125.6 (d, CH_{arom}), 126.1 (d, CH_{arom}), 128.9 (d, CH_{arom}), 140.2 (s, C_{qarom}), 165.2 (s, C-3).

exo-1-Ethyl-5-methoxy-1,3-dimethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-11d). From propionaldehyde (0.29 g, 5 mmol) and 2,4-dimethyl-5-methoxyoxazole (**2**) (0.64 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.78 g of the oxetane **11d** as a pale yellow oil. 84 %, $^1\text{H-NMR}$ δ 0.84 (t, J = 7.2 Hz, 3H, CH_3), 1.80 (s, 3H, CH_3), 1.55 (dt, J = 7.8, 7.2 Hz, 2H, CH_2), 1.95 (s, 3H, CH_3), 3.41 (s, 3H, OCH_3), 4.05 (dd, J = 7.8, 6.3 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 8.7 (q, CH_3), 11.8 (q, CH_3), 14.7 (q, CH_3), 25.1 (t, CH_2), 51.0 (q, OCH_3), 73.2 (s, C-1), 90.1 (d, C-7), 124.3 (s, C), 164.7 (s, C-3). HRMS: ($\text{C}_9\text{H}_{15}\text{NO}_3$, M = 185.11 g/mol) Calcd: 185.1048, Found: 185.1039.

exo-1-Diethyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-13d). From propionaldehyde (0.29 g, 5 mmol) and 4-ethyl-2-methyl-5-methoxyoxazole (**3**) (0.71 g, 5 mmol) in benzene (50 mL). Distillation of the solvent under vacuum afforded 0.9 g of the oxetane **13d** as a pale yellow oil. 90 %, $^1\text{H-NMR}$ δ 0.76 (t, J = 7.5 Hz, 3H, CH_3), 0.82 (t, J = 7.4 Hz, 3H, CH_3), 1.55 (q, J = 7.5 Hz, 2H, CH_2), 1.58 (m, 2H, CH_2), 1.93 (s, 3H, CH_3), 3.58 (s, 3H, OCH_3), 4.03 (dd, J = 5.7, 8.3 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 7.8 (q, CH_3), 8.9 (q, CH_3), 14.7 (q, CH_3), 19.0 (t, CH_2), 24.9 (t, CH_2), 50.8 (q, OCH_3), 76.7 (s, C-1), 90.4 (d, C-7), 124.3 (s, C-5), 164.8 (s, C-3).

exo-7-Ethyl-5-methoxy-3-methyl-1-propyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-15d). From propionaldehyde (0.29 g, 5 mmol) and 2-methyl-4-propyl-5-methoxyoxazole (**4**)

(0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.92 g of the oxetane **15d** as a pale yellow oil. 86 %, IR: (film) $\tilde{\nu}$ (cm⁻¹) = 2998, 1615, 1440, 1110, 967. ¹H-NMR δ 0.81 (t, J = 7.1 Hz, 3H, CH₃), 0.84 (t, J = 7.5 Hz, 3H, CH₃), 0.85 (t, J = 7.1 Hz, 2H, CH₂), 1.21 (m, 2H, CH₂), 1.65 (m, 2H, CH₂), 1.96 (s, 3H, CH₃), 3.41 (s, 3H, OCH₃), 4.07 (dd, J = 6.0, 8.0 Hz, 1H, 7-H). ¹³C-NMR δ 8.9 (q, CH₃), 14.2 (q, CH₃), 16.9 (t, CH₂), 24.9 (t, CH₂), 28.3 (t, CH₂), 50.8 (q, OCH₃), 76.4 (s, C-1), 90.5 (d, C-7), 124.3 (s, C-5), 164.5 (s, C-3). MS: (EI, 70 eV) m/z (%) = 184 (M⁺-Et, 12), 172 (M⁺-CH₃CN, 16), 155 (80), 144 (75), 126 (34), 112 (43), 99 (20), 95 (15), 86 (55), 68 (88), 57 (100).

exo-7-Ethyl-1-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-17d). From propionaldehyde (0.29 g, 5 mmol) and 4-isopropyl-2-methyl-5-methoxyoxazole (**5**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.88 g of the oxetane **17d** as a pale yellow oil. 83 %, ¹H-NMR δ 0.71 (d, J = 6.8 Hz, 3H, CH₃), 0.87 (t, J = 7.5 Hz, 3H, CH₃), 1.65 (m, 2H, CH₂), 1.97 (s, 3H, CH₃), 2.15 (s, 3H, CH₃), 3.42 (s, 3H, OCH₃), 4.10 (dd, J = 9.4, 4.4 Hz, 1H, 7-H). ¹³C-NMR δ 9.0 (q, CH₃), 14.9 (q, CH₃), 16.6 (q, CH₃), 17.7 (q, CH₃), 21.5 (d, CH), 24.7 (t, CH₂), 50.7 (q, OCH₃), 80.0 (s, C-1), 90.8 (d, C-7), 124.5 (s, C-5), 164.9 (s, C-3).

exo-7-Ethyl-1-isobutyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-19d). From propionaldehyde (0.29 g, 5 mmol) and 4-isobutyl-2-methyl-5-methoxyoxazole (**6**) (0.84 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.99 g of the oxetane **19d** as a pale yellow oil. 87 %, ¹H-NMR δ 0.75 (d, J = 6.6 Hz, 3H, CH₃), 0.86 (d, J = 7.4 Hz, 3H, CH₃), 1.05 (m, 1H, CH), 1.54 (t, J = 7.5 Hz, 3H, CH₃), 1.67 (m, 2H, CH₂), 1.88 (m, 2H, CH₂), 1.95 (s, 3H, CH₃), 3.42 (s, 3H, OCH₃), 4.05 (dd, J = 7.6, 6.2 Hz, 1H, 7-H). ¹³C-NMR δ 8.9 (q, CH₃), 13.9 (q, CH₃), 14.8 (q, CH₃), 20.8 (d, CH), 22.9 (q, CH₃), 27.6 (t, CH₂), 34.1 (t, CH₂), 50.8 (q, OCH₃), 76.9 (s, C-1), 90.9 (d, C-7), 124.4 (s, C-5), 164.0 (s, C-3). HRMS: (C₁₂H₂₁NO₃, M = 227.15 g/mol) Calcd: 227.1516, Found: 227.1512.

exo-7-Isopropyl-5-methoxy-1,3-dimethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-11e). From isobutyraldehyde (0.36 g, 5 mmol) and 2,4-dimethyl-5-methoxyoxazole (**2**) (0.64 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.93 g of the oxetane **11e** as a pale yellow oil. 93 %, ¹H-NMR: (300 MHz, CDCl₃) δ 0.82 (d, J = 7.5 Hz, 3H, CH₃), 0.85 (d, J = 7.2 Hz, 3H, CH₃), 0.87 (m, 1H, CH), 1.31 (s, 3H, CH₃), 2.10 (s, 3H, CH₃), 3.48 (s, 3H, OCH₃), 4.28 (d, J = 3.4 Hz, 1H, 7-H). ¹³C-NMR δ 12.4 (q, CH₃), 14.9 (q, CH₃), 17.7 (q, CH₃), 19.3 (q, CH₃), 23.8 (d, CH), 52.5 (q, OCH₃), 73.0 (s, C-1), 93.9 (d, C-7), 124.6 (s, C-5), 165.0 (s, C-3).

exo-1-Ethyl-7-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-13e). From isobutyraldehyde (0.36 g, 5 mmol) and 4-ethyl-2-methyl-5-methoxyoxazole (**3**) (0.71 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.93 g of the oxetane **13e** as a pale yellow oil. 87 %, ¹H-NMR δ 0.75 (d, J = 7.2 Hz, 6H, 2CH₃), 0.85 (m, 1H, CH), 1.00 (t, J = 7.5 Hz, 3H, CH₃), 1.58 (m, 2H, CH₂), 2.15 (s, 3H, CH₃), 3.60 (s, 3H, OCH₃), 4.04 (d, J = 8.4 Hz, 1H, CH). ¹³C-NMR δ 8.7 (q, CH₃), 15.0 (q, CH₃), 18.3 (q, CH₃), 18.9 (q, CH₃), 21.2 (t, CH₂), 27.8 (d, CH), 50.8 (q, OCH₃), 78.3 (s, C-1), 85.1 (d, C-7), 124.1 (s, C-5), 164.1 (s, C-3).

exo-7-Isopropyl-5-methoxy-3-methyl-1-propyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-15e). From isobutyraldehyde (0.36 g, 5 mmol) and 2-methyl-4-propyl-5-methoxyoxazole (**4**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.98 g of the oxetane **15e** as a pale yellow oil. 86 %, ¹H-NMR δ 0.79 (d, J = 6.8 Hz, 6H, 2CH₃), 0.82 (t, J = 7.2 Hz, 3H, CH₃), 1.12 (m, 2H, CH₂), 1.23 (m, 2H, CH₂), 1.75 (m, 2H, CH₂), 1.78 (m, 1H, CH), 2.21 (s, 3H, CH₃), 3.44 (s, 3H, OCH₃), 3.66 (d, J = 12.2 Hz, 1H, 7-H). ¹³C-NMR δ 13.5 (q, CH₃), 13.9 (q, CH₃), 14.3 (q, CH₃), 16.7 (q, CH₃), 17.7 (d, CH), 26.3 (t, CH₂), 28.6 (t, CH₂), 50.8 (q, OCH₃), 90.7 (s, C-1), 94.1 (d, C-7), 124.1 (s, C-5), 164.8 (s, C-3). HRMS: (C₁₂H₂₁NO₃, M = 227.15 g/mol) Calcd: 227.1516, Found: 227.1509.

***exo*-1,7-Diisopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-17e).** From isobutyraldehyde (0.36 g, 5 mmol) and 4-isopropyl-2-methyl-5-methoxyoxazole (**5**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.92 g of the oxetane **17e** as a pale yellow oil. 81 %, $^1\text{H-NMR}$ δ 0.78 (d, J = 6.6 Hz, 6H, 2CH₃), 0.82 (m, 1H, CH), 0.85 (d, J = 7.2 Hz, 3H, CH₃), 0.92 (t, J = 7.4 Hz, 3H, CH₃), 1.43 (m, 1H, CH), 1.85 (m, 2H, CH₂), 2.00 (s, 3H, CH₃), 3.62 (s, 3H, OCH₃), 4.18 (d, J = 8.4 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 8.7 (q, CH₃), 11.3 (q, CH₃), 11.9 (t, CH₂), 14.5 (q, CH₃), 19.3 (q, CH₃), 19.5 (q, CH₃), 22.3 (d, CH), 27.2 (d, CH), 52.0 (q, OCH₃), 82.3 (s, C-1), 88.1 (d, C-7), 124.3 (s, C-5), 165.7 (s, C-3).

***exo*-1-Isobutyl-7-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-19e).** From isobutyraldehyde (0.36 g, 5 mmol) and 4-isobutyl-2-methyl-5-methoxyoxazole (**6**) (0.84 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.1 g of the oxetane **19e** as a pale yellow oil. 91 %, $^1\text{H-NMR}$ δ 0.82 (d, J = 6.8 Hz, 6H, 2CH₃), 0.85 (d, J = 7.2 Hz, 6H, 2CH₃), 1.12 (m, 2H, 2CH), 1.78 (m, 2H, CH₂), 1.98 (s, 3H, CH₃), 3.46 (s, 3H, OCH₃), 3.79 (d, J = 12.1 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 14.1 (q, CH₃), 14.9 (q, CH₃), 17.7 (q, CH₃), 18.6 (q, CH₃), 18.9 (q, CH₃), 29.7 (d, CH), 34.4 (d, CH), 40.6 (t, CH₂), 50.8 (q, OCH₃), 91.5 (s, C-1), 94.1 (d, C-7), 124.3 (s, C-5), 164.5 (s, C-3).

***exo*-7-Isobutyl-5-methoxy-1,3-dimethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-11f).** From 3-methylbutyraldehyde (0.43 g, 5 mmol) and 2,4-dimethyl-5-methoxyoxazole (**2**) (0.64 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.89 g of the oxetane **11f** as a pale yellow oil. 84 %, $^1\text{H-NMR}$ δ 0.83 (d, J = 6.6 Hz, 3H, CH₃), 0.86 (d, J = 6.6 Hz, 3H, CH₃), 0.92 (m, 1H, CH), 1.23 (s, 3H, CH₃), 1.43 (m, 2H, CH₂), 2.02 (s, 3H, CH₃), 3.47 (s, 3H, OCH₃), 4.32 (dd, J = 4.0, 8.5 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 12.2 (q, CH₃), 14.9 (q, CH₃), 21.9 (q, CH₃), 23.2 (d, CH), 24.4 (q, CH₃), 40.8 (t, CH₂), 51.3 (q, OCH₃), 73.6 (s, C-1), 82.8 (d, C-7), 124.6 (s, C-5), 165.0 (s, C-3). HRMS: (C₁₁H₁₉NO₃, M = 213.14 g/mol) Calcd: 213.1360, Found: 213.1353.

***exo*-1-Ethyl-7-isobutyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-13f).** From 3-methylbutyraldehyde (0.43 g, 5 mmol) and 4-ethyl-2-methyl-5-methoxyoxazole (**3**) (0.64 g, 5 mmol). Distillation of the solvent under vacuum afforded 0.98 g of the oxetane **13f** as a pale yellow oil. 86 %, $^1\text{H-NMR}$ δ 0.88 (d, J = 6.8 Hz, 3H, CH₃), 0.92 (d, J = 6.6 Hz, 3H, CH₃), 1.02 (t, J = 7.5 Hz, 3H, CH₃), 1.23 (m, 1H, CH), 1.43 (m, 2H, CH₂), 1.54 (m, 2H, CH₂), 1.54 (m, 2H, CH₂), 2.03 (s, 3H, CH₃), 3.78 (s, 3H, OCH₃), 4.12 (dd, J = 9.2, 2.1 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 9.0 (q, CH₃), 15.4 (q, CH₃), 19.2 (q, CH₃), 19.7 (q, CH₃), 23.1 (t, CH₂), 27.3 (d, CH), 34.1 (t, CH₂), 50.2 (q, OCH₃), 79.2 (s, C-1), 88.3 (d, C-7), 124.2 (s, C-5), 164.7 (s, C-3).

***exo*-7-Isobutyl-5-methoxy-3-methyl-1-propyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-15f).** From 3-methylbutyraldehyde (0.43 g, 5 mmol) and 2-methyl-4-propyl-5-methoxyoxazole (**4**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.0 g of the oxetane **15f** as a pale yellow oil. 83 %, $^1\text{H-NMR}$ δ 0.80 (d, J = 7.1 Hz, 6H, 2CH₃), 0.83 (t, J = 7.2 Hz, 3H, CH₃), 1.23 (m, 2H, CH₂), 1.65 (m, 2H, CH₂), 1.73 (m, 1H, CH), 1.98 (s, 3H, CH₃), 3.41 (s, 3H, OCH₃), 4.27 (dd, J = 10.2, 4.9 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 14.3 (q, CH₃), 14.9 (q, CH₃), 16.9 (t, CH₂), 21.7 (d, CH), 23.2 (q, CH₃), 24.2 (q, CH₃), 28.4 (t, CH₂), 40.4 (t, CH₂), 50.9 (q, OCH₃), 76.5 (s, C-1), 87.8 (d, C-7), 124.4 (s, C-5), 164.7 (s, C-3). HRMS: (C₁₃H₂₃NO₃, M = 241.17 g/mol) Calcd: 241.1764, Found: 241.1762.

***exo*-7-Isobutyl-1-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-17f).** From 3-methylbutyraldehyde (0.43 g, 5 mmol) and 4-isopropyl-2-methyl-5-methoxyoxazole (**5**) (0.75 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.1 g of the oxetane **17f** as a pale yellow oil. 87 %, $^1\text{H-NMR}$ δ 0.72 (d, J = 6.6 Hz, 3H, CH₃), 0.81 (d, J = 6.5 Hz, 3H, CH₃), 0.88 (d, J = 6.8 Hz, 6H, 2CH₃), 1.35 (m, 2H, 2CH), 1.76 (m, 2H, CH₂), 1.98 (s, 3H, CH₃), 3.42 (s, 3H, OCH₃), 4.29 (dd, J = 11.2, 2.3 Hz, 1H, 7-H). $^{13}\text{C-NMR}$ δ 14.8 (q, CH₃), 16.6 (q, CH₃), 17.7 (q, CH₃), 21.3 (q, CH₃), 23.5 (q, CH₃), 23.9 (d, CH), 24.7

(d, CH), 40.1 (t, CH₂), 50.7 (q, OCH₃), 80.0 (s, C-1), 87.9 (d, C-7), 124.4 (s, C-5), 164.9 (s, C-3).

exo-1,7-Diisobutyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (exo-19f). From 3-methylbutyraldehyde (0.43 g, 5 mmol) and 4-isobutyl-2-methyl-5-methoxyoxazole (6) (0.84 g, 5 mmol). Distillation of the solvent under vacuum afforded 1.12 g of the oxetane **19f** as a pale yellow oil. 88 %, ¹H-NMR δ 0.81 (d, J = 6.8 Hz, 3H, CH₃), 0.83 (d, J = 6.7 Hz, 3H, CH₃), 0.85 (d, J = 7.1 Hz, 3H, CH₃), 0.89 (d, J = 7.5 Hz, 3H, CH₃), 1.53 (m, 2H, CH₂), 1.54 (m, 2H, CH₂), 1.56 (m, 2H, CH₂), 2.00 (s, 3H, CH₃), 3.46 (s, 3H, OCH₃), 4.27 (dd, J = 1.3, 7.7 Hz, 1H, 7-H). ¹³C-NMR δ 15.0 (q, CH₃), 21.6 (q, CH₃), 23.1 (q, CH₃), 23.4 (q, CH₃), 23.9 (q, CH₃), 24.2 (d, CH), 24.4 (d, CH), 34.3 (t, CH₂), 40.7 (t, CH₂), 51.0 (q, OCH₃), 76.3 (s, C-1), 88.3 (d, C-7), 124.6 (s, C-5), 164.2 (s, C-3). HRMS: (C₁₄H₂₅NO₃, M = 255.18 g/mol) Calcd: 255.1828, Found: 255.1822.

6) Yields and spectroscopic data for C3-quaternary *erythro* (2S*,3S*)- α -acetamido- α -alkylated- β -hydroxy esters.

erythro-Methyl (2S*,3S*) 2-(N-acetylamino)-2-ethyl-3-hydroxy-3-phenylpropanoate (erythro-14a). Following the general procedure, oxetane **13a** (0.49 g, 2 mmol) was hydrolyzed in 5h. Preparative chromatography yielded 0.39 g of **14a** as a colorless oil. 73 %, ¹H-NMR: δ 1.08 (t, J = 7.5 Hz, 3H, CH₃), 1.35 (q, J = 7.5 Hz, 2H, CH₂), 1.99 (s, 3H, CH₃CO), 3.89 (s, 3H, OCH₃), 4.35 (s, 1H, CHOH), 7.26-7.34 (m, 5H, H_{arom}). ¹³C-NMR: (75.5 MHz, CDCl₃) δ 8.8 (q, CH₃), 24.1 (q, CH₃), 25.4 (t, CH₂), 53.0 (q, OCH₃), 63.2 (s, C-2), 70.2 (d, C-3), 126.1 (d, CH_{arom}), 128.1 (d, CH_{arom}), 129.3 (d, CH_{arom}), 134.5 (s, C_{qarom}), 170.3 (s, CON), 171.5 (s, COO). HRMS: (C₁₄H₁₉NO₄, M = 265.13 g/mol) Calcd: 265.1314, Found: 265.1310.

erythro-Methyl (2S*,3S*) 2-(N-acetylamino)-3-hydroxy-2-propyl-3-phenylpentanoate (erythro-16a). Following the general procedure, oxetane **15a** (0.52 g, 2 mmol) was hydrolyzed in 6h. Preparative chromatography yielded 0.32 g of **16a** as a colorless oil. 58 %, ¹H-NMR δ 0.88 (t, J = 7.4 Hz, 3H, CH₃), 1.27 (m, 2H, CH₂), 1.51-1.81 (m, 2H, CH₂), 1.98 (s, 3H, CH₃CO), 3.78 (s, 3H, OCH₃), 4.02 (s, 1H, CHOH), 6.21 (bs, 1H, NH), 7.22-7.26 (m, 5H, H_{arom}). ¹³C-NMR δ 13.6 (q, CH₃), 18.5 (q, CH₃), 23.1 (t, CH₂), 34.6 (t, CH₂), 52.2 (q, OCH₃), 70.1 (s, C-2), 78.0 (d, C-3), 125.7 (d, CH_{arom}), 126.9 (d, CH_{arom}), 128.1 (d, CH_{arom}), 136.4 (s, C_{qarom}), 169.8 (s, CON), 173.3 (s, COO). MS (EI, 70 eV) m/z (%) = 278 (M⁺-1, 5), 262 (M⁺-OH, 7), 202 (8), 220 (M⁺-CO₂Me, 10), 219 (55), 155 (78), 127 (100), 105 (43), 91 (15), 77 (30), 59 (13). Anal: (C₁₅H₂₁NO₄, M = 279.33 g/mol) Calcd: C 64.50 H 7.58 N 5.01, Found: C 64.72 H 7.42 N 5.16.

Methyl (2S*,3S*) 2-(N-acetylamino)-2-(hydroxy-phenyl-methyl)-3-methylbutanoate (18a). Following the general procedure, oxetane **17a** (0.52 g, 2 mmol) were hydrolyzed in 4h. Preparative chromatography yielded 0.27 g of the *erythro*-isomer and 0.12 g of the *threo*-isomer, both as a colorless oils. **erythro-diastereoisomer:** 48 %, TLC: R_f = 0.47 (ethylacetate/n-hexane 1: 4), ¹H-NMR δ 0.76 (d, J = 6.9 Hz, 3H, CH₃), 1.23 (d, J = 6.8 Hz, 3H, CH₃), 1.43 (sextet, J = 6.8 Hz, 1H, CH), 2.13 (s, 3H, CH₃CO), 3.78 (s, 3H, OCH₃), 4.14 (s, 1H, CHOH), 7.32-7.34 (m, 5H, H_{arom}). ¹³C-NMR δ 17.5 (q, CH₃), 18.2 (q, CH₃), 23.7 (q, CH₃), 30.3 (d, CH), 52.8 (q, OCH₃), 75.1 (s, C-2), 75.6 (d, C-3), 125.8 (d, CH_{arom}), 127.8 (d, CH_{arom}), 128.8 (d, CH_{arom}), 141.2 (s, C_{qarom}), 170.2 (s, CON), 171.4 (s, COO). Anal: (C₁₅H₂₁NO₄, M = 279.33 g/mol) Calcd: C 64.50 H 7.58 N 5.01, Found: C 64.65 H 7.38 N 4.98.

threo-diastereoisomer: 25 %, TLC: R_f = 0.31 (ethylacetate/n-hexane 1: 4), ¹H-NMR δ 0.95 (d, J = 6.8 Hz, 3H, CH₃), 0.99 (d, J = 6.6 Hz, 3H, CH₃), 2.19 (s, 3H, CH₃CO), 2.33 (sextet, J = 6.8 Hz, 1H, CH), 3.09 (s, 3H, OCH₃), 3.83 (s, 1H, CHOH), 7.14-7.26 (m, 5H, H_{arom}). ¹³C-NMR δ 13.8 (q, CH₃), 15.7 (q, CH₃), 18.2 (q, CH₃), 35.9 (d, CH), 51.3 (q, OCH₃), 86.2 (s, C-2), 87.2 (d, C-3), 125.8 (d, CH_{arom}), 127.8 (d, CH_{arom}), 128.8 (d, CH_{arom}), 137.4 (s, C_{qarom}),

165.9 (s, CON), 171.3 (s, COO). HRMS ($C_{15}H_{21}NO_4$, $M = 279.15$ g/mol) Calcd: 279.1465, Found: 279.1460.

Methyl (2S*,3S*) 2-(N-acetylamino)-2-(hydroxy-phenyl-methyl)-4-methylpentanoate (20a). Following the general procedure, oxetane **19a** (0.55 g, 2 mmol) were hydrolyzed in 3h. Preparative chromatography yielded 0.34 g of the *erythro*-isomer and 0.13 g of the *threo*-isomer, both as a colorless oils. ***erythro*-diastereoisomer:** 58 %, TLC: $R_f = 0.39$ (ethylacetate/n-hexane 1: 3), 1H -NMR δ 0.81 (d, $J = 6.6$ Hz, 3H, CH_3), 0.83 (d, $J = 6.6$ Hz, 3H, CH_3), 0.90 (m, 1H, CH), 1.89 (dd, $J = 13.4, 6.7$ Hz, 1H, CH), 1.92 (s, 3H, CH_3CO), 2.72 (dd, $J = 13.4$ Hz, 4.5 Hz, 1H, CH), 3.80 (s, 3H, OCH_3), 3.82 (s, 1H, $CHOH$), 7.25-7.38 (m, 5H, H_{arom}). ^{13}C -NMR δ 1.7 (q, CH_3), 21.9 (q, CH_3), 23.9 (q, CH_3), 24.6 (q, CH_3), 40.8 (t, CH_2), 52.8 (q, OCH_3), 68.3 (s, C-2), 71.0 (d, C-3), 126.1 (d, CH_{arom}), 129.8 (d, CH_{arom}), 137.2 (s, C_{qarom}), 169.6 (s, CON), 172.9 (s, COO). MS: (EI, 70 eV) m/z (%) = 278 ($M^+ - 1$, 5), 261 ($M^+ - H_2O$, 8), 218 (10), 202 (12), 156 (15), 155 (100), 140 (38), 127 (80), 112 (45), 105 (65), 84 (27), 77 (30), 55 (8). ***threo*-diastereoisomer:** 24 %, TLC: $R_f = 0.33$ (ethylacetate/n-hexane 1: 3), 1H -NMR δ 0.86 (d, $J = 6.8$ Hz, 3H, CH_3), 1.00 (d, $J = 6.6$ Hz, 3H, CH_3), 1.33 (m, 1H, CH), 1.70 (dd, $J = 13.5, 5.6$ Hz, 1H, CH), 2.17 (s, 3H, CH_3CO), 2.80 (dd, $J = 13.5, 5.6$ Hz, 1H, CH), 3.09 (s, 3H, OCH_3), 3.77 (s, 1H, $CHOH$), 7.23-7.36 (m, 5H, H_{arom}). ^{13}C -NMR 14.2 (q, CH_3), 22.9 (q, CH_3), 24.3 (q, CH_3), 24.5 (q, CH_3), 24.9 (t, CH_2), 52.9 (q, OCH_3), 83.3 (s, C-2), 90.0 (d, C-3), 125.8 (d, CH_{arom}), 126.1 (d, CH_{arom}), 128.6 (d, CH_{arom}), 140.6 (s, C_{qarom}), 165.7 (s, CON), 171.5 (s, COO). HRMS: ($C_{16}H_{23}NO_4$, $M = 293.16$ g/mol) Calcd: 293.1627, Found: 293.1632.

***erythro*-Methyl (2S*,3S*) 2-(N-acetylamino)-3-hydroxy-2-methyl-3-naphth-2-yl-propanoate (*erythro*-12b).** Following the general procedure, oxetane **11b** (0.57 g, 2 mmol) was hydrolyzed in 7h. Preparative chromatography yielded 0.29 g of **12b** as a colorless oil. 48 %, 1H -NMR δ 1.25 (s, 3H, CH_3), 2.17 (s, 3H, CH_3CO), 3.82 (s, 3H, OCH_3), 4.22 (s, 1H, $CHOH$), 7.46-7.92 (m, 7H, H_{arom}). ^{13}C -NMR δ 13.5 (q, CH_3), 23.5 (q, CH_3), 47.9 (s, C-2), 49.3 (d, C-3), 53.0 (q, OCH_3), 126.3 (d, CH_{arom}), 127.8 (d, CH_{arom}), 128.3 (d, CH_{arom}), 129.3 (d, CH_{arom}), 133.2 (s, C_{arom}), 136.9 (s, C_{qarom}), 166.9 (s, CON), 180.0 (s, COO). MS: (EI, 70 eV) m/z (%) = 299 ($M^+ - H_2$, 5), 242 (4), 241 (50), 207 (28), 206 (30), 182 (19), 181 (100), 140 (30), 139 (48), 102 (10), 89 (5), 63 (4). HRMS: ($C_{17}H_{19}NO_4$, $M = 301.12$ g/mol) Calcd: 301.1314, Found: 301.1320.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2-methyl-5-phenylpentanoate (*erythro*-12c).** Following the general procedure, oxetane **11c** (0.52 g, 2 mmol) was hydrolyzed in 4h. Preparative chromatography yielded 0.31 g of **12c** as a colorless oil. 55 %, IR: (Film) $\tilde{\nu}$ (cm^{-1}) = 3420, 3270, 2986, 1735, 1690, 1600, 1550, 1340, 1062, 957. 1H -NMR δ 1.35 (s, 3H, CH_3), 1.94 (dd, $J = 6.3, 14.0$ Hz, 2H, CH_2), 2.03 (s, 3H, CH_3CO), 2.68-2.92 (dd, $J = 7.4, 14.0$ Hz, 2H, CH_2), 3.76 (s, 3H, OCH_3), 4.66 (dd, $J = 7.4, 6.3$ Hz, 1H, $CHOH$), 5.30 (s, 1H, NH), 7.21-7.31 (m, 5H, H_{arom}). ^{13}C -NMR δ 14.2 (q, CH_3), 19.6 (q, CH_3), 31.9 (t, CH_2), 32.9 (t, CH_2), 52.6 (q, OCH_3), 75.0 (s, C-2), 83.9 (d, C-3), 126.1 (d, C_{arom}), 128.3 (d, C_{arom}), 129.3 (d, C_{arom}), 141.0 (s, C_{qarom}), 165.2 (s, CON), 174.3 (s, COO). HRMS: ($C_9H_{17}NO_4$, $M = 203.12$ g/mol) Calcd: 203.1157, Found: 203.1161.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2-isopropyl-5-phenylpentanoate (*erythro*-18c).** Following the general procedure, oxetane **17c** (0.58 g, 2 mmol) was hydrolyzed in 5h. Preparative chromatography yielded 0.34 g of **18c** as a colorless oil. 55 %, 1H -NMR δ 0.85 (d, $J = 6.8$ Hz, 3H, CH_3), 1.23 (m, 1H, CH), 2.02 (s, 3H, CH_3CO), 2.13 (dd, $J = 7.4, 14.0$ Hz, 2H, CH_2), 2.63-2.92 (dd, $J = 6.4, 14.0$ Hz, 2H, CH_2), 3.68 (s, 3H, OCH_3), 4.32 (dd, $J = 10.4, 3.3$ Hz, 1H, $CHOH$), 7.21-7.33 (m, 5H, H_{arom}). ^{13}C -NMR δ 14.1 (q, CH_3), 17.5 (q, CH_3), 18.8 (q, CH_3), 30.7 (d, CH), 31.3 (t, CH_2), 33.5 (t, CH_2), 52.1 (q, OCH_3), 67.5 (s, C-2), 75.1 (d, C-3), 126.1 (d, CH_{arom}), 128.3 (d, CH_{arom}), 129.3 (d, CH_{arom}), 141.0 (s, C_{qarom}), 169.5 (s, CON), 171.3 (s, COO). MS: (EI, 70 eV) m/z (%) = 307 (M^+ , 5), 290 ($M^+ - OH$, 4),

289 ($M^+ - H_2O$, 10), 246 ($M^+ - CO_2Me$, 8), 231 (15), 230 (100), 214 (20), 188 (13), 141 (10), 133 (28), 130 (40), 105 (60), 91 (78), 79 (7), 70 (10), 55 (8). HRMS: ($C_{17}H_{25}NO_4$, $M = 307.18$ g/mol) Calcd: 307.1784, Found: 307.1790.

***erythro*-Methyl (2S*,3S*) 2-acetyl-amino-3-hydroxy-2-methylpentanoate (*erythro*-12d).**

Following the general procedure, oxetane **11d** (0.37 g, 2 mmol) was hydrolyzed in 3h.

Preparative chromatography yielded 0.28 g of **12d** as a colorless oil. 70 %, IR: (film) $\tilde{\nu}$ (cm^{-1}) = 3330, 3260, 2983, 1720, 1685, 1448, 1345, 1042, 987. 1H -NMR δ 1.03 (t, $J = 7.2$ Hz, 3H, CH_3), 1.55 (s, 3H, CH_3), 1.76 (dq, $J = 2.1, 7.2$ Hz, 2H, CH_2), 1.96 (s, 3H, CH_3CO), 3.71 (s, 3H, OCH_3), 4.15 (dd, $J = 11.1, 2.1$ Hz, 1H, $CHOH$), 6.22 (bs, 1H, NH). ^{13}C -NMR δ 11.8 (q, CH_3), 17.2 (q, CH_3), 23.3 (q, CH_3), 26.2 (t, CH_2), 52.9 (q, OCH_3), 63.5 (s, C-2), 69.9 (d, C-3), 169.4 (s, CON), 171.7 (s, COO). HRMS: ($C_9H_{17}NO_4$, $M = 203.12$ g/mol) Calcd: 203.1248, Found: 203.1247.

***erythro*-Methyl (2S*,3S*) 2-acetyl-amino-2-ethyl-3-hydroxypentanoate (*erythro*-14d).**

Following the general procedure, oxetane **13d** (0.4 g, 2 mmol) was hydrolyzed in 3h.

Preparative chromatography yielded 0.28 g of **14d** as a colorless oil. 65 %, 1H -NMR δ 0.75 (t, $J = 7.4$ Hz, 3H, CH_3), 1.03 (t, $J = 7.4$ Hz, 3H, CH_3), 1.50 (m, 2H, CH_2), 2.07 (s, 3H, CH_3CO), 2.13 (sextet, $J = 7.4$ Hz, 1H, CH), 3.78 (s, 3H, OCH_3), 4.23 (dd, $J = 11.6, 2.1$ Hz, 1H, $CHOH$), 6.40 (bs, 1H, NH). ^{13}C -NMR δ 9.0 (q, CH_3), 12.0 (q, CH_3), 24.5 (t, CH_2), 26.7 (t, CH_2), 53.2 (q, OCH_3), 69.9 (s, C-2), 70.2 (d, C-3), 169.5 (s, CON), 172.2 (s, COO). Anal: ($C_{10}H_{19}NO_4$, $M = 217.3$ g/mol) Calcd: C 55.28 H 8.81 N 6.46 Found: C 55.85 H 8.22 N 6.41

Methyl (2S*,3S*) 2-acetyl-amino-3-hydroxy-2-propylpentanoate (16d**).** Following the general procedure, oxetane **15d** (0.43 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.35 g of the *erythro*-isomer and 0.08 g of the *threo*-isomer, both as a colorless oils. ***erythro*-diastereoisomer:** 75 %, TLC: $R_f = 0.19$ (ethylacetate/n-hexane 1: 4), 1H -NMR δ 0.85 (t, $J = 6.9$ Hz, 3H, CH_3), 1.00 (t, $J = 7.2$ Hz, 3H, CH_3), 1.53 (m, 2H, CH_2), 1.64 (dd, $J = 7.2, 2.1$ Hz, 1H, CH), 1.76 (ddd, $J = 7.2, 4.6, 3.2$ Hz, 1H, CH), 2.00 (s, 3H, CH_3CO), 2.15 (m, 1H, CH), 2.85 (m, 1H, CH), 3.77 (s, 3H, OCH_3), 4.21 (dd, $J = 11.6, 2.1$ Hz, 1H, $CHOH$), 6.41 (bs, 1H, NH). ^{13}C -NMR δ 12.0 (q, CH_3), 13.9 (q, CH_3), 18.1 (q, CH_3), 24.5 (t, CH_2), 26.6 (t, CH_2), 33.4 (t, CH_2), 53.2 (q, OCH_3), 69.2 (s, C-2), 70.3 (d, C-3), 169.5 (s, CON), 172.3 (s, COO). HRMS: ($C_{11}H_{21}NO_4$, $M = 231.15$ g/mol) Calcd: 231.1465, Found: 231.1461. ***threo*-diastereoisomer:** 12 %, TLC: $R_f = 0.15$ (ethylacetate/n-hexane 1: 4), 1H -NMR δ 0.84 (t, $J = 7.2$ Hz, 3H, CH_3), 1.02 (t, $J = 7.4$ Hz, 3H, CH_3), 1.23 (m, 2H, CH_2), 1.48 (m, 2H, CH_2), 1.67 (m, 2H, CH_2), 1.95 (s, 3H, CH_3CO), 3.69 (s, 3H, OCH_3), 4.33 (dd, $J = 10.1, 3.5$ Hz, 1H, $CHOH$), 6.41 (bs, 1H, NH). ^{13}C -NMR δ 11.3 (q, CH_3), 14.2 (q, CH_3), 14.3 (q, CH_3), 17.9 (t, CH_2), 22.9 (t, CH_2), 36.0 (t, CH_2), 52.3 (q, OCH_3), 78.4 (s, C-2), 86.9 (d, C-3), 165.1 (s, CON), 174.5 (s, COO).

***erythro*-Methyl (2S*,3S*) 2-acetyl-amino-3-hydroxy-2-isopropylpentanoate (*erythro*-18d).**

Following the general procedure, oxetane **17d** (0.43 g, 2 mmol) was hydrolyzed in 3h.

Preparative chromatography yielded 0.31 g of **18d** as a colorless oil. 67 %, TLC: $R_f = 0.47$ (ethylacetate/n-hexane 1: 4), IR: (film) $\tilde{\nu}$ (cm^{-1}) = 3339, 3220, 2998, 1740, 1665, 1450, 1355, 1042, 980. 1H -NMR δ 0.83 (d, $J = 6.9$ Hz, 3H, CH_3), 1.00 (d, $J = 7.1$ Hz, 3H, CH_3), 1.05 (t, $J = 7.2$ Hz, 3H, CH_3), 1.56 (ddq, $J = 7.2, 7.1, 4.4$ Hz, 1H, CH), 2.01 (s, 3H, CH_3CO), 2.14 (dd, $J = 7.2, 1.6$ Hz, 1H, CH), 2.71 (sextet, $J = 6.9$ Hz, 1H, CH), 3.79 (s, 3H, OCH_3), 4.51 (dd, $J = 11.7, 1.7$ Hz, 1H, $CHOH$), 6.45 (bs, 1H, NH). ^{13}C -NMR δ 12.1 (q, CH_3), 17.2 (q, CH_3), 18.6 (q, CH_3), 25.1 (q, CH_3), 26.8 (t, CH_2), 52.9 (q, OCH_3), 69.9 (s, C-2), 73.5 (d, C-3), 169.7 (s, CON), 171.8 (s, COO). MS: (EI, 70 eV) m/z (%) = 215 ($M^+ - H_2O$, 27), 214 ($M^+ - OH$, 21), 172 (29), 164 (42), 154 (50), 130 (100), 112 (48), 98 (25), 95 (30), 70 (87), 60 (67), 57 (50). HRMS: ($C_{11}H_{21}NO_4$, $M = 231.15$ g/mol) Calcd: 231.1465, Found: 231.1463

***threo*-Methyl (2S*,3R*) 2-acetyl-amino-3-hydroxy-2-isobutylpentanoate (*threo*-20d).**

Following the general procedure, oxetane **19d** (0.45 g, 2 mmol) was hydrolyzed in 5h.

Preparative chromatography yielded 0.34 g of **20d** as a colorless oil. 70 %, TLC: R_f = 0.40 (ethylacetate/n-hexane 1: 3), IR: (Film) $\tilde{\nu}$ (cm⁻¹) = 3335, 3210, 2978, 1745, 1675, 1453, 1355, 1042, 980. ¹H-NMR δ 0.80 (d, J = 6.6 Hz, 3H, CH₃), 0.88 (d, J = 6.6 Hz, 3H, CH₃), 1.02 (t, J = 7.4 Hz, 3H, CH₃), 1.41 (dd, J = 13.4, 5.0 Hz, 1H, CH), 1.62 (m, 2H, CH₂), 1.87 (m, 1H, CH), 1.96 (s, 3H, CH₃CO), 2.03 (dd, J = 13.9, 3.5 Hz, 1H, CH), 3.64 (s, 3H, OCH₃), 4.20 (dd, J = 10.1, 3.5 Hz, 1H, CHOH). ¹³C-NMR δ 11.4 (q, CH₃), 14.3 (q, CH₃), 22.7 (q, CH₃), 23.7 (q, CH₃), 24.4 (d, CH), 24.6 (t, CH₂), 42.0 (t, CH₂), 52.3 (q, OCH₃), 77.9 (s, C-2), 87.8 (d, C-3), 164.9 (s, CON), 174.9 (s, COO). Anal: (C₁₂H₂₃NO₄, M = 245.16 g/mol) Calcd: C 58.75 H 9.45 N 5.71, Found: C 58.52 H 9.27 N 5.49.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2,4-dimethylpentanoate (*erythro*-12e).** Following the general procedure, oxetane **11e** (0.4 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.3 g of **12e** as a colorless oil. 70 %, TLC: R_f = 0.51 (ethylacetate/n-hexane 1: 3), ¹H-NMR δ 0.88 (d, J = 6.6 Hz, 3H, CH₃), 0.90 (d, J = 6.8 Hz, 3H, CH₃), 1.25 (m, 1H, CH), 1.43 (s, 3H, CH₃), 2.10 (s, 3H, CH₃CO), 3.78 (s, 3H, OCH₃), 4.43 (d, J = 9.2 Hz, 1H, CHOH). ¹³C-NMR δ 18.7 (q, CH₃), 18.9 (q, CH₃), 19.3 (q, CH₃), 23.9 (q, CH₃), 27.3 (d, CH), 52.3 (q, OCH₃), 68.1 (s, C-2), 78.1 (d, C-3), 169.1 (s, CON), 170.3 (s, COO). HRMS: (C₁₀H₁₉NO₄, M = 217.12 g/mol) Calcd: 217.1314, Found: 217.13079.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-2-ethyl-3-hydroxy-4-methylpentanoate (*erythro*-14e).** Following the general procedure, oxetane **13e** (0.43 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.33 g of **14e** as a colorless oil. 71 %, ¹H-NMR δ 0.90 (t, J = 7.4 Hz, 3H, CH₃), 0.95 (d, J = 6.9 Hz, 6H, 2CH₃), 1.31 (m, 1H, CH), 1.59 (m, 2H, CH₂), 1.99 (s, 3H, CH₃CO), 3.74 (s, 3H, OCH₃), 4.16 (d, J = 8.4 Hz, 1H, CHOH). ¹³C-NMR δ 8.9 (q, CH₃), 14.1 (q, CH₃), 19.8 (q, CH₃), 19.9 (q, CH₃), 25.6 (t, CH₂), 27.9 (d, CH), 52.4 (q, OCH₃), 78.6 (s, C-2), 90.8 (d, C-3), 165.2 (s, CON), 174.5 (s, COO). Anal: (C₁₁H₂₁NO₄, M = 231.29 g/mol) Calcd: C 57.12 H 9.12 N 6.06, Found: C 56.92 H 8.96 N 5.87.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-4-methyl-2-propylpentanoate (*erythro*-16e).** Following the general procedure, oxetane **15e** (0.45 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.36 g of **16e** as a colorless oil. 73 %, ¹H-NMR δ 0.85 (d, J = 6.6 Hz, 3H, CH₃), 0.87 (d, J = 6.8 Hz, 3H, CH₃), 1.00 (t, J = 7.5 Hz, 3H, CH₃), 1.21 (m, 1H, CH), 1.24 (m, 2H, CH₂), 1.54 (m, 2H, CH₂), 2.05 (s, 3H, CH₃CO), 3.74 (s, 3H, OCH₃), 4.34 (d, J = 8.9 Hz, 1H, CHOH). ¹³C-NMR δ 8.7 (q, CH₃), 23.7 (q, CH₃), 24.5 (q, CH₃), 24.7 (q, CH₃), 27.1 (t, CH₂), 28.1 (d, CH), 39.2 (t, CH₂), 52.0 (q, OCH₃), 68.2 (s, C-2), 74.1 (d, C-3), 169.1 (s, CON), 172.1 (s, COO). HRMS: (C₁₂H₂₃NO₄, M = 245.16 g/mol) Calcd: 245.1627, Found: 245.1631.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2-isopropyl-4-methylpentanoate (*erythro*-18e)** Following the general procedure, oxetane **17e** (0.45 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.29 g of **18e** as a colorless oil. 59 %, ¹H-NMR δ 0.85 (d, J = 6.8 Hz, 6H, 2CH₃), 0.89 (d, J = 6.5 Hz, 3H, CH₃), 0.93 (d, J = 6.6 Hz, 3H, CH₃), 1.03 (m, 2H, 2CH), 2.04 (s, 3H, CH₃CO), 3.79 (s, 3H, OCH₃), 4.44 (d, J = 8.4 Hz, 1H, CHOH). ¹³C-NMR δ 17.0 (q, CH₃), 18.7 (q, CH₃), 19.2 (q, CH₃), 19.7 (q, CH₃), 20.3 (q, CH₃), 23.8 (d, CH), 27.2 (d, CH), 52.7 (q, OCH₃), 66.3 (s, C-2), 73.5 (d, C-3), 170.1 (s, CON), 171.8 (s, COO). HRMS: (C₁₂H₂₃NO₄, M = 245.16 g/mol) Calcd: 245.1627, Found: 245.1618.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2-isobutyl-4-methylpentanoate (*erythro*-20e)** Following the general procedure, oxetane **19e** (0.48 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.34 g of **20e** as a colorless oil. 65 %, ¹H-NMR δ 0.65 (dd, J = 6.6, 6.6 Hz, 6H, 2CH₃), 0.75 (dd, J = 7.5, 7.5 Hz, 6H, 2CH₃), 1.30 (m, 1H, CH), 1.65 (dd, J = 1.6, 3.4 Hz, 1H, CH), 1.94 (s, 3H, CH₃CO), 2.42 (dd, J = 7.5, 1.5 Hz, 1H, CH), 3.50 (m, 1H, CH), 3.72 (s, 3H, OCH₃), 3.92 (dd, J = 4.5, 1.6 Hz, 1H, CHOH), 6.97 (bs, 1H, NH). ¹³C-NMR δ 19.8 (q, CH₃), 23.5 (q, CH₃), 23.6 (q, CH₃), 23.7 (q, CH₃), 24.4 (d, CH),

24.6 (d, CH), 41.1 (t, CH₂), 52.8 (q, OCH₃), 69.0 (s, C-2), 78.0 (d, C-3), 171.3 (s, CON), 174.5 (s, COO). HRMS: (C₁₃H₂₅NO₄, M = 259.18 g/mol) Calcd: 259.1777, Found: 259.1773. ***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2,5-dimethylhexanoate (*erythro*-12f).** Following the general procedure, oxetane **13f** (0.43 g, 2 mmol) was hydrolyzed in 3h.

Preparative chromatography yielded 0.25 g of **12f** as a colorless oil. 55 %, ¹H-NMR δ 0.83 (d, J = 6.5 Hz, 3H, CH₃), 0.91 (d, J = 6.5 Hz, 3H, CH₃), 0.94 (m, 1H, CH), 1.24 (dd, J = 1.9, 1.8 Hz, 1H, CH), 1.54 (s, 3H, CH₃), 1.87 (m, 1H, CH), 1.96 (s, 3H, CH₃CO), 3.71 (s, 3H, OCH₃), 4.25 (dd, J = 11.4, 1.8 Hz, 1H, CHOH), 6.20 (bs, 1H, NH). ¹³C-NMR δ 17.1 (q, CH₃), 20.4 (q, CH₃), 23.3 (q, CH₃), 23.5 (q, CH₃), 25.1 (d, CH), 41.6 (t, CH₂), 52.8 (q, OCH₃), 63.4 (s, C-2), 65.9 (d, C-3), 169.2 (s, CON), 171.5 (s, COO). HRMS: (C₁₁H₂₁NO₄, M = 231.15 g/mol) Calcd: 231.1470, Found: 231.1466.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-2-ethyl-3-hydroxy-5-methylhexanoate (*erythro*-14f).** Following the general procedure, the bicyclic oxetane **13f** (0.45 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.32 g of **14f** as a colorless oil. 66 %, ¹H-NMR δ 0.75 (t, J = 7.5 Hz, 3H, CH₃), 0.84 (d, J = 6.6 Hz, 3H, CH₃), 0.92 (d, J = 6.8 Hz, 3H, CH₃), 1.23 (m, 1H, CH), 1.63 (ddd, J = 10.1, 2.1, 2.1 Hz, 2H, CH₂), 1.85 (m, 1H, CH), 2.12 (s, 3H, CH₃CO), 2.80 (sextet, J = 7.5 Hz, 1H, CH), 3.76 (s, 3H, OCH₃), 4.40 (dd, J = 11.5, 2.1 Hz, 1H, CHOH), 6.39 (bs, 1H, NH). ¹³C-NMR δ 9.0 (q, CH₃), 20.5 (q, CH₃), 23.7 (q, CH₃), 24.4 (q, CH₃), 24.5 (t, CH₂), 25.1 (d, CH), 42.0 (t, CH₂), 53.1 (q, OCH₃), 65.7 (s, C-2), 69.7 (d, C-3), 169.3 (s, CON), 172.2 (s, COO). HRMS: (C₁₂H₂₃NO₄, M = 245.16 g/mol) Calcd: 245.1621, Found: 245.1618.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-5-methyl-2-propylhexanoate (*erythro*-16f).** Following the general procedure, oxetane **15f** (0.48 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.35 g of **16f** as a colorless oil. 68 %, ¹H-NMR δ 0.83 (d, J = 6.5 Hz, 3H, CH₃), 0.86 (t, J = 4.3 Hz, 3H, CH₃), 0.91 (d, J = 6.6 Hz, 3H, CH₃), 1.52 (dddd, J = 3.1, 3.2, 3.1, 3.2 Hz, 2H, CH₂), 1.65 (ddd, J = 6.5, 2.1, 1.9 Hz, 1H, CH), 1.81 (m, 2H, CH₂), 2.00 (s, 3H, CH₃CO), 3.77 (s, 3H, OCH₃), 4.39 (dd, J = 11.5, 2.1 Hz, 1H, CHOH), 6.39 (bs, 1H, NH). ¹³C-NMR δ 13.9 (q, CH₃), 18.1 (q, CH₃), 20.5 (q, CH₃), 23.7 (q, CH₃), 25.1 (t, CH₂), 33.4 (t, CH₂), 41.9 (t, CH₂), 53.1 (q, OCH₃), 65.9 (s, C-2), 69.0 (d, C-3), 169.3 (s, CON), 172.3 (s, COO). Anal: (C₁₃H₂₅NO₄, M = 259.18 g/mol) Calcd: C 60.21 H 9.72 N 5.40, Found: C 59.98 H 9.61 N 5.32.

***erythro*-Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2-isopropyl-5-methylhexanoate (*erythro*-18f).** Following the general procedure, oxetane **17f** (0.48 g, 2 mmol) was hydrolyzed in 3h. Preparative chromatography yielded 0.32 g of **18f** as a colorless oil. 62 %, IR: (film) $\tilde{\nu}$ (cm⁻¹) = 3396, 2919, 1738, 1668, 1471, 1056, 1031, 978, 680. ¹H-NMR δ 0.83 (d, J = 6.8 Hz, 3H, CH₃), 0.89 (d, J = 6.5 Hz, 3H, CH₃), 0.93 (d, J = 6.6 Hz, 3H, CH₃), 1.00 (d, J = 7.1 Hz, 3H, CH₃), 1.70 (m, 2H, CH₂), 1.86 (m, 1H, CH), 2.04 (s, 3H, CH₃CO), 3.57 (sextet, J = 6.9 Hz, 1H, CH), 3.79 (s, 3H, OCH₃), 4.69 (dd, J = 11.2, 2.4 Hz, 1H, CHOH), 6.43 (bs, 1H, NH). ¹³C-NMR δ 17.0 (q, CH₃), 18.7 (q, CH₃), 20.5 (q, CH₃), 23.8 (q, CH₃), 25.1 (d, CH), 28.1 (d, CH), 41.6 (t, CH₂), 52.9 (q, OCH₃), 65.9 (s, C-2), 73.5 (d, C-3), 169.6 (s, CON), 171.8 (s, COO). Anal: (C₁₃H₂₅NO₄, M = 259.3 g/mol) Calcd: C 60.21 H 9.72 N 5.40, Found: C 59.89 H 9.64 N 5.27.

Methyl (2S*,3S*) 2-acetylamino-3-hydroxy-2-isobutyl-5-methylhexanoate (20f).

Following the general procedure, oxetane **19f** (0.51 g, 2 mmol) was hydrolyzed in 3h.

Preparative chromatography yielded 0.26 g of the *erythro*-isomer and 0.09 g of the *threo*-isomer, both as a colorless oil. ***erythro*-diastereoisomer:** 45 %, TLC: R_f = 0.33

(ethylacetate/n-hexane 1: 3), ¹H-NMR δ 0.74 (d, J = 6.5 Hz, 3H, CH₃), 0.81 (d, J = 6.6 Hz, 3H, CH₃), 0.86 (d, J = 6.6 Hz, 3H, CH₃), 0.95 (d, J = 6.8 Hz, 3H, CH₃), 1.51 (m, 2H, CH₂), 1.57 (m, 2H, CH₂), 1.94 (s, 3H, CH₃CO), 2.21 (m, 1H, CH), 2.26 (dd, J = 13.5, 3.5 Hz, 1H, CH), 2.81 (dd, J = 13.5, 1.8 Hz, 1H, CH), 3.78 (s, 3H, OCH₃), 4.43 (dd, J = 11.6, 1.8 Hz, 1H,

CHOH), 6.50 (bs, 1H, NH). ^{13}C -NMR δ 14.4 (q, CH_3), 20.5 (q, CH_3), 21.5 (q, CH_3), 23.2 (q, CH_3), 25.1 (d, CH), 28.1 (d, CH), 37.9 (t, CH_2), 52.8 (q, OCH_3), 68.0 (s, C-2), 73.9 (d, C-3), 169.3 (s, CON), 172.9 (s, COO). MS: (EI, 70 eV) m/z (%) = 271 ($\text{M}^+ - \text{H}_2$, 5), 256 ($\text{M}^+ - \text{H}_2\text{O}$, 7), 214 ($\text{M}^+ - \text{CO}_2\text{Me}$, 12), 196 (22), 187 (60), 170 (20), 154 (30), 112 (26), 102 (100), 85 (70), 57 (52). ***threo*-diastereoisomer:** 13 %, TLC: R_f = 0.23 (ethylacetate/n-hexane 1: 3), ^1H -NMR δ 0.81 (d, J = 6.8 Hz, 3H, CH_3), 0.88 (d, J = 6.6 Hz, 3H, CH_3), 0.93 (dd, J = 7.5, 7.4 Hz, 6H, 2CH_3), 1.00 (dd, J = 6.9 Hz, 1H, CH), 1.12 (dd, J = 7.0, 1.8 Hz, 1H, CH), 1.53 (m, 1H, CH), 2.00 (s, 3H, CH_3CO), 2.43 (dd, J = 7.5, 1.5 Hz, 1H, CH), 3.67 (s, 3H, OCH_3), 4.00 (dd, J = 8.5, 1.4 Hz, 1H, CHOH), 6.97 (bs, 1H, NH). ^{13}C -NMR δ 14.0 (q, CH_3), 20.0 (q, CH_3), 23.3 (q, CH_3), 23.8 (q, CH_3), 24.7 (d, CH), 27.9 (d, CH), 40.6 (t, CH_2), 52.3 (q, OCH_3), 80.0 (s, C-2), 91.5 (d, C-3), 164.9 (s, CON), 174.9 (s, COO). Anal: ($\text{C}_{14}\text{H}_{27}\text{NO}_4$, M = 273.2 g/mol) Calcd: C 61.51 H 9.96 N 5.12, Found: C 61.62 H 9.86 N 4.97.

7) ^{13}C -NMR-spectra of the bicyclic oxetanes. **8c**, **8d**, **8e**, **11b**, **11e**, **13a**, **13f**, **15d**, **17a**, **17d**, **19a**, and **19e**. These compounds were isolated as mixtures with unreacted oxazole and used for hydrolytic ring-opening without purification.

exo-5-Methoxy-3-methyl-7-phenethyl-4,6-dioxa-2-aza-bicyclo[3.2.0]hept-2-ene (***exo*-8c**):

exo-7-Ethyl-5-methoxy-3-methyl-4,6-dioxa-2-aza-bicyclo[3.2.0]hept-2-ene (***exo*-8d**):

exo-7-Isopropyl-5-methoxy-3-methyl-4,6-dioxa-2-aza-bicyclo[3.2.0]hept-2-ene (***exo*-8e**):

exo-5-Methoxy-1,3-dimethyl-7-naphth-2-yl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-11b):

exo-7-Isopropyl-5-methoxy-1,3-dimethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-11e):

exo-1-Ethyl-5-methoxy-3-methyl-7-phenyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (***exo*-13a**):

exo-1,7-Diethyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (***exo*-13d**):

exo-1-Ethyl-7-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-13e):

exo-1-Ethyl-7-isobutyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-13f):

exo-5-Methoxy-3-methyl-7-phenyl-1-propyl-4,6-dioxa-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-15a):

exo-7-Ethyl-5-methoxy-3-methyl-1-propyl-4,6-dioxa-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-15d):

1-Isopropyl-5-methoxy-3-methyl-7-phenyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (**17a**):

exo-1-Isopropyl-5-methoxy-3-methyl-7-phenylethyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-**17c**):

exo-7-Ethyl-1-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-17d):

exo-1,7-Diisopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-17e):

exo-7-Isobutyl-1-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene
(*exo*-17f):

exo-1-Isobutyl-5-methoxy-3-methyl-7-phenyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene
(*exo*-19a):

exo-1-Isobutyl-7-isopropyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-19e):

exo-1-*sec*-Butyl-7-ethyl-5-methoxy-3-methyl-4,6-dioxo-2-aza-bicyclo[3.2.0]hept-2-ene (*exo*-22) 1:1 mixture of 2 diastereoisomers:

9) X-ray crystallographic data for compound **23**:

data_SBO341

```

_audit_creation_method          SHELXL-97
_chemical_formula_sum           'C12 H23 N O4'
_chemical_formula_weight        245.31

loop_
  _atom_type_symbol
  _atom_type_description
  _atom_type_scatter_dispersion_real
  _atom_type_scatter_dispersion_imag
  _atom_type_scatter_source
  'C' 'C' 0.0033 0.0016
  'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
  'H' 'H' 0.0000 0.0000
  'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
  'N' 'N' 0.0061 0.0033
  'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
  'O' 'O' 0.0106 0.0060
  'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'

_symmetry_cell_setting          orthorhombic
_symmetry_space_group_name_H-M P212121

loop_
  _symmetry_equiv_pos_as_xyz
  'x, y, z'
  '-x+1/2, -y, z+1/2'
  'x+1/2, -y+1/2, -z'
  '-x, y+1/2, -z+1/2'

_cell_length_a                  6.434(1)
_cell_length_b                  14.945(1)
_cell_length_c                  15.721(1)
_cell_angle_alpha               90.00
_cell_angle_beta                90.00
_cell_angle_gamma               90.00
_cell_volume                    1511.7(3)
_cell_formula_units_Z           4
_cell_measurement_temperature   293(2)
_cell_measurement_reflns_used   3219
_cell_measurement_theta_min     1.88
_cell_measurement_theta_max     26.90

_exptl_crystal_description      thin needle
_exptl_crystal_colour           colourless
_exptl_crystal_size_max         0.15
_exptl_crystal_size_mid         0.08
_exptl_crystal_size_min         0.05
_exptl_crystal_density_meas     ?
_exptl_crystal_density_diffn    1.078
_exptl_crystal_density_method   'not measured'
_exptl_crystal_F_000            536
_exptl_absorpt_coefficient_mu   0.080
_exptl_absorpt_correction_type   'none'
_exptl_absorpt_correction_T_min ?
_exptl_absorpt_correction_T_max ?

```

```

_exptl_absorpt_process_details      ?

_exptl_special_details
;
?
;

_diffn_ambient_temperature          293(2)
_diffn_radiation_wavelength          0.71073
_diffn_radiation_type                MoK\alpha
_diffn_radiation_source              'fine-focus sealed tube'
_diffn_radiation_monochromator        graphite
_diffn_measurement_device_type        'Nonius KappaCCD'
_diffn_measurement_method            'Phi/Omega-Scans'
_diffn_detector_area_resol_mean      ?
_diffn_standards_number              ?
_diffn_standards_interval_count      ?
_diffn_standards_interval_time       ?
_diffn_standards_decay_%             ?
_diffn_reflns_number                 3219
_diffn_reflns_av_R_equivalents        0.0000
_diffn_reflns_av_sigmaI/netI         0.0669
_diffn_reflns_limit_h_min            -8
_diffn_reflns_limit_h_max            8
_diffn_reflns_limit_k_min            -18
_diffn_reflns_limit_k_max            18
_diffn_reflns_limit_l_min            -19
_diffn_reflns_limit_l_max            19
_diffn_reflns_theta_min              1.88
_diffn_reflns_theta_max              26.90
_reflns_number_total                 3219
_reflns_number_gt                    1762
_reflns_threshold_expression          >2sigma(I)

_computing_data_collection            'Nonius COLLECT'
_computing_cell_refinement            'Nonius DENZO'
_computing_data_reduction             'Nonius DENZO'
_computing_structure_solution         'SHELXS-97 (Sheldrick, 1997)'
_computing_structure_refinement       'SHELXL-97 (Sheldrick, 1997)'
_computing_molecular_graphics         ?
_computing_publication_material       ?

_refine_special_details
;
Refinement of F2 against ALL reflections. The weighted R-factor wR and
goodness of fit S are based on F2, conventional R-factors R are based
on F, with F set to zero for negative F2. The threshold expression of
F2 > 2sigma(F2) is used only for calculating R-factors(gt) etc. and is
not relevant to the choice of reflections for refinement. R-factors based
on F2 are statistically about twice as large as those based on F, and R-
factors based on ALL data will be even larger.
;

_refine_ls_structure_factor_coef      Fsqd
_refine_ls_matrix_type                full
_refine_ls_weighting_scheme           calc
_refine_ls_weighting_details          'calc w=1/[s2(Fo2)+(0.0460P)2+0.2800P] where P=(Fo2+2Fc2)/3'
_atom_sites_solution_primary          direct
_atom_sites_solution_secondary        difmap
_atom_sites_solution_hydrogens        geom
_refine_ls_hydrogen_treatment         refall

```



```

_refine_ls_extinction_method      none
_refine_ls_extinction_coef        ?
_refine_ls_abs_structure_details
'Flack H D (1983), Acta Cryst. A39, 876-881'
_refine_ls_abs_structure_Flack    0.9(18)
_refine_ls_number_reflns          3219
_refine_ls_number_parameters      247
_refine_ls_number_restraints      0
_refine_ls_R_factor_all           0.1396
_refine_ls_R_factor_gt            0.0605
_refine_ls_wR_factor_ref          0.1308
_refine_ls_wR_factor_gt          0.1063
_refine_ls_goodness_of_fit_ref    1.045
_refine_ls_restrained_S_all       1.045
_refine_ls_shift/su_max           0.020
_refine_ls_shift/su_mean          0.002

```

```

loop_

```

```

_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_U_iso_or_equiv
_atom_site_adp_type
_atom_site_occupancy
_atom_site_symmetry_multiplicity
_atom_site_calc_flag
_atom_site_refinement_flags
_atom_site_disorder_assembly
_atom_site_disorder_group
C1 C 0.9309(6) 0.6966(2) 0.4679(3) 0.0638(9) Uani 1 1 d . . .
H1A H 1.073(6) 0.675(2) 0.447(2) 0.095(12) Uiso 1 1 d . . .
H1B H 0.819(7) 0.708(2) 0.424(3) 0.110(15) Uiso 1 1 d . . .
H1C H 0.954(5) 0.752(3) 0.498(2) 0.089(12) Uiso 1 1 d . . .
C2 C 0.9495(5) 0.5972(2) 0.58821(19) 0.0534(7) Uani 1 1 d . . .
C3 C 0.8300(4) 0.52612(19) 0.64233(19) 0.0510(7) Uani 1 1 d . . .
C4 C 0.7866(4) 0.44039(19) 0.5837(2) 0.0581(8) Uani 1 1 d . . .
H4 H 0.730(4) 0.3957(17) 0.6275(16) 0.047(7) Uiso 1 1 d . . .
C5 C 0.9932(7) 0.4049(3) 0.5440(4) 0.0918(12) Uani 1 1 d . . .
H5A H 1.101(8) 0.387(3) 0.593(3) 0.140(18) Uiso 1 1 d . . .
H5B H 0.966(8) 0.344(3) 0.507(3) 0.158(19) Uiso 1 1 d . . .
H5C H 1.034(6) 0.446(3) 0.500(3) 0.101(15) Uiso 1 1 d . . .
C6 C 0.6147(6) 0.4528(2) 0.5143(3) 0.0698(10) Uani 1 1 d . . .
H6A H 0.674(4) 0.484(2) 0.466(2) 0.067(10) Uiso 1 1 d . . .
H6B H 0.486(5) 0.486(2) 0.543(2) 0.079(10) Uiso 1 1 d . . .
C7 C 0.5369(8) 0.3620(3) 0.4784(3) 0.0858(12) Uani 1 1 d . . .
H7A H 0.433(6) 0.373(2) 0.432(3) 0.101(14) Uiso 1 1 d . . .
H7B H 0.656(7) 0.326(3) 0.450(3) 0.128(16) Uiso 1 1 d . . .
H7C H 0.477(6) 0.324(3) 0.522(3) 0.100(13) Uiso 1 1 d . . .
C8 C 0.9412(5) 0.4409(2) 0.7764(2) 0.0632(9) Uani 1 1 d . . .
C9 C 1.1284(7) 0.4126(4) 0.8308(3) 0.0836(13) Uani 1 1 d . . .
H9A H 1.100(8) 0.393(3) 0.885(4) 0.15(2) Uiso 1 1 d . . .
H9B H 1.213(8) 0.382(3) 0.796(3) 0.14(2) Uiso 1 1 d . . .
H9C H 1.210(9) 0.464(4) 0.844(4) 0.18(3) Uiso 1 1 d . . .
C10 C 0.6235(5) 0.5709(2) 0.6781(2) 0.0561(8) Uani 1 1 d . . .
H10 H 0.555(4) 0.6023(17) 0.6261(16) 0.042(6) Uiso 1 1 d . . .
C11 C 0.6702(6) 0.6421(3) 0.7480(3) 0.0753(10) Uani 1 1 d . . .
H11A H 0.798(6) 0.681(3) 0.729(2) 0.109(14) Uiso 1 1 d . . .
H11B H 0.731(5) 0.613(2) 0.797(2) 0.080(11) Uiso 1 1 d . . .
C12 C 0.4762(10) 0.6946(4) 0.7760(4) 0.1011(15) Uani 1 1 d . . .
H12A H 0.518(7) 0.735(3) 0.819(3) 0.115(15) Uiso 1 1 d . . .

```

```

H12B H 0.405(7) 0.720(3) 0.728(3) 0.14(2) Uiso 1 1 d . . .
H12C H 0.359(9) 0.663(4) 0.796(4) 0.17(3) Uiso 1 1 d . . .
N1 N 0.9799(4) 0.49968(16) 0.71189(16) 0.0564(7) Uani 1 1 d . . .
H1N H 1.127(6) 0.511(2) 0.702(2) 0.091(11) Uiso 1 1 d . . .
O1 O 0.8292(3) 0.63516(13) 0.52804(13) 0.0586(6) Uani 1 1 d . . .
O2 O 1.1348(4) 0.61505(16) 0.59765(16) 0.0848(8) Uani 1 1 d . . .
O3 O 0.7611(4) 0.41201(17) 0.79469(16) 0.0816(7) Uani 1 1 d . . .
O4 O 0.4742(3) 0.50531(16) 0.70829(16) 0.0710(7) Uani 1 1 d . . .
H4O H 0.538(8) 0.464(3) 0.747(3) 0.143(19) Uiso 1 1 d . . .

```

```

loop_
  _atom_site_aniso_label
  _atom_site_aniso_U_11
  _atom_site_aniso_U_22
  _atom_site_aniso_U_33
  _atom_site_aniso_U_23
  _atom_site_aniso_U_13
  _atom_site_aniso_U_12
C1 0.064(2) 0.062(2) 0.065(2) 0.017(2) 0.008(2) -0.0014(18)
C2 0.0491(19) 0.0519(16) 0.0592(18) 0.0053(15) -0.0033(16) 0.0046(15)
C3 0.0448(17) 0.0490(16) 0.0593(18) 0.0101(15) -0.0007(15) 0.0008(13)
C4 0.0546(19) 0.0510(17) 0.069(2) 0.0031(17) 0.0078(16) -0.0010(15)
C5 0.081(3) 0.084(3) 0.109(4) -0.020(3) 0.009(3) 0.017(2)
C6 0.071(2) 0.069(2) 0.070(2) -0.005(2) -0.0025(19) -0.0080(19)
C7 0.088(3) 0.081(3) 0.089(3) -0.018(3) 0.001(3) -0.015(3)
C8 0.058(2) 0.0645(19) 0.067(2) 0.0135(18) 0.0029(17) 0.0032(17)
C9 0.068(3) 0.107(4) 0.076(3) 0.036(3) 0.002(2) 0.020(3)
C10 0.0469(18) 0.063(2) 0.058(2) 0.0084(17) 0.0048(15) 0.0020(16)
C11 0.074(3) 0.076(2) 0.076(3) -0.008(2) -0.002(2) 0.012(2)
C12 0.108(4) 0.117(4) 0.079(3) -0.018(3) 0.005(3) 0.034(3)
N1 0.0477(15) 0.0605(16) 0.0611(16) 0.0188(13) 0.0019(13) 0.0028(12)
O1 0.0503(12) 0.0621(12) 0.0634(12) 0.0208(11) -0.0044(11) -0.0018(10)
O2 0.0480(13) 0.1030(19) 0.1036(19) 0.0486(15) -0.0160(13) -0.0167(13)
O3 0.0590(14) 0.0929(17) 0.0929(17) 0.0397(14) 0.0055(13) -0.0073(13)
O4 0.0492(13) 0.0794(16) 0.0844(16) 0.0202(14) 0.0098(13) -0.0019(12)

```

```
_geom_special_details
```

```
;
```

All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic)

treatment of cell esds is used for estimating esds involving l.s. planes.

```
;
```

```

loop_
  _geom_bond_atom_site_label_1
  _geom_bond_atom_site_label_2
  _geom_bond_distance
  _geom_bond_site_symmetry_2
  _geom_bond_publ_flag
C1 O1 1.471(4) . ?
C2 O2 1.231(3) . ?
C2 O1 1.348(3) . ?
C2 C3 1.562(4) . ?
C3 N1 1.510(4) . ?
C3 C10 1.590(4) . ?
C3 C4 1.603(4) . ?
C4 C5 1.561(5) . ?
C4 C6 1.564(5) . ?

```

C6 C7 1.552(5) . ?
 C8 O3 1.269(4) . ?
 C8 N1 1.365(4) . ?
 C8 C9 1.536(5) . ?
 C10 O4 1.452(3) . ?
 C10 C11 1.560(5) . ?
 C11 C12 1.538(6) . ?

loop_

_geom_angle_atom_site_label_1
 _geom_angle_atom_site_label_2
 _geom_angle_atom_site_label_3
 _geom_angle
 _geom_angle_site_symmetry_1
 _geom_angle_site_symmetry_3
 _geom_angle_publ_flag
 O2 C2 O1 123.3(3) . . ?
 O2 C2 C3 123.9(3) . . ?
 O1 C2 C3 112.7(2) . . ?
 N1 C3 C2 105.0(2) . . ?
 N1 C3 C10 112.8(2) . . ?
 C2 C3 C10 108.5(2) . . ?
 N1 C3 C4 108.6(2) . . ?
 C2 C3 C4 108.4(2) . . ?
 C10 C3 C4 113.2(2) . . ?
 C5 C4 C6 111.3(3) . . ?
 C5 C4 C3 110.7(3) . . ?
 C6 C4 C3 115.4(3) . . ?
 C7 C6 C4 112.2(3) . . ?
 O3 C8 N1 123.6(3) . . ?
 O3 C8 C9 119.7(3) . . ?
 N1 C8 C9 116.6(3) . . ?
 O4 C10 C11 110.9(3) . . ?
 O4 C10 C3 112.6(2) . . ?
 C11 C10 C3 112.1(2) . . ?
 C12 C11 C10 113.2(4) . . ?
 C8 N1 C3 126.2(3) . . ?
 C2 O1 C1 117.3(3) . . ?

_diffn_measured_fraction_theta_max	0.993
_diffn_refl_theta_full	26.90
_diffn_measured_fraction_theta_full	0.993
_refine_diff_density_max	0.116
_refine_diff_density_min	-0.134
_refine_diff_density_rms	0.035