A SYNTHETIC APPROACH TO SUBSTITUTED CYCLOPROPANES
BASED ON THE COUPLING REACTION OF LITHIATED
CHLOROALKYLOXAZOLINES WITH FISCHER CARBENE
COMPLEXES

*Saverio Florio,^{*a} Filippo M. Perna,^a Renzo Luisi,^a José Barluenga,^{*b} Félix Rodríguez,^b and
Francisco J. Fañanás,^b*

^aDipartimento Farmaco-Chimico, University of Bari, Via E. Orabona 4, I-70126-Bari, Italy

C.N.R., Istituto di Chimica dei Composti Organometallici “ICCOM”, Sezione di Bari,

^b Instituto Universitario de Química Organometálica “Enrique Moles”, Unidad Asociada al C.S.I.C.
Julián Clavería, 8, Universidad de Oviedo, 33006 Oviedo, Spain

Experimental Section

General. Tetrahydrofuran (THF) and diethyl ether (Et₂O) were freshly distilled under a nitrogen atmosphere over sodium/benzophenone ketyl. All other chemicals were of commercial grade and used without further purification. Commercial solutions of *n*-BuLi (2.5 M solution in hexanes) were titrated by using *N*-pivaloyl-*o*-toluidine prior to use.¹ For the ¹H and ¹³C NMR spectra (300, 500 MHz; ¹³C NMR 75.4, 125 MHz), CDCl₃ was used as the solvent. Carbon multiplicities were assigned by DEPT techniques. GC-MS spectrometry analyses were performed on a gas chromatograph HP 5890 II (dimethylsilicon capillary column, 30 m, 0.25 mm i.d.) equipped with a mass selective detector operating at 70 eV (EI). Elemental analyses were carried out on a Perkin-Elmer 2400 and Carlo Erba 1108 microanalyzers. Melting points were uncorrected. TLC was performed on Merck silica gel plates with F-254 indicator; visualization was accomplished by UV light (254 nm) or by exposing to I₂ vapor. All reactions involving air-sensitive reagents were performed under nitrogen in oven-dried glassware using syringe-septum cap technique. The compounds **3a-e** were prepared according to the reported procedure.²

Preparation of carbene complexes **4** and cyclopropanecarboxylates **5**, **6**, and **7**. General Procedure.

To a solution of LDA, generated by reaction of *N,N*-diisopropylamine (1.2 mmol) and BuLi (1.2 mmol, 0.48 mL of a 2.5N solution in hexanes) in THF (5 mL) was added a solution of the corresponding oxazoline **1** (1.0 mmol) in THF (2 mL) at -98° C under a nitrogen atmosphere. The mixture was stirred for 10 min and then a solution of the corresponding carbene complex **3** in THF (5 mL) was added dropwise. The reaction mixture was kept at -98 °C for other 30 min, then allowed to warm to room temperature and, once checked the consumption of **3a**, filtered through

¹ Suffert, J. *J. Org. Chem.* **1989**, *54*, 509.

² Aumann, R.; Heinen, H. *Chem. Ber.* **1987**, *120*, 537

celite and silica gel. The solvent was removed under reduced pressure and the crude residue was purified by flash column chromatography (ethyl acetate/hexane: 1/9) to give the cyclopropylcarbene complexes **4**.

In order to obtain cyclopropanecarboxylate derivatives **5**, **6**, and **7**, the solution of the corresponding cyclopropylcarbene complex in hexane (30 mL) was subjected to the air oxidation under sun light. After 16 h the mixture was filtered off on celite, the solvent removed and the residue purified by column chromatography (AcOEt 1/ petroleum ether 9) to give the cyclopropane carboxylates **5**, **6**, and **7**. Alternatively, the oxidation can be carried out with pyridine N-oxide in THF.

Pentacarbonyl-[(1*R,2*R**,3*S**)-2-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-2-methyl-3-phenylcyclopropyl]methoxymethylene}tungsten (**4d**):** yellow solid, mp 43-44 °C, 80 %. ¹H NMR (300 MHz, C₆D₆) δ 1.01(s, 3 H), 1.11 (s, 3 H), 1.44 (s, 3 H), 3.20 (d, *J* = 6.2 Hz, 1 H), 3.39 and 3.49 (2 × d, AB system, *J* = 8.1 Hz, 2 H), 3.92 (s, 3 H), 5.23 (d, *J* = 6.2 Hz, 1 H), 7.04–7.24 (m, 5 H); ¹³C NMR (75.4MHz, C₆D₆) δ 16.5 (CH₃), 28.4 (CH₃), 28.5 (CH₃), 40.8 (CH), 41.2 (*C_q*), 60.1 (CH₃), 68.1 (CH), 69.7 (*C_q*), 79.9 (CH₂), 127.4 (CH), 128.1 (2 × CH), 128.3 (2 × CH), 138.1 (*C_q*), 163.7 (C=N), 198.4 (4 × C=O), 204.8 (C=O), 327.5 (C=W); GC-MS (70 eV) *m/z* (%) 271 (M⁺ – WC₅O₅, 4), 228 (100), 200 (6), 157 (11), 77(21); FT-IR (film, cm⁻¹): 2967, 2066, 1931, 1662 (s, C=N), 752, 699. Anal. calcd for C₂₂H₂₁NO₇W: C, 44.39; H, 3.56; N, 2.35. Found: C, 44.73; H, 3.62; N, 2.30

Pentacarbonyl-[(1*R,2*R**,3*S**)-2-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-3-(4-methoxyphenyl)-2-methylcyclopropyl]methoxymethylene}tungsten (**4e**):** yellow solid, mp 48-49°C, 85 %. ¹H NMR (300 MHz, C₆D₆) δ 0.92 (s, 3 H), 1.09 (s, 3 H), 1.48 (s, 3 H), 3.22 (d, *J* = 6.8 Hz, 1 H), 3.33 and 3.49 (2 × d, AB system, *J* = 8.1 Hz, 2 H), 3.36 (s, 3 H), 3.92 (s, 3 H), 5.21 (d, *J* = 6.8 Hz, 1 H), 6.81 (d, *J* = 8.9 Hz, 2 H), 7.25 (d, *J* = 8.9 Hz, 2 H); ¹³C NMR (75.4MHz, C₆D₆) δ 16.6 (CH₃), 28.7 (CH₃), 28.9 (CH₃), 40.9 (CH), 41.2 (*C_q*), 55.4 (CH₃), 60.0 (CH₃), 68.1 (CH), 69.7 (*C_q*), 80.1 (CH₂), 114.4 (2 × CH), 128.8 (2 × CH), 131.0 (*C_q*), 159.9 (*C_q*), 163.8 (C=N), 198.4 (4 × C=O), 204.8

(C=O), 327.1(C=W); GC-MS (70 eV) m/z (%), 301 ($M^+ - WC_5O_5$, 6), 258 (100), 230 (8), 187 (9), 159 (9), 127 (8), 115 (5); FT-IR (film, cm^{-1}): 2967, 2066, 1931, 1660 (s, C=N), 1230, 1108. Anal. calcd. for $C_{23}H_{23}NO_8W$: C, 44.18; H, 3.71; N, 2.24. Found: C, 44.02; H, 3.51; N, 2.02.

Methyl-(1*R,2*R**,3*S**)-2-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-2-methyl-3-**

phenylcyclopropanecarboxylate (5a): white solid, mp 53-54 °C, 72 %. 1H NMR (300 MHz) δ 0.81 (s, 3 H), 1.06 (s, 3 H), 1.62 (s, 3 H), 2.86 (d, $J = 6.6$ Hz, 1 H), 3.10 (d, $J = 6.6$ Hz, 1 H), 3.46 and 3.55 (2 $\times$ d, AB system, $J = 7.9$ Hz, 2 H), 3.74 (s, 3 H), 7.22–7.30 (m, 5 H); ^{13}C NMR (75.4 MHz) δ 15.6 (CH_3), 28.1 (CH_3), 28.2 (CH_3), 29.6 (CH), 31.1 (C_q), 36.9 (CH), 52.0 (CH_3), 66.9 (C_q), 78.9 (CH_2), 127.3 (CH), 128.1 (2 $\times$ CH), 129.2 (2 $\times$ CH), 135.3 (C_q), 163.4 (C=N), 171.8 (C=O); GC-MS (70 eV) m/z (%) 287 (M^+ , 5), 229 (28), 228 (100), 200 (13), 157 (13), 129 (20), 115 (8); FT-IR (film, cm^{-1}): 2957, 2927, 1732 (s, C=O), 1660 (s, C=N), 1450, 1302. Anal. calcd. for $C_{17}H_{21}NO_3$: C, 71.06; H, 7.37; N, 4.87. Found: C, 71.42; H, 7.50; N, 4.80.

Methyl-(1*R,2*R**,3*S**)-2-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-3-(4-methoxyphenyl)-2-**

methylcyclopropanecarboxylate (5b): yellow solid, 57-59 °C, 85 %. 1H NMR (300 MHz) δ 0.85 (s, 3 H), 1.09 (s, 3 H), 1.60 (s, 3 H), 2.79 (d, $J = 6.6$ Hz, 1 H), 3.03 (d, $J = 6.6$ Hz, 1 H), 3.48 and 3.58 (2 $\times$ d, AB system, $J = 7.8$ Hz, 2 H), 3.74 (s, 3 H), 3.75 (s, 3 H), 6.76 (d, $J = 8.9$ Hz, 2 H), 7.12 (d, $J = 8.9$ Hz, 2 H); ^{13}C NMR (75.4 MHz) δ 15.7 (CH_3), 27.9 (CH_3), 28.2 (CH_3), 29.3 (CH), 30.8 (C_q), 36.5 (CH), 52.0 (CH_3), 55.2 (CH_3), 66.8 (C_q), 78.9 (CH_2), 113.2 (2 $\times$ CH), 127.5 (2 $\times$ CH), 129.7 (C_q), 158.4 (C_q), 164.0 (C=N), 171.3 (C=O); GC-MS (70 eV) m/z (%) 317 (M^+ , 12), 286 (7), 258 (100), 230 (16), 187 (17), 159 (15), 127 (12); FT-IR (film, cm^{-1}): 2967, 1732 (s, C=O), 1658 (s, C=N), 1441, 1408, 1171, 810. Anal. calcd. for $C_{18}H_{23}NO_4$: C, 68.12; H, 7.30; N, 4.41. Found: C, 68.44; H, 7.56; N, 4.22.

Methyl-(1*R,2*R**,3*S**)-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-3-(2-furyl)-2-**

methylcyclopropanecarboxylate (5c): colorless oil, 83 %. 1H NMR (300 MHz) δ 0.99 (s, 3 H),

1.15 (s, 3 H), 1.55 (s, 3 H), 2.72 (d, $J = 6.4$ Hz, 1 H), 3.02 (d, $J = 6.4$ Hz, 1 H), 3.66 and 3.71 ($2 \times$ d, AB system, $J = 7.9$ Hz, 2 H), 3.73 (s, 3 H), 6.04–6.08 (m, 1 H), 6.20–6.25 (m, 1 H), 7.24–7.30 (m, 1 H); ^{13}C NMR (75.4 MHz) δ 15.2 (CH_3), 28.0 (CH_3), 28.1 (CH_3), 29.6 (CH), 29.7 (CH), 30.4 (C_q), 52.0 (CH_3), 66.9 (C_q), 79.4 (CH_2), 107.3 (CH), 110.1 (CH), 141.5 (CH), 150.1 (C_q), 163.9 (C=N), 170.5 (C=O); GC-MS (70 eV) m/z (%) 277 (M^+ , 5), 246 (3), 218 (100), 190 (11), 147 (10), 129 (20), 91 (9); FT-IR (film, cm^{-1}): 2956, 2920, 1732 (s, C=O), 1649 (s, C=N), 1436, 1378, 1205, 919. Anal. calcd. for $\text{C}_{15}\text{H}_{19}\text{NO}_4$: C, 64.97; H, 6.91; N, 5.05. Found: C, 65.29; H, 7.02; N, 4.99.

Methyl-2-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-3-phenylcyclopropanecarboxylate (6a): colorless oil, 49 % inseparable mixture of two diastereoisomers, dr (1*R**,2*R**,3*S**)/(1*R**,2*S**,3*S**) 1/1. ^1H NMR (300 MHz) δ 0.86 (s, 3 H), 1.15 (s, 3 H), 1.24 (s, 3 H), 1.27 (s, 3 H), 2.50–2.61 (m, 2 H), 2.74–2.85 (m, 2 H), 2.91–3.06 (m, 2 H), 3.50 (s, 3 H), 3.52 (s, 3 H), 3.58 and 3.67 ($2 \times$ d, AB system, $J = 8.0$ Hz, 2 H), 3.99 (s, 2 H), 7.02–7.40 (m, 10 H); ^{13}C NMR (75.4 MHz) δ , 24.6 (CH), 24.9 (CH), 27.8 ($2 \times \text{CH}_3$), 27.9 (CH_3), 28.0 (CH_3), 29.1 (CH), 29.2 (CH), 31.2 (CH), 31.5 (CH), 51.8 (CH_3), 52.2 (CH_3), 66.9 (C_q), 67.1 (C_q), 79.0 (CH_2), 79.2 (CH_2), 127.0 (CH), 127.3 (CH), 128.1 ($2 \times \text{CH}$), 128.2 ($2 \times \text{CH}$), 129.1 ($2 \times \text{CH}$), 129.2 ($2 \times \text{CH}$), 134.8 (C_q), 135.3 (C_q), 163.4 (C=N), 163.7 (C=N), 171.2 (C=O), 172.5 (C=O); GC-MS (70 eV) m/z (%) diastereoisomer with a lower t_r , 273 (M^+ , 8), 215 (30), 204 (100), 158 (16), 143 (16), 77 (10); diastereoisomer with a higher t_r , 273 (M^+ , 10), 242 (8), 215 (29), 204 (100), 186 (13), 115 (16), 77 (19); FT-IR (film, cm^{-1}): 2965, 1731 (s, C=O), 1660 (s, C=N), 1652, 1249, 1175, 750, 700. Anal. Calcd for: $\text{C}_{16}\text{H}_{19}\text{NO}_3$: C, 70.31; H, 7.01; N, 5.12. Found: C, 70.53; H, 7.29; N, 5.03.

Methyl-2-(4,5-dihydro-4,4-dimethyl-2-oxazolyl)-3-(4-

methoxyphenyl)cyclopropanecarboxylate 6b: colorless oil, 53 % inseparable mixture of two diastereoisomers, dr (1*R**,2*R**,3*S**)/(1*R**,2*S**,3*S**) 1/1. ^1H NMR (300 MHz) δ 0.91 (s, 3 H), 1.15 (s, 3 H), 1.27 (s, 6 H), 2.56 (dd, $^3J = 5.1$ Hz, $^3J' = 5.3$ Hz, 2 H), 2.82 (dd, $^3J = 5.3$ Hz, $^3J' = 5.4$ Hz, 2 H), 2.92–3.04 (m, 2 H), 3.50 (s, 3 H), 3.58 and 3.67 ($2 \times$ d, AB system, $J = 8.1$ Hz, 2 H), 3.76 (s, 3

H), 3.79 (6 H), 3.95 (s, 2 H), 6.77–6.86 (m, 4 H), 7.16–7.24 (m, 4 H); ^{13}C NMR (75.4 MHz) δ 24.7 (CH), 25.0 (CH), 27.9 (CH₃), 28.1 (CH₃), 28.3 (CH₃), 28.9 (CH₃), 29.3 (CH), 29.5 (CH), 31.1 (CH), 31.4 (CH), 51.8 (CH₃), 52.2 (CH₃), 55.1 (CH₃), 55.2 (CH₃), 66.9 (*C_q*), 67.1 (*C_q*), 79.0 (CH₂), 79.2 (CH₂), 113.3 (2 $\times$ CH), 113.5 (2 $\times$ CH), 129.8 (2 $\times$ CH), 129.9 (2 $\times$ CH), 137.3 (*C_q*), 137.4 (*C_q*), 158.6 (2 $\times$ *C_q*), 163.5 (C=N), 163.7 (C=N) 172.2 (C=O), 172.5 (C=O); GC-MS (70 eV) *m/z* (%) diastereoisomer with a lower *t_r*, 303 (*M*⁺, 8), 272 (8), 245 (30), 244 (100), 216 (13), 188 (16), 173 (16), 145 (14), 113 (17); diastereoisomer with a higher *t_r*, 303 (*M*⁺, 10), 272 (8), 245 (29), 244 (100), 216 (13), 188 (19), 173 (16), 145 (16), 113 (19); FT-IR (film, cm⁻¹): 2965, 1733 (s, C=O), 1663 (s, C=N), 1657 (s, C=N) 1518, 1249, 1175. Anal. Calcd. for C₁₇H₂₁NO₄: C, 67.31; H, 6.98; N, 4.62. Found: C, 67.58; H, 7.21; N, 4.60.

Methyl-(1*R*,2*R*,3*S*,4'*S*)- and (1*S*,2*S*,3*R*,4'*S*)-2-[4,5-dihydro-4-isopropyl-2-oxazolyl]-2-methyl-3-phenylcyclopentanecarboxylate (7a). Major diastereoisomer: colorless oil, 45 %. ^1H NMR (300 MHz) δ 0.60 (d, *J* = 6.5 Hz, 3 H), 0.74 (d, *J* = 6.5 Hz, 3 H), 1.25–1.56 (m, 1 H), 1.66 (s 3 H), 2.90 (d, *J* = 6.5 Hz, 1 H), 3.15 (d, *J* = 6.5 Hz, 1H), 3.41–3.92 (m, 3 H), 3.80 (s, 3 H), 7.20–7.37 (m, 5 H); ^{13}C NMR (75.4 MHz) δ 15.8 (CH₃), 18.1 (CH₃), 18.9 (CH₃), 29.5 (CH), 31.1 (*C_q*), 32.7 (CH), 37.4 (CH), 52.0 (CH₃), 69.6 (CH), 70.6 (CH₂), 127.8 (CH), 128.4 (2 $\times$ CH), 128.7 (2 $\times$ CH), 135.4 (*C_q*), 163.3 (C=N), 171.2 (C=O); GC-MS (70 eV) *m/z* (%), 301 (*M*⁺,11), 233 (38), 232 (100), 204 (8), 77 (10); FT-IR (film, cm⁻¹): 2961, 1729 (s, C=O), 1653 (s, C=N), 1444, 1153, 754, 699. Anal. calcd for C₁₈H₂₃NO₃: C, 71.73; H, 7.69; N, 4.65. Found: C, 71.95; H, 7.53; N, 4.37. **Minor diastereoisomer:** colorless oil, 10 %. ^1H NMR (300 MHz) δ 0.67 (d, *J* = 6.6 Hz, 3 H), 0.82 (d, *J* = 6.6 Hz, 3 H), 1.21–1.53 (m, 1 H), 1.68 (s 3 H), 2.90 (d, *J* = 6.5 Hz, 1 H), 3.09 (d, *J* = 6.5 Hz, 1 H), 3.38–3.90 (m, 3 H), 3.82 (s, 3 H), 7.23–7.43 (m, 5 H); ^{13}C NMR (75.4 MHz) δ 15.7 (CH₃), 18.1 (CH₃), 18.9 (CH₃), 29.5 (CH), 31.2 (*C_q*), 32.7 (CH), 37.4 (CH), 51.9 (CH₃), 69.6 (CH), 72.0 (CH₂), 126.8 (CH), 128.4 (2 $\times$ CH), 128.6 (2 $\times$ CH), 132.6 (*C_q*), 163.0 (C=N), 171.1 (C=O); GC-MS (70

eV) m/z (%), 301 (M^+ , 5), 233 (20), 232 (100), 204 (3), 77 (15); FT-IR (film, cm^{-1}): 2964, 1727 (s, C=O), 1655 (s, C=N), 1440, 1153, 750, 702. Anal. calcd for $\text{C}_{18}\text{H}_{23}\text{NO}_3$: C, 71.73; H, 7.69; N, 4.65. Found: C, 72.07; H, 7.74; N, 4.41.

Methyl-(1*R*,2*R*,3*S*,4'*S*)- and (1*S*,2*S*,3*R*,4'*S*)-2-[(4,5-dihydro-4-isopropyl-2-oxazolyl]-3-(4-methoxyphenyl)-2-methyl-cyclopopanecarboxylate (7b). Major diastereoisomer : white solid, mp 47-48 °C, 49 %. ^1H NMR (300 MHz) δ 0.62 (d, 6.5 Hz, 3 H), 0.75 (d, 6.5 Hz, 3 H), 1.22–1.55 (m, 1 H), 1.63 (s, 3 H), 2.84 (d, J = 6.5 Hz, 1 H), 3.08 (d, J = 6.5 Hz, 1H), 3.48–3.92 (m, 3 H), 3.78 (s, 3 H), 3.80 (s, 3 H), 6.70–6.85 (m, 2 H), 7.15–7.20 (m, 2 H); ^{13}C NMR (75.4 MHz) δ 16.2 (CH_3), 18.7 (CH_3), 18.9 (CH_3), 29.5 (CH), 31.1 (C_q), 32.6 (CH), 36.6 (CH), 51.9 (CH_3), 55.1 (CH_3), 69.4 (CH), 71.9 (CH), 113.3 ($2 \times \text{CH}$), 127.5 ($2 \times \text{CH}$), 129.6 (C_q), 158.3 (C_q), 165.0 (C=N), 172.2 (C=O); GC-MS (70 eV) m/z (%), 331 (M^+ , 11), 273 (38), 272 (100), 244 (3), 159 (12), 127 (8); FT-IR (film, cm^{-1}): 2957, 1732 (s, C=O), 1661 (s, C=N), 1516, 1443, 1250, 1173, 818. Anal calcd. for $\text{C}_{19}\text{H}_{25}\text{NO}_4$: C, 68.86; H, 7.60; N, 4.23. Found: C, 69.21; H, 7.51; N, 4.10. **Minor diastereoisomer:** white solid, mp 45-46 °C, 11 %. ^1H NMR (300 MHz) δ 0.69 (d, J = 6.6 Hz, 3 H), 0.84 (d, J = 6.6 Hz, 3 H), 1.24–1.56 (m, 1 H), 1.65 (s 3 H), 2.84 (d, J = 6.5 Hz, 1 H), 3.03 (d, J = 6.5 Hz, 1 H), 3.46–3.92 (m, 3 H), 3.79 (s, 3 H), 3.81 (s, 3H), 6.68–6.83 (m, 2 H), 7.14–7.20 (m, 2 H); ^{13}C NMR (75.4 MHz) δ 16.1 (CH_3), 18.0 (CH_3), 18.8 (CH_3), 29.5 (CH), 32.1 (C_q), 32.6 (CH), 36.4 (CH), 51.8 (CH_3), 55.2 (CH_3), 69.8 (CH), 72.3 (CH_2), 113.1 ($2 \times \text{CH}$), 127.5 ($2 \times \text{CH}$), 129.4 (C_q), 158.2 (C_q), 164.9 (C=N), 171.5 (C=O); GC-MS (70 eV) m/z (%), 331 (M^+ , 5), 273 (24), 272 (100), 244 (3), 159 (8), 127 (6); FT-IR (film, cm^{-1}): 2957, 1732 (s, C=O), 1661 (s, C=N), 1516, 1443, 1250, 1173, 818. Anal calcd. for $\text{C}_{19}\text{H}_{25}\text{NO}_4$: C, 68.86; H, 7.60; N, 4.23. Found: C, 69.16; H, 7.82; N, 4.16.

Methyl-(1*R*,2*R*,3*S*,4*S*,5*S*)- and (1*S*,2*S*,3*R*,4*S*,5*S*)-2-[4,5-dihydro-4-methoxymethyl-5-phenyl-2-oxazolyl]-3-(4-methoxyphenyl)-2-methylcyclopopanecarboxylate (7c): white solid, mp 62-64 °C, 55 %, inseparable mixture of two diastereoisomers (dr = 3/2). ^1H NMR (300 MHz) δ 1.48 (s, 3 H, minor), 1.50 (s, 3 H, major), 2.64 (d, J = 6.4 Hz, 1 H, minor), 2.70 (d, J = 6.4 Hz, 1 H, major),

3.00 (d, $J = 6.4$ Hz, 1 H major) 3.12 (d, $J = 6.4$ Hz, 1 H minor), 3.32 (s, 3 H, major), 3.34 (s, 3 H, minor), 3.40–3.62 (m, 2 H major and 2 H minor), 3.52 (s, 3 H major and 3 H minor), 3.70–3.95 (m, 1 H major and 1 H minor), 3.76 (s, 3 H, major), 3.80 (s, 3 H, minor), 4.82 (d, $J = 7.7$ Hz, 1 H, major), 4.86 (d, $J = 7.7$ Hz, 1 H, minor), 6.74–6.89 (m, 2 H major and 2 H minor), 7.12–7.24 (m, 2 H major and 2 H minor), 7.21–7.44 (m, 5 H major and 5 H minor); ^{13}C NMR (75.4 MHz) δ 21.2 (CH_3 , minor), 21.3 (CH_3 , major), 29.9 (CH , major), 30.0 (CH , minor), 32.1 (CH , major), 32.3 (CH , minor), 39.6 (C_q , major), 39.7 (C_q , minor), 51.9 (CH_3 , minor), 52.2 (CH_3 , major), 55.4 (CH_3 , major), 55.6 (CH_3 , minor), 59.2 (CH_3 , major), 59.3 (CH_3 , minor), 73.6 (CH_2 , major), 73.7 (CH_2 , minor), 74.2 (CH , major), 74.6 (CH , minor), 84.1 (CH , minor), 84.2 (CH , major), 114.3 ($2 \times \text{CH}$, minor), 114.5 ($2 \times \text{CH}$, major), 125.3 ($2 \times \text{CH}$, minor), 125.6 ($2 \times \text{CH}$, major), 128.1 (CH , major), 128.2 (CH , minor), 128.6 ($2 \times \text{CH}$, major), 128.7 ($2 \times \text{CH}$, minor), 129.1 ($2 \times \text{CH}$, minor), 129.3 ($2 \times \text{CH}$, major), 131.0 (C_q , major), 131.6 (C_q , minor), 140.2 (C_q , major), 140.3 (C_q , minor), 158.8 (C_q , minor), 159.4 (C_q , major), 166.2 ($\text{C}=\text{N}$, minor), 166.5 ($\text{C}=\text{N}$, major), 171.0 ($\text{C}=\text{O}$, minor), 171.3 ($\text{C}=\text{O}$, major); GC-MS (70 eV) m/z (%) diastereoisomer minor, 409 (M^+ , 1), 380 (10), 337 (24), 336 (100), 308 (3), 159 (8), 77 (6); diastereoisomer major, 409 (M^+ , 1), 364 (10), 337 (20), 336 (100), 223 (8), 77 (6); FT-IR (film, cm^{-1}): 2967, 1741 (s, $\text{C}=\text{O}$), 1661, 1655 (s, $\text{C}=\text{N}$), 1443, 1250, 1173, 818, 703. Anal. Calcd. for $\text{C}_{24}\text{H}_{27}\text{NO}_5$: C, 70.40; H, 6.65; N, 3.42. Found: C, 70.12; H, 6.52; N, 3.56.