Tandem Oxidation Processes for the Preparation of Functionalised Cyclopropanes

Magalie F. Oswald, Steven A. Raw, Richard J. K. Taylor*
Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK
rjkt1@york.ac.uk

Supplementary Information

Compounds **3a**, ^{4c}**3b**, ^{4a}**3c**, ^{4b}**3g**, ^{4d}**3h**, ^{4e}**3i**, ^{4f}**3o**, ^{4g}**3p**, ^{4h}**3r**⁴ⁱ and **3t**⁹ are known compounds and the data correspond with those reported in the literature.

To a solution of allylic alcohol **1a** (36 mg, 0.50 mmol) in dichloromethane (5 mL) was added powdered 4Å molecular sieves (0.50 g), (benzoylmethylene)dimethylsulfurane **2b** (108 mg, 0.60 mmol) and manganese dioxide (435 mg, 5.0 mmol). The mixture was heated at reflux for 2 hours then cooled to room temperature. The crude mixture was then filtered through celite[®] and the residue washed with dichloromethane (50 mL) giving a pale yellow solution. After removal of the solvent, the residue was purified by flash column chromatography (19 :1 to 4 :1 PE :EtOAc) to give the *title compound* **3d** (50 mg, 53%) as a colorless oil, ~4.2:1 *trans:cis* by ¹H NMR integration. $\nu_{\max}$ (film) 2935, 1712, 1665, 1596, 1449, 1379, 1305, 1225 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) *trans*: δ 1.26 (3 H, s, Me), 1.59 (1 H, dd, *J* 4.9 Hz, *J* 8.2 Hz), 1.91 (1 H, dd, *J* 4.9 Hz, *J* 7.0 Hz), 3.20 (1 H, dd, *J* 8.2 Hz, *J* 7.0 Hz), 7.47 (2 H, dd (app. t), *J* 7.6 Hz, arom.), 7.58 (1 H, dd (app. t), *J* 7.6 Hz, arom.), 7.90 (1 H, d, *J* 7.6 Hz, arom.), 9.09 (1 H, s, CHO) *cis*: δ 1.44 (3 H, s, Me), 1.54 (1 H, dd, *J* 4.9 Hz, *J* 7.6 Hz), 2.39 (1 H, dd, *J* 4.9 Hz, *J* 7.0 Hz), 3.01 (1 H, dd, *J* 7.6 Hz, *J* 7.0 Hz), 7.47 (2 H, dd (app. t), *J* 7.6 Hz, arom.), 7.58 (1 H, dd (app. t), *J* 7.6 Hz, arom.), 7.93 (1 H, d, *J* 7.6 Hz, arom.), 9.11 (1 H, s, CHO); ¹³C NMR (100 MHz, CDCl₃) *trans*: δ 10.3 (CH₃), 18.6 (CH₂), 29.9 (CH), 37.4 (C), 128.3 (CH), 128.9 (CH), 133.6 (CH), 137.8 (C), 194.9 (C), 200.0 (C) *cis*: δ 17.5 (CH₃), 21.2 (CH₂), 35.0 (CH), 37.8 (C), 128.4 (CH), 128.9 (CH), 133.7 (CH), 137.2 (C), 195.6 (C), 200.4 (C); *m/z* (CI) 189 (MH⁺) [HRMS (CI) [MH⁺] C₁₂H₁₃O₂ requires 189.0916. Found: 189.0912 (2.1 ppm error).

To a solution of allylic alcohol **1c** (101 mg, 0.50 mmol) in dichloromethane (5 mL) was added powdered 4Å molecular sieves (0.50 g), (carbethoxymethylene)dimethylsulfurane **2a** (90 mg, 0.60 mmol) and manganese dioxide (430 mg, 4.9 mmol). The mixture was heated at reflux for 18 hours, cooled to room temperature and stirred for a further 8 hours. The crude mixture was then filtered through celite[®] and the residue washed with dichloromethane (50 mL) giving a pale yellow solution. After removal of the solvent, the residue was purified by flash column chromatography (100% petroleum ether) to give the *title compound* **3e** (100 mg, 59%) as a colorless oil, ~3.5:1 *trans:cis* by ¹H NMR integration. $\nu_{\max}$ (film) 779, 839, 1097, 1176, 1257,

1704, 1732, 2856, 2931, 2953 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) **trans**: δ 0.02 (3 H, s, SiMe), 0.05 (3 H, s, SiMe); 0.84 (9 H, s, SiCMe₃), 1.23-1.28 (3 H, m, $\text{CO}_2\text{CH}_2\text{CH}_3$), 1.45 (1 H, dd, J 4.5 Hz, J 8.5 Hz), 1.53 (1 H, dd, J 4.5 Hz, J 6.7 Hz), 2.31 (1 H, dd, J 8.5 Hz, J 6.7 Hz), 3.86 (1 H, d, J 11.1 Hz), 4.03-4.20 (2 H, m, $\text{CO}_2\text{CH}_2\text{CH}_3$), 4.32 (1 H, d, J 11.1 Hz), 9.52 (1 H, s, CHO) **cis**: δ 0.02 (3 H, s, SiMe), 0.05 (3 H, s, SiMe); 0.84 (9 H, s, SiCMe₃), 1.23-1.28 (3 H, m, $\text{CO}_2\text{CH}_2\text{CH}_3$), 1.63 (1 H, dd, J 4.8 Hz, J 8.2 Hz), 1.91 (1 H, dd, J 4.8 Hz, J 6.7 Hz), 2.33 (1 H, dd, J 8.2 Hz, J 6.7 Hz), 3.93 (2 H, s), 4.03-4.20 (2 H, m, $\text{CO}_2\text{CH}_2\text{CH}_3$), 9.38 (1 H, s, CHO); ^{13}C NMR (67.5 MHz, CDCl_3) **trans**: δ -5.5 (CH₃), -5.4 (CH₃), 14.3 (CH₃), 18.3 (C), 18.9 (CH), 25.8 (CH₃), 26.4 (CH), 40.6 (C), 59.4 (CH₂), 61.3 (CH₂), 170.1 (C), 200.2 (C) **cis**: δ -5.5 (CH₃), -5.4 (CH₃), 14.3 (CH₃), 16.1 (CH₂), 18.3 (C), 25.2 (CH), 25.8 (CH₃), 40.3 (C), 59.1 (CH₂), 61.5 (CH₂), 171.0 (C), 200.4 (C); m/z (CI) 287 (MH^+) [HRMS (CI) [MH^+] $\text{C}_{14}\text{H}_{27}\text{O}_4\text{Si}$ requires 287.1679. Found: 287.1681 (0.2 ppm error).

To a solution of allylic alcohol **1c** (51 mg, 0.25 mmol) in dichloromethane (5 mL) was added powdered 4Å molecular sieves (0.25 g), (benzoylmethylene)dimethylsulfurane **2b** (54 mg, 0.30 mmol) and manganese dioxide (217 mg, 2.5 mmol). The mixture was heated at reflux for 4.5 hours then cooled to room temperature. The crude mixture was then filtered through celite[®] and the residue washed with dichloromethane. After removal of the solvent, the residue was purified by flash column chromatography (19 :1 to 9 :1 PE :EtOAc) to give the *title compound* **3f** (58 mg, 73%) as a colorless oil, ~2.0:1 *trans*:*cis* by ^1H NMR integration. ν_{max} (film) 2954, 2929, 2857, 1716, 1675, 1451, 1255, 1223, 1090 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) **trans**: δ -0.29 (3 H, s, SiMe), -0.09 (3 H, s, SiMe); 0.62 (9 H, s, SiCMe₃), 1.49 (1 H, dd, J 4.1 Hz, J 7.9 Hz), 1.91 (1 H, dd, J 4.1 Hz, J 6.7 Hz), 3.35 (1 H, dd, J 7.9 Hz, J 6.7 Hz), 3.58 (1 H, d, J 11.3 Hz, OCHa), 4.35 (1 H, d, J 11.3 Hz, OCHb), 7.39-7.49 (2 H, m, arom.), 7.50-7.62 (2 H, m, arom.), 7.94 (1 H, d, J 7.3 Hz, arom.), 9.76 (1 H, s, CHO) **cis**: δ 0.11 (3 H, s, SiMe), 0.12 (3 H, s, SiMe); 0.92 (9 H, s, SiCMe₃), 1.68 (1 H, dd, J 4.6 Hz, J 7.6 Hz), 2.27 (1 H, dd, J 4.6 Hz, J 6.7 Hz), 3.35 (1 H, dd, J 7.6 Hz, J 6.7 Hz), 3.79 (1 H, d, J 10.8 Hz, OCHa), 4.30 (1 H, d, J 10.8 Hz, OCHb), 7.39-7.49 (2 H, m, arom.), 7.50-7.62 (2 H, m, arom.), 8.03 (1 H, d, J 7.6 Hz, arom.), 9.12 (1 H, s, CHO); ^{13}C NMR (100 MHz, CDCl_3) **trans**: δ -6.0 (CH₃), -5.8 (CH₃), 18.1 (C), 18.6 (CH₂), 25.6 (CH₃), 30.2 (CH), 43.7 (C), 59.2 (CH₂), 128.5 (CH), 128.6 (CH), 133.2 (CH), 137.7 (C), 194.9 (C), 200.7 (C) **cis**: δ -5.4 (CH₃), -5.3 (CH₃), 16.6 (CH₂), 18.4 (C), 25.9 (CH₃), 30.1 (CH), 42.8 (C), 60.3 (CH₂), 128.6 (CH), 128.8 (CH), 133.6 (CH), 137.3 (C), 195.9 (C), 199.4 (C); m/z (CI) 319 (MH^+) [HRMS (CI) [MH^+] $\text{C}_{18}\text{H}_{27}\text{O}_3\text{Si}$ requires 319.1729. Found: 319.1730 (0.1 ppm error).

To a solution of alcohol **1f** (50 μL , 0.51 mmol) in dichloromethane (10 mL) was added powdered 4Å molecular sieves (0.50 g), (carbethoxymethylene)dimethylsulfurane **2a** (166 mg, 1.12 mmol) and manganese dioxide (443 mg, 5.1 mmol). The solution was heated at reflux for 4 hours. The crude mixture was cooled and filtered through celite[®] and the residue washed with dichloromethane (50 mL). After removal of the solvent, the crude material was purified by flash column chromatography (5:1 PE:EtOAc) to give the *title compound* **3j** (78

mg, 60%) as a colourless wax: $\nu_{\max}$ (film) 1186, 1334, 1369, 1692, 1731, 2983 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) δ 1.26 (6 H, t, J 7.0 Hz), 1.40-1.46 (4 H, m), 2.16-2.23 (2 H, m), 2.58-2.65 (2 H, m), 4.14 (4 H, q, J 7.0 Hz); ^{13}C NMR (67.5 MHz, CDCl_3) δ 14.3 (CH_3), 17.8 (CH), 24.9 (CH), 29.9 (CH_2), 61.3 (CH_2), 171.9 (C), 204.7 (C); m/z (CI) 255 (MH^+) 272 (MNH_4^+) [HRMS (CI) $[\text{MNH}_4^+]$ $\text{C}_{13}\text{H}_{22}\text{NO}_5$ requires 272.1498. Found: 272.1500 (0.7 ppm error).

To a solution of alcohol **1f** (49 μL , 0.50 mmol) in dichloromethane (5 mL) was added powdered 4Å molecular sieves (0.50 g), (benzoylmethylene)dimethylsulfurane **2b** (216 mg, 1.20 mmol) and manganese dioxide (435 mg, 5.0 mmol). The solution was heated at reflux for 2.5 hours. The crude mixture was cooled and filtered through celite® and the residue washed with dichloromethane (50 mL). After removal of the solvent, the crude material was purified by flash column chromatography (19:1 to 4:1 PE :EtOAc) to give the *title compound* **3k** (110 mg, 69%) as a colourless solid: m.p. 148-150 $^\circ\text{C}$; $\nu_{\max}$ (film) 1660, 1349, 1225 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 1.58-1.78 (4 H, m), 2.84-2.97 (2 H, m), 3.23-3.38 (2 H, m), 7.38-7.51 (4 H, m, arom.), 7.52-7.64 (2 H, m, arom.), 7.91-8.06 (2 H, m, arom.); ^{13}C NMR (100 MHz, CDCl_3) δ 20.2/20.3 (CH_2), 28.4/28.5 (CH), 32.3/32.4 (CH), 128.3/128.4 (CH), 128.7/128.8 (CH), 133.4/133.5 (CH), 136.9 (C), 196.9 (C), 205.8 (C); m/z (CI) 319 (MH^+) 336 (MNH_4^+) [HRMS (CI) $[\text{MH}^+]$ $\text{C}_{21}\text{H}_{19}\text{O}_3$ requires 319.1334. Found: 319.1330 (1.3 ppm error).

To a solution of allylic alcohol **1g** (76 mg, 0.50 mmol) in dichloromethane (5 mL) was added powdered 4Å molecular sieves (0.50 g), (benzoylmethylene)dimethylsulfurane **2b** (108 mg, 0.60 mmol) and manganese dioxide (435 mg, 5.0 mmol). The mixture was heated at reflux for 16 hours then cooled to room temperature. The crude mixture was then filtered through celite® and the residue washed with dichloromethane. After removal of the solvent, the residue was purified by flash column chromatography (19:1 to 4:1 PE :EtOAc) to give the *title compound* **3l** (102 mg, 76%) as a colorless oil, ~5.0:1 mixture of isomers by ^1H NMR integration. $\nu_{\max}$ (film) 2924, 1703, 1664, 1597, 1449, 1385, 1326, 1289, 1220, 1033 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) **major**: δ 0.94 (3 H, s, Me), 1.01 (1 H, d, J 10.7 Hz); 1.28 (3 H, s, Me), 1.65 (1 H, dd, J 6.4 Hz, J 3.4 Hz), 1.72 (1 H, dd, J 7.9 Hz, J 3.4 Hz), 1.93 (1 H, t, J 6.7 Hz), 2.10-2.19 (1 H, m), 2.30-2.39 (1 H, m), 2.56 (1 H, dd, J 19.2 Hz, J 2.8 Hz), 2.71 (1 H, dt, J 19.2 Hz, J 2.5 Hz), 3.27 (1 H, t, J 7.3 Hz), 7.39-7.45 (2 H, m, arom.), 7.49-7.55 (2 H, m, arom.), 7.91 (1 H, d, J 7.6 Hz, arom.) **minor**: δ 0.56 (3 H, s, Me), 1.13 (3 H, s, Me), 1.37 (1 H, d, J 10.5 Hz), 1.52-1.59 (2 H, m), 1.93 (1 H, obscured), 2.10-2.19 (2 H, m), 2.54 (1 H, dd, J 19.3 Hz, J 2.8 Hz), 2.65 (1 H, dt, J 19.3 Hz, J 2.8 Hz), 3.56 (1 H, t, J 7.3 Hz), 7.39-7.45 (2 H, m, arom.), 7.49-7.55 (2 H, m, arom.), 7.93 (1 H, d, J 7.3 Hz, arom.); ^{13}C NMR (100 MHz, CDCl_3) **major**: δ 21.1 (CH_3), 22.5 (CH_2), 26.2 (CH_3), 32.2 (CH), 32.6 (CH_2), 38.6 (CH), 40.3 (CH), 40.4 (C), 43.5 (CH_2), 45.3 (CH), 128.2 (CH), 128.6 (CH), 133.2 (CH), 137.9 (C), 195.8 (C), 210.5 (C) **minor**: δ 21.4 (CH_3), 22.1 (CH_2), 25.9 (CH_3), 32.6 (CH_2), 33.4 (CH), 38.7 (CH), 39.5 (C), 40.0 (CH), 43.4 (CH_2), 45.6 (C), 128.4 (CH), 128.6 (CH), 133.1 (CH), 137.8

(C), 195.8 (C), 210.4 (C); m/z (CI) 269 (MH^+) [HRMS (CI) [MH^+] $C_{18}H_{21}O_2$ requires 269.1542. Found: 269.1537 (1.7 ppm error).

Ethyl 2-benzoyl-3-(4-chlorophenyl)cyclopropanecarboxylate, 3q.

To a solution of alcohol **1j** (113 mg, 0.46 mmol) in CH_2Cl_2 (10 mL) was added 4Å molecular sieves (0.48 g), (carbethoxymethylene)dimethylsulfurane **2a** (85 mg, 0.57 mmol) and manganese dioxide (435 mg, 5 mmol). The mixture was heated at reflux for 14 hours. The crude mixture was filtered through celite[®] and the residue was washed with dichloromethane (50 mL) giving a pale yellow solution. After removal of the solvent, the crude material was purified by flash column chromatography (10:1 PE:EtOAc) to yield the *title compound* **3q** (116 mg, 80%) as a mixture of isomers, partially separable by chromatography. The first eluting material was shown to be **A-3q**, a colorless oil; ν_{max} (film) 1014, 1191, 1372, 1450, 1494, 1680, 1730, 2980 cm^{-1} ; 1H NMR (270 MHz, $CDCl_3$) δ 1.11 (3 H, t, J 7.2 Hz, $CO_2CH_2CH_3$), 2.82 (1 H, dd, J 4.8 Hz, J 10.0 Hz), 3.21 (1 H, dd, J 6.3 Hz, J 10.0 Hz), 3.80 (1 H, dd, J 4.8 Hz, J 6.3 Hz), 4.01 (2 H, q, J 7.2 Hz, CO_2CH_2), 7.27 (4 H, br s, chlorophenyl H's), 7.48-7.56 (2 H, m, ArCO meta-H's), 7.61-7.67 (1 H, m, ArCO para-H), 8.10 (2 H, d, J 7.2 Hz, ArCO ortho-H's); ^{13}C NMR (67.5 MHz, $CDCl_3$) δ 14.3, 29.8, 32.3, 34.3, 61.3, 128.6, 128.7, 130.4, 132.2, 133.3, 133.4, 133.8, 137.0, 168.8, 196.5. The next eluting material was an amorphous, colorless solid; m.p. 102-105 °C, shown to be a mixture of compounds **B-3q** and **C-3q**, ~3.6:1 **C:B** (by 1H NMR integration); ν_{max} (film) 1093, 1185, 1372, 1453, 1496, 1680, 1729, 2977 cm^{-1} ; 1H NMR (270 MHz, $CDCl_3$) **B-3q**: 1.30 (3 H, t, J 7.2 Hz, $CO_2CH_2CH_3$), 3.15 (1 H, dd, J 5.2 Hz, J 6.3 Hz), 3.26 (1 H, dd, J 6.3 Hz, J 10.4 Hz), 3.54 (1 H, dd, J 4.8 Hz, J 10.0 Hz), 4.19 (2 H, q, J 7.2 Hz, CO_2CH_2), 7.09-7.17 (2 H, m, ArCl H's), 7.23-7.33 (2 H, m, ArCl H's), 7.40-7.49 (2 H, m, ArCO meta-H's), 7.50-7.60 (1 H, m, ArCO ortho-H's), 7.90 (2 H, d, J 8.4 Hz, ArCO ortho-H's). **C-3q**: 1.10 (3 H, t, J 7.2 Hz, $CO_2CH_2CH_3$), 2.58 (1 H, dd, J 5.9 Hz, J 9.7 Hz), 3.02 (1 H, dd, J 6.3 Hz, J 9.7 Hz), 3.31 (1 H, t, J 6.1 Hz), 4.04 (2 H, q, J 7.2 Hz, CO_2CH_2), 7.09-7.17 (2 H, m, ArCl H's), 7.23-7.33 (2 H, m, ArCl H's), 7.40-7.49 (2 H, m, ArCO meta-H's), 7.50-7.60 (1 H, m, ArCO ortho-H's), 7.98 (2 H, d, J 8.1 Hz, ArCO ortho-H's); ^{13}C NMR (67.5 MHz, $CDCl_3$) δ **C-3q**: 14.3 (CH_3), 29.0 (CH), 31.7 (CH), 34.1 (CH), 61.3 (CH_2), 128.1 (CH), 128.5 (CH), 128.8 (CH), 129.0 (CH), 133.0 (C), 133.5 (CH), 136.7 (C), 136.9 (C), 169.0 (C), 193.5 (C); m/z (CI) 328 (MH^+) [HRMS (CI) [MH^+] $C_{19}H_{18}O_3^{35}Cl$ requires 329.0944. Found 329.0947 (0.8 ppm error).

Ethyl 2-benzoyl-3-(4-nitrophenyl)cyclopropanecarboxylate, 3s.

To a solution of alcohol **1n** (71 mg, 0.28 mmol) in THF (10 mL) was added 4Å molecular sieves (0.30 g), (carbethoxymethylene)dimethylsulfurane **2a** (82 mg, 0.56 mmol) and manganese dioxide (242 mg, 2.78 mmol). The mixture was heated at reflux for 14 hours. The crude mixture was filtered through celite[®] and the residue was washed with dichloromethane (50 mL) giving a pale yellow solution. After removal of the solvent, the

crude material was purified by flash column chromatography (5:1 PE:EtOAc) to yield the *title compound* **3s** (85 mg, 90%) as a mixture of isomers, partially separable by chromatography. The first eluting material was shown to be compound **A-3s**, a pale yellow oil ; $\nu_{\max}$ (film) 1010, 1344, 1515, 1597, 2852, 2923, 3390 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) δ 1.12 (3 H, t, J 7.1 Hz, $\text{CO}_2\text{CH}_2\text{CH}_3$), 2.88 (1 H, dd, J 4.8 Hz, J 10.0 Hz), 3.32 (1 H, dd, J 6.3 Hz, J 9.7 Hz), 3.88 (1 H, dd, J 4.8 Hz, J 6.3 Hz), 4.02 (2 H, q, J 7.1 Hz, CO_2CH_2), 7.45-7.58 (4 H, m), 7.60-7.69 (1 H, m), 8.07-8.12 (2 H, m), 8.13-8.20 (2 H, m); ^{13}C NMR (67.5 MHz, CDCl_3) δ 14.2 (CH_3), 29.9 (CH), 32.4 (CH), 33.9 (CH), 61.5 (CH_2), 123.5 (CH), 128.5 (CH), 129.0 (CH), 130.0 (CH), 134.0 (CH), 136.7 (C), 142.5 (C), 147.3 (C), 168.5 (C), 195.9 (C); m/z (CI) 240 (MH^+) 257 (MNH_4^+) [HRMS (CI) [MNH_4^+] $\text{C}_{19}\text{H}_{21}\text{N}_2\text{O}_5$ requires 357.1450. Found: 357.1448 (0.2 ppm error). The next eluting material was a pale yellow oil, shown to be compounds **B-3s** and **C-3s**, ~1.6:1 **C:B** (by ^1H NMR integration); $\nu_{\max}$ (film) 1184, 1347, 1520, 1601, 1679, 1730, 2983 cm^{-1} ; ^1H NMR (270 MHz, CDCl_3) δ **B-3s**: 1.34 (3 H, t, J 7.0 Hz, $\text{CO}_2\text{CH}_2\text{CH}_3$), 3.25 (1 H, dd, J 5.0 Hz, J 6.3 Hz), 3.38 (1 H, dd, J 10.0 Hz, J 6.3 Hz), 3.66 (1 H, dd, J 5.0 Hz, J 10.0 Hz), 4.24 (2 H, q, J 7.0 Hz), 7.34-7.43 (2 H, m), 7.44-7.53 (2 H, m), 7.53-7.64 (1 H, m), 7.89-7.96 (2 H, m), 7.99-8.10 (2 H, m) **C-3s**: 1.12 (3 H, t, J 7.4 Hz, $\text{CO}_2\text{CH}_2\text{CH}_3$), 2.72 (1 H, dd, J 5.9 Hz, J 9.7 Hz), 3.16 (1 H, dd, J 6.3 Hz, J 9.7 Hz), 3.46 (1 H, dd, J 5.9 Hz, J 6.3 Hz), 4.07 (2 H, q, J = 7.4 Hz, CO_2CH_2), 7.34-7.43 (2 H, m), 7.44-7.53 (2 H, m), 7.53-7.64 (1 H, m), 7.99-8.10 (2 H, m), 8.17-8.23 (2 H, m); ^{13}C NMR (67.5 MHz, CDCl_3) δ (There was difficulty in assigning signals to a given isomer, but the following were observed) 14.1 (CH_3), 14.3 (CH_3), 26.7 (CH), 28.9 (CH), 32.3 (CH), 34.9 (CH), 35.0 (CH), 35.6 (CH), 61.6 (CH_2), 61.8 (CH_2), 123.5 (ArC), 124.2 (ArC), 127.5 (ArC), 128.3 (ArC), 128.5 (ArC), 128.8 (ArC), 128.9 (ArC), 129.9 (ArC), 133.8 (ArC), 136.6 (C), 136.7 (C), 141.1 (C), 146.0 (C), 147.2 (C), 168.4 (C), 171.5 (C), 192.8 (C); m/z (CI) 240 (MH^+) [HRMS (CI) [MH^+] $\text{C}_{19}\text{H}_{18}\text{NO}_5$ requires 340.1184. Found : 340.1184 (0.0 ppm error).

Representative ^1H NMR Spectra

2-Benzoyl-1-methylcyclopropanecarboxaldehyde, **3d**

Title: k3867sar

Converted from "D:\POSTDOC5\PD479_1\K3867SAR.GXD"

LS

2-Benzoyl-1-methylcyclopropanecarboxaldehyde, **3d**

Title: k3868sar

Converted from "E:\POSTDOC5\PD479_1\K3868S-1.GXD"

Ethyl 2-(tert-butyltrimethylsilyloxymethyl)-2-formyl-cyclopropane-carboxylate, **3e**

Title: b6882mfo

Converted from "N:\SAR113\MAGALIE\MO29\1H\B6882MFO.GXD"

Ethyl 2-(tert-butyltrimethylsilyloxymethyl)-2-formyl-cyclopropane-carboxylate, 3e

Title: b6883mfo

Converted from "E:\NMRDAT~1\MO29\13C\B6883MFO.GXD"

2-Benzoyl-1-(tert-butyltrimethylsilyloxymethyl)cyclopropanecarboxaldehyde, **3f**

Title: k3843sar

Converted from "D:\POSTDOC5\PD481_1\IH\K3843SAR.GXD"

Title: k3844sar

Converted from "E:\POSTDOC5\PD481_1\13C\K3844SAR.GXD"

Diethyl 2,2'-carbonyldicyclopropanecarboxylate, **3j**.

Title: b6003mfo

Converted from "N:\SAR113\MAGALIE\MO16\1H\B6003MFO.GXD"

S13

Diethyl 2,2'-carbonyldicyclopropanecarboxylate, **3j**.

Title: b6004mfo

Converted from "E:\NMRDAT~1\MO16\13C\B6004M~1.GXD"

{2-[(2-Benzoylcyclopropyl)carbonyl]cyclopropyl} (phenyl)methan-one, **3k**

Title: k4010sar

Converted from "D:\POSTDOC5\PD492_1\1H\K4010SAR.GXD"

{2-[(2-Benzoylcyclopropyl)carbonyl]cyclopropyl} (phenyl)methan-one, **3k**

Title: k4011sar

Converted from "E:\POSTDOC5\PD492_1\13C\K4011SAR.GXD"

(1*R*,3*R*,6*S*,8*S*)-1-Benzoyl-7,7-dimethyl-6,8-methylene-spiro[2.5]octan-4-one
and (1*S*,3*S*,6*S*,8*S*)-isomer, **31**

Title: k3869sar

Converted from "D:\POSTDOC5\PD484_1\K3869SAR.GXD"

L1S

(1*R*,3*R*,6*S*,8*S*)-1-Benzoyl-7,7-dimethyl-6,8-methylene-spiro[2.5]octan-4-one
and (1*S*,3*S*,6*S*,8*S*)-isomer, **3l**

Title: k3870sar

Converted from "E:\POSTDOC5\PD484_1\K3870S-1.GXD"

81S

Ethyl 2-benzoyl-3-(4-chlorophenyl)cyclopropanecarboxylate, A-3q

Title: b6251mfo

Converted from "E:\NMRDAT~1\MO23\MO23-1\1H\B6251MFO.GXD"

Ethyl 2-benzoyl-3-(4-chlorophenyl)cyclopropanecarboxylate, A-3q

Title: b6252mfo

Converted from "E:\NMRDAT~1\MO23\MO23-1\13C\B6252MFO.GXD"

Ethyl 2-benzoyl-3-(4-chlorophenyl)cyclopropanecarboxylate, **B-3q/C-3q**

Title: b6253mfo

Converted from "E:\NMRDAT~1\MO23\MO23-3\1H\B6253MFO.GXD"

Ethyl 2-benzoyl-3-(4-chlorophenyl)cyclopropanecarboxylate, B-3q/C-3q

Title: k0395mfo

Converted from "E:\NMRDAT~1\MO23\MO23-3\13CCOR~1\K0395M~1.GXD"

S22

\\NMRDAT~1\MO27\3-6\1H\B6586MFO.GXD

Ethyl 2-benzoyl-3-(4-nitrophenyl)cyclopropanecarboxylate, **A-3s**

\\NMRDAT-1\MO27\3-6\13C\B6587MFO.GXD

MO27\10-17\1H\B6588M-1.GX1

Title: b6589mfo

Converted from "E:\NMRDAT~1\MO27\10-17\13C\B6589M~1.GXD"