Tandem Oxidation of Allylic and Benzylic Alcohols to Esters Catalyzed by *N*-Heterocyclic Carbenes

Brooks E. Maki, Audrey Chan, Eric M. Phillips, Karl A. Scheidt*

Department of Chemistry, Northwestern University,
2145 Sheridan Road, Evanston, Illinois, 60208

Supporting Information

1. General methods	S1-S2
2. Procedure and characterization data for reactions with alcohol as solvent	S2-S4
3. Procedure and characterization data for reactions with toluene as solvent	S4-S5
4. Procedure and characterization data for asymmetric reactions and catalyst	S5-S7
5. Selected NMR Spectra	S8-S17

General Information. All reactions were carried out under a nitrogen atmosphere in flame-dried glassware with magnetic stirring. Toluene, MeOH, and CH₂Cl₂ were purified by passage through a bed of activated alumina.¹ Reagents were purified prior to use unless otherwise stated following the guidelines of Perrin and Armarego.² Purification of reaction products was carried out by flash chromatography using EM Reagent silica gel 60 (230-400 mesh). Analytical thin layer chromatography was performed on EM Reagent 0.25 mm silica gel 60-F plates. Visualization was accomplished with UV light and anisaldehyde, ceric ammonium nitrate stain, potassium permanganate, or phosphomolybdic acid followed by heating. Infrared spectra were recorded on a Perkin Elmer 1600 series FT-IR spectrometer. ¹H-NMR spectra were recorded on a Varian Inova 500 (500 MHz) or Mercury 400 (400 MHz) spectrometer and are reported in ppm using solvent as an internal standard (CDCl₃ at 7.26 ppm). Data are reported as (ap = apparent, s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, b = broad; coupling constant(s) in Hz; integration. Proton-decoupled ¹³C-NMR spectra were recorded on a Varian Inova 500 (125 MHz) or Mercury 400 (100 MHz) spectrometer and are reported in ppm using solvent as an

-
1. Pangborn, A. B.; Giardello, M. A.; Grubbs, R. H.; Rosen, R. K.; Timmers, F. J. *Organometal.* **1996**, *15*, 1518-1520.
 2. Perrin, D. D. and Armarego, W. L. *Purification of Laboratory Chemicals*; 3rd Ed., Pergamon Press, Oxford. 1988.

internal standard (CDCl_3 at 77.0 ppm). Mass spectra data were obtained on a Varian 1200 Quadrupole Mass Spectrometer and Micromass Quadro II Spectrometer.

3-Phenyl-butyn-1-ol was purchased from GFS Chemicals. Manganese(IV) oxide and all other alcohols were commercially available from Sigma-Aldrich Chemical Company. Triazolium salt **21** was prepared using the method of Rovis, et al.³ with mesitylhydrazine prepared using the method of Carlin and Moores.⁴

General Procedure for triazolium-catalyzed oxidation with alcohol as solvent:

A flame-dried round bottom flask was charged with triazolium salt (22 mg, 0.10 mmol). The flask was sealed with a septum and put under positive pressure of nitrogen. The allylic alcohol (1 mmol), and the nucleophilic alcohol (5 mL) were added. The septum was removed and to the flask was added MnO_2 (1.304 g, 15 mmol). The flask was then sealed with septum and DBU (0.015 mL, 0.10 mmol) was added *via* syringe. The reaction stirred at ambient temperature until allylic alcohol and aldehyde were consumed (monitored by GC). The mixture was filtered through a thin pad of Celite. The Celite was washed with ethyl acetate (15 mL). The filtrate was then concentrated *in vacuo*. The resulting residue was purified by flash chromatography on silica gel.

***n*-Butyl *trans*-cinnamate (6a):** Filtrate was washed with H_2O (2x), sat. NaCl , dried with Na_2SO_4 . Volatiles were removed yielding 190 mg (93%) of **6a** as a colorless oil. Spectral data matched those found in the literature.⁵

***n*-Butyl 2-methyl-3-phenyl-propenoate (6b):** 25 mol % triazolium salt used. Purified by column chromatography (8% ethyl acetate/hexanes) yielding 199 mg (91%) of **6b** as a clear oil. R_f = 0.65 (17:3 hexanes/EtOAc); IR (film) 3175, 3038, 2963, 2873, 1710, 1636, 1449 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.69 (s, 1H, $\text{HC}=\text{C}$); 7.40 (m, 4H); 7.32 (m, 1H); 4.23 (t, 2H, OCH_2); 2.12 (s, 3H, $\text{C}=\text{CCH}_3$); 1.72 (m, 2H, CH_2); 1.46 (m, 2H, CH_2); 0.98 (t, 3H, CH_3); ^{13}C NMR (500 MHz, CDCl_3) δ 168.9, 138.8, 136.1, 129.8, 128.8, 128.5, 128.4, 65.0, 31.0, 19.5, 14.2; LRMS (electrospray): Mass calculated for $\text{C}_{14}\text{H}_{18}\text{O}_2$ $[\text{3M}+\text{Na}]^+$ 677. Found 676.8

***n*-Butyl 1-naphthoate (7):** Filtrate was washed with H_2O (2x), sat. NaCl , dried with Na_2SO_4 . Volatiles were removed yielding 204 mg

3. Kerr, M. S.; de Alaniz, J. R.; Rovis, T. *J. Org. Chem.* **2005**, 70, 5725-5728.

4. Carlin, R. B.; Moores, M. S. *J. Am. Chem. Soc.* **1962**, 84, 4107-4112.

5. Calo, V.; Nacci, A.; Monopoli, A.; Ieva, E.; Cioffi, N. *Org. Lett.* **2005**, 7, 617-620.

(90%) of **7** as a white solid. Spectral data matched those found in the literature.⁶

***n*-Butyl 2-naphthoate (8):** Filtrate was washed with H₂O (2x), sat. NaCl, dried with Na₂SO₄. Volatiles were removed yielding 207 mg (91%) of **8** as a white solid. *R*_f = 0.6 (8:2 hexanes/EtOAc); IR (film) 3054, 2962, 1712, 1264 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 8.21 (s, 1H); 8.08 (m, 1H); 7.87 (m, 2H); 7.55 (m, 2H); 4.40 (t, *J*=7, 2H); 1.82 (m, 2H); 1.53 (m, 2H); 1.02 (t, *J*=7, 3H); ¹³C NMR (500 MHz, CDCl₃) δ 167.11, 135.73, 132.76, 131.21, 131.18, 129.60, 128.41, 128.35, 128.01, 126.85, 125.52, 65.26, 31.11, 19.60, 14.09; LRMS (electrospray): Mass calculated for C₁₅H₁₆O₂ [2M]⁺ 456. Found 456.2

***n*-Butyl 3-phenylpropiolate (9):** Purified by column chromatography (15% ether/hexanes) yielding 171 mg (85%) of **9** as a yellow oil. IR (film) 3054, 2963, 2221, 1706, 1273 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.3-7.6 (m, 5H); 4.23 (t, 2H); 1.69 (m, 2H); 1.42 (m, 2H); 0.95 (t, 3H); ¹³C NMR (500 MHz, CDCl₃) δ 154.2, 132.9, 130.8, 128.5, 119.6, 86.0, 80.6, 65.9, 30.4, 19.0, 14.1; LRMS (electrospray): Mass calculated for C₁₃H₁₄O₂ [2M+Na]⁺ 427. Found 427.8

***n*-Butyl furan-2-carboxylate (10):** Purified by column chromatography (20:1 toluene/EtOAc) yielding 123 mg (73%) of **10** as a colorless oil. Spectral data matched those found in the literature.⁷

***n*-Butyl pent-2-enoate (11):** Purified by column chromatography (15% ether/hexanes) yielding 136 mg (87%) of **11** as a colorless oil. IR (film) 3054, 2963, 2873, 1723, 1654, 1262 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.15 (dt, *J*=15.9, 6.4, 1H, CH₂CH); 5.81 (d, *J*=15.9, 1H, CH); 4.13 (t, 2H, OCH₂); 2.22 (m, 2H, CH₂); 1.64 (m, 2H, CH₂); 1.40 (m, 2H, CH₂); 1.07 (t, 3H, CH₃); 0.94 (t, 3H, CH₃); ¹³C NMR (500 MHz, CDCl₃) δ 167.0, 150.6, 120.3, 64.1, 30.7, 25.3, 19.2, 13.7, 12.1; LRMS (electrospray): Mass calculated for C₉H₁₆O₂ [2M+Na]⁺ 335. Found 335.6

***n*-Butyl ethyl fumarate (12):** Purified by column chromatography (10% ether/hexanes) yielding 130 mg (65%) of **12** as colorless oil. Spectral data matched those found in the literature.⁸

6. Cai, M. Z.; Song, C. S.; Huang, X. *J. Chem. Soc. Perkin Trans. I* **1997**, 2273-2274.

7. Pittelkow, M.; Kamounah, F. S.; Boas, U.; Pedersen, B.; Christensen, J. B. *Synthesis* **2004**, 2485-2492.

8. Zaderenko, P.; Lopez, M. C.; Ballesteros, P. *J. Org. Chem.* **1996**, 61, 6825-6828.

literature.⁹

***n*-Butyl benzoate (13):** The standard procedure was followed with 2 mmol benzyl alcohol and 50 mol% triazolium and DBU. Purified by column chromatography (20:1 toluene/EtOAc) to yield 313 mg (88%) of **13** as colorless oil. Spectral data matched those found in the

¹³C NMR (500 MHz, CDCl₃) δ 166.4, 135.3, 134.3, 132.4, 127.9, 127.1, 121.6, 65.5, 30.6, 19.3, 13.7; LRMS (electrospray): Mass calculated for C₁₁H₁₃O₂Br [2M+Na]⁺ 537.2 Found 537.7

***n*-Butyl 2-bromobenzoate (14):** 25 mol% of triazolium salt used. Purified by column chromatography (15% ether/hexanes) yielding 180 mg (70%) of **14** as a yellow oil. IR (film) 3054, 2962, 1729, 1696, 1271, 705 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.3-7.9 (m, 4H); 4.34 (t, 2H, OCH₂); 1.75 (m, 2H, CH₂); 1.48 (m, 2H, CH₂); 0.97 (t, 3H, CH₃);

General Procedure for triazolium-catalyzed oxidation with toluene as solvent:

A flame-dried round bottom flask was charged with triazolium salt (34 mg, 0.15 mmol). The flask was sealed with a septum and put under positive pressure of nitrogen. Toluene (5 ml), the allylic alcohol (1 mmol), and the nucleophilic alcohol (5 mmol) were added. The septum was removed and to the flask was added MnO₂ (1.304 g, 15 mmol). The flask was then sealed with septum and DBU (0.165 ml, 1.1 mmol) was added *via* syringe. The reaction stirred at ambient temperature until allylic alcohol and aldehyde were consumed (monitored by GC). Upon consumption of allylic alcohol and aldehyde, methanol (10 mL) was added to the reaction. The mixture was then quickly filtered through a thin pad of celite (prolonged stirring with methanol resulted in transesterification to the methyl ester). The Celite was washed with ethyl acetate (15 ml). The filtrate was then concentrated *in vacuo*. The resulting residue was purified by flash chromatography on silica gel.

Methyl *trans*-cinnamate (5): Filtration through Celite and concentration *in vacuo* yielded 154 mg (95%) of **5** as a white solid. Spectral data matched those found in the literature.¹⁰

2-Propyl *trans*-cinnamate (15): Cinnamyl alcohol combined with MnO₂ in toluene and stirred for 30 minutes (until complete consumption of alcohol) at ambient temperature. Triazolium salt, DBU, and 2-propanol were added and stirred. Reaction mixture stirred with methanol and quickly filtered through Celite. Filtrate was concentrated *in vacuo* yielding 169 mg (89%) of **15** as a yellow oil. Spectral data matched those found in the literature.¹¹

9. Hans, J. J.; Driver, R. W.; Burke, S. D. *J. Org. Chem.* **2000**, 65, 2114-2121.

10. Crosignani, S.; White, P. D.; Linclau, B. *Org. Lett.* **2002**, 4, 2961-2963.

2,2,2-Trichloroethyl *trans*-cinnamate (16): Purified by column chromatography (20:1 toluene/EtOAc) yielding 228 mg (82%) of **16** as colorless crystals. Spectral data matched those found in the literature.¹²

2-Methoxyethyl *trans*-cinnamate (17): Purified by column chromatography (20:1 toluene/EtOAc) yielding 169 mg (82%) of **17** as a yellow oil. $R_f = 0.33$ (8:2 hexanes/EtOAc); IR (film) 3054, 2986, 2932, 1710, 1633, 1300, 1250, 1200, 1180 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.72 (d, $J=16$, 1H); 7.52 (m, 2H); 7.38 (m, 3H); 6.50 (d, $J=16$, 1H); 4.37 (t, $J=4$, 2H); 3.67 (t, $J=4$, 2H); 3.43 (s, 3H); ^{13}C NMR (500 MHz, CDCl_3) δ 167.24, 145.40, 134.60, 130.59, 129.15, 128.37, 118.03, 70.82, 63.84, 59.32; LRMS (electrospray): Mass calculated for $\text{C}_{12}\text{H}_{14}\text{O}_3$ $[2\text{M} + \text{Na}]^+$ 435. Found 435.2

2-(Trimethylsilyl)ethyl *trans*-cinnamate (18): Purified by column chromatography (15% ether/hexanes) yielding 183 mg (74%) of **18** as a yellow oil. IR (film) 3054, 2956, 1706, 1638, 1268 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.68 (d, $J=15.9$, 1H, CH); 7.3-7.5 (m, 5H); 6.43 (d, $J=15.9$, 1H, CH); 4.31 (t, $J=8.5$, 2H, OCH_2); 1.08 (t, $J=8.5$, 2H, CH_2Si); 0.08 (s, 9H, $\text{Si}(\text{CH}_3)_3$); ^{13}C NMR (500 MHz, CDCl_3) δ 167.0, 144.4, 134.4, 130.1, 128.8, 128.0, 118.4, 62.6, 17.3, -1.4, -1.6, -1.9; LRMS (electrospray): Mass calculated for $\text{C}_{14}\text{H}_{20}\text{O}_2\text{Si}$ $[2\text{M} + \text{Na}]^+$ 519.8 Found 519.4

Methyl 3-phenylpropionate (19): A flame-dried round bottom flask was charged with triazolium salt (22 mg, 0.10 mmol). The flask was sealed with a septum and put under positive pressure of nitrogen. Hydrocinnamaldehyde (0.5 mmol), and methanol (2.5 mL) were added. The septum was removed and to the flask was added MnO_2 (652 mg, 7.5 mmol). The flask was then sealed with septum and DBU (0.015 mL, 0.10 mmol) was added *via* syringe. The reaction stirred at ambient temperature until aldehyde was consumed (monitored by TLC). The mixture was filtered through a thin pad of Celite. The Celite was washed with ethyl acetate (15 mL). The filtrate was then concentrated *in vacuo* to afford 75 mg (91%) **19** as a yellow oil. Spectral data matched those found in the literature.¹³

General Procedure for NHC-catalyzed desymmetrization of meso diols:

11. Shono, T.; Ishige, O.; Uyama, H.; Kashimura, S. *J. Org. Chem.* **1986**, *51*, 546-549.

12. Dambacher, J.; Zhao, W.; El-Batta, A.; Anness, R.; Jiang, C. C.; Bergdahl, M. *Tetrahedron Lett.* **2005**, *46*, 4473-4477.

13. Semmelhack, M. F.; Stauffer, R. D.; Yamashita, A. *J. Org. Chem.* **1977**, *42*, 3180-3188.

A flame-dried 10 ml round bottom flask equipped with stirbar was charged with the chiral triazolium salt **E** (50 mg, 0.119 mmol), K₂CO₃ (16.4 mg, 0.119 mmol), and 18-crown-6 (15.7 mg, 0.06 mmol). The flask was sealed with a septum and placed under a positive pressure of nitrogen. Into it was then added the *meso* diol (115 mg, 0.99 mmol), proton sponge (85 mg, 0.397 mmol), and MnO₂ (517 mg, 5.95 mmol). Dichloromethane (1.6 mL, 0.25 M) was added resulting in a black suspension. Lastly, cinnamaldehyde (50 μ L, 0.397 mmol) was added via syringe. The reaction was allowed to stir at -30 °C under a positive pressure of nitrogen until the aldehyde was consumed as determined by TLC. Upon consumption of the aldehyde, the mixture was filtered through a thin pad of celite and washed with ethyl acetate (25 mL). The filtrate was then concentrated *in vacuo* and the resulting residue was purified by flash chromatography on silica gel.

(E)-(1S,2R)-2-hydroxycyclohexyl cinnamate (21): Purified by column chromatography (25% EtOAc/hexane) yielding 57 mg (58%) of **21** as a white solid. Mp: 61-63 °C; IR (film) 3464, 3048, 2936, 2861, 2351, 1701, 1281 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.10 (d, *J*=15.8, 1H, CH); 7.53 (m, 2H); 7.39 (m, 3H); 6.58 (d, *J*=16.4, 1H, CH); 5.07 (m, 1H); 3.95 (m, 1H); 1.98-1.95 (m, 2H); 1.84-1.82 (m, 1H); 1.69-1.65 (m, 3H); 1.57 (s, 1H); 1.43-1.39 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 166.9, 145.4, 134.5, 130.6, 129.1, 128.3, 118.3, 74.5, 69.5, 30.6, 27.3, 22.3, 21.3; LRMS (electrospray): Mass calculated for C₁₅H₁₈O₃ [2M+Na]⁺ 515.6 Found 515.4

Determination of Enantiomeric excess:

The desymmetrization of *cis*-1,2-cyclohexanediol was carried out using cinnamaldehyde and the chiral catalyst **21** to yield **20** with an enantiomeric excess of 80%. Enantiomeric excess determined by HPLC on a Chiralcel AD-H column. 10% IPA/Hexanes, 1mL/min.

Racemic product: =====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,4 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.778	BB	0.4061	4483.20117	163.41731	50.5538
2	16.976	PB	0.4681	4384.97656	138.98174	49.4462

Totals : 8868.17773 302.39905

Optically enriched product: =====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: DAD1 A, Sig=254,4 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	15.333	MM	0.3977	168.55333	7.06430	9.7477
2	17.353	MM	0.5499	1560.61377	47.29740	90.2523

Totals : 1729.16710 54.36171

Selected NMR Spectra:

11