Tandem Sonogashira Coupling: an efficient tool for the synthesis of diarylalkynes

Zoltán Novák, Péter Nemes, András Kotschy

*Department of General and Inorganic Chemistry,
Eötvös Loránd University,
H-1117 Pázmány Péter sétány 1/A
Budapest, Hungary
e-mail: kotschy@para.chem.elte.hu*

Supporting Information

Contents

Experimental procedures	2
Characterization data	4
References	9
NMR spectra	10

General

Unless otherwise indicated, all starting materials were obtained from commercial suppliers (Aldrich, Fisher, Merck) and were used without further purification including, dichloromethane, diisopropylamine, n-hexane, toluene, ethyl acetate, PPh_3 , CuI , $\text{PdCl}_2(\text{PPh}_3)_2$, 3-bromopyridine, 3-iodotoluene, iodobenzene, 4-iodoanisole, 2-bromochlorobenzene, 2-methyl-3-butyn-2-ol, MgSO_4 , NaH (55% in mineral oil), KOH . Toluene and diisopropylamine were purged with argon prior to use. All reactions were performed under an atmosphere of argon. Analytical thin-layer chromatography (TLC) was performed on Polygram SIL G/ UV 254 Pre-coated plastic TLC plates with 0.25mm silica gel from Macherey-Nagel + Co. Visualization was performed with a 254 nm UV lamp. Silica gel column chromatography was carried out with Flash silica gel (0.040 – 0.063 mm) from Merck. Melting points were determined using a Büchi melting point apparatus. All melting point values are reported as the midpoint of the thermal transition. The ^1H and ^{13}C NMR spectra were recorded on a Bruker DRX-250 spectrometer in CDCl_3 . Chemical shifts are expressed in parts per million (δ) using residual solvent protons as internal standards: chloroform (δ 7.26 for ^1H , δ 77.00 for ^{13}C). Coupling constants (J) are reported in Hertz (Hz). Splitting patterns are designated as s (singlet), d (doublet), t (triplet), m (multiplet), br s (broad singlet), dd (doublet doublet). Gas chromatographic measurements were performed with a Chrompack CP9001 Gas Chromatograph (10 m x 0.25 mm column with 0.12 μm CP-SIL 5CB coating, H_2 carrier gas). Combination gas chromatography and low resolution mass spectrometry was obtained on a Hewlett-Packard 5790A Gas Chromatograph (30 m x 0.25 mm column with 0.25 μm RH-5 MS+ coating, He carrier gas) and VG 12-250 Mass Spectrometer (Ion source: EI+, 70eV, 250°C; interface: 250°C). In the GC measurements di(ethylene glycol) dibutyl ether (40 μL per reaction) was used as internal standard.

Method A for the syntheses of diarylakynes: 10 mmol aryl halide, 351 mg $\text{PdCl}_2(\text{PPh}_3)_2$ (0,5 mmol, 5%) and 95 mg CuI (0,5 mmol, 5%) were placed into a flame dried Schlenk flask. 20 mL diisopropylamine was added to the flask followed by 1260 μL 2-methyl-3-butyn-2-ol (13 mmol, 1091mg). The reaction mixture was stirred under argon for one hour at 50°C. Then 4,48 g KOH (80 mmol), 351 mg $\text{PdCl}_2(\text{PPh}_3)_2$ (0.5 mmol,

5%), 95 mg CuI (0.5 mmol, 5%) and 10 mmol aryl halide were added and the reaction mixture was heated for five hours in a 110 °C oil bath. After cooling to room temperature the KOH was neutralized with 1M HCl, and then the mixture was extracted with DCM. The combined organic phases were dried over MgSO₄ then evaporated. The crude product was purified by column chromatography using hexane-ethyl acetate mixtures.

Method B for the synthesis of diarylalkynes: 5 mmol aryl halide, 100 mg PdCl₂(PPh₃)₂ (0.14 mmol, 2.8%) and 28 mg CuI (0.14 mmol, 2.8%) were placed into a flame dried Schlenk flask. 12 mL toluene and 1200 µL diisopropylamine were added to the flask followed by 630 µL 2-methyl-3-butyne-2-ol (6.5 mmol, 546 mg). The reaction mixture was stirred at 80 °C under argon until the first coupling was complete. Then the temperature of the oil bath was increased to 110 °C and 400 mg NaH (55% dispersion, 9.16 mmol) was added slowly to the mixture. After 5 min stirring 5 mmol of the appropriate aryl halide was added to the reaction mixture, which was followed after 25 min by the careful addition of another 100 mg portion of NaH (1.83 mmol). The reaction was stirred at 110 °C until completion (ca. 1 hour). After cooling to room temperature the suspension was filtered, and the separated amine-hydrochloride was washed with toluene. Evaporation of the combined toluene solutions gave a crude product, which was purified by column chromatography using hexane-ethyl acetate mixtures.

Combinatorial synthesis by tandem Sonogashira coupling

One-pot library synthesis

180 mg PdCl₂(PPh₃)₂ (0.25 mmol, 5%) and 47 mg CuI (0.25 mmol, 5%) were placed into a flame dried Schlenk flask followed by 12 mL diisopropylamine. Then 790 mg (5 mmol) 3-bromopyridine and 532 µL 2-methyl-3-butyne-2-ol (5.5 mmol, 462 mg) was added to the solution. The reaction mixture was stirred at 70 °C under argon until the first coupling was complete (1 hour). Then 180 mg PdCl₂(PPh₃)₂ (0.25 mmol, 5%), 47 mg CuI (0.25 mmol, 5%), 186 µL (1.66 mmol, 339 mg) iodobenzene, 213 µL (1.66 mmol, 362 mg) 3-iodotoluene, and 388 mg (1.66 mmol) 4-iodoaniline was added to the mixture followed by 2.24 g KOH. The Schlenk flask was purged again with argon, sealed and heated in an oil bath at 110 °C until completion.

Two-step library synthesis

90 mg $\text{PdCl}_2(\text{PPh}_3)_2$ (0.0125 mmol, 5%) and 24 mg CuI (0.0125 mmol, 5%) were placed into a flame dried Schlenk flask followed by 6 mL diisopropylamine, 402 mg (2.5 mmol) 4-(3'-pyridyl)-2-methyl-3-butyn-2-ol, 93 μL (0.83 mmol, 169.5 mg) iodobenzene, 107 μL (0.83 mmol, 181 mg) 3-iodotoluene, 194 mg (0.83 mmol) 4-iodoanazole and 1.12 g KOH (8eq.) under argon. Then the Schlenk flask was placed into the oil bath which was preheated to 110 $^\circ\text{C}$ and the reaction was stirred until completion. Samples were taken off, diluted with ether and examined with GC.

Characterization of the products

Diphenylacetylene (4a)¹: Yield: Method A: 1317mg (7.4 mmol, 74%); Method B: 498 mg (2.8 mmol, 56%) white crystal. m.p.: 60-61 $^\circ\text{C}$. ^1H NMR (CDCl_3 ; 250 MHz): δ 7.59-7.55 (m, 4H), 7.39-7.36 (m, 6H); ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 131.6, 128.3, 128.2, 123.2, 89.4; GC-MS (EI, 70eV) m/z $\text{C}_{14}\text{H}_{10}$ (% relative intensity, ion): 178 (100, $[\text{M}]^+$), 176 (61), 152 (40), 126 (29), 89 (23), 76 (68).

1-Methoxy-4-(phenylethynyl)-benzene (4b)²: Yield: Method A: 603 mg (2.9 mmol, 29%), 1331 mg (6.4 mmol, 64%), Method B: 582 mg (2.8 mmol, 56%), 707 mg (3.4 mmol, 68%) white crystal. m.p.: 79-81 $^\circ\text{C}$. ^1H NMR (CDCl_3 ; 250 MHz): δ 7.56-7.48 (m, 4H), 7.36-7.33 (m, 3H), 6.80 (dd, 1H, $J = 4.2, 2.0$ Hz), 3.83 (s, 3H). ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 159.2, 133.0, 131.4, 128.3, 127.9, 123.5, 115.2, 113.9, 89.3, 88.0, 55.2 MS(EI, 70eV) m/z $\text{C}_{15}\text{H}_{12}\text{O}$ (% relative intensity, ion): 208 (100, $[\text{M}]^+$), 193 (82), 165 (68), 139 (25).

1-Methyl-3-(phenylethynyl)-benzene (4c)³: Yield: Method B: 662 mg (3.45 mmol, 69%); 806 mg (4.2 mmol, 84%) yellow oil. ¹H NMR (CDCl₃; 250 MHz): δ 7.44-7.40 (m, 2H), 7.26-7.20 (m, 5H), 7.11 (t, 1H, *J* = 8.0 Hz), 7.03 (d, 1H, *J* = 8.0 Hz), 2.23 (s, 3H). ¹³C NMR (CDCl₃, 62.5 MHz): δ 137.8, 131.6, 129.1, 128.7, 128.6, 128.4, 128.3, 128.2, 123.4, 123.0, 89.6, 89.0, 21.2. MS(EI, 70eV) *m/z* C₁₅H₁₂ (% relative intensity, ion): 192 (100, [M]⁺), 165 (18), 139 (7), 139 (8), 115 (10).

1-Chloro-2-(phenylethynyl)-benzene (4d)²: Yield: Method A: 806 mg (3.8 mmol, 38%); Method B: 508 mg (2.4 mmol, 47 %), 763 mg (3.6 mmol, 71 %) yellow oil. ¹H NMR (CDCl₃; 250 MHz): δ 7.50-7.43 (m, 3H), 7.29-7.22 (m, 3H), 7.14-7.10 (m, 3H). ¹³C NMR (CDCl₃, 62.5 MHz): δ 135.8, 133.2, 131.7, 131.5, 129.22, 129.20, 128.6, 128.32, 128.29, 126.4, 123.1, 122.8, 94.5, 86.1 MS(EI, 70eV) *m/z* C₁₄H₉Cl (% relative intensity, ion): 212(100, [M]⁺), 153(11), 126(21).

3-Phenylethynyl-pyridine (4e)⁴: Yield: Method A: 1002 mg (5.6 mmol, 56%) yellow crystal. m.p.: 50-51 °C. ¹H NMR (CDCl₃; 250 MHz): δ 8.7 (d, 1H, *J* = 1.5 Hz), 8.49 (dd, 1H, *J* = 5.0, 1.5 Hz), 7.77 (dt, 1H, *J* = 1.75, 8.0 Hz), 7.50-7.46 (m, 2H), 7.33-7.28 (m, 3H), 7.25-7.19 (m, 1H). ¹³C NMR (CDCl₃, 62.5 MHz): δ 151.9, 148.3, 138.6, 131.7,

128.8, 128.4, 123.1, 122.4, 92.8, 85.8 MS(EI, 70eV) m/z $C_{13}H_9N$ (% relative intensity, ion): 179(100, $[M]^+$), 153(11), 126(21).

1-Methoxy-4-(4'-methoxy-phenylethynyl)-benzene (4f)¹: Yield: Method A: 603.2 mg (2.9 mmol, 29%); Method B: 785 mg (3.3 mmol, 66 %), yellow crystal. m.p.: 140-142 °C. 1H NMR ($CDCl_3$; 250 MHz): δ 7.39-7.36 (m, 4H), 6.89-6.69 (m, 4H) 3.74 (s, 6H). ^{13}C NMR ($CDCl_3$, 62.5 MHz): δ 159.3, 133.7, 114.8, 113.1, 88.0, 56.4 MS(EI, 70eV) m/z $C_{16}H_{14}O_2$ (% relative intensity, ion): 238(100), $[M]^+$, 223(84), 180(16), 151(17), 119(20).

1-Methyl-3-(4'-methoxyphenyl)-benzene (4g): Yield: Method B: 877 mg (3.95 mmol, 79%), 733 mg (3.3 mmol, 66%) yellow crystal. m.p.:43-45°C. 1H NMR ($CDCl_3$; 250 MHz): δ 7.47 (d, 2H, J = 8.75 Hz), 7.33 (d, 2H, J = 8.5 Hz), 7.21 (t, 1H, J = 6.75 Hz), 7.13 (d, 1H, J = 7.5 Hz), 6.88 (d, 2H, J = 8.75 Hz), 3.80 (s, 3H), 2.34 (s, 3H). ^{13}C NMR ($CDCl_3$, 62.5 MHz): δ 159.5, 137.9, 133.0, 132.0, 128.8, 128.5, 128.2, 123.4, 115.5, 114.0, 89.0, 88.2, 55.2, 21.2. MS(EI, 70eV) m/z $C_{16}H_{14}O$ (% relative intensity, ion): 222(100, $[M]^+$), 207(48), 178(31), 151(9).

1-Methoxy-4-(2'-chloro- phenylethynyl)-benzene (4h): Yield: Method A: 1645 mg (6.8 mmol, 68%); Method B: 509 mg (2.1 mmol, 41 %), 921 mg (3.8 mmol, 75 %) yellow crystal. m.p.: 52-54 °C. ^1H NMR (CDCl_3 ; 250 MHz): δ 7.45-7.41 (m, 3H), 7.35 (d, 1H, $J = 9.5$ Hz), 7.16 (d, 2H, $J = 9.5$ Hz), 6.82 (d, 2H, $J = 8.75$ Hz) 3.74 (s, 3H). ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 159.9, 133.2, 133.0, 129.2, 128.9, 127.1, 126.4, 123.5, 114.9, 114.0, 94.6, 84.9, 55.3. MS(EI, 70eV) m/z $\text{C}_{15}\text{H}_{11}\text{OCl}$ (% relative intensity, ion): 244 ((31), $[\text{M}]^+$), 242 ((100), $[\text{M}]^+$), 226(59), 198 (47). Analysis ($\text{C}_{15}\text{H}_{11}\text{ClO}$) calculated: C, 74.23; H, 4.57; Found: C, 74.11; H, 4.40.

1-Methoxy-4-(3'-pyridynethynyl)-benzene (4i): Yield: Method A: 1359 mg (6.5 mmol, 65%) yellow crystal. m.p.: 46-48°C. ^1H NMR (CDCl_3 ; 250 MHz): δ 8.67 (s, 1H), 8.44 (d, 1H, $J = 3.75$ Hz), 7.71 (dt, 1H, $J = 8.0, 2.0$ Hz) 7.43-7.37 (m, 2H), 7.20-7.17 (m, 1H), 6.83-6.79 (m, 2H). ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 159.9, 151.9, 148.1, 138.2, 133.1, 123.0, 114.4, 114.0, 92.7, 84.6, 55.2. MS(EI, 70eV) m/z $\text{C}_{14}\text{H}_{11}\text{NO}$ (% relative intensity, ion): 209 (100), $[\text{M}]^+$, 194 (91), 166 (46), 139 (60), 113 (25). Analysis ($\text{C}_{14}\text{H}_{11}\text{NO}$) calculated: C, 80.36; H, 5.30; N, 6.69. Found: C, 80.45; H, 5.29; N, 6.84.

1-Methyl-3-(3'-methylphenyl)-benzene (4j)⁵: Yield: Method B: 803 mg (3.9 mmol, 78%) white crystal. m.p.: 57-59°C. ^1H NMR (CDCl_3 ; 250 MHz): δ 7.34 (d, 4H, $J = 7.75$ Hz), 7.20 (t, 2H, $J = 7.5$ Hz), 7.12 (d, 2H, $J = 7.5$ Hz), 2.32 (s, 6H). ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 137.9, 132.1, 129.0, 128.6, 128.2, 123.1, 89.2, 21.2 MS(EI, 70eV) m/z $\text{C}_{16}\text{H}_{14}$ (% relative intensity, ion): 206 (100, $[\text{M}]^+$), 189 (25), 178 (10), 165 (9), 139 (8).

1-Methyl-3-(2'-chloro-phenylethynyl)-benzene (4k): Yield: Method B: 727 mg (2.95 mmol, 59%), 493 mg (2.0 mmol, 40%) yellow crystal. m.p.: 39-41°C. ^1H NMR (CDCl_3 ; 250 MHz): δ 7.53 (t, 1H, $J = 5.0$ Hz), 7.42-7.36 (m, 3H), 7.26-7.12 (m, 4H), 2.33 (s, 3H). ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 138.0, 135.8, 133.1, 132.2, 129.5, 129.2, 128.8, 126.3, 123.2, 122.6, 94.7, 85.8, 21.2. MS(EI, 70eV) m/z $\text{C}_{15}\text{H}_{11}\text{Cl}$ (% relative intensity, ion): 228 (25, $[\text{M}]^+$), 226 (100, $[\text{M}]^+$), 189 (50), 165 (11), 113 (10), 94 (18).

1-Chloro-2-(2'-chlorophenyl-ethynyl)-benzene (4l): Yield: Method A: 1641 mg (6.7 mmol, 67%); Method B: 395 mg (1.6 mmol, 32%) yellow crystal. m.p.: 83-85°C. ^1H NMR (CDCl_3 ; 250 MHz): δ 7.54-7.51 (m, 2H), 7.38-7.34 (m, 2H), 7.21-7.16 (m, 4H). ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 136.0, 133.4, 129.6, 129.3, 126.4, 122.8, 91.1. MS(EI, 70eV) m/z $\text{C}_{14}\text{H}_8\text{Cl}_2$ (% relative intensity, ion): 248 (30, $[\text{M}]^+$), 245 (100, $[\text{M}]^+$), 211 (7), 209 (13), 176 (15), 175 (17), 174 (10), 150 (119), 123 (16).

2-(3'-Pyridylethynyl)-chlorobenzene (4m): Yield: Method A: 1320 mg (6.2 mmol, 62%) yellow oil. ^1H NMR (CDCl_3 ; 250 MHz): δ 8.7 (d, 1H, $J = 1.75$ Hz), 8.46 (dd, 1H, $J = 5.0, 1.75$ Hz), 7.74 (dt, 1H, $J = 7.25, 2.5$ Hz), 7.47-7.43 (m, 1H), 7.34-7.30 (m, 1H), 7.20-7.10 (m, 3H). ^{13}C NMR (CDCl_3 , 62.5 MHz): δ 152.0, 148.7, 138.4, 135.8, 133.1, 129.7,

129.2, 126.4, 122.9, 122.3, 119.9, 90.8, 89.2. MS(EI, 70eV) m/z $C_{13}H_8ClN$ (% relative intensity, ion): 215 ((32), $[M]^+$), 213 ((100), $[M]^+$), 178 (23), 151 (29), 150 (17), 125 (20). Analysis ($C_{13}H_8ClN$) calculated: C, 73.08; H, 3.77; N, 6.56; Found: C, 73.21; H, 3.73; N, 6.32;

3-(3'-Pyridylethynyl)-pyridine (4n)⁶: Yield: Method A: 1512 mg (8.4 mmol, 84%) yellow crystal. m.p.: °C. 1H NMR ($CDCl_3$; 250 MHz): δ 8.72 (d, 2H, $J = 1.75$ Hz), 8.53 (dd, 2H, $J = 5.0, 1.75$ Hz), 7.79 (dt, 2H, $J = 7.25, 2.5$ Hz) 7.28-7.22 (m, 2H). ^{13}C NMR ($CDCl_3$, 62.5 MHz): δ 152.1, 148.9, 138.6, 123.1, 119.7, 89.1. MS(EI, 70eV) m/z $C_{12}H_8N_2$ (% relative intensity, ion): 180 ((100), $[M]^+$), 152 (15), 127 (11), 100 (18).

References

- 1 Mio, M. J.; Kopel, L. C.; Braun, J. B.; Gidzikwa, T. L.; Hull, K. L.; Brisbois, R. G.; Markworth, C. J.; Grieco, P. A.; *Org. Lett.* **2002**, 4, 3199.
- 2 Mouriés, V.; Waschbüsch, R.; Carran, J.; Savignac, P.; *Synthesis* **1998**, 271.
- 3 Eberhard, M. R.; Wang, Z.; Jensen, C. M.; *Chem. Comm.* **2002**, (8), 818.
- 4 Mori, Y.; Seki, M.; *J. Org. Chem.* **2003**, 68, 1517.
- 5 Newkome, G. R.; Roper, J. M.; Robinson, J. M.; *J. Org. Chem.* **1980**, 45, 4380.
- 6 Negishi, E.-I.; Caiding, X.; Ze, T.; Martin, K. *Heterocycles* **1997**, 46, 209.

25

7.339
7.308
7.233
7.203
7.172
7.118
7.089

2.320

6.0

3.9

2.0.1

7.548
7.537
7.530
7.510
7.429
7.416
7.389
7.359
7.256
7.243
7.224
7.213
7.202
7.194
7.184
7.152
7.123

2.333

30

4.0

3.0

1.0

3.0