Supporting Information

Belonging the manuscript

The first 1-alkyl-3-perfluoroalkyl-4, 5-dimethyl-1, 2, 4-triazolium salts

Hong Xue, Brendan Twamley, and Jean'ne M. Shreeve*

Department of Chemistry, University of Idaho, Moscow, Idaho 83843-2343, U.S.A.

jshreeve@uidaho.edu

Table of Contents

S-2 General methods.

S-3 Figure 1. ORTEP diagram for **11c**.

S-4 Table 1. Crystal data and structure refinement for **11c**.

S-5 Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for **11c**.

S-7 Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for **11c**.

S-9 Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for **11c**.

S-11 Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for **11c**.

S-12 Table 6. Intermolecular close contact lengths [$\text{\AA}$] for **11c**.

General Methods. ^1H , ^{13}C and ^{19}F NMR spectra were recorded on a 300 MHz spectrometer operating at 300.13, 75.48, 282.37 MHz respectively, chemical shifts were reported relative to Me_4Si or CFCl_3 . Acetone- d_6 was the solvent unless otherwise specified. GC/MS spectra were determined using an appropriate instrument. Mass spectra for ionic salts were determined by insertion using solid probe. M^+ is the mass of the cation. The melting and decomposition points were recorded on a differential scanning calorimeter at a scan rate of 10 $^\circ\text{C}/\text{min}$. IR spectra were recorded using NaCl plates for neat liquids and KBr pellets for solids. Elemental analyses were performed by professional analysts.

Figure 1. ORTEP drawing of compound **11c**

Table 1. Crystal data and structure refinement for [C₈F₁₇-Me₃-triazole][BF₄]

Identification code	1,4,5-Trimethyl-3-perfluorooctyl-1,2,4-triazolium tetrafluoroborate (11c)	
Empirical formula	C ₁₃ H ₉ F ₂₁ N ₃ B	
Formula weight	617.04	
Temperature	84(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2(1)/c	
Unit cell dimensions	a = 25.843(5) Å	α = 90°.
	b = 6.3886(13) Å	β = 90.20(3)°.
	c = 12.629(3) Å	γ = 90°.
Volume	2085.0(8) Å ³	
Z	4	
Density (calculated)	1.966 Mg/m ³	
Absorption coefficient	0.250 mm ⁻¹	
F(000)	1208	
Crystal size	0.61 x 0.18 x 0.02 mm ³	
Crystal color and habit	colorless plate	
Diffractionmeter	Bruker/Siemens SMART APEX	
Theta range for data collection	2.36 to 25.00°.	
Index ranges	-30 ≤ h ≤ 30, -7 ≤ k ≤ 7, -15 ≤ l ≤ 15	
Reflections collected	20320	
Independent reflections	3675 [R(int) = 0.0460]	
Completeness to theta = 25.00°	99.8 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.9950 and 0.8626	
Solution method	XS, Bruker SHELXTL v. 6.12	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	3675 / 0 / 346	
Goodness-of-fit on F ²	1.078	
Final R indices [I > 2σ(I)]	R1 = 0.0471, wR2 = 0.1070	
R indices (all data)	R1 = 0.0592, wR2 = 0.1132	
Largest diff. peak and hole	0.346 and -0.320 e.Å ⁻³	

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for **11c**. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
B(1)	5899(1)	9786(5)	3758(3)	19(1)
C(3)	3994(1)	6824(4)	3926(2)	16(1)
C(5)	3211(1)	7601(4)	3466(2)	16(1)
C(6)	2641(1)	7495(4)	3257(2)	19(1)
C(7)	2511(1)	6263(4)	2233(2)	18(1)
C(8)	1942(1)	6368(4)	1861(2)	18(1)
C(9)	1798(1)	4611(4)	1053(2)	18(1)
C(10)	1304(1)	5080(4)	395(2)	18(1)
C(11)	1079(1)	3134(4)	-185(2)	20(1)
C(12)	716(1)	3685(5)	-1124(2)	22(1)
C(13)	365(1)	1867(5)	-1493(2)	26(1)
C(21)	4419(1)	9965(4)	3138(2)	22(1)
C(31)	4460(1)	5795(4)	4343(2)	20(1)
C(41)	3345(1)	4196(4)	4517(3)	24(1)
F(1)	2452(1)	9460(2)	3141(1)	24(1)
F(2)	2396(1)	6590(3)	4083(1)	25(1)
F(3)	2812(1)	7023(3)	1458(1)	24(1)
F(4)	2641(1)	4251(2)	2410(1)	24(1)
F(5)	1862(1)	8234(2)	1396(1)	25(1)
F(6)	1628(1)	6207(3)	2696(1)	25(1)
F(7)	2195(1)	4342(3)	385(1)	23(1)
F(8)	1723(1)	2826(2)	1593(1)	24(1)
F(9)	1416(1)	6539(3)	-333(1)	25(1)
F(10)	940(1)	5846(3)	1044(1)	23(1)
F(11)	812(1)	1985(3)	521(1)	28(1)
F(12)	1469(1)	1967(3)	-554(2)	32(1)
F(13)	408(1)	5284(3)	-840(1)	26(1)
F(14)	1004(1)	4286(3)	-1944(1)	34(1)
F(15)	155(1)	2360(3)	-2420(1)	38(1)
F(16)	-12(1)	1526(3)	-814(2)	31(1)

F(17)	640(1)	133(3)	-1604(2)	39(1)
F(18)	5586(1)	8158(3)	4134(2)	32(1)
F(19)	5586(1)	11520(3)	3558(2)	40(1)
F(20)	6269(1)	10243(3)	4509(1)	36(1)
F(21)	6133(1)	9159(3)	2829(1)	27(1)
N(1)	3493(1)	9117(3)	3088(2)	18(1)
N(2)	3980(1)	8600(3)	3382(2)	16(1)
N(4)	3504(1)	6155(3)	3997(2)	16(1)

Table 3. Bond lengths [Å] and angles [°] for **11c**.

B(1)-F(20)	1.377(4)	C(13)-F(15)	1.326(4)
B(1)-F(21)	1.382(4)	C(21)-N(2)	1.464(4)
B(1)-F(19)	1.394(4)	C(21)-H(21A)	0.9800
B(1)-F(18)	1.401(4)	C(21)-H(21B)	0.9800
C(3)-N(2)	1.327(3)	C(21)-H(21C)	0.9800
C(3)-N(4)	1.340(4)	C(31)-H(31A)	0.9800
C(3)-C(31)	1.469(4)	C(31)-H(31B)	0.9800
C(5)-N(1)	1.303(4)	C(31)-H(31C)	0.9800
C(5)-N(4)	1.368(4)	C(41)-N(4)	1.473(3)
C(5)-C(6)	1.498(4)	C(41)-H(41A)	0.9800
C(6)-F(2)	1.352(3)	C(41)-H(41B)	0.9800
C(6)-F(1)	1.355(3)	C(41)-H(41C)	0.9800
C(6)-C(7)	1.550(4)	N(1)-N(2)	1.352(3)
C(7)-F(3)	1.344(3)		
C(7)-F(4)	1.347(3)	F(20)-B(1)-F(21)	109.8(3)
C(7)-C(8)	1.542(4)	F(20)-B(1)-F(19)	111.0(2)
C(8)-F(6)	1.337(3)	F(21)-B(1)-F(19)	109.4(2)
C(8)-F(5)	1.345(3)	F(20)-B(1)-F(18)	109.0(2)
C(8)-C(9)	1.561(4)	F(21)-B(1)-F(18)	109.1(2)
C(9)-F(7)	1.342(3)	F(19)-B(1)-F(18)	108.5(3)
C(9)-F(8)	1.343(3)	N(2)-C(3)-N(4)	106.5(2)
C(9)-C(10)	1.551(4)	N(2)-C(3)-C(31)	126.1(3)
C(10)-F(9)	1.342(3)	N(4)-C(3)-C(31)	127.4(2)
C(10)-F(10)	1.342(3)	N(1)-C(5)-N(4)	111.9(2)
C(10)-C(11)	1.554(4)	N(1)-C(5)-C(6)	121.3(2)
C(11)-F(12)	1.339(3)	N(4)-C(5)-C(6)	126.7(2)
C(11)-F(11)	1.347(3)	F(2)-C(6)-F(1)	108.1(2)
C(11)-C(12)	1.551(4)	F(2)-C(6)-C(5)	110.3(2)
C(12)-F(14)	1.333(3)	F(1)-C(6)-C(5)	109.4(2)
C(12)-F(13)	1.345(3)	F(2)-C(6)-C(7)	108.9(2)
C(12)-C(13)	1.545(4)	F(1)-C(6)-C(7)	107.6(2)
C(13)-F(16)	1.318(4)	C(5)-C(6)-C(7)	112.4(2)
C(13)-F(17)	1.324(4)	F(3)-C(7)-F(4)	108.7(2)

F(3)-C(7)-C(8)	108.5(2)	F(13)-C(12)-C(11)	109.0(2)
F(4)-C(7)-C(8)	109.2(2)	C(13)-C(12)-C(11)	114.4(2)
F(3)-C(7)-C(6)	107.4(2)	F(16)-C(13)-F(17)	109.1(3)
F(4)-C(7)-C(6)	107.0(2)	F(16)-C(13)-F(15)	108.3(2)
C(8)-C(7)-C(6)	115.9(2)	F(17)-C(13)-F(15)	108.8(2)
F(6)-C(8)-F(5)	108.6(2)	F(16)-C(13)-C(12)	111.2(2)
F(6)-C(8)-C(7)	109.7(2)	F(17)-C(13)-C(12)	110.3(2)
F(5)-C(8)-C(7)	108.5(2)	F(15)-C(13)-C(12)	109.0(3)
F(6)-C(8)-C(9)	108.4(2)	N(2)-C(21)-H(21A)	109.5
F(5)-C(8)-C(9)	108.4(2)	N(2)-C(21)-H(21B)	109.5
C(7)-C(8)-C(9)	113.1(2)	H(21A)-C(21)-H(21B)	109.5
F(7)-C(9)-F(8)	108.8(2)	N(2)-C(21)-H(21C)	109.5
F(7)-C(9)-C(10)	108.5(2)	H(21A)-C(21)-H(21C)	109.5
F(8)-C(9)-C(10)	108.4(2)	H(21B)-C(21)-H(21C)	109.5
F(7)-C(9)-C(8)	108.8(2)	C(3)-C(31)-H(31A)	109.5
F(8)-C(9)-C(8)	108.2(2)	C(3)-C(31)-H(31B)	109.5
C(10)-C(9)-C(8)	114.0(2)	H(31A)-C(31)-H(31B)	109.5
F(9)-C(10)-F(10)	108.6(2)	C(3)-C(31)-H(31C)	109.5
F(9)-C(10)-C(9)	108.7(2)	H(31A)-C(31)-H(31C)	109.5
F(10)-C(10)-C(9)	108.7(2)	H(31B)-C(31)-H(31C)	109.5
F(9)-C(10)-C(11)	108.4(2)	N(4)-C(41)-H(41A)	109.5
F(10)-C(10)-C(11)	108.5(2)	N(4)-C(41)-H(41B)	109.5
C(9)-C(10)-C(11)	113.8(2)	H(41A)-C(41)-H(41B)	109.5
F(12)-C(11)-F(11)	108.4(2)	N(4)-C(41)-H(41C)	109.5
F(12)-C(11)-C(12)	108.3(2)	H(41A)-C(41)-H(41C)	109.5
F(11)-C(11)-C(12)	108.7(2)	H(41B)-C(41)-H(41C)	109.5
F(12)-C(11)-C(10)	109.2(2)	C(5)-N(1)-N(2)	103.8(2)
F(11)-C(11)-C(10)	108.4(2)	C(3)-N(2)-N(1)	112.0(2)
C(12)-C(11)-C(10)	113.7(2)	C(3)-N(2)-C(21)	126.8(2)
F(14)-C(12)-F(13)	108.8(2)	N(1)-N(2)-C(21)	121.2(2)
F(14)-C(12)-C(13)	108.1(2)	C(3)-N(4)-C(5)	105.8(2)
F(13)-C(12)-C(13)	107.7(2)	C(3)-N(4)-C(41)	124.5(2)
F(14)-C(12)-C(11)	108.8(2)	C(5)-N(4)-C(41)	129.6(2)

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for **11c**. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
B(1)	28(2)	13(2)	16(2)	3(1)	1(1)	-5(1)
C(3)	25(2)	12(1)	11(1)	-4(1)	0(1)	1(1)
C(5)	26(2)	10(1)	12(1)	-4(1)	0(1)	1(1)
C(6)	22(2)	16(1)	18(1)	-2(1)	2(1)	1(1)
C(7)	23(2)	13(1)	18(2)	1(1)	3(1)	2(1)
C(8)	22(2)	14(1)	17(1)	-2(1)	3(1)	2(1)
C(9)	22(2)	14(1)	18(2)	1(1)	3(1)	0(1)
C(10)	21(1)	19(1)	15(1)	2(1)	2(1)	-1(1)
C(11)	19(2)	20(2)	19(2)	1(1)	1(1)	-1(1)
C(12)	24(2)	25(2)	16(1)	-2(1)	2(1)	1(1)
C(13)	26(2)	28(2)	23(2)	-4(1)	-5(1)	3(1)
C(21)	24(2)	15(1)	26(2)	2(1)	-2(1)	-3(1)
C(31)	23(2)	13(1)	24(2)	2(1)	-2(1)	1(1)
C(41)	29(2)	14(1)	29(2)	7(1)	1(1)	-3(1)
F(1)	26(1)	18(1)	29(1)	-5(1)	-5(1)	7(1)
F(2)	24(1)	34(1)	18(1)	0(1)	4(1)	-1(1)
F(3)	23(1)	31(1)	16(1)	-4(1)	4(1)	-7(1)
F(4)	27(1)	15(1)	29(1)	-4(1)	-7(1)	5(1)
F(5)	32(1)	13(1)	31(1)	0(1)	-9(1)	3(1)
F(6)	22(1)	34(1)	18(1)	-6(1)	3(1)	-1(1)
F(7)	20(1)	27(1)	23(1)	-8(1)	2(1)	2(1)
F(8)	31(1)	14(1)	26(1)	5(1)	-8(1)	-3(1)
F(9)	28(1)	23(1)	22(1)	7(1)	-4(1)	-6(1)
F(10)	19(1)	28(1)	22(1)	-5(1)	-1(1)	4(1)
F(11)	34(1)	25(1)	26(1)	8(1)	-9(1)	-11(1)
F(12)	26(1)	30(1)	40(1)	-18(1)	-7(1)	8(1)
F(13)	30(1)	23(1)	25(1)	1(1)	-6(1)	5(1)
F(14)	35(1)	49(1)	16(1)	1(1)	4(1)	-11(1)
F(15)	44(1)	46(1)	24(1)	-4(1)	-16(1)	-5(1)
F(16)	25(1)	35(1)	33(1)	-1(1)	-3(1)	-5(1)

F(17)	32(1)	30(1)	55(1)	-20(1)	-12(1)	7(1)
F(18)	34(1)	27(1)	37(1)	14(1)	2(1)	-8(1)
F(19)	51(1)	22(1)	46(1)	10(1)	11(1)	9(1)
F(20)	43(1)	39(1)	24(1)	-11(1)	-1(1)	-13(1)
F(21)	43(1)	23(1)	16(1)	-2(1)	3(1)	-3(1)
N(1)	24(1)	13(1)	17(1)	-2(1)	-3(1)	2(1)
N(2)	21(1)	10(1)	17(1)	-2(1)	-2(1)	-2(1)
N(4)	21(1)	11(1)	15(1)	-1(1)	0(1)	0(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for **11c**.

	x	y	z	U(eq)
H(21A)	4551	10592	3793	32
H(21B)	4306	11074	2653	32
H(21C)	4693	9141	2803	32
H(31A)	4550	4606	3889	31
H(31B)	4394	5297	5064	31
H(31C)	4748	6797	4354	31
H(41A)	3284	3117	3980	36
H(41B)	3026	4436	4918	36
H(41C)	3619	3731	5001	36

Table 6. Intermolecular close contact lengths [$\text{\AA}$] for **11c**.

C(3) – F(20)#1	2.809(3)
C(3) – F(21)#2	2.812(3)
N(2) – F(19)#2	3.007(3)
N(2) – F(20)#1	2.840(3)
N(4) – F(21)#2	2.800(3)

Symmetry transformations used to generate equivalent atoms:

#1 1-x,2-y,1-z #2 1-x, -0.5+y, 0.5-z