The First Non-thiolic, Odorless 1,3-Propanedithiol Equivalent and Its Application in Thioacetalization

Qun Liu, Guangbo Che, Haifeng Yu, Yingchun Liu,*

Jingping Zhang, Qian Zhang and Dewen Dong

Department of Chemistry, Northeast Normal University, Changchun, 130024, P. R. China

liuqun@nenu.edu.cn

Supporting Information

Table of contents

Materials and Reagents	S2
Instrumentation	S2
Synthesis and analytical data of compound 1	S2
Synthesis and analytical data of compound 4	S3
Analytical data of intermediates 2 , 5i , 6 , 7ai	S3
¹ H NMR spectra for compounds 1 , 4d , 5i , 6 and 7ai	S4-S8
X-ray crystal structure of 4d	S9

Materials and Reagents. All reagents were purchased from the Aldrich Chemical Co. and used without further treatment unless otherwise noted. All solvents were purchased from Tianjin Chemical Co. and silica gel (300-400 mesh) was purchased from Qingdao Ocean Chemical Co. Absolute solvents, such as methanol was prepared by distillation from sodium and 99% methanol was obtained by adding water into absolute methanol.

Instrumentation. Infrared (IR) spectra were obtained on a Magna-560 FT-IR spectrometer and were recorded as KBr mulls. ^1H -NMR spectra were obtained on a Unity-400 (400 MHz) spectrometer unless otherwise noted. ^1H -NMR spectra were measured in CDCl_3 and chemical shifts are reported in ppm. Mass spectra were recorded on a HP GC 6890/MS 5973 mass spectrometer.

Synthesis and analytical data for compound 1. The Vilsmeier reagent was prepared by mixing ice cold DMF (10 ml) and POCl_3 (0.62 ml, 5.0 mmol) and then kept at 0 °C until further use. Compound **1** was synthesized by chlorination of 3-(1,3)-dithian-ylidene-pentane-2, 4-dione **2** with Vilsmeier reagent. To the above Vilsmeier reagent in a 100 mL three-necked flask equipped with condenser, N_2 inlet and outlet, a solution of **2** (1.1g, 5.0 mmol) in DMF (2 mL) was added. The mixture was stirred under N_2 at room temperature for 4 hrs. Then the mixture was poured over ice (5 g) and the pH was adjusted to approximate 8.0 by adding sodium hydroxide solution (1.0 N). The crude product precipitated from the solution as colourless crystal in 99% yield, mp: 79-81 °C. ^1H -NMR (ppm): 2.20 (m, 2 H), 2.98 (t, J = 6.4 Hz, 4 H), 5.35 (d, J = 2.0 Hz, 2 H), 5.71 (d, J = 2.0 Hz,

2 H). IR (KBr, cm^{-1}) 2927, 1616, 1508. EIMS (20 *ev*) *m/z*: 253 (M^+). Calculated for $\text{C}_9\text{H}_{10}\text{Cl}_2\text{S}_2$: C, 42.69, H3.98; Found, C, 42.61, H, 3.93.

Synthesis and analytical data for compound 4. The dithiane **4** were prepared by the thioacetalization reaction between **1** and aldehydes/ketones **3** via a simple procedure. The preparation of dithiane **4d** was described below as an example. Piperonal **3d** (2.0 mmol), compound **1** (2.2 mmol) and methanol (99%, 15 mL) were added into a flask equipped with a condenser, N_2 inlet and outlet. The mixture was heated at reflux temperature and stirred under N_2 . Once the aldehyde was consumed, as indicated by TLC, the resulting mixture was worked up and separated *via* silica gel chromatography (eluent: petroleum ether) to afford **4d** 466 mg (97% yield). With the same precusure, compounds **4a-4q** were synthesized with yields ranging from 79% to 97%. All of the dithianes **4a-4q** are known compounds and were identified by ^1H -NMR, IR spectral and m.p. (for all solids). All the analytical data for **4a-q** were in good agreement with that in the literature.

Analytical data for intermediates 2, 5i, 6, 7ai

3-(1,3)-dithian-ylidene-pentane-2,4-dione **2** and Dimethylacetal **5i** are known compounds. All the analytical data for **2** and **5i** are in good agreement with that in literature.

Dithio orthoester **6**. Colorless viscous liquid. ^1H -NMR: 1.84 (s, 3H), 1.87 (m, 1H), 2.09 (m, 1H), 2.72 (m, 1H), 2.76 (m, 1H), 3.01 (m, 1H), 3.04 (m, 1H), 3.05 (s, 2H), 3.70 (s, 3H). IR: (KBr, cm^{-1}) 1736, 1435, 1328, 1278, 1199, 1076, 1010, 908. EIMS (20 *ev*) *m/z*: 206 (M^+). Calculated for $\text{C}_8\text{H}_{14}\text{O}_2\text{S}_2$: C, 46.57, H, 6.84; Found, C, 46.70, H, 6.79.

Monothioacetal **7ai**. Colorless viscous liquid. **7ai** is unstable and quickly transfers to **4i**.

^1H -NMR: 1.77 (s, 3H), 1.83 (m, 1H), 2.05 (m, 1H), 2.75 (m, 2H), 3.03 (m, 2H), 3.05 (s, 2H), 3.34 (s, 6H), 3.70 (s, 3H), 5.48 (s, 1H), 7.65 (d, $J = 8.8\text{Hz}$, 2H), 8.23 (d, $J = 8.8\text{Hz}$, 2H). IR (KBr, cm^{-1}): 1736, 1609, 1525, 1437, 1345.

1

6

7ai

X-ray crystal structure of **4d** with $F(000) = 1008$, $R_1 = 0.0793$, $wR_2 = 0.2526$.