The Oxidative Mannich Reaction Catalyzed by Dirhodium Caprolactamate

Arthur J. Catino, Jason M. Nichols, Brian J. Nettles, and Michael P. Doyle

Department of Chemistry and Biochemistry
The University of Maryland
College Park, MD 20742

Supporting Information
(17 pages)

General. All products were characterized and in agreement with those previously described. Yields reported are for isolated yields unless otherwise noted. *tert*-Butyl hydroperoxide (T-HYDRO®, 70 wt. % in H₂O) was obtained from Aldrich and used as received. The preparation of dirhodium(II) caprolactamate [Rh₂(cap)₄·2CH₃CN] has been previously described.¹ 2-Triisopropylsiloxyfurans were prepared according to the procedure of Martin.² 5-Allyl-2-triisopropylsiloxyfuran was prepared according to the procedure of Liras.³ Amines (Table 2, entries 2, 3, and 7) were prepared via reductive amination with NaCNBH₃-ZnCl₂ according to the procedure of Kim, *et al.*⁴ 2-*N,N*-Dimethylaminostyrene was prepared via Wittig reaction from the known 2-*N,N*-dimethylaminobenzaldehyde.⁵ *N*-Ethyl-*N*-methylaniline (Table 2, entry 6) was obtained from the NaBH₄-I₂ reduction of *N*-methylacetanilide.⁶ *N*-Phenyl-1,2,3,4-tetrahydroisoquinoline was prepared according to the procedure of Quach.⁷ Preparative chromatographic purification was performed using SiliCycle (60Å, 40-63 mesh) silica gel according to the method of Still.⁸ Thin layer chromatography (TLC) was performed on Merck 0.25 mm silica gel 60 F₂₅₄ plates with visualization by fluorescence quenching. ¹H NMR (400 MHz) and ¹³C NMR (100 MHz) spectra were obtained on a Bruker DRX-400 NMR as solutions in CDCl₃. Chemical shifts are reported in parts per million (ppm), δ downfield from Me₄Si (TMS); coupling constants are reported in Hertz (Hz).

Representative Procedure. T-HYDRO® (1.2 equiv) was added in one portion to a stirring solution of amine (2.0 equiv), 2-triisopropoxysilylfuran (1.0 equiv), and Rh₂(cap)₄ (1.0 mol%) in MeOH (0.27 M/[siloxyfuran]). The reaction mixture was heated at 60 °C for 3-5 hr (or 16 hr with 0.1 mol % catalyst). The solvent was then evaporated and the product was purified directly on silica gel.

¹ Doyle, M. P.; Westrum, L. J.; Wolthuis, W. N. E.; See, M. M.; Boone, W. P.; Bagheri, V.; Pearson, M. M. *J. Am. Chem. Soc.* **1993**, *115*, 958.

² Bur, S. K.; Martin, S. F. *Tetrahedron* **2001**, *57*, 3221 and references therein.

³ Liras, S.; Davoren, J. E.; Bordner, J. *Org. Lett.* **2001**, *3*, 703.

⁴ Kim, S.; Oh, C. H.; Ko, J. S.; Ahn, K. H.; Kim, Y. J. *J. Org. Chem.* **1985**, *50*, 1927.

⁵ Stanetty, P.; Rodler, I. K.; Krumpak, B. *J. Prakt. Chem.* **1994**, *336*, 333.

⁶ Prasad, A. S. B.; Kanth, J. V. B.; Periasamy, M. *Tetrahedron* **1992**, *48*, 4623.

⁷ Quach, T. D.; Batey, R. A. *Org. Lett.* **2003**, *5*, 4397.

⁸ Still, W. C.; Kahn, M.; Mitra, A. J. *J. Org. Chem.* **1978**, *43*, 2923.

Scheme 1, **4**: Purified by chromatography on silica gel (5:1 → 1:1 hexanes/EtOAc); orange oil: TLC R_f = 0.30 (1:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.47 (dd, J = 5.8, 1.5 Hz, 1H), 7.25 (t, J = 8.0 Hz, 2H), 6.77 (t, J = 7.3 Hz, 1H), 6.72 (d, J = 8.0 Hz, 2H), 6.13 (dd, J = 5.6, 2.1 Hz, 1H), 5.27 (tt, J = 5.8, 1.8 Hz, 1H), 3.69 (d, J = 6.0 Hz, 2H), 3.02 (s, 3H); ^{13}C NMR (100 MHz) δ 172.6, 154.4, 148.2, 129.4, 122.2, 117.4, 112.3, 81.9, 55.0, 39.5; IR (neat) 1755 (C=O) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{12}\text{H}_{14}\text{NO}_2$ 204.1025, found 204.1026 ($\text{M}+\text{H}$).

Table 2, entry 1 (R = CH₃): Purified by chromatography on silica gel (5:1 → 1:1 hexanes/EtOAc); orange oil: TLC R_f = 0.50 (1:1 hexanes/EtOAc); ¹H NMR (400 MHz, CDCl₃) δ 7.47 (d, J = 5.6 Hz, 1H), 7.06 (d, J = 7.4 Hz, 2H), 6.64 (d, J = 8.1, 2H), 6.11 (d, J = 5.4 Hz, 1H), 5.26 – 5.24 (m, 1H), 3.64 (d, J = 5.8 Hz, 2H), 2.99 (s, 3H), 2.26 (s, 3H); ¹³C NMR (100 MHz) δ 172.6, 154.5, 146.2, 129.9, 126.7, 122.0, 112.6, 82.0, 55.3, 39.6, 20.1; IR (neat) 1751 (C=O) cm⁻¹; HRMS (EI) calcd for C₁₃H₁₅NO₂ 218.1181, found 218.1176 (M+H).

Table 2, entry 1 (R = *t*-Bu): Purified by chromatography on silica gel (5:1 → 1:1 hexanes/EtOAc); orange oil: TLC R_f = 0.25 (2:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.48 (d, J = 5.6 Hz, 1H), 7.26 (d, J = 8.7 Hz, 2H), 6.66 (d, J = 8.7 Hz, 2H), 6.10 - 6.08 (m, 1H), 5.25 - 5.22 (m, 1H), 3.67 - 3.56 (m, 2H), 2.97 (s, 3H), 1.28 (s, 9H); ^{13}C NMR (100 MHz) δ 172.5, 154.6, 145.9, 139.8, 125.9, 121.8, 111.9, 81.9, 55.1, 39.3, 33.5, 31.3; IR (neat) 1756 (C=O) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{16}\text{H}_{22}\text{NO}_2$ 260.1651, found 260.1646 (M+H).

Table 2, entry 1 (R = Br): Purified by chromatography on silica gel (5:1 → 1:1 hexanes/EtOAc); yellow oil: TLC R_f = 0.22 (2:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.45 (dd, J = 5.8, 1.3 Hz, 1H), 7.32 (d, J = 8.9 Hz, 2H), 6.58 (d, J = 8.9 Hz, 2H), 6.14 (dd, J = 5.7, 1.9 Hz, 1H), 5.26 – 5.23 (m, 1H), 3.70 (dd, J = 15.3, 5.6 Hz, 1H), 3.63 (dd, J = 15.5, 6.0 Hz, 1H), 3.00 (s, 3H); ^{13}C NMR (100 MHz) δ 172.4, 153.9, 147.2, 132.0, 122.4, 113.9, 109.3, 81.8, 54.8, 39.6; IR (neat) 1755 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{12}\text{H}_{13}\text{BrNO}_2$ 282.0130, found 282.0130 ($\text{M}+\text{H}$).

Table 2, entry 2: Purified by chromatography on silica gel (5:1 $\rightarrow$ 1:1 hexanes/EtOAc); orange oil: TLC R_f = 0.14 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.48 (dd, J = 5.8, 1.6 Hz, 1H), 7.00 (d, J = 8.1 Hz, 1H), 6.54 – 6.54 (m, 1H), 6.50 – 6.47 (m, 1H), 6.12 (dd, J = 5.8, 1.6 Hz, 1H), 5.25 (tt, J = 5.8, 1.6 Hz, 1H), 3.69 – 3.58 (m, 2H), 2.94 (s, 3H), 2.24 (s, 3H), 2.17 (s, 3H); ^{13}C NMR (100 MHz) δ 172.6, 154.6, 145.6, 137.4, 130.3, 125.5, 121.9, 114.2, 110.1, 82.0, 55.3, 39.6, 20.3, 18.5; IR (neat) 1751 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{14}\text{H}_{18}\text{NO}_2$ 232.1338, found 232.1331 ($\text{M}+\text{H}$).

Table 2, entry 3: Purified by chromatography on silica gel (5:1 $\rightarrow$ 1:1 hexanes/EtOAc); orange oil: TLC R_f = 0.36 (1:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.47 (dd, J = 5.6, 1.6 Hz, 1H), 6.72 (d, J = 8.3 Hz, 1H), 6.38 (d, J = 2.6 Hz, 1H), 6.15 (d, J = 2.6 Hz, 1H), 6.15 – 6.13 (m, 1H), 5.89 (s, 2H), 5.24 (tt, J = 5.8 Hz, 1.6 Hz, 1H), 3.64 – 3.53 (m, 2H), 2.95 (s, 3H); ^{13}C NMR (100 MHz) δ 172.6, 154.4, 148.6, 144.6, 139.8, 122.1, 108.5, 105.1, 100.8, 96.2, 81.9, 56.3, 40.3; IR (neat) 1751 (C=O) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{14}\text{NO}_4$ 248.0923, found 248.0927 ($\text{M}+\text{H}$).

Table 2, entry 4: Purified by chromatography on silica gel (8:1 → 2:1 hexanes/EtOAc); bright yellow oil: TLC R_f = 0.35 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.48 (dd, J = 7.7, 1.5 Hz, 1H), 7.34 (dd, J = 5.7, 1.6 Hz, 1H), 7.28 – 7.24 (m, 1H), 7.14 – 7.01 (comp, 3H), 6.07 (dd, J = 5.6, 2.1 Hz, 1H) 5.68 (dd, J = 17.7, 1.4 Hz, 1H), 5.27 (dd, J = 10.9, 1.4 Hz, 1H), 5.11 (tt, J = 5.6, 1.6 Hz, 1H), 3.33 (d, J = 6.0 Hz, 2H), 2.86 (s, 3H); ^{13}C NMR (100 MHz) δ 172.8, 155.1, 149.5, 134.1, 133.0, 128.6, 127.0, 123.9, 121.7, 120.6, 114.3, 82.2, 58.4, 43.4; IR (neat) 1746 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{14}\text{H}_{16}\text{NO}_2$ 230.1181, found 230.1187 ($\text{M}+\text{H}$).

Table 2, entry 5: Purified by chromatography on silica gel (2:1 $\rightarrow$ 1:2 hexanes/EtOAc); yellow oil: TLC R_f = 0.25 (1:2 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 9.78 (s, 1H), 7.77 (d, J = 8.9 Hz, 2H), 7.50 (dd, J = 5.8, 1.4 Hz, 1H), 6.76 (d, J = 8.9 Hz, 2H), 6.19 (dd, J = 5.5, 2.0 Hz, 1H), 5.32 – 5.29 (m, 1H), 3.89 (dd, J = 15.5, 5.0 Hz, 1H), 3.74 (dd, J = 15.5, 6.2 Hz, 1H), 3.14 (s, 3H); ^{13}C NMR (100 MHz) δ 190.2, 172.1, 153.4, 152.7, 132.0, 126.1, 122.7, 111.2, 81.7, 54.1, 39.8; IR (neat) 1746, 1588 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{14}\text{NO}_3$ 232.0974, found 232.0974 ($\text{M}+\text{H}$).

Table 2, entry 6: Purified by chromatography on silica gel (5:1 $\rightarrow$ 2:1 hexanes/EtOAc); yellow oil: TLC R_f = 0.22 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.50 (d, J = 5.8 Hz, 1H), 7.27 – 7.23 (m, 2H), 6.73 – 6.70 (comp, 3H), 6.16 – 6.14 (m, 1H), 5.27 – 5.34 (m, 1H), 3.70 (dd, J = 15.2, 5.8 Hz, 1H), 3.57 (dd, J = 15.3, 6.4 Hz, 1H), 3.52 – 3.34 (m, 2H), 1.17 (t, J = 7.0 Hz, 3H); ^{13}C NMR (100 MHz) δ 172.6, 154.7, 146.9, 129.5, 122.1, 117.1, 112.5, 81.8, 52.9, 45.8, 11.9; IR (neat) 1751 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{16}\text{NO}_2$ 218.1181, found 218.1184 ($\text{M}+\text{H}$).

Table 2, entry 7: Purified by chromatography on silica gel (6:1 $\rightarrow$ 3:1 hexanes/EtOAc); yellow oil: TLC R_f = 0.26 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.42 (dd, J = 5.6, 1.4 Hz, 1H), 7.30 – 7.17 (comp, 7H), 6.73 (t, J = 7.2 Hz, 1H), 6.63 (d, J = 8.1 Hz, 2H), 6.10 (dd, J = 5.8, 1.9 Hz, 1H), 5.21 – 5.18 (m, 1H), 3.65 (dd, J = 15.2, 6.0 Hz, 1H), 3.56 (dd, J = 15.2, 6.2 Hz, 1H), 3.44 – 3.27 (m, 2H), 2.64 (t, J = 7.6 Hz, 2H), 1.92 (p, J = 7.6 Hz, 2H); ^{13}C NMR (100 MHz) δ 172.5, 154.5, 146.9, 141.2, 129.4, 128.4, 128.2, 126.0, 122.0, 117.2, 112.7, 81.6, 53.5, 51.0, 33.0, 28.1; IR (neat) 1765 (C=O) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{20}\text{H}_{22}\text{NO}_2$ 308.1651, found 308.1643 ($\text{M}+\text{H}$).

Table 2, entry 8: Purified by chromatography on silica gel (5:1 → 1:1 hexanes/EtOAc); orange oil: TLC R_f = 0.26 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.52 (d, J = 5.8 Hz, 1H), 7.38 (d, J = 5.8 Hz, 1H), 7.33 – 7.18 (comp, 6H), 7.00 (d, J = 8.3 Hz, 1H), 6.89 (d, J = 8.3 Hz, 1H), 6.86 – 6.80 (m, 1H), 6.14 – 6.12 (m, 1H), 5.94 – 5.92 (m, 1H), 5.46 – 5.45 (m, 1H), 5.36 – 5.34 (m, 1H), 5.18 (d, J = 4.2 Hz, 1H), 4.91 (d, J = 6.2 Hz, 1H), 3.82 – 3.76 (m, 1H), 3.67 – 3.55 (comp, 2H), 3.47 – 3.41 (m, 1H), 3.10 – 2.91 (comp, 4H); ^{13}C NMR (100 MHz) δ 172.5, 172.4, 154.6, 153.6, 148.9, 148.8, 135.7, 135.3, 132.4, 131.9, 129.4, 129.4, 128.6, 128.4, 128.1, 127.8, 127.7, 127.5, 126.3, 125.9, 122.5, 122.2, 118.8, 118.6, 114.5, 114.4, 85.9, 85.4, 61.7, 60.5, 44.0, 43.4, 28.3, 27.2; IR (neat) 1756 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{19}\text{H}_{18}\text{NO}_2$ 292.1338, found 292.1331 ($\text{M}+\text{H}$).

Table 2, entry 9: Purified by chromatography on silica gel (5:1 → 1:1 hexanes/EtOAc); yellow oil: TLC R_f = 0.22 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.46 – 7.40 (comp, 2H), 7.30 – 7.22 (comp, 4H), 6.80 – 6.72 (comp, 4H), 6.59 (d, J = 8.1 Hz, 2H), 6.19 – 6.16 (comp, 2H), 5.40 – 5.39 (m, 1H), 5.03 – 5.02 (m, 1H), 4.40 – 4.37 (m, 1H), 3.86 (t, J = 7.2 Hz, 1H), 3.64 – 3.55 (comp, 2H), 3.24 – 3.16 (comp, 2H), 2.17 – 1.68 (comp, 8H); ^{13}C NMR (100 MHz) δ 172.8, 172.7, 155.9, 153.6, 146.7, 146.8, 129.4, 129.2, 122.7, 121.6, 117.2, 117.0, 112.6, 112.4, 84.0, 82.4, 60.5, 59.1, 49.4, 49.1, 27.8, 25.2, 24.1, 23.2; IR (neat) 1756 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{14}\text{H}_{16}\text{NO}_2$ 230.1181, found 230.1177 ($\text{M}+\text{H}$).

Table 2, entry 10: Purified by chromatography on silica gel (5:1 $\rightarrow$ 1:1 hexanes/EtOAc); yellow oil: TLC R_f = 0.27 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.26 – 7.22 (m, 2H), 7.05 – 7.04 (m, 1H), 6.76 – 6.70 (comp, 3H), 5.12 – 5.09 (m, 1H), 3.65 (dd, J = 15.3, 5.3 Hz, 1H), 3.55 (dd, J = 15.3, 5.9 Hz, 1H), 3.00 (s, 3H), 1.89 (s, 3H); ^{13}C NMR (100 MHz) δ 173.7, 148.3, 146.7, 130.6, 129.2, 117.1, 112.2, 79.6, 55.3, 39.4, 10.6; IR (neat) 1751 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{16}\text{NO}_2$ 218.1181, found 218.1181 ($\text{M}+\text{H}$).

Table 2, entry 11: Purified by chromatography on silica gel (5:1 → 1:1 hexanes/EtOAc); light yellow solid (mp = 116 – 117 °C): TLC R_f = 0.44 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.36 (d, J = 5.6 Hz, 1H), 7.23 – 7.19 (m, 2H), 6.72 (t, J = 7.2 Hz, 1H), 6.65 (d, J = 8.1 Hz, 2H), 5.92 (d, J = 5.6 Hz, 1H), 3.69 (d, J = 15.5 Hz, 1H), 3.63 (d, J = 15.7 Hz, 1H), 2.97 (s, 3H), 1.51 (s, 3H); ^{13}C NMR (100 MHz) δ 171.9, 157.8, 148.8, 129.1, 121.2, 117.0, 112.0, 90.5, 58.8, 40.0, 21.8; IR (neat) 1751 (C=O) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{13}\text{H}_{16}\text{NO}_2$ 218.1181, found 218.1183 (M+H).

Table 2, entry 12: Purified by chromatography on silica gel (5:1 $\rightarrow$ 1:1 hexanes/EtOAc); yellow oil: TLC R_f = 0.35 (3:1 hexanes/EtOAc); ^1H NMR (400 MHz, CDCl_3) δ 7.31 (d, J = 5.8 Hz, 1H), 7.23 – 7.19 (m, 2H), 6.72 (t, J = 7.3 Hz, 1H), 6.64 (d, J = 8.3 Hz, 2H), 5.95 (d, J = 5.6 Hz, 1H), 5.73 – 5.63 (m, 1H), 5.20 – 5.15 (comp, 2H), 3.78 (d, J = 15.7 Hz, 1H), 3.66 (d, J = 15.7 Hz, 1H), 2.96 (s, 3H), 2.66 – 2.54 (m, 2H); ^{13}C NMR (100 MHz) δ 171.9, 156.3, 148.8, 130.1, 129.1, 122.3, 120.5, 117.0, 112.0, 92.2, 57.6, 40.1, 39.6; IR (neat) 1756 ($\text{C}=\text{O}$) cm^{-1} ; HRMS (EI) calcd for $\text{C}_{15}\text{H}_{18}\text{NO}_2$ 244.1338, found 244.1333 ($\text{M}+\text{H}$).