Supporting information for

The Reactivity of Epoxides with Lithium 2,2,6,6-Tetramethylpiperidide in Combination with Organolithiums or Grignard Reagents

David M. Hodgson,* Matthew J. Fleming and Steven J. Stanway

david.hodgson@chem.ox.ac.uk

Contents

(I)	Initial investigations	S1
(II)	General directions	S6
(III)	General procedures	S7
(IV)	Experimental data	S8
(V)	¹ H and ¹³ C NMR spectra of all major products	S34

(I) Initial investigations

We first evaluated a variety of organolithiums to ascertain which were (in)capable of deprotonating terminal epoxides. Mioskowski and co-workers demonstrated that reductive alkylation in THF was highly efficient ($\geq 91\%$ yield) for terminal epoxides with alkyllithiums:¹ secondary and tertiary alkyllithiums giving exclusively *E*-isomers, while primary alkyllithiums gave *E:Z* mixtures. However, the addition of excess of an aryllithium (PhLi) to a terminal epoxide gave only 20% yield of the corresponding *E*-alkene.^{1a} We repeated this latter reaction and found that the major product was alcohol **16b** formed in 69% yield in THF. A similar result was obtained when the reaction was carried in hexane, where alcohol **16b** was obtained in 65%

(1) (a) Mioskowski, C.; Dechoux, L.; Doris, E. *Tetrahedron Lett.* **1994**, 35, 7943–7946. see also: (b) Concellon, J. M.; Suarez, J. R.; Del Solar, V. *Org. Lett.* **2006**, 8, 349–351.

yield along with alkene **14b** in 10% yield (Table 13, entry 1). Evidently, nucleophilic ring-opening of the epoxide by PhLi out-competed deprotonation of the epoxide by PhLi. A variety of other organolithiums were also screened with 1,2-epoxydodecane **13** to investigate whether alkene **14** or alcohol **16** was the predominant product. Use of 2-furanyl- and 2-thienyllithium gave only the corresponding secondary alcohols **16d** and **16e** in 98% and 91% yield, respectively (entries 2 and 3). Vinylolithium gave 72% yield of homoallylic alcohol **16f** and trace amounts of diene **14f** (entry 4; 81% of **16f** obtained in THF). (*E*)-Propenyllithium gave 99% yield of (*E*)-homoallylic alcohol **16h**, while (*Z*)-propenyllithium gave 90% yield of (*Z*)-**16h**. LiCH₂SiMe₃ gave predominantly secondary alcohol **16l** in 41% yield (entry 7; 84% yield of **16l** obtained in THF), along with allylsilane **14l** (14% yield, mixture of stereoisomers) and some recovered epoxide (22%). The more substituted α -silyllithium, 1-(trimethylsilyl)hexyllithium, also predominantly gave the corresponding secondary alcohol **16m** in 70% yield, along with *E*-allylsilane **14m** in 17% yield (entry 8). 1-Hexynyllithium was unreactive giving $\geq 90\%$ recovered epoxide (entry 9).

Table 13. Reactions of organolithiums with epoxide **13**

$ \begin{array}{c} \text{C}_{10}\text{H}_{21} \text{---} \text{epoxide} \text{---} \text{C}_{10}\text{H}_{21} \\ \textbf{13} \end{array} \xrightarrow[\text{hexane, 0 }^{\circ}\text{C to rt, 14 h}]{\text{RLi (3 equiv)}} \begin{array}{c} \text{C}_{10}\text{H}_{21} \text{---} \text{CH(OH)---R} \\ \textbf{16} \end{array} + \begin{array}{c} \text{C}_{10}\text{H}_{21} \text{---} \text{CH=CH---R} \\ \textbf{14} \end{array} + \text{recovered epoxide } \textbf{13} $						
Products						
Entry	RLi	Alcohol 16	Yield, % ^a	Alkene 14 ^b	Yield, % ^a	Recovered 13 , %
1	Li-Ph		65		10	—
2			98		—	—
3			91		—	—
4	Li-CH=CH ₂		72		3	22
5	Li-CH=CH-CH=CH ₂		99		—	—
6	Li-CH=CH-CH=CH-CH=CH ₂		90		—	—
7	LiCH ₂ SiMe ₃		41		14 ^c	15
8			70 ^d		17	—
9	Li-C≡C-C ₄ H ₉		—		—	>90

^a Isolated yield. ^b *E*:*Z* > 98:2 (determined by GCMS) unless stated otherwise. ^c *E*:*Z* = 73:27. ^d *dr* = 50:50.

As the alkynyllithium was unreactive with 1,2-epoxydodecane in hexane (Table 13, entry 9), this organolithium was considered as an ideal starting point to try to develop a modified reductive alkylation procedure, as any competing nucleophilic ring-opening of the epoxide would be absent. Initial experiments aimed to synthesize enyne **14a** by adding a base to a solution of 1,2-epoxydodecane **13** and 1-hexynyllithium (Scheme 7). It was envisaged that the base could deprotonate the epoxide to generate α -lithiated epoxide **53**, which would then be

trapped out by the excess alkynyllithium and then undergo the usual process of reductive alkylation (*cf* **54** to **14a**) to give enyne **14a**.

Scheme 7. Possible enyne **14a** formation from epoxide **13** and 1-hexynyllithium and a base

As *s*-BuLi is known to deprotonate terminal epoxides efficiently in a regio- and stereoselective manner,¹ it was chosen as the base to generate the α -lithiated epoxide in the presence of the alkynyllithium. The general procedure was to add 1 equiv of *s*-BuLi (1.6 M in hexanes) dropwise over 1 h to a solution of the epoxide (0.2 M) and excess alkynyllithium (4 equiv) in THF. The temperature at which *s*-BuLi was added was varied from -90 to 0 °C (Table 14, entries 1–5). After addition of *s*-BuLi, the reaction was left at that temperature for 1 h and then allowed to warm to room temperature overnight. The results of these reactions were essentially identical, in that 35–42% of undesired *E*-alkene **14p** was produced, *via* the reductive alkylation of the epoxide by *s*-BuLi and 50–65% of the epoxide was recovered. None of the desired enyne **14a** was observed.

Table 14. Attempted enyne **14a** formation using *s*-BuLi as the base^a

Entry	Solvent	Temp, °C	14p yield, %	13 recovered, %
1	THF	-90	35	65
2	THF	-78	40	50
3	THF	-50	42	53
4	THF	-20	39	50
5	THF	0	41	52
6 ^b	THF	-50	38	50
7 ^b	Et ₂ O	-50	40	58
8 ^b	hexane	-50	42	51

^a 1.6 M *s*-BuLi unless otherwise indicated. ^b 0.2 M *s*-BuLi used.

As 1-hexynyllithium proved to be unsuccessful it was next investigated whether more nucleophilic 2-thienyllithium² (*cf.* Table 13) could undergo this chemistry while using *s*-BuLi as the base. Following a similar procedure as before, a solution of *s*-BuLi (0.2 M in hexane) was added dropwise (over 1 h) to a mixture of 1,2-epoxydodecane and 2-thienyllithium (3 equiv) in various solvents and at different temperatures with the aim of producing arylated alkene **14e**. However, only unwanted *E*-alkene **14p** (27–41% yield) and secondary alcohol **16e** (36–55% yield) were isolated (Table 15, entries 1–4). These reactions possibly failed because deprotonation of the epoxide and insertion of the resulting carbenoid into the organolithium could be essentially intramolecular events (due to the aggregate state of *s*-BuLi).³ An additional experiment was performed wherein 1,2-epoxydodecane (**13**) was added to a mixture of *s*-BuLi (1 equiv) and 2-thienyllithium (2.5 equiv) in hexane at –50 °C. The two organolithiums were stirred together for 12 h prior to addition of the epoxide to maximize possible formation of mixed aggregates. This reaction generated for the first time the desired arylated alkene **14e**, albeit in only 7% yield (Table 15, entry 5). The secondary alcohol **16e** and alkene **14p** were isolated as the main products. Attempts to improve the yield of alkene **14e** by varying temperature, solvent and number of equivalents of organolithium used all failed.

Table 15. Attempted modified reductive alkylation with 2-thienyllithium^a

Entry	Solvent	Temp °C	14e yield, % ^b	16e yield, % ^b	14p yield, % ^{b,c}
1	THF	–78	–	38	41
2	THF	–50	–	51	33
3	Et ₂ O	–50	–	36	30
4	hexane	0	–	55	27
5 ^d	THF	–50	7	21	48

^a Reactions left at specified temperature for 1 h then allowed to warm to rt and left for 3 h.

^b Isolated yields. ^c *E*-isomer only. ^d Organolithiums left for 14 h prior to addition of epoxide.

The above studies indicated that it was possible to generate a modified reductive alkylation procedure by deprotonation of an epoxide with one type of organolithium and then ring-open the resulting lithiated epoxide with another type of organolithium, however the yield

(2) 62% of alcohol **16e** is formed within 2 h when 2-thienyllithium is added to epoxide **13** compared with 23% of alcohol **16b** when PhLi is added to epoxide **13** under same reaction conditions (3 equiv organolithium, hexane, 0 °C, 0.2 M)

(3) Brown, J. L. *Acc. Chem. Res.* **1968**, *1*, 23–32.

of the corresponding alkene (Table 15, entry 5) was very low. So it was next investigated whether changing the base (to LTMP) would influence the reaction.

(II) General directions

All reactions required anhydrous conditions and were conducted in flame-dried glassware under an atmosphere of argon. All solvents were distilled under an atmosphere of argon; ethers from sodium benzophenone ketyl and hydrocarbons from CaH_2 . 2,2,6,6-Tetramethylpiperidine (TMP) and 1,2-epoxydodecane were distilled from CaH_2 prior to use. Solutions of commercially available and prepared organolithiums and Grignard reagents were titrated prior to use using *s*-BuOH and 1,10-phenanthroline.⁴ Petrol refers to the fraction of light petroleum boiling at 30–40 °C. Reactions were monitored by thin layer chromatography using commercially available aluminium backed plates pre-coated with silica containing a fluorescent indicator (0.2 mm, Merck 60 F₂₅₄) which were developed using phosphomolybdic acid solution. Column chromatography was performed on silica gel (Kieselgel 60, 40–63 μm) or octadecyl-functionalized (C₁₈) silica gel. Infra-red spectra were obtained as liquid films of the compounds on NaCl discs. Peak intensities are specified as strong (s), medium (m), weak (w) or broad (br). Only selected absorbencies (ν_{max}) are reported. ¹H and ¹³C NMR spectra were recorded in CDCl_3 at 25 °C. Data are expressed as chemical shifts in parts per million (ppm) relative to residual chloroform (¹H δ 7.27) or CDCl_3 (¹³C δ 77.0) as the internal standard on the δ scale. The multiplicity of each signal is designated by the following abbreviations; s, singlet; d, doublet; t, triplet; br, broad. Coupling constants *J* are given in Hz. GCMS analyses were performed on a spectrometer with a BPX5 column - HP 6890 (dimethylsilicon capillary column, 30 m, 0.25 mm i.d.) equipped with a mass selective detector operating at 60 eV (CI). Flow rate (He) = 1 mL/min.

(4) Schlosser, M. *Organometallics in Synthesis*, 2nd ed. Wiley, New York, **2002**, p. 671.

(III) General procedures

General Procedure A: Optimized conditions for alkene formation *via* LTMP-modified reductive alkylation using preformed or commercially available organolithiums, or Grignard reagents.

To a solution of 2,2,6,6-tetramethylpiperidine (1.5–2 equiv) in the stated solvent at 0 °C was added *n*-BuLi (1.5–2 equiv) dropwise. The reaction mixture was stirred at this temperature for 10 min and then a solution of the appropriate organometallic reagent (1.3–1.5 equiv) was added dropwise. The reaction was left for a further 5 min and then the epoxide (1 equiv) was added. After 2 h the reaction was quenched with sat. brine solution (10 mL) and the layers were separated. The aqueous layer was extracted with Et₂O (2 x 10 mL), the combined organic layers were dried (MgSO₄) and solvent removed *in vacuo*. The residue was purified by column chromatography (petrol/Et₂O) to give the alkene.

General Procedure B: Optimized conditions for alkene formation *via* LTMP-modified reductive alkylation using organolithiums generated *in situ* *via* halogen-lithium exchange.

To a solution of the organohalide (1.3 equiv) in the stated solvent at –78 °C was added *t*-BuLi (2.6 equiv) dropwise. The reaction mixture was stirred at this temperature for 30 min and then warmed to 0 °C. After 5 min a solution of LTMP [prepared by the addition of *n*-BuLi (2 equiv) to 2,2,6,6-tetramethylpiperidine (2 equiv) in the stated solvent at 0 °C] was added dropwise. The reaction was left for a further 5 min at 0 °C and then the appropriate epoxide (1 equiv) was added. After 2 h the reaction was quenched with sat. brine solution (10 mL) and the layers were separated. The aqueous layer was extracted with Et₂O (2 x 10 mL), the combined organic layers were dried (MgSO₄) and solvent evaporated *in vacuo*. The residue was purified by column chromatography (Et₂O/petrol) to give the alkene.

General Procedure C: Optimized conditions for allylsilane formation *via* LTMP modified reductive alkylation using 1-(trimethylsilyl)hexyllithium generated *in situ*.

To a solution of vinyltrimethylsilane (84 mg, 84 mmol, 1.55 equiv) in THF (1 mL) at –78 °C was added *n*-BuLi (1.6 M in hexanes; 0.44 mL, 0.70 mmol, 1.3 equiv) dropwise. The reaction mixture was stirred at this temperature for 2 h and then at –30 °C for 2 h. The reaction mixture was warmed to 0 °C and a solution of LTMP [prepared from the addition of *n*-BuLi (1.6 M in hexanes; 0.44 mL, 0.70 mmol, 1.3 equiv) to 2,2,6,6-tetramethylpiperidine (99 mg, 0.70 mmol, 1.3 equiv) in hexane (7 mL) at 0 °C] was added dropwise. After 5 min the epoxide (0.54 mmol, 1 equiv) was added and the reaction mixture was left stirring for 2 h at 0 °C. After quenching

with sat. brine solution (10 mL), the layers were separated. The aqueous layer was extracted with Et₂O (2 x 10 mL), the combined organic layers were dried (MgSO₄) and solvent evaporated *in vacuo*. The residue was purified by (reverse phase) column chromatography (petrol/Et₂O/MeCN/CH₂Cl₂) to give the corresponding allylsilane.

General Procedure D: Secondary/tertiary alcohol formation generated by addition of an epoxide to excess organolithium.

The appropriate epoxide (1 equiv) was added dropwise to the organolithium (3 equiv) in the stated solvent at 0 °C. The ice bath was removed and the reaction was left stirring for 14 h. After quenching with sat. brine solution (10 mL), the layers were separated. The aqueous layer was extracted with Et₂O (2 x 20 mL), the combined organic layers were dried (MgSO₄) and solvent evaporated *in vacuo*. The residue was purified by column chromatography (Et₂O/petrol) to give the secondary/tertiary alcohol.

(IV) Experimental data

(*E*)- and (*Z*)-7-Octadecen-5-yne **14a** (Table 1, entry 14 conditions)

To a solution of 2,2,6,6-tetramethylpiperidine (153 mg, 1.08 mmol) and 1-hexyne (133 mg, 1.62 mmol) in hexane (8 mL) at 0 °C was added *n*-BuLi (1.6 M in hexanes; 1.69 mL, 2.70 mmol) dropwise. The reaction mixture was left at this temperature for 15 min and then 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) was added. After 2 h the reaction was quenched with sat. brine solution (10 mL) and the layers were separated. The aqueous layer was extracted with Et₂O (2 x 20 mL) and the combined organic layers were dried (MgSO₄) and solvent evaporated *in vacuo*. The residue was purified by column chromatography (100% petrol) to give *enyne* **14a** (108 mg, 80%, *E*:*Z* = 57:43 by GCMS analysis, *t_R* *E*-**14a** 9.43 min, *t_R* *Z*-**14a** 9.95 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.60 (100% petrol); IR (neat)/cm⁻¹ 3020w, 2957s, 2924s, 2855s, 1466m, 1378w, 1327w, 1260w, 1104w; ¹H NMR (400 MHz) δ 6.05 (dt, *J* = 16, 7, 0.6H), 5.45–5.41 (m, 0.6H), 2.35 (td, *J* = 7, 2, 1.2H), 2.09–2.05 (m, 1.2H) (*E*-isomer), 5.81 (dt, *J* = 10.5, 7.5, 0.4H), 5.46–5.42 (m, 0.4H), 2.32–2.28 (m, 1.6H) (*Z*-isomer), 1.58–1.22 (m, 20H), 0.95–0.92 (m, 6H) (*E*- and *Z*-isomers); ¹³C NMR (400 MHz) δ 143.4, 142.7, 109.7, 109.2, 99.6, 94.4, 88.6, 79.2, 33.0, 31.9, 31.9, 31.0, 30.9, 30.0, 29.6, 29.6, 29.5, 29.4, 29.3, 29.2, 29.1, 28.9, 28.8, 22.7, 22.0, 21.9, 19.2, 19.0, 14.1, 13.6 (*E*- and *Z*-

isomers); MS m/z (CI) 266 ($M+NH_4^+$, 75), 249 (100), 248 (33), 206 (51), 191 (43), 177 (30), 163 (34), 135 (62), 121 (75), 107 (60), 93 (96), 80 (26); HRMS calcd for $C_{18}H_{36}N$ ($M+NH_4^+$) 266.2848, found 266.2844.

(*E*)-1-Phenyl-1-dodecene⁵ **14b**

Following general procedure A, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in hexane (8 mL)] and PhLi (2.0 M in *n*-Bu₂O; 0.35 mL, 0.70 mmol) gave after workup and column chromatography (100% petrol) *alkene* **14b** (123 mg, 93%, *E*:*Z* = 98:2, by GCMS analysis, t_R **Z-14b** 13.02 min, t_R **E-14b** 14.22 min, initial temp. 80 °C, max. temp. 280 °C, rate 10 °C/min) as a colorless oil; R_f 0.48 (100% petrol); IR (neat)/cm⁻¹ 3025m, 2956s, 2925s, 2853s, 1625w, 1599w, 1494w, 1466m, 927s; ¹H NMR (400 MHz) δ 7.38–7.29 (m, 4H), 7.21 (tt, J = 7.5, 1.5, 1H), 6.41 (d, J = 16, 1H), 6.26 (dt, J = 16, 7, 1H), 2.22 (dt, J = 7, 7, 2H), 1.53–1.45 (m, 2H), 1.40–1.25 (m, 14H), 0.92 (t, J = 6.5, 3H); ¹³C NMR (100 MHz) δ 138.0, 131.3, 129.7, 128.5, 126.7, 125.9, 33.1, 32.0, 29.7, 29.6, 29.4, 29.4, 29.3, 22.7, 14.1; MS m/z (CI) 245 ($M+H^+$, 18), 244 (57), 117 (47), 104 (100), 91 (14); HRMS calcd for $C_{18}H_{29}$ ($M+H^+$) 245.2269, found 245.2267.

(*E*)-(1-Dodec-1-enyl)-4-methoxybenzene⁶ **14c**

Following general procedure B, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in hexane (8 mL)] and (4-methoxyphenyl)lithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 0.94 mL, 1.40 mmol) to 4-bromoanisole (131 mg, 0.70 mmol) in THF (1 mL)] gave after column chromatography (2% Et₂O in petrol) *alkene* **14c** (104 mg, 70%, *E*:*Z* = 98:2 by GCMS analysis, t_R **Z-14c** 14.35 min, t_R **E-14c** 16.78 min, initial temp. 100 °C, max. temp. 280 °C, rate 10 °C/min) as a white solid; R_f 0.48 (10% Et₂O in petrol); mp 36–39 °C; IR (CHCl₃)/cm⁻¹ 2954m, 2920s, 2850m, 1608w, 1513w, 1465w, 1251w, 1176w, 1023w, 962w; ¹H NMR (400 MHz) δ 7.31 (d, J = 9, 2H), 6.87 (d, J = 9, 2H), 6.35 (d, J = 16, 1H), 6.12 (dt, J = 16, 7, 1H), 3.82 (s, 3H), 2.21 (q, J = 7, 2H), 1.52–1.44 (m, 2H), 1.41–1.25 (m, 14H), 0.92 (t, J = 7, 3H); ¹³C NMR (400 MHz) δ 158.6, 130.8, 129.1, 129.0, 127.0, 113.9, 55.2,

(5) Terao, J.; Watabe, H.; Miyamoto, M.; Kambe, N. *Bull. Chem. Soc. Jpn.* **2003**, 76, 2209–2214.

(6) Hodgson, D. M.; Fleming, M. J.; Stanway, S. J. *J. Am. Chem. Soc.* **2004**, 126, 12250–12251.

33.1, 31.9, 29.7, 29.6, 29.6, 29.4, 29.3, 22.7, 14.1; MS m/z (CI) 275 ($M+H^+$, 100), 274 (69), 147 (63), 134 (18), 121 (25); HRMS calcd for $C_{19}H_{31}O$ ($M+H^+$) 275.2375, found 275.2373.

(*E*)-2-(Dodec-1-enyl)furan **14d**

Following general procedure A, the addition of 1,2-epoxydodecane **13** (308 mg, 1.67 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 2.09 mL, 3.34 mmol) to TMP (472 mg, 3.34 mmol) in hexane (24 mL)] and 2-furanyllithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 1.45 mL, 2.17 mmol) to furan (148 mg, 2.17 mmol) at 0 °C in Et₂O (2 mL)] gave after workup and column chromatography (100% petrol) *alkene* **14d** (125 mg, 32%, *E:Z* = 98:2 by GCMS analysis, t_R **Z-14d** 8.88 min, t_R **E-14d** 9.27 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; R_f 0.40 (100% petrol); IR (neat)/cm⁻¹ 2925s, 2854s, 1466m, 1378w, 1254w, 1152m, 1075m, 1012m, 959m, 926w, 884w, 727s; ¹H NMR (400 MHz) δ 7.31 (d, J = 2, 1H), 6.35 (dd, J = 3.5, 2, 1H), 6.22–6.19 (m, 2H), 6.13 (d, J = 3.5, 1H), 2.20–2.17 (m, 2H), 1.49–1.42 (m, 2H), 1.37–1.22 (m, 14H), 0.89 (t, J = 7, 3H); ¹³C NMR (100 MHz) δ 153.4, 141.1, 130.3, 118.4, 111.0, 105.8, 32.8, 31.9, 29.6, 29.6, 29.5, 29.3, 29.2, 29.2, 22.7, 14.1; MS m/z (CI) 235 ($M+H^+$, 100), 234 (28), 107 (24), 94 (28); HRMS calcd for $C_{16}H_{27}O$ ($M+H^+$) 235.2062, found 235.2052.

(*E*)-2-(Dodec-1-enyl)thiophene **14e**

Following general procedure A, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in hexane (8 mL)] and 2-thienyllithium (1.0 M in THF; 0.70 mL, 0.70 mmol) gave after workup and column chromatography (100% petrol) *alkene* **14e** (53 mg, 39%, *E:Z* = 99:1, by GCMS analysis, t_R **Z-14e** 7.41 min, t_R **E-14e** 8.02 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; R_f 0.53 (100% petrol); IR (neat)/cm⁻¹ 2923s, 2854s, 1465m, 1437m, 1378w, 1284w, 1239w, 1204w, 1041w, 953s; ¹H NMR (400 MHz) δ 7.08 (d, J = 5, 1H), 6.94 (dd, J = 5, 3.5, 1H), 6.87 (d, J = 3.5, 1H), 6.50 (d, J = 15.5, 1H), 6.08 (dt, J = 15.5, 7, 1H), 2.18 (dt, J = 7, 1.5, 2H), 1.49–1.42 (m, 2H), 1.35–1.21 (m, 14H), 0.89 (t, J = 7, 3H); ¹³C NMR (100 MHz) δ 143.2, 131.3, 127.2, 124.0, 122.9, 122.9, 32.8, 31.9, 29.6, 29.5, 29.5, 29.3, 29.2, 29.2, 22.7, 14.1; MS m/z (CI) 251 ($M+H^+$, 100), 250 (44), 149 (8), 132 (13), 129 (10), 71 (17); HRMS calcd for $C_{16}H_{27}S$ ($M+H^+$) 251.1833, found 251.1824.

(E)-1,3-Tetradecadiene⁷ **14f**

Following general procedure A, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in hexane (8 mL)] and vinylolithium⁸ (1.5 M in Et₂O; 0.47 mL, 0.70 mmol) gave after workup and column chromatography (100% petrol) *diene* **14f** (77 mg, 73%, *E:Z* = 98:2 by GCMS analysis, *t_R* **Z-14f** 6.83 min, *t_R* **E-14f** 7.05 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.58 (100% petrol); IR (neat)/cm⁻¹ 3008m, 2925s, 2855s, 1794w, 1654m, 1603m, 1466s, 1378w, 1001s, 949m, 895s; ¹H NMR (400 MHz) δ 6.33 (ddd, *J* = 17, 10, 10, 1H), 6.06 (dd, *J* = 15, 10, 1H), 5.72 (dt, *J* = 15, 7, 1H), 5.10 (d, *J* = 17, 1H), 4.96 (d, *J* = 10, 1H), 2.08 (dt, *J* = 7, 7, 2H), 1.40–1.22 (m, 16H), 0.90 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 137.4, 135.6, 130.8, 114.5, 32.6, 31.9, 29.6, 29.6, 29.5, 29.3, 22.7, 14.1; MS *m/z* (CI) 212 (M+NH₄⁺, 60), 193 (9), 195 (100), 123 (8), 109 (13), 95 (28), 68 (9); HRMS calcd for C₁₄H₃₀N (M+NH₄⁺) 212.2378, found 212.2379.

(E)-2,3-Dimethyl-2,4-pentadecadiene⁶ **14g**

Following general procedure B, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in hexane (8 mL)] and (3-methylbut-2-en-2-yl)lithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 0.94 mL, 1.40 mmol) to 2-bromo-3-methyl-2-butene (105 mg, 0.70 mmol) in THF (1.5 mL)] gave after workup and column chromatography (100% petrol) *diene* **14g** (109 mg, 85%, *E*-only by GCMS analysis, *t_R* **E-14g** 7.10 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.56 (100% petrol); IR (neat)/cm⁻¹ 2923s, 2854s, 1465m, 1375m, 1170w, 959s; ¹H NMR (400 MHz) δ 6.05 (d, *J* = 15.5, 1H), 5.59 (dt, *J* = 15.5, 7, 1H), 2.12 (dt, *J* = 7, 7, 2H), 1.82 (s, 3H), 1.78 (s, 3H), 1.76 (s, 3H), 1.33–1.24 (m, 16H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 129.1, 128.5, 128.4, 126.1, 33.4, 31.9, 30.0, 29.6, 29.6, 29.4, 29.3, 22.7, 21.6, 20.2, 14.4, 14.1; MS *m/z* (CI) 237 (M+H⁺, 100), 109 (21); HRMS calcd for C₁₇H₃₃ (M+H⁺) 237.2582, found 237.2585.

(7) Suzuki, K.; Hasegawa, T.; Imai, T.; Maeta, H.; Ohba, S. *Tetrahedron* **1995**, 51, 4483–4494.

(8) Seyferth, D.; Weiner, M. *J. Am. Chem. Soc.* **1961**, 83, 3583–3586.

(2*E*,4*E*)-2,4-Pentadecadiene⁶ **14h**

Following general procedure A, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in hexane (8 mL)] and (*E*)-propenyllithium⁹ (2.0 M in Et₂O; 0.35 mL, 0.70 mmol) gave after workup and column chromatography (100%, petrol) *diene* **14h** (92 mg, 82%, *E,E*:*Z,E* = 98:2 by GCMS analysis, *t_R* *Z,E*-**14h** 10.82 min, *t_R* *E,E*-**14h** 10.98 min, initial temp. 100 °C, max. temp. 280 °C, rate 8 °C/min) as a colorless oil; *R_f* 0.64 (100% petrol); IR (neat)/cm⁻¹ 3016m, 2957s, 2925s, 2854s, 1466m, 1378w, 986s; ¹H NMR (400 MHz) δ 6.07–5.95 (m, 2H), 5.62–5.52 (m, 2H), 2.05 (dt, *J* = 7, 7, 2H), 1.74 (d, *J* = 6.5, 3H), 1.40–1.22 (m, 16H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 132.3, 131.7, 130.2, 126.6, 32.6, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 22.7, 18.0, 14.1; MS *m/z* (CI) 226 (M+NH₄⁺, 33), 208 (100), 207 (17), 95 (22), 81 (48); HRMS calcd for C₁₅H₃₂N (M+NH₄⁺) 226.2535, found 226.2535.

(7*E*,9*E*)-7,9-Icosadiene⁶ **14i**

Following general procedure B, the addition of 1,2-epoxydodecane **13** (80 mg, 0.43 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.54 mL, 0.87 mmol) to TMP (123 mg, 0.87 mmol) in hexane (7 mL)] and (*E*)-1-octenyllithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 0.74 mL, 1.12 mmol) to (*E*)-1-iodo-1-octene¹⁰ (133 mg, 0.56 mmol) in Et₂O (1.0 mL)] gave after workup and column chromatography (100% petrol) *diene* **14i** (101 mg, 84%, *E,E*:*Z,E* = 99:1, by GCMS analysis, *t_R* *Z,E*-**14i** 16.68 min, *t_R* *E,E*-**14i** 17.07 min, initial temp. 80 °C, max. temp. 280 °C, rate 10 °C/min) as a colorless oil; *R_f* 0.59 (100% petrol); IR (neat)/cm⁻¹ 3014w, 2957s, 2925s, 2854s, 1460m, 1378w, 986s; ¹H NMR (400 MHz) δ 6.03–5.98 (m, 2H), 5.57 (dt, *J* = 14.5, 7, 2H), 2.05 (dt, *J* = 7, 7, 4H), 1.42–1.21 (m, 24H), 0.89 (t, *J* = 7, 6H); ¹³C NMR (100 MHz) δ 132.4, 130.3, 32.6, 31.9, 31.7, 29.6, 29.5, 29.4, 29.4, 29.3, 29.2, 22.7, 22.6, 14.1, 14.1; MS *m/z* (CI) 296 (M+NH₄⁺, 30), 279 (63), 278 (100), 277 (56), 276 (20), 275 (17), 138 (13), 95 (16), 81 (15); HRMS calcd for C₂₀H₄₂N (M+NH₄⁺) 296.3317, found 296.3310.

(1*E*,3*E*)-1-Phenyl-1,3-tetradecadiene⁶ **14j**

Following general procedure A, the addition of 1,2-epoxydodecane (230 mg, 1.25 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 1.56 mL, 2.50 mmol)

(9) Seyferth, D.; Vaughan, L. G. *J. Am. Chem. Soc.* **1964**, 86, 883–890.

(10) Schwartz, J.; Hart, D. W.; Blackburn, T. F. *J. Am. Chem. Soc.* **1975**, 97, 679–680.

to TMP (353 mg, 2.50 mmol) in hexane (19 mL)] and (*E*)-2-phenylethenyllithium¹¹ (1.1 M in Et₂O; 1.48 mL, 1.63 mmol) gave after workup and reverse phase column chromatography (C₁₈ silica, MeCN)¹² diene **14j** (243 mg, 72%, *E,E:Z,E* = 98:2, by GCMS analysis, *t_R* *Z,E*-**14j** 11.55 min, *t_R* *E,E*-**6g** 11.83 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colourless oil; *R_f* 0.26 (C₁₈ silica, MeCN). IR (neat)/cm⁻¹ 3022m, 2955s, 2925s, 2853s, 1644w, 1597w, 1495w, 1466w, 1448w, 986s; ¹H NMR (400 MHz) δ 7.40–7.37 (m, 2H), 7.33–7.28 (m, 2H), 7.20 (tt, *J* = 7.5, 1.5, 1H), 6.76 (dd, *J* = 15.5, 10.5, 1H), 6.44 (d, *J* = 15.5, 1H), 6.20 (dd, *J* = 15, 10.5, 1H), 5.83 (dt, *J* = 15, 7, 1H), 2.14 (dt, *J* = 7, 7, 2H), 1.47–1.22 (m, 16H), 0.88 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 137.7, 136.1, 130.4, 129.9, 129.5, 128.5, 127.0, 126.1, 32.9, 31.9, 29.6, 29.5, 29.4, 29.3, 29.2, 22.7, 14.1; MS *m/z* (CI) 271 (M+H⁺ 100), 270 (64), 143 (25), 128 (14); HRMS calcd for C₂₀H₃₁ (M+H⁺) 271.2426, found 271.2429.

(*2Z,4E*)-2,4-Pentadecadiene⁶ **14h**

Following general procedure A, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in hexane (8 mL)] and (*Z*)-propenyllithium⁹ (1.4 M in Et₂O; 0.50 mL, 0.70 mmol) gave after workup and column chromatography (100% petrol) diene-**14h** (79 mg, 70%, *Z,E:E,E* = 90:10 by GCMS analysis, *t_R* *Z,E*-**14h** = 10.82 min, *t_R* *E,E*-**14h** = 10.98 min, initial temp. 100 °C, max. temp. 280 °C, rate 8 °C/min) as a colorless oil; *R_f* 0.64 (100% petrol); IR (neat)/cm⁻¹ 3015m, 2960s, 2920s, 2844s, 1466m, 1349w, 970s; ¹H NMR (400 MHz) δ 6.33 (ddq, *J* = 11, 10, 1.5, 1H), 6.02–5.95 (m, 1H), 5.67 (dt, *J* = 15, 7.5, 1H), 5.38 (dq, *J* = 11, 7, 1H), 2.15–2.07 (m, 2H), 1.75 (dd, *J* = 7, 1.5, 3H), 1.44–1.20 (m, 16H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 134.7, 129.5, 125.3, 123.8, 32.9, 31.9, 29.6, 29.5, 29.4, 29.4, 29.3, 22.7, 14.1, 13.3; MS *m/z* (CI) 226 (M+NH₄⁺, 52), 208 (100), 207 (11), 166 (87), 95 (10), 81 (22); HRMS calcd for C₁₅H₃₂N (M+NH₄⁺) 226.2535, found 226.2541.

(*7Z,9E*)-7,9-Icosadiene¹³ **14i**

Following general procedure B, the addition of 1,2-epoxydodecane **13** (80 mg, 0.43 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.54 mL, 0.87 mmol)

(11) Prepared by the addition of (*E*)-β-bromostyrene to Li metal in Et₂O at 0 °C.

(12) Farina, V. *J. Org. Chem.* **1991**, 56, 4985–4987.

(13) Bestmann, H. J.; Vostrowsky, O.; Paulus, H.; Billmann, W.; Stransky, W. *Tetrahedron Lett.* **1977**, 18, 121–124.

to TMP (123 mg, 0.87 mmol) in hexane (7 mL)] and (Z)-1-octenyllithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 0.74 mL, 1.12 mmol) to (Z)-1-iodo-1-octene¹⁴ (133 mg, 0.56 mmol) in Et₂O (1.0 mL)] gave after workup and column chromatography (100% petrol) *diene* **14i** (96 mg, 80%, *Z,E:E,E*:other isomer = 91:5:4 by GCMS analysis, *t_R* *Z,E*-**14i** 16.68 min, *t_R* other isomer 16.90, *t_R* *E,E*-**14i** 17.07 min, initial temp. 80 °C, max. temp. 280 °C, rate 10 °C/min) as a colorless oil; *R_f* 0.59 (100% petrol); IR (neat)/cm⁻¹ 3019m, 2926s, 2854s, 1466s, 1378s, 982m, 946m; ¹H NMR (400 MHz) δ 6.31 (dd, *J* = 15, 11, 1H), 5.97 (dd, *J* = 11, 10.5, 1H), 5.67 (dt, *J* = 15, 7.5, 1H), 5.31 (dt, *J* = 10.5, 7.5, 1H), 2.20–2.02 (m, 4H), 1.43–1.21 (m, 24H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 134.7, 130.1, 128.6, 125.6, 32.9, 31.9, 31.8, 29.7, 29.6, 29.5, 29.4, 29.3, 29.2, 28.9, 27.7, 22.7, 22.6, 14.1, 14.1; MS *m/z* (CI) 296 (M+NH₄⁺, 31), 279 (75), 278 (100), 277 (31), 276 (13), 166 (10), 138 (14), 96 (14), 81 (19); HRMS calcd for C₂₀H₄₂N (M+NH₄⁺) 296.3317, found 296.3325.

(5*E*,7*E*)-5-Methyl-5,7-octadecadiene⁶ **14k**

Following general procedure B, the addition of 1,2-epoxydodecane **13** (125 mg, 0.68 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.85 mL, 1.36 mmol) to TMP (192 mg, 1.36 mmol) in hexane (10 mL)] and (*E*)-(2-methyl-1-hexenyl)lithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 1.18 mL, 1.77 mmol) to (*E*)-1-iodo-2-methyl-1-hexene¹⁵ (198 mg, 0.88 mmol) in Et₂O (2 mL)] gave after workup and column chromatography (100% petrol) *diene* **14k** (153 mg, 85% *E,E:Z,E* = 91:9 by GCMS analysis, *t_R* *Z,E*-**14k** 9.78 min, *t_R* *E,E*-**14k** 10.00 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.69 (100% petrol); IR (neat)/cm⁻¹ 3020m, 2924s, 2854s, 1642w, 1466s, 1378m, 963s; ¹H NMR (400 MHz) δ 6.25 (dd, *J* = 15, 11, 1H), 5.81 (d, *J* = 11, 1H), 5.59 (dt, *J* = 15, 7, 1H), 2.09 (dt, *J* = 7, 7, 2H), 2.04 (t, *J* = 7.5, 2H), 1.74 (s, 3H), 1.44–1.26 (m, 20H), 0.93–0.88 (m, 6H); ¹³C NMR (100 MHz) δ 136.7, 132.4, 126.6, 124.5, 39.5, 33.0, 31.9, 30.1, 29.6, 29.6, 29.4, 29.3, 29.3, 22.7, 22.4, 16.4, 14.1, 14.0; MS *m/z* (CI) 265 (M+H⁺, 100), 264 (29), 263 (10), 137 (10); HRMS calcd for C₁₉H₃₇ (M+H⁺) 265.2895, found 265.2889.

(*E*)-1-Trimethylsilyl-2-tridecene¹⁶ **14l** (from Table 4, entry 4 conditions)

Following general procedure A, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol)

(14) Brown, H. C.; Blue, C. D.; Nelson, D. J.; Bhat, N. G. *J. Org. Chem.* **1989**, *54*, 6064–6067.

(15) Negishi, E.; Van Horn, D. E.; Yoshida, T. *J. Am. Chem. Soc.* **1985**, *107*, 6639–6647.

(16) Masao, A.; Yamaguchi, H.; Fujita, E.; Nagao, Y.; Ochiai, M. *Chem. Pharm. Bull.* **1989**, *37*, 3221–3324.

to TMP (153 mg, 1.08 mmol) in Et₂O (8 mL)] and (trimethylsilylmethyl)lithium (1.0 M in pentane; 0.70 mL, 0.70 mmol) gave after workup and column chromatography (100% petrol) *allylsilane* **14l** (89 mg, 65%, *E:Z* = 97:3, by GCMS analysis, *t_R* *E*-**14l** 8.22 min, *t_R* *Z*-**14l** 8.30 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.63 (petrol); IR (neat)/cm⁻¹ 3007m, 2928s, 2854s, 1659w, 1466s, 1416m, 1378w, 1248s, 1155s, 934s, 857s; ¹H NMR (400 MHz) δ 5.42–5.33 (m, 1H), 5.29–5.23 (m, 1H), 2.29 (dt, *J* = 7, 7, 2H), 1.41 (dd, *J* = 7.5, 1, 2H), 1.33–1.27 (m, 16H), 0.90 (t, *J* = 7, 3H), 0.00 (s, 9H); ¹³C NMR (100 MHz) δ 129.1, 125.9, 32.8, 31.9, 30.0, 29.7, 29.6, 29.5, 29.4, 29.1, 22.7, 22.6, 14.1, –2.0; MS *m/z* (CI) 255 (*M*+H⁺, 25), 90 (100), 73(10); HRMS calcd for C₁₆H₃₅Si (*M*+H⁺) 255.2508, found 255.2501.

When *Z*-isomer **14l** was formed in appreciable amounts along with the *E*-isomer (Table 4, entries 1–3) it was identified from the allylic protons and allylic/vinylic/–SiMe₃ carbons: ¹H NMR (400 MHz) δ 1.48 (d, *J* = 8, 2H); ¹³C NMR (100 MHz) δ 127.8, 125.2, 27.4, –1.8.

(*E*)-(1-Pentyl-1-tridecen-1-yl)trimethylsilane⁶ **14m**

Following general procedure C, using 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) gave after workup and reverse phase column chromatography¹² (C₁₈ silica, 15% Et₂O in MeCN) *allylsilane* **14m** (130 mg, 74%, *E*-only by GCMS analysis, *E*-**14m** 10.18 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.13 (C₁₈ silica, 15% Et₂O in MeCN); IR (neat)/cm⁻¹ 2924s, 2854s, 1467m, 1378w, 1258m, 969m, 861m; ¹H NMR (400 MHz) δ 5.24–5.12 (m, 2H), 1.99 (dt, *J* = 6.5, 6.5, 2H), 1.42–1.16 (m, 25 H), 0.91–0.86 (m, 6H), –0.01 (s, 9H); ¹³C NMR (100 MHz) δ 131.5, 128.2, 32.9, 32.9, 31.9, 31.8, 30.1, 29.7, 29.6, 29.5, 29.4, 29.1, 29.0, 28.9, 22.7, 22.6, 14.2, 14.1, –3.1; MS *m/z* (CI) 325 (*M*+H⁺, 34), 251 (13), (8), 250 (48), 90 (100), 73 (25); HRMS calcd for C₂₁H₄₅Si (*M*+H⁺) 325.3291, found 325.3287.

(*E*)-5-Hexadecene⁶ **14n** (Table 10, entry 3 conditions)

Following general procedure A (except reaction left for 3 h), the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.68 mL, 1.08 mmol) to TMP (153 mg, 1.08 mmol) in Et₂O (8 mL)] and *n*-BuMgCl (2.0 M in Et₂O; 0.38 mL, 0.76 mmol) gave after workup and column chromatography (100% petrol) *alkene* **14n** (84 mg, 69%, *E:Z* = 96:4, by ¹H NMR analysis of vinylic protons in the δ 6.00–5.00 region) as a colorless oil; *R_f* 0.70 (100% petrol); IR (neat)/cm⁻¹ 2924s, 2854s 1466m, 1378w, 966m; ¹H NMR (500 MHz) δ 5.43–5.83 (m, 2H), 2.03–1.93 (m, 4H), 1.39–1.22 (m,

20H), 0.93–0.87 (m, 6H); ^{13}C NMR (125 MHz) δ 130.3, 130.2, 32.6, 32.2, 31.9, 31.8, 29.6, 29.5, 29.3, 29.1, 22.6, 22.1, 14.0, 13.9; MS m/z (EI) 224 (M^+ , 21), 111 (39), 98 (29), 97 (88), 84 (47), 84 (47), 83 (100), 71 (32), 70 (59), 69 (92), 67 (41), 57 (30), 55 (65); HRMS calcd for $\text{C}_{16}\text{H}_{32}$ (M^+) 224.2504, found 224.2514.

When *Z*-isomer **14n** was formed in appreciable amounts along with the *E*-isomer (Table 9, entries 1–5) it was identified from the vinylic protons and carbons: ^1H NMR (500 MHz) δ 5.36 (dt, $J = 7.5, 6, 2\text{H}$); ^{13}C NMR (125 MHz) δ 129.9, 129.8.

(*E*)-2-Tridecene⁶ **14o** (Table 11, entry 4 conditions)

Following general procedure A (except reaction left for 3 h), the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.51 mL, 0.81 mmol) to TMP (114 mg, 0.81 mmol) in hexane (8 mL)] and MeMgCl (2.2 M in THF; 0.45 mL, 0.98 mmol) gave after workup and column chromatography (100% petrol) *alkene* **14o** (70 mg, 71%, *E:Z* = 98:2 by GCMS analysis, t_R *E*-**14o** 9.10 min, t_R *Z*-**14o** 9.25 min, initial temp. 80 °C, max. temp. 280 °C, rate 10 °C/min) as a colorless oil; R_f 0.66 (100% petrol); IR (neat)/ cm^{-1} 2925s, 2855s, 1465m, 1378w, 964m; ^1H NMR (400 MHz) δ 5.44–5.40 (m, 2H), 1.99–1.94 (m, 2H), 1.66–1.64 (m, 3H), 1.35–1.24 (m, 16H), 0.89 (t, $J = 7, 3\text{H}$); ^{13}C NMR (100 MHz) δ 131.7, 124.5, 32.6, 31.9, 29.6, 29.5, 29.4, 29.2, 22.7, 17.9, 14.1; MS m/z (EI) 182 (M^+ , 41), 111 (32), 98 (34), 97 (87), 84 (61), 83 (94), 71 (37), 70 (83), 69 (100), 67 (38), 55 (71); HRMS calcd for $\text{C}_{13}\text{H}_{26}$ (M^+) 182.2035, found 182.2035.

When *Z*-isomer **14o** was formed in appreciable amounts along with the *E*-isomer (Table 8, entries 1, 3–6) it was identified from the allylic protons and vinylic carbons:

^1H NMR (400 MHz) δ 1.61 (dd, $J = 6, 1, 3\text{H}$); ^{13}C NMR (100 MHz) δ 130.9, 123.5.

When *E*-alkene **14p** was observed it was identified as follows:

(*E*)-3-Methyl-4-pentadecene **14p**

Colorless oil; R_f 0.75 (100% petrol); IR (neat)/ cm^{-1} 2959s, 2924s, 2854s, 1458s, 1378m, 1158w; ^1H NMR (400 MHz) δ 5.36 (dt, $J = 15.5, 6.5, 1\text{H}$), 5.25 (dd, $J = 15.5, 7.5, 1\text{H}$), 2.07–1.90 (m, 3H), 1.40–1.20 (m, 18H), 0.96 (d, $J = 7, 3\text{H}$), 0.89 (t, $J = 7, 3\text{H}$), 0.85 (t, $J = 7.5, 3$); ^{13}C NMR (100 MHz) δ 136.1, 128.7, 38.4, 32.6, 31.9, 29.9, 29.7, 29.6, 29.5, 29.3, 29.1, 22.7, 20.5, 14.1, 11.8; MS m/z (CI) 242 ($\text{M}+\text{NH}_4^+$, 5), 224 (48), 195 (30), 125 (24), 111 (35), 97 (49), 83 (100); HRMS calcd for $\text{C}_{16}\text{H}_{36}\text{N}$ ($\text{M}+\text{NH}_4^+$) 242.2848, found 242.2850.

When aldehyde **15** was observed it was identified as follows:

Dodecanal¹⁷ **15**

Colourless oil; R_f 0.36 (5% Et₂O in petrol); ¹H NMR (400 MHz) δ 9.72 (t, J = 2, 1H), 2.38 (td, J = 7, 2, 2H), 1.70–1.48 (m, 2H), 1.40–1.08 (m, 16H), 0.84 (t, J = 6.5, 3H); ¹³C NMR (100 MHz) δ 203.0, 44.3, 31.9, 29.6, 29.4, 29.3, 29.3, 29.2, 22.7, 22.1, 14.1.

1-Phenyldodecan-2-ol⁶ **16b**

Following general procedure D, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to PhLi (2.0 M in *n*-Bu₂O; 0.82 mL, 1.63 mmol) in hexane (8 mL) gave after workup and column chromatography (15% Et₂O in petrol) *alcohol* **16b** (98 mg, 69%) as a white solid; R_f 0.16 (10% Et₂O in petrol); mp 40–41 °C; IR (CHCl₃)/cm⁻¹ 3367br, 3027m, 2919s, 1496w, 1466s, 1350w; ¹H NMR (400 MHz) δ 7.35–7.31 (m, 2H), 7.27–7.22 (m, 3H), 3.85–3.79 (m, 1H), 2.84 (dd, J = 13.5, 4.5, 1H), 2.65 (dd, J = 13.5, 8.5, 1H), 1.52–1.46 (m, 3H), 1.35–1.24 (m, 16H), 0.90 (t, J = 7, 3H); ¹³C NMR (100 MHz) δ 138.7, 129.4, 128.5, 126.4, 72.7, 44.1, 36.9, 31.9, 29.7, 29.6, 29.3, 25.8, 22.7, 14.1; MS m/z (CI) 280 (M+NH₄⁺, 100), 262 (6), 244 (36), 169 (12), 117 (27), 104 (64), 92 (76), 91 (40); HRMS calcd for C₁₈H₃₄NO (M+NH₄⁺) 280.2640, found 280.2632.

1-(4-Methoxy-phenyl)dodecanol-2-ol **16c**

Following general procedure D, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to (4-methoxyphenyl)lithium [prepared by the addition of *t*-BuLi (1.5 M in hexanes; 2.17 mL, 3.26 mmol) to 4-bromoanisole (305 mg, 1.63 mmol) at -78 °C in THF (2.5 mL)] in hexane (7 mL) gave after column chromatography (10% Et₂O in petrol) *alcohol* **16c** (132 mg, 83%) as a white solid; R_f 0.11 (10% Et₂O in petrol); mp 55–56 °C; IR (KBr)/cm⁻¹ 3418br, 2924s, 1460s, 1378s, 1032w; ¹H NMR (400 MHz) δ 7.14 (d, J = 8.5, 2H), 6.87 (d, J = 8.5, 2H), 3.81 (s, 3H), 3.80–3.74 (m, 1H), 2.79 (dd, J = 13.5, 4.5, 1H), 2.58 (dd, J = 13.5, 8.5, 1H), 1.58–1.22 (m, 19H), 0.89 (t, J = 7, 3H); ¹³C NMR (400 MHz) δ 158.2, 130.3, 114.0, 99.6, 72.8, 55.2, 43.1, 36.7, 31.9, 29.7, 29.6, 29.3, 25.8, 22.7, 14.1; MS m/z (CI) 310 (M+NH₄⁺, 100), 275 (11), 122 (38), 121 (12); HRMS calcd for C₁₉H₃₆NO₂ (M+NH₄⁺) 310.2741, found 310.2742.

(17) Spectra matched commercially available (Aldrich, product no. D222003) material.

1-(Furan-2-yl)dodecan-2-ol **16d**

Following general procedure D, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to 2-furanyllithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 1.08 mL, 1.62 mmol) to furan (110 mg, 1.62 mmol) at 0 °C in Et₂O (2 mL)] in hexane (8 mL) gave after workup and column chromatography (10% Et₂O in petrol) *alcohol* **16d** (134 mg, 98%) as a white solid; *R*_f 0.10 (15% Et₂O in petrol); mp 46–47 °C; IR (CHCl₃)/cm⁻¹ 3357m, 2930s, 2855s, 1597w, 1507w, 1466w, 1147w, 1081w, 1009w; ¹H NMR (400 MHz) δ 7.35 (dd, *J* = 2, 1, 1H), 6.32 (dd, *J* = 3, 2, 1H), 6.12 (dd, *J* = 3, 1, 1H), 3.91–3.88 (m, 1H), 2.83 (dd, *J* = 15, 4, 1H), 2.72 (dd, *J* = 15, 8, 1H), 1.76 (d, *J* = 4, 1H), 1.52–1.46 (m, 2H), 1.35–1.24 (m, 16H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 152.9, 141.6, 110.3, 107.0, 70.5, 36.7, 36.1, 31.9, 29.6, 29.3, 25.6, 22.7, 14.1; MS *m/z* (CI) 270 (M+NH₄⁺, 6), 268 (24), 253 (58), 235 (33), 233 (22), 170 (13), 98 (13), 82 (14); HRMS calcd for C₁₆H₂₉O₂ (M+H⁺) 253.2168, found 253.2176.

1-(Thiophen-2-yl)dodecan-2-ol **16e**

Following general procedure D, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to 2-thienyllithium (1.0 M in THF; 1.62 mL, 1.62 mmol) in hexane (8 mL) gave after workup and column chromatography (15% Et₂O in petrol) *alcohol* **16e** (131 mg, 91%) as a white solid; *R*_f 0.15 (10% Et₂O in petrol); mp 45–46 °C; IR (CDCl₃)/cm⁻¹ 3368s, 2923s, 2853s, 1640w, 1534w, 1466s, 1439s, 1078m, 1045m, 1021m; ¹H NMR (400 MHz) δ 7.18 (d, *J* = 5.5, 1H), 6.97 (dd, *J* = 5.5, 3.5, 1H), 6.88 (d, *J* = 3.5, 1H), 3.85–3.79 (m, 1H), 3.04 (dd, *J* = 15, 4, 1H), 2.89 (dd, *J* = 15, 8, 1H), 1.69 (s, 1H), 1.57–1.44 (m, 2H), 1.35–1.22 (m, 16H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 140.6, 127.0, 126.0, 124.1, 72.4, 37.9, 36.5, 31.9, 29.6, 29.3, 25.7, 22.7, 14.1; MS *m/z* (CI) 286 (M+NH₄, 100), 269 (11), 98 (11); HRMS calcd for C₁₆H₃₂NOS (M+NH₄⁺) 286.2199, found 286.2201.

1-Tetradecen-4-ol¹⁸ **16f**

Following general procedure D, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to vinylolithium⁸ (1.5 M in Et₂O; 1.09 mL, 1.63 mmol) in hexane (8 mL) gave after workup and column chromatography (15% Et₂O in petrol) *alcohol* **16f** (83 mg, 72%) as a white solid; *R*_f

(18) Chang, Y. H.; Uang, B. J.; Wu, C. M.; Yu, T. H. *Synthesis* **1990**, 1033–1034.

0.20 (15% Et₂O in petrol); mp 45–46 °C; IR (CHCl₃)/cm⁻¹ 3341br, 2925s, 2854s, 1641w, 1466w, 994w, 912w; ¹H NMR (400 MHz) δ 5.89–5.78 (m, 1H), 5.17–5.14 (m, 1H), 5.13–5.11 (m, 1H), 3.68–3.61 (m, 1H), 2.34–2.27 (m, 1H), 2.18–2.10 (m, 1H), 1.63 (br, 1H), 1.48–1.22 (m, 18H), 0.88 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 134.9, 118.0, 70.7, 41.9, 36.8, 31.9, 29.7, 29.6, 29.3, 25.7, 22.7, 14.1; MS *m/z* (CI) 230 (M+NH₄⁺, 100), 211 (34), 169 (25), 97 (9); HRMS calcd for C₁₄H₃₂NO (M+NH₄⁺) 230.2484, found 230.2478.

(*E*)-2-Pentadecen-5-ol⁶ **16h**

Following general procedure D, the addition of 1,2-epoxydodecane (100 mg, 0.54 mmol) to (*E*)-propenyllithium⁹ (2.0 M in Et₂O; 0.81 mL, 1.63 mmol) in hexane (8 mL) gave after workup and column chromatography (15% Et₂O in petrol) *alcohol* **16h** (122 mg, 99%, *E:Z* = 97:3 by GCMS analysis, *t_R* *E*-**16h** 12.77 min, *t_R* *Z*-**16h** 12.95 min, initial temp. 100 °C, max. temp. 280 °C, rate 8 °C/min) as a colourless oil; *R_f* 0.16 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3351br, 2925m, 2855m, 1466w, 967w; ¹H NMR (400 MHz) δ 5.56 (dq, *J* = 15, 6, 1H), 5.48–5.40 (m, 1H), 3.61–3.54 (m, 1H), 2.27–2.19 (m, 1H), 2.09–2.01 (m, 1H), 1.70 (dd, *J* = 6.3, 1.1, 3H), 1.60 (br, 1H), 1.47–1.23 (m, 18H), 0.88 (t, *J* = 6.8, 3H); ¹³C NMR (100 MHz) δ 128.9, 127.2, 70.9, 40.7, 36.8, 31.9, 29.7, 29.6, 29.3, 25.7, 22.7, 18.1, 14.1; MS *m/z* (CI) 244 (M+NH₄⁺, 100), 242 (63), 227 (35), 226 (37), 225 (63), 207 (20), 169 (41); HRMS calcd for C₁₅H₃₄NO (M+NH₄⁺) 244.2640, found 244.2644.

(*Z*)-2-Pentadecen-5-ol⁶ **16h**

Following general procedure D, the addition of 1,2-epoxydodecane (100 mg, 0.54 mmol) to (*Z*)-propenyllithium⁹ (1.4 M in Et₂O; 1.16 mL, 1.63 mmol) in hexane (8 mL) gave after column chromatography (15% Et₂O in petrol) *alcohol* **16h** (111 mg, 90%, *Z:E* = 99:1 by GCMS analysis, *t_R* *E*-**16h** 12.77 min, *t_R* *Z*-**16h** 12.95 min, initial temp. 100 °C, max. temp. 280 °C, rate 8 °C/min) as a colourless oil; *R_f* 0.16 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3349br, 3017m, 2923s, 2854s, 1657w, 1467s, 1077m, 1027m; ¹H NMR (400 MHz) δ 5.66 (dq, *J* = 11, 7, 1H), 5.49–5.40 (m, 1H), 3.64 (quint, *J* = 6, 1H), 2.23 (t, *J* = 6, 2H), 1.65 (d, *J* = 7, 3H), 1.57 (br, 1H), 1.52–1.20 (m, 18H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 127.3, 126.2, 71.5, 36.9, 35.0, 31.9, 29.7, 29.6, 29.3, 25.8, 22.7, 14.1, 13.0; MS *m/z* (CI) 244 (M+NH₄⁺, 100), 242 (81), 227 (30), 226 (40), 225 (69), 208 (25), 169 (19); HRMS calcd for C₁₅H₃₄NO (M+NH₄⁺) 244.2640, found 244.2639.

1-(Trimethylsilyl)tridecan-3-ol⁶ **16l**

Following general procedure D, the addition of 1,2-epoxydodecane **13** (100 mg, 0.54 mmol) to (trimethylsilylmethyl)lithium (1.0 M in pentane, 1.63 mL, 1.63 mmol) in THF (8 mL) gave after column chromatography (10% Et₂O in petrol) *alcohol* **16l** (123 mg, 84%) as a colorless oil; *R_f* 0.20 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3346br, 2925s, 2854s, 1466m, 1378m, 1248s, 1179m, 1127m, 1076m, 1007m; ¹H NMR (400 MHz) δ 3.50–3.44 (m, 1H), 1.74 (br, 1H), 1.48–1.20 (m, 20H), 0.86 (t, *J* = 7, 3H), 0.59 (ddd, *J* = 14, 14, 4.5, 1H) 0.40 (ddd, *J* = 14, 13, 5, 1H), –0.02 (s, 9H); ¹³C NMR (100 MHz) δ 74.1, 36.7, 31.9, 31.5, 29.8, 29.6, 29.3, 25.7, 22.7, 14.1, 12.0, –1.8; MS *m/z* (CI) 290 (M+NH₄⁺, 100), 288 (18), 272 (25), 92 (26), 91 (34), 90 (83); HRMS calcd for C₁₆H₄₀NOSi (M+NH₄⁺) 290.2874, found 290.2877.

6-Trimethylsilyl-octadecan-8-ol⁶ **16m**

Following general procedure D, the addition of 1,2-epoxydodecane (100 mg, 0.54 mmol) to 1-(trimethylsilyl)hexyllithium [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 1.02 mL, 1.63 mmol) to trimethylvinylsilane (196 mg, 1.96 mmol) at –78 °C in THF (2 mL)] in hexane (8 mL) gave after column chromatography (10% Et₂O in petrol) *alcohol* **16m** (130 mg, 70% 50:50 diastereoisomers by GC analysis, *t_R* 11.23 min, *t_R* 11.28 min, initial temp. 80 °C, max. temp. 280 °C, rate = 20 °C/min) as a colorless wax; *R_f* 0.24 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3360br, 2925s, 2855s, 1730w, 1467m, 1378w, 1248s, 834m; ¹H NMR (400 MHz) δ 3.70–3.60 (m, 1H), 1.60 (br, 1H), 1.58–1.20 (m, 27H), 0.88 (t, *J* = 6.5, 6H), 0.81–0.75 (m, 1H), 0.69–0.62 (m, 1H), –0.01 (s, 9H); ¹³C NMR (100 MHz) δ 71.5, 70.0, 38.2, 37.7, 37.3, 32.5, 32.3, 31.9, 29.8, 29.7, 29.7, 29.6, 29.3, 29.0, 28.8, 25.8, 25.6, 22.7, 22.6, 22.6, 22.2, 21.9, 14.1, –2.2, –2.3; MS *m/z* (CI) 360 (M+H⁺, 52), 325 (5), 90 (100); HRMS: *m/z* calcd for C₂₁H₅₀SiNO (M+NH₄⁺) 360.3656, found 360.3651.

When alcohol **16n** was observed it was identified as follows:

Hexadecan-6-ol¹⁹ **16n**

White solid; mp 41–42 °C; *R_f* 0.13 (10% Et₂O in petrol); IR (CHCl₃)/cm⁻¹ 3340br, 2920w, 2859w, 1467w; ¹H NMR (400 MHz) δ 3.63–3.56 (m, 1H), 1.51–1.19 (m, 27H), 0.93–0.83 (m,

(19) Ohta, H.; Fujiwara, H.; Tsuchihashi, G. *Agric. Biol. Chem.* **1984**, 48, 1509–1516.

6H); ^{13}C NMR (100 MHz) δ 72.0, 37.5, 37.4, 31.9, 29.7, 29.6, 29.6, 29.3, 25.6, 25.3, 22.7, 22.6, 14.1, 14.0; MS m/z (CI) 260 ($\text{M}+\text{NH}_4^+$, 100), 258 (83), 152 (10); HRMS calcd for $\text{C}_{16}\text{H}_{38}\text{NO}$ ($\text{M}+\text{NH}_4^+$) 260.2945, found 260.2945.

When alcohol **16o** was observed it was identified as follows:

Tridecan-3-ol²⁰ **16o**

Colorless oil; R_f 0.15 (10% Et_2O in petrol); IR (neat)/ cm^{-1} 3347br, 2925s, 2855s, 1464w, 967w; ^1H NMR (400 MHz) δ 3.56–3.50 (m, 1H), 1.58–1.20 (m, 21H), 0.95 (t, $J = 7.5$, 3H), 0.89 (t, $J = 7$, 3H); ^{13}C NMR (100 MHz) δ 73.3, 36.9, 31.9, 30.1, 29.7, 29.6, 29.3, 25.7, 22.7, 14.1, 9.9; MS m/z (CI) 218 ($\text{M}+\text{NH}_4^+$, 34), 216 (100), 152 (30), 90 (38), 60 (88); HRMS calcd for $\text{C}_{13}\text{H}_{28}\text{NO}$ ($\text{M}+\text{NH}_4^+$) 218.2478, found 218.2479.

(3*E*,5*Z*)-3,5-*tert*-Butyldodecadienyl(oxydimethyl)silane **19**

Following general procedure B, the addition of *tert*-butyldimethyl(2-oxiranyloxy)silane²¹ **17** (241 mg, 1.19 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 1.49 mL, 2.38 mmol) to TMP (336 mg, 2.38 mmol) in hexane (11 mL)] and (*Z*)-1-octenyllithium [prepared by the addition of *t*-BuLi (1.5 M in pentane; 2.06 mL, 3.09 mmol) to (*Z*)-1-iodo-1-octene¹⁴ (368 mg, 1.55 mmol) in Et_2O (2 mL)] gave after workup and column chromatography (100% petrol $\rightarrow$ 3% Et_2O in petrol) *diene* **19** (247 mg, 70%, *E,Z:E,E*:other isomer = 92:5:3 by GCMS analysis; t_R other isomer 14.73 min, t_R *E,Z*-**19** 15.58 min, t_R *E,E*-**19** 15.93 min, initial temp. 100 $^\circ\text{C}$, max. temp. 280 $^\circ\text{C}$, rate 8 $^\circ\text{C}/\text{min}$) as a colorless oil; R_f 0.18 (100% petrol); IR (neat)/ cm^{-1} 2928s, 2857s, 1471w, 1256m, 1103s, 836s; ^1H NMR (400 MHz) δ 6.37 (ddd, $J = 15, 11, 1.5$, 1H), 5.95 (dd, $J = 11, 11$, 1H), 5.65 (dt, $J = 15, 7$, 1H), 5.34 (dt, $J = 11, 7$, 1H), 3.66 (t, $J = 7$, 2H), 2.33 (t, $J = 7$, 2H), 2.16 (t, $J = 7$, 2H), 1.42–1.25 (m, 8H), 0.91–0.88 (m, 12H), 0.06 (s, 6H); ^{13}C NMR (100 MHz) δ 130.8, 130.3, 128.4, 127.6, 63.0, 36.5, 31.8, 29.7, 29.0, 27.7, 25.9, 22.6, 18.3, 14.1, -5.2 ; MS m/z (CI) 297 ($\text{M}+\text{H}^+$, 16), 256 (15), 239 (56), 165 (100) 74 (18); HRMS calcd for $\text{C}_{18}\text{H}_{37}\text{OSi}$ ($\text{M}+\text{H}^+$) 297.2614, found 297.2613.

(20) Yosioka, I. *Chem. Pharm. Bull.* **1974**, 22, 1943–1945.

(21) (a) Pirrung, M. C.; Webster, M. C.; Nicholas, J. G. *J. Org. Chem.* **1987**, 52, 3603–3613. (b) Ficini, J.; Barbara, C.; Desmaele, D.; Overfelli, O. *Heterocycles* **1987**, 25, 329–332.

(3*E*,5*Z*)-3,5-dodecadienyl acetate²² **20**

Diene **19** (200 mg, 0.67 mmol) was added to a solution of TBAF (1.0 M in THF; 0.67 mL, 0.67 mmol) and stirred for 1 h. After the addition of Et₂O (3 mL) and sat. brine solution (3 mL), the layers were separated. The aqueous layer was extracted with Et₂O (2 x 3 mL), the combined organic layers were dried (MgSO₄) and solvent evaporated *in vacuo* to give the crude *alcohol*. To this was added Ac₂O (0.11 mL, 1.1 mmol), pyridine (0.16 mL, 2.01 mmol) and DMAP (15 mg, 0.12 mmol) and the reaction was left stirring for 3 h at rt. After addition of 1 M HCl solution (4 mL), the layers were separated. The aqueous layer was extracted with Et₂O (5 mL), the combined organic layers were dried (MgSO₄) and solvent evaporated *in vacuo*. The residue was purified by column chromatography (3% Et₂O in petrol) to give *diene* **20** (120 mg, 80%, *E,Z:E,E*:other isomer = 92:5:3 by GCMS analysis; *t_R* other isomer **20** 10.02 min, *t_R* *E,Z*-**20** 10.48 min, *t_R* *E,E*-**20** 10.68 min, initial temp. 100 °C, max. temp. 280 °C, rate 8 °C/min) as a colorless oil; *R_f* 0.15 (3% Et₂O in petrol). Further purification on AgNO₃ impregnated silica²³ (3% Et₂O in petrol) gave *diene* **20** (96 mg, 64%, *E,Z:E,E* = 99:1 by GCMS analysis); IR (neat)/cm⁻¹ 2927s, 2856m, 1744s, 1458m, 1364m, 1237s, 1035m; ¹H NMR (400 MHz) δ 6.39 (ddd, *J* = 15, 11, 1.5, 1H), 5.95 (dd, *J* = 11, 11, 1H), 5.60 (dt, *J* = 15, 7, 1H), 5.37 (dt, *J* = 11, 7, 1H), 4.12 (t, *J* = 7, 2H), 2.43 (q, *J* = 7, 2H), 2.16 (q, *J* = 7, 2H), 2.06 (s, 3H) 1.42–1.22 (m, 8H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 171.1, 131.6, 128.6, 128.2, 128.0, 63.8, 32.1, 31.7, 29.6, 28.9, 27.7, 22.6, 21.0, 14.1; MS *m/z* (CI) 242 (M+NH₄⁺, 100), 165 (92), 164 (85), 93 (29), 80 (94); HRMS calcd for C₁₄H₂₈NO₂ (M+NH₄⁺) 242.2120, found 242.2122.

(*E*)-(4-Methyl-2-pentenyl)trimethylsilane²⁴ **21** (from Table 4, entry 6 conditions)

Following general procedure A, the addition of isopropylloxirane (123 mg, 1.43 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 1.79 mL, 2.86 mmol) to TMP (404 mg, 2.86 mmol) in THF (21 mL)] and (trimethylsilylmethyl)lithium (1.0 M in pentane; 1.86 mL, 1.86 mmol) gave after workup and column chromatography (100% petrol) *allylsilane* **21** (159 mg, 71%, *E:Z* = 98:2, by GCMS analysis, *t_R* *E*-**21** 3.15 min, *t_R* *Z*-**21** 3.23, min initial temp. 60 °C, max. temp. 240 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.66 (100% petrol); IR (neat)/cm⁻¹ 2957s, 2926s, 2871m, 1466w, 1248s, 859s, 840s; ¹H NMR (400 MHz) δ

(22) Chong, J. M.; Heaft, M. A. *Tetrahedron* **1999**, *55*, 14243–14250.

(23) Williams, C. M.; Mander, L. N. *Tetrahedron* **2001**, *57*, 425–447.

(24) Shimizu, N.; Imazu, S.; Shibata, F.; Tsuno, Y. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 1122–1128.

5.39–5.30 (m, 1H), 5.23 (dd, $J = 15, 7, 1\text{H}$), 2.24 (d, sept, $J = 7, 7, 1\text{H}$), 1.39 (d, $J = 7.5, 2\text{H}$), 0.97 (d, $J = 7, 6\text{H}$), 0.00 (s, 9H); ^{13}C NMR (100 MHz) δ 136.4, 122.8, 31.3, 23.0, 22.4, -2.1 ; MS m/z (CI) 157 ($\text{M}+\text{H}^+$, 12), 91 (12), 90 (100); HRMS calcd for $\text{C}_9\text{H}_{22}\text{Si}$ ($\text{M}+\text{H}^+$) 157.1413, found 157.1416.

When *Z*-isomer **21** was formed in appreciable amounts along with the *E*-isomer (Table 4, entry 5) it was identified from the vinylic/allylic protons and allylic carbons: ^1H NMR (400 MHz) δ 5.10 (dd, $J = 10, 9.5, 1\text{H}$), 2.53 (d, sept, $J = 7, 7, 1\text{H}$), 1.48 (d, $J = 8, 2\text{H}$); ^{13}C NMR (100 MHz) δ 18.4.

(*E*)-(4,4-Dimethyl-2-pentenyl)trimethylsilane²⁴ **22** (from Table 4, entry 9 conditions)

Following general procedure A, the addition of isobutyloxirane (123 mg, 1.23 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 1.16 mL, 1.85 mmol) to TMP (261 mg, 1.85 mmol) in THF (18 mL)] and (trimethylsilylmethyl)lithium (1.0 M in pentane; 1.85 mL, 1.85 mmol) gave after workup and column chromatography (100% petrol) *allylsilane* **22** (132 mg, 63%, *E*-only by GCMS analysis, t_R *E*-**22** 4.33 min, initial temp. 50 °C, max. temp. 240 °C, rate 20 °C/min) as a colorless oil; R_f 0.63 (100% petrol); IR (neat)/ cm^{-1} 2925s, 2854s, 2361w, 1464w, 1378m, 1248w, 1156m, 969m, 851m; ^1H NMR (400 MHz) δ 5.30–5.25 (m, 2H), 1.39 (dd, $J = 5, 2, 2\text{H}$), 0.99 (s, 9H), -0.01 (s, 9H); ^{13}C NMR (100 MHz) δ 140.2, 120.4, 32.9, 30.0, 22.4, -2.1 ; MS m/z (CI) 171 ($\text{M}+\text{H}^+$, 27), 155 (9), 148 (10), 91 (12), 90 (100), 73 (23); HRMS calcd for $\text{C}_{10}\text{H}_{23}\text{Si}$ ($\text{M}+\text{H}^+$) 171.1569, found 171.1561.

When *Z*-isomer **22** was formed in appreciable amounts along with the *E*-isomer (Table 4, entry 8) it was identified as follows: ^1H NMR (400 MHz) δ 5.25–5.15 (m, 2H), 1.61 (dd, $J = 5, 2\text{ Hz}$, 2H), 1.11 (s, 9H), 0.05 (s, 9H); ^{13}C NMR (100 MHz) δ 137.6, 124.0, 31.3, 22.3, 19.5, -2.3 .

When alcohol **23** was observed it was identified as follows:

4-Methyl-(1-trimethylsilyl)pentan-3-ol **23**

Colorless oil; R_f 0.13 (10% Et_2O in petrol); IR (neat)/ cm^{-1} 3368br, 2958s, 2875s, 1470m, 1248s, 864s, 836s; ^1H NMR (400 MHz) δ 3.27 (dt, $J = 8, 5, 1\text{H}$), 1.75–1.65 (m, 1H), 1.54–1.30 (m, 3H), 0.93 (d, $J = 7, 3\text{H}$), 0.91 (d, $J = 7, 3\text{H}$), 0.65 (dtd, $J = 13, 4.5, 1.5, 1\text{H}$), 0.41 (dtd, $J = 12.5, 5, 1.5, 1\text{H}$), 0.00 (s, 9H); ^{13}C NMR (100 MHz) δ 79.0, 32.7, 28.2, 19.1, 17.0, 12.2, -1.8 ; MS m/z

(CI) 192.2 ($M+NH_4^+$, 13), 157.1 (30), 90.1 (100), 91.1 (28); HRMS calcd for $C_9H_{26}NOSi$ ($M+NH_4^+$) 192.1778, found 192.1778.

When alcohol **24** was observed it was identified as follows:

4,4-Dimethyl-(1-trimethylsilyl)pentan-3-ol²⁵ **24**

Colorless oil; R_f 0.26 (15% Et_2O in petrol); IR (neat)/ cm^{-1} 3368br, 2958s, 2875s, 1470m, 1250s; 1H NMR (400 MHz) δ 3.09 (dd, $J = 10.5, 2$, 1H), 1.55 (dddd, $J = 14, 13, 10, 4$, 1H), 1.46 (br, 1H), 1.19 (dddd, $J = 14, 12, 4.8, 1.8$, 1H), 0.89 (s, 9H), 0.80 (ddd, $J = 14, 13, 4.5$, 1H), 0.36 (ddd, $J = 14, 12, 5$, 1H), 0.00 (s, 9H); ^{13}C NMR (100 MHz) δ 82.6, 35.0, 25.8, 25.4, 13.6, -1.7; MS m/z (CI) 206 ($M+NH_4^+$, 50); 90 (100); HRMS calcd for $C_{10}H_{28}NOSi$ ($M+NH_4^+$) 206.1940, found 206.1941.

When aldehyde **25** was observed it was identified as follows:

3-Methylbutanal²⁶ **25**

Colorless oil; R_f 0.40 (5% Et_2O in petrol); 1H NMR (400 MHz) δ 9.70 (t, $J = 1.5$, 1H), 2.28 (dd, $J = 7, 1.5$, 2H), 2.21 (d, sept, $J = 7, 6.5$, 1H), 0.97 (d, $J = 6.5$, 6H); ^{13}C NMR (100 MHz) δ 205.3, 52.7, 24.8, 22.5, 22.1.

When aldehyde **26** was observed it was identified as follows:

3,3-Dimethylbutanal²⁷ **26**

Colorless oil; R_f 0.50 (10% Et_2O in petrol); IR (neat)/ cm^{-1} 2960s, 2905s, 1723w, 1455w; 1H NMR (400 MHz) δ 9.90 (t, $J = 1.5$, 1H), 2.22 (d, $J = 1.5$, 2H), 1.06 (s, 9H); ^{13}C NMR (100 MHz) δ 205.3, 56.2, 31.4, 30.0; MS m/z (CI) 118 ($M+NH_4^+$, 100), HRMS calcd for $C_6H_{16}NO$ ($M+NH_4^+$) 118.1232, found 118.1233.

(25) Mizojiri, R.; Urabe, H.; Sato, F. *J. Org. Chem.* **2000**, 65, 6217-6222.

(26) Spectra matched commercially available (Aldrich, product no. 59820) material.

(27) Guo, Z.; Sawyer, R.; Prakash, I. *Synth. Commun.* **2001**, 3395-3400.

When allylic alcohol **27** was observed it was identified as follows:

1-Tridecen-3-ol²⁸ **27**

Colorless oil; R_f 0.13 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3355s, 1645m, 1465w, 990s; ¹H NMR (400 MHz) δ 5.92–5.83 (m, 1H), 5.16 (dq, J = 17, 1.5, 1H), 5.11 (dq, J = 10.5, 1.5, 1H), 4.10 (q, J = 6, 1H), 1.60 (br, 1 H), 1.55–1.21 (m, 18 H), 0.88 (t, J = 6.5, 3H); ¹³C NMR (100 MHz) δ 141.3, 114.5, 73.3, 37.0, 31.9, 29.6, 29.5, 29.3, 25.3, 22.7, 14.1; MS m/z (CI) 216 (M+NH₄⁺, 61), 198 (100), 180 (84), 81 (42); HRMS calcd for C₁₃H₃₀NO (M+NH₄⁺) 216.2327, found 216.2323.

When allylic alcohol **28** was observed it was identified as follows:

4-Methyl-1-penten-3-ol²⁹ **28**

Colorless oil; R_f 0.23 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3361br, 3052s, 1717w; ¹H NMR (400 MHz) δ 5.91 (ddd, J = 6.5, 10.5, 17, 1H), 5.30 (ddd, J = 17, 1.5, 1, 1H), 5.13 (ddd, J = 10.5, 1.5, 1, 1H), 4.05 (dddd, J = 7, 6.5, 1, 1, 1H), 2.6 (br, 1H), 1.76 (d, sept, J = 7, 6.5, 1H), 1.24 (d, J = 6.5, 3H), 0.96 (d, J = 6.5, 3H); ¹³C NMR (100 MHz) δ 139.3, 115.5, 78.1, 33.4, 17.7, 15.1; MS m/z (CI) 118 (M+NH₄⁺, 100), HRMS calcd for C₆H₁₆NO (M+NH₄⁺) 118.1232, found 118.1234.

When allylic alcohol **29** was observed it was identified as follows:

4,4-Dimethyl-1-penten-3-ol³⁰ **29**

Colorless oil; R_f 0.30 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3404br, 2956s, 1365s, 1073m; ¹H NMR (400 MHz) δ 5.91 (ddd, J = 16.5, 9.5, 6.5, 1H), 5.18 (ddd, J = 16.5, 1.5, 1, 1H), 5.12 (ddd, J = 9.5, 1.5, 1, 1H), 3.70 (ddd, J = 6.5, 1, 1, 1H), 2.20 (br, 1H), 0.91 (s, 9H); ¹³C NMR (100 MHz) δ 139.3, 115.5, 78.1, 25.4, 13.6; MS m/z (CI) 132 (M+NH₄⁺, 100), HRMS calcd for C₇H₁₈NO (M+NH₄⁺) 132.1388, found 132.1395.

(28) Yazbak, A.; Sinha, S. C.; Keinan, E. *J. Org. Chem.* **1998**, 63, 5863–5868.

(29) Concellón, J. M.; Llavon, L.; Bernard Jr., P. L. *Tetrahedron* **1995**, 51, 5573–5584.

(30) Craig, D.; Pennington, M. W.; Warner, P. *Tetrahedron* **1999**, 55, 13495–13512.

(E)-(1-Pentyl-2,10-undecadien-1-yl)trimethylsilane⁶ **30**

Following general procedure C, using 1,2-epoxy-9-decene (83 mg, 0.54 mmol) gave after workup and reverse phase column chromatography¹² (C₁₈ silica, MeCN) *allylsilane* **30** (120 mg, 75%, *E*-only by GCMS analysis, *E*-**30** 11.57 min, initial temp. 80 °C, max. temp. 280 °C, rate 15 °C/min) as a colorless oil; *R*_f 0.13 (C₁₈ silica, MeCN); IR (neat)/cm⁻¹ 3078s, 2926s, 2854s, 1642w, 1464w, 1247s, 836s; ¹H NMR (400 MHz) δ 5.87–5.77 (m, 1H), 5.24–5.12 (m, 2H), 5.03–4.92 (m, 2H), 2.05 (dt, *J* = 7, 7, 2H), 1.99 (dt, *J* = 6.5, 6.5, 2H), 1.44–1.14 (m, 17H), 0.89 (t, *J* = 7, 3H), –0.05 (s, 9H); ¹³C NMR (100 MHz) δ 139.3, 131.5, 128.1, 114.1, 33.8, 32.9, 32.8, 31.8, 30.0, 29.0, 28.9, 28.9, 28.8, 22.7, 14.1, –3.2; MS *m/z* (CI) 295 (M+H⁺, 40), 220 (29), 90 (100); HRMS calcd for C₁₉H₃₉Si (M+H⁺) 295.2821, found 295.2822.

(E)-(1-pentyl-5-phenyl-2-penten-1-yl)trimethylsilane⁶ **31**

Following general procedure C, using 1,2-epoxy-4-phenylbutane³¹ (80 mg, 0.54 mmol) gave after workup and column chromatography (100% petrol) *allylsilane* **31** (109 mg, 70%, *E*:*Z* = 98:2 by GCMS analysis, *t*_R *Z*-**31** 12.33 min, *t*_R *E*-**31** 12.43 min, initial temp. 80 °C, max. temp. 280 °C, rate 15 °C/min) as a colorless oil; *R*_f 0.48 (100% petrol); IR (neat)/cm⁻¹ 3064w, 3027m, 2925s, 2853s, 1605w, 1496m, 1454m, 1247s; ¹H NMR (400 MHz) δ 7.37–7.28 (m, 2H), 7.24–7.18 (m, 3H), 5.31–5.20 (m, 2H), 2.70 (t, *J* = 7.5, 2H), 2.40 (dt, *J* = 7.5, 7.5, 2H), 1.47–1.13 (m, 9H), 0.92 (t, *J* = 7, 3H), –0.04 (s, 9H); ¹³C NMR (100 MHz) δ 142.2, 132.4, 128.5, 128.2, 127.0, 125.6, 36.6, 34.7, 33.0, 31.8, 29.0, 28.8, 22.6, 14.2, –3.2; MS *m/z* (CI) 289 (M+H⁺, 100) 215 (19), 214 (91), 197 (29), 123 (23), 90 (40); HRMS calcd for C₁₉H₃₃Si (M+H⁺) 289.2352, found 289.2344.

(E)-1-(*tert*-Butyldimethylsilyloxy)-6-(trimethylsilyl)-4-undecene⁶ **32**

Following general procedure C, using 1-*tert*-butyldimethylsilyloxy-4,5-epoxypentane³² (117 mg, 0.54 mmol) gave after workup and column chromatography (100% petrol) *allylsilane* **32** (126 mg, 65%, *E*-only by GCMS analysis, *E*-**32** 10.17 min, initial temp. 80 °C, max. temp. 280 °C, rate 15 °C/min) as a colorless oil; *R*_f 0.20 (100% petrol); IR (neat)/cm⁻¹ 2929m, 2857m, 1471w, 1388w, 1248m, 1105m, 969w, 836m; ¹H NMR (400 MHz) δ 5.25–5.14 (m, 2H), 3.62 (t, *J* = 7,

(31) Elings, J. A.; Downing, R. S.; Sheldon, R. A. *Eur. J. Org. Chem.* **1999**, 837–846.

(32) Yang, L.; Weber, A. E.; Greenlee, W. J.; Patchett, A. A. *Tetrahedron Lett.* **1993**, 34, 7035–7038.

2H), 2.05 (dt, $J = 7, 7$, 2H), 1.58 (dd, $J = 7, 7$, 2H), 1.47–1.12 (m, 9H), 0.91 (s, 9H), 0.88 (t, $J = 7$, 3H), 0.06 (s, 6H), –0.05 (s, 9H); ^{13}C NMR (100 MHz) δ 131.9, 127.5, 62.8, 33.2, 33.0, 31.8, 29.1, 29.0, 28.8, 26.0, 22.6, 18.4, 14.2, –3.1, –5.3; MS m/z (CI) 357 ($\text{M}+\text{H}^+$, 31), 299 (32), 269 (21), 242 (16), 228 (19), 225 (77), 164 (43), 147 (100), 90 (46), 73 (55); HRMS calcd for $\text{C}_{21}\text{H}_{45}\text{Si}$ ($\text{M}+\text{H}^+$) 357.3009, found 357.3011.

(*E*)-(3-Cyclohexyl-1-pentyl-allyl)trimethylsilane **33**

Following general procedure C, using 2-cyclohexyloxirane³³ (69 mg, 0.54 mmol) gave after workup and reverse phase column chromatography¹² (C_{18} silica, 10% CH_2Cl_2 in MeCN) *allylsilane* **33** (115 mg, 80%, *E*-only by GCMS analysis, *E*-**33** 12.63 min, initial temp. 80 °C, max. temp. 280 °C, rate 10 °C/min) as colorless oil; R_f 0.22 (C_{18} silica, 10% CH_2Cl_2 in MeCN); IR (neat)/ cm^{-1} 2920s, 2852s, 1449s, 1247s, 968s, 836; ^1H NMR (400 MHz) δ 5.20–5.09 (m, 2H), 1.95–1.88 (m, 1H), 1.72–1.62 (m, 6H), 1.42–1.01 (m, 13H), 0.88 (t, $J = 7$, 3H), –0.05 (s, 9H); ^{13}C NMR (100 MHz) δ 134.4, 128.8, 41.1, 33.8, 33.7, 32.8, 31.7, 28.9, 28.8, 26.3, 26.2, 22.6, 14.1, –3.2; MS m/z (CI) 267 ($\text{M}+\text{H}^+$, 45), 193 (23), 192 (100), 90 (61), 73 (69); HRMS calcd for $\text{C}_{17}\text{H}_{35}\text{Si}$ ($\text{M}+\text{H}^+$) 267.2508, found 267.2507.

(*E*)-1-(3-Cyclohexyl-1-pentyl-allyl)dimethylphenylsilane **34**

Following general procedure C, using 2-cyclohexyloxirane³³ (69 mg, 0.54 mmol) and dimethylphenylvinylsilane (136 mg, 0.84 mmol) instead of vinyltrimethylsilane gave after workup and reverse phase column chromatography¹² (C_{18} silica, 100% MeCN) *allylsilane* **34** (128 mg, 72%, *E*-only by GCMS analysis, *E*-**34** 14.45 min, initial temp. 80 °C, max. temp. 280 °C, rate 15 °C/min) as colorless oil; R_f 0.39 (C_{18} silica, 100% MeCN); IR (neat)/ cm^{-1} 3070m, 2923s, 2852s, 1449m, 1428m, 1247m, 1113m, 969m; ^1H NMR (400 MHz) δ 7.51–7.48 (m, 2H), 7.36–7.35 (m, 3H), 5.19–5.08 (m, 2H), 1.95–1.87 (m, 1H), 1.72–1.60 (m, 6H), 1.41–1.03 (m, 13H), 0.85 (t, $J = 7$, 3H), 0.26 (s, 3H), 0.24 (3H); ^{13}C NMR (100 MHz) δ 138.4, 135.1, 134.1, 128.7, 128.2, 127.5, 41.1, 33.7, 33.6, 32.3, 31.6, 28.8, 28.8, 26.3, 26.1, 22.6, 14.1, –4.3, –5.1;

(33) Michnick, T. J.; Matteson, D. S. *Synlett* **1991**, 631–632.

MS m/z (CI) 329 ($M+H^+$, 6), 193 (10), 192 (38), 152 (100), 135 (68); HRMS calcd for $C_{22}H_{37}Si$ ($M+H^+$) 329.2665, found 329.2667.

(*E*)-Dimethyl(1-pentyl-2-pentenyl)phenylsilane **35**

Following general procedure C, using 1,2-epoxybutane (39 mg, 0.54 mmol) and dimethylphenylvinylsilane (136 mg, 0.84 mmol) instead of vinyltrimethylsilane gave after workup, and reverse phase column chromatography¹² (C_{18} silica, 100% MeCN) *allylsilane* **35** (107 mg, 72%, $E:Z$ = 95:5, by GCMS analysis, t_R *E*-**35** 11.28 min, t_R *Z*-**35** 11.43 min initial temp. 80 °C, max. temp. 280 °C, rate 15 °C/min) as a colorless oil; R_f 0.35 (C_{18} silica, 100% MeCN); IR (neat)/ cm^{-1} 3051m, 2958s, 2924s, 2852, 1460m, 1428m, 1248s, 1113s, 969s; 1H NMR (400 MHz) δ 7.53–7.48 (m, 2H), 7.36–7.30 (m, 3H), 5.29–5.22 (m, 1H), 5.18–5.11 (m, 1H), 2.00 (quint, J = 7.5, 2H), 1.65–1.58 (m, 1H), 1.44–1.06 (m, 8H), 0.95 (t, J = 7, 3H), 0.85 (t, J = 7, 3H), 0.26 (s, 3H), 0.25 (s, 3H); ^{13}C NMR (100 MHz) δ 138.4, 134.1, 130.5, 129.9, 128.7, 127.5, 32.3, 31.6, 31.5, 28.9, 25.9, 22.6, 14.5, 14.1, –4.3, –5.1; MS m/z (CI) 275 ($M+H^+$, 6), 153 (15), 152 (100), 135 (59); HRMS calcd for $C_{18}H_{31}Si$ ($M+H^+$) 275.2195, found 275.2188.

(*E*)-Trimethyl(1-pentyl-2-pentenyl)silane³⁴ **36**

Following general procedure C, using 1,2-epoxybutane (39 mg, 0.54 mmol) gave after workup, and reverse phase column chromatography¹² (C_{18} silica, 15% Et_2O in MeCN) *allylsilane* **36** (88 mg, 77%, $E:Z$ = 94:6, by GCMS analysis, t_R *E*-**36** 6.93 min, t_R *Z*-**36** 7.17 min, initial temp. 80 °C, max. temp. 280 °C, rate 10 °C/min) as a colorless oil; R_f 0.34 (C_{18} silica, 15% Et_2O in MeCN); IR (neat)/ cm^{-1} 2959s, 2925s, 2873s, 1461m, 1259m, 1247s, 968m, 836s; 1H NMR (400 MHz) δ 5.31–5.13 (m, 2H), 2.01 (quint, J = 7, 2H), 1.45–1.21 (m, 9H), 0.97 (t, J = 7, 3H), 0.89 (t, J = 7, 3H), –0.05 (s, 9H); ^{13}C NMR (100 MHz) δ 130.5, 129.8, 32.9, 31.8, 29.0, 28.9, 26.0, 22.6, 14.6, 14.1, –3.2; MS m/z (CI) 213 ($M+H^+$, 59), 138 (61), 90 (100); HRMS calcd for $C_{13}H_{29}Si$ ($M+H^+$) 213.2039, found 213.2043.

(34) Matsuda, I.; Kato, T.; Sato, S.; Izumi, Y. *Tetrahedron Lett.* **1986**, 27, 5747–5750.

Cis-1-Deuterio-1,2-epoxydecane³⁵ **37**

To a solution of 1-decyne (5.0 g, 36.2 mmol) in hexane (100 mL) at 0 °C was added *n*-BuLi (1.6 M in hexanes; 27.1 mL, 43.4 mmol) dropwise. The reaction mixture was stirred at this temperature for 5 min and then D₂O (25 mL) added and the reaction was stirred for a further 10 min at this temperature. The organic layer was separated, dried (MgSO₄) and solvent removed *in vacuo* to yield the crude deuterated alkyne (5.0 g). To a solution of deuterated alkyne (5.0 g) in pentane (20 mL) at rt was added DIBAL-H (1 M in hexanes; 43.4 mL, 43.4 mmol) dropwise. The reaction mixture was heated to 50 °C and stirred at this temperature for 3 h. After cooling to rt the reaction mixture was diluted with pentane (60 mL) then H₂O (10 mL) was added dropwise to the vigorously stirred solution which was then left for 12 h. The reaction mixture was filtered and solvent removed *in vacuo* to yield the crude *Z*-deuterated alkene (4.1 g). To a solution of the *Z*-deuterated alkene (4.1 g) in CH₂Cl₂ (100 mL) at 0 °C was added a solution of *m*CPBA (5.17 g, 30 mmol in 100 mL CH₂Cl₂) dropwise. The reaction was left at rt for 14 h then quenched with sat. aq. NaSO₃ solution (50 mL). The layers were separated and the organic layer was washed with sat. aq. NaHCO₃ solution (4 x 50 mL), then dried (MgSO₄) and solvent evaporated *in vacuo*. The residue was purified by column chromatography (5% Et₂O in petrol) to give *Z*-epoxide **37** (3.45 g, 61%, 97% D incorporation determined by mass spectrometry) as a colorless oil; *R*_f 0.29 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 2927s, 2857s, 2228w, 1467m, 1428m, 836m; ¹H NMR (400 MHz) δ 2.93–2.89 (m, 1H), 2.74 (d, *J* = 4, 1H), 1.55–1.25 (m, 14H), 0.89 (t, *J* = 7, 3H); ¹³C NMR (100 MHz) δ 52.3, 46.7, 32.4, 31.8, 29.5, 29.4, 29.2, 25.9, 22.6, 14.0; MS *m/z* (CI) 175 (100, M+H⁺), 158(37), 157 (27); HRMS calcd for C₁₀H₂₀DO (M+H⁺), 158.1655, found 158.1658.

Trans-1-Deuterio-1,2-epoxydecane³⁵ **37**

To a solution of 1-decyne (5.0 g, 36.2 mmol) in pentane (10 mL) at rt was added DIBAL-H (1 M in hexanes; 43.4 mL, 43.4 mmol) dropwise. The reaction mixture was warmed to 50 °C and stirred at this temperature for 3 h. After cooling to rt the reaction mixture was diluted with pentane (50 mL) then D₂O (10 mL) was added dropwise to the vigorously stirred solution which was left stirring for 12 h. The reaction mixture was filtered and solvent removed *in vacuo* to yield the crude *trans*-deuterated alkene (4.2 g). To a solution of the *trans*-deuterated alkene (4.2

(35) (a) Yanagisawa, H.; Katsutaka, Y.; Yamamoto, H. *J. Chem. Soc., Chem. Commun.* **1994**, 2103–2104.

(b) Patrick, D.W.; Truesdale, L. K.; Biller, S. A.; Sharpless, K. B. *J. Org. Chem.* **1978**, 43, 2628–2638.

g) in CH_2Cl_2 (100 mL) at 0 °C was added a solution of *m*CPBA (5.30 g, 30.7 mmol in 100 mL CH_2Cl_2) dropwise. The reaction was left at rt for 14 h then quenched with sat. aq. NaSO_3 solution (50 mL). The layers were separated and the organic layer was washed with sat. aq. NaHCO_3 solution (4 x 50 mL), then dried (MgSO_4) and solvent evaporated *in vacuo*. The residue was purified by column chromatography (5% Et_2O in petrol) to give the *E*-epoxide **37** (3.67 g, 65%, 96% D incorporation determined by mass spectrometry) as a colourless oil; R_f 0.20 (5% Et_2O in petrol); IR (neat)/ cm^{-1} 2927s, 2856s, 2242w, 1466m, 1431m, 901m; ^1H NMR (400 MHz) δ 2.90 (td, $J = 3, 5.5$, 1H), 2.45 (d, $J = 3$, 1H), 1.55–1.25 (m, 14), 0.89 (t, $J = 7$ Hz, 3H); ^{13}C NMR (100 MHz) δ 52.6, 46.8, 32.4, 31.8, 29.5, 29.4, 29.2, 25.9, 22.6, 14.0; MS m/z (CI) 175 (100, $\text{M}+\text{H}^+$), 158 (18), 157 (11); HRMS calcd for $\text{C}_{10}\text{H}_{20}\text{DO}$ ($\text{M}+\text{H}^+$) 158.1655, found 158.1653.

General procedure A was used for alkene **38** formation using *cis/trans*-deuterated epoxides **37**. The *E:Z* ratio was determined by GCMS analysis. The deuterium incorporation of each isomer was determined by analysis of the molecular ion peak in the mass spectrum and then verified by analysis of the ^1H NMR spectrum.

Benzylidenecyclododecane³⁶ **42** and (*Z*)-cyclododecenylmethanol³⁷

Following general procedure A, the addition of 1-oxa-spiro[2.11]tetradecane³³ **41** (100 mg, 0.51 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.48 mL, 0.77 mmol) to TMP (109 mg, 0.77 mmol) in hexane (7 mL)] and PhLi (2.0 M in *n*-Bu₂O; 0.38 mL, 0.77 mmol) gave after workup and column chromatography (100% petrol→10% Et_2O in petrol) alkene **42** (55 mg, 42%) as a colorless oil; R_f 0.58 (100% petrol); IR (neat)/ cm^{-1} 3452br, 3027w, 2936s, 2862s, 1600w, 1494s, 1470m, 1347w; ^1H NMR (400 MHz) δ 7.35–7.31 (m, 2H), 7.26–7.18 (m, 3H), 6.34 (s, 1H), 2.28 (t, $J = 7$, 2H), 2.23 (t, $J = 7$, 2H), 1.69–1.53 (m, 4H), 1.47–1.34 (m, 14H); ^{13}C NMR (100 MHz) δ 142.7, 138.9, 128.6, 128.0, 125.8, 125.5, 32.6, 27.9, 25.2, 24.9, 24.4, 24.2, 24.2, 24.1, 23.9, 23.0, 22.4; MS m/z (CI) 257 ($\text{M}+\text{H}^+$, 25), 256 (100), 148 (28); HRMS calcd for $\text{C}_{19}\text{H}_{29}$ ($\text{M}+\text{H}^+$) 257.2269, found 257.2258.

Also isolated was (*Z*)-cyclododecenylmethanol³⁷ (47 mg, 47%) as a colorless oil; R_f 0.10 (10% Et_2O in petrol); IR (neat)/ cm^{-1} 3320br, 2925s, 1461s, 1247m, 1013s; ^1H NMR (400 MHz) δ 5.51 (t, $J = 8$, 1H), 4.18 (s, 2H), 2.23–2.19 (m, 2H), 2.17–2.12 (m, 2H), 1.52–1.45 (m, 4H),

(36) Petasis, N. A.; Bzowej, E. I. *J. Org. Chem.* **1992**, 57, 1327–1330.

(37) Yasuda, A.; Yamamoto, H.; Nozaki, H. *Bull. Chem. Soc. Jpn.* **1979**, 52, 1705–1708.

1.38–1.22 (m, 13H); ^{13}C NMR (100 MHz) δ 136.9, 131.8, 59.2, 33.5, 27.4, 26.8, 26.6, 25.6, 25.6, 24.4, 24.4, 24.4, 23.5; MS m/z (CI) 197 ($\text{M}+\text{H}^+$, 3), 196 (25), 179 (100); HRMS calcd for $\text{C}_{13}\text{H}_{25}\text{O}$ ($\text{M}+\text{H}^+$) 197.1905, found 197.1900. The geometry of the double bond was determined by a NOSEY experiment.

(2-Cyclododecylideneethyl)trimethylsilane **43**

Following general procedure A, the addition of 1-oxa-spiro[2.11]tetradecane³³ **41** (236 mg, 1.02 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.96 mL, 1.53 mmol) to TMP (216 mg, 1.53 mmol) in hexane (8 mL)] and (trimethylsilyl)methylolithium (1.0 M in pentane; 1.53 mL, 1.53 mmol) gave after workup and column chromatography (100% petrol) *alkene* **43** (109 mg, 40%) as a colorless oil; R_f 0.63 (100% Et_2O in petrol); IR (neat)/ cm^{-1} 3006m, 2930s, 2854s, 1658w, 1466s, 1416m, 1380w, 1241s, 1152s; ^1H NMR (400 MHz) δ 5.23 (t, $J = 8$, 1H), 2.02 (q, $J = 7$, 4H), 1.57–1.26 (m, 20H), -0.01 (s, 9H); ^{13}C NMR (100 MHz) δ 135.6, 120.3, 32.1, 27.6, 24.7, 24.6, 24.5, 24.4, 24.3, 24.0, 23.9, 23.9, 23.1, 22.3, 18.3, -1.7 ; MS m/z (CI) 267 ($\text{M}+\text{H}^+$, 100), 266 (52), 192 (27), 90 (100); HRMS calcd for $\text{C}_{17}\text{H}_{35}\text{Si}$ ($\text{M}+\text{H}^+$) 267.2508, found 267.2496.

(*E*)- and (*Z*)-1,2-Diphenyl-1-propene³⁸ **45**

Following general procedure A, the addition of 2-methyl-2-phenyloxirane³³ **44** (100 mg, 0.75 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.71 mL, 1.13 mmol) to TMP (159 mg, 1.13 mmol) in hexane (8 mL)] and PhLi (2.0 M in *n*-Bu₂O; 0.57 mL, 1.13 mmol) gave after workup and column chromatography (100% petrol) *alkene* **45** (60 mg, 41%, *E*:*Z* = 84:16 by GCMS analysis, t_R *Z*-**45** 8.07 min, t_R *E*-**45** 9.02 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a white solid; R_f 0.25 (100% petrol); mp 80–81 °C; IR (KBr)/ cm^{-1} 3453s, 1625w, 1443m, 1079m; ^1H NMR (400 MHz) δ 6.72 (d, $J = 1.5$, 0.8H), 2.17 (d, $J = 1.5$, 2.4H) (*E*-isomer), 6.35 (d, $J = 1.5$, 0.2H), 2.10 (d, $J = 1.5$, 0.6H) (*Z*-isomer), 7.42–7.09 (m, 10H) (*E*- and *Z*-isomers); ^{13}C NMR (100 MHz) δ 144.5, 144.0, 138.9, 138.4, 128.0, 137.5, 129.7, 129.2, 129.0, 128.9, 128.7, 128.5, 128.4, 128.2, 127.9, 127.7, 127.2, 127.0, 126.5, 126.0, 18.0, 17.5 (*E*- and *Z*-isomers); MS m/z (CI) 195 ($\text{M}+\text{H}^+$, 88), 194 (100), 179 (40), 178 (28), 115 (15); HRMS calcd for $\text{C}_{15}\text{H}_{15}$ ($\text{M}+\text{H}^+$) 195.1174, found 195.1175.

(38) Tingoli, M.; Tiecco, M.; Testaferri, L.; Temperini, A.; Pelizzi, G.; Bacchi, A. *Tetrahedron* **1995**, *51*, 4691–4700.

(*E*)- and (*Z*)-1,2-Diphenyl-1-butene³⁹ **47**

Following general procedure A, the addition of 2-ethyl-2-phenyloxirane³³ **46** (100 mg, 0.68 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.64 mL, 1.02 mmol) to TMP (144 mg, 1.02 mmol) in hexane (8 mL)] and PhLi (2.0 M in *n*Bu₂O; 0.51 mL, 1.02 mmol) gave after workup and column chromatography (100% petrol) *alkene* **47** (88 mg, 62%, *E*:*Z* = 82:18 by GCMS analysis, *t_R* *Z*-**47** 7.04 min, *t_R* *E*-**47** 7.62 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.28 (100% petrol); IR (neat)/cm⁻¹ 3023m, 2966m, 2874m, 1599m, 1495m, 1445m, 1376m; ¹H NMR (400 MHz) δ 6.74 (s, 0.8H), 2.79 (q, *J* = 7.5, 1.6H) (*E*-isomer), 6.47 (s, 0.2H), 2.55 (q, *J* = 7.5, 0.4H) (*Z*-isomer), 7.66–7.07 (m, 10H), 1.11 (t, *J* = 7.5, 3H) (*E*- and *Z*-isomers); ¹³C NMR (100 MHz) δ 145.0, 144.5, 142.7, 141.5, 138.3, 137.6, 129.0, 128.8, 128.6, 128.5, 128.4, 128.3, 127.8, 127.6, 127.2, 126.8, 126.7, 126.6, 126.1, 125.1, 33.6, 23.3, 13.5, 12.9 (*E*- and *Z*-isomers); MS *m/z* (CI) 226 (M+NH₄⁺, 33), 209 (39), 208 (100), 193 (17), 178 (11); HRMS calcd for C₁₆H₂₀N (M+NH₄⁺) 226.1596, found 226.1601.

(*E*)- and (*Z*)-Trimethyl(3-phenyl-2-butenyl)silane⁴⁰ **48**

Following general procedure A, the addition of 2-methyl-2-phenyloxirane³³ **44** (100 mg, 0.75 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.71 mL, 1.13 mmol) to TMP (159 mg, 1.13 mmol) in hexane (8 mL)] and (trimethylsilyl)methylolithium (1.0 M in pentane; 1.13 mL, 1.13 mmol) gave after workup and column chromatography (100% petrol) *alkene* **48** (64 mg, 42%, *E*:*Z* = 76:24 by GCMS analysis, *t_R* *Z*-**48** 4.79 min, *t_R* *E*-**48** 5.66 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C/min) as a colorless oil; *R_f* 0.52 (100% petrol); IR (neat)/cm⁻¹ 3028w, 2955m, 1689w, 1599w, 1494m, 1446m, 1248s; ¹H NMR (400 MHz) δ 5.92 (tq, *J* = 9, 1.5, 0.8H), 2.03–2.02 (m, 2.4H), 1.70 (dd, *J* = 9, 1, 1.6H), 0.09 (s, 7.2H) (*E*-isomer), 5.52 (tq, *J* = 8.5, 1.5, 0.2H), 2.06–2.05 (m, 0.6H), 1.40 (dd, *J* = 8.5, 1, 0.4H), 0.01 (s, 1.8H) (*Z*-isomer), 7.41–7.31 (m, 4H), 7.27–7.31 (m, 1H) (*E*- and *Z*-isomers); ¹³C NMR (100 MHz) δ 144.4, 142.6, 134.2, 132.0, 128.3, 128.1, 128.1, 126.1, 126.0, 125.4, 124.9, 123.0, 25.9, 20.4, 19.3, 15.6, –1.5, –1.6 (*E*- and *Z*-isomers); MS *m/z* (CI) 205 (M+H⁺, 100), 90 (100), 73 (29); HRMS calcd for C₁₃H₂₁Si (M+H⁺) 205.1413, found 205.1409. The geometry of the double bond was determined by a NOSEY experiment.

(39) Stüdetmann, T.; Ibrahim-Ouali, M.; Knochel, P. *Tetrahedron* **1998**, *54*, 1299–1316.

(40) Shimizu, N.; Imazu, S.; Shibata, F.; Tsuno, Y. *Bull. Chem. Soc. Jpn.* **1991**, *64*, 1122–1128.

(E)- and *(Z)*-Trimethyl(3-phenyl-2-pentenyl)silane **49**

Following general procedure A, the addition of 2-ethyl-2-phenyloxirane³³ **46** (100 mg, 0.68 mmol) to a solution of LTMP [prepared by the addition of *n*-BuLi (1.6 M in hexanes; 0.64 mL, 1.02 mmol) to TMP (144 mg, 1.02 mmol) in hexane (8 mL)] and (trimethylsilyl)methylolithium (1.0 M in pentane; 1.02 mL, 1.02 mmol) gave after workup and column chromatography (100% petrol) *alkene* **49** (65 mg, 44% *E:Z* = 53:47 by GCMS analysis, t_R **Z-49** 4.12 min, t_R **E-49** 5.03 min, initial temp. 80 °C, max. temp. 280 °C, rate 20 °C) as a colorless oil; R_f 0.53 (100% petrol); IR (neat)/cm⁻¹ 3023w, 2958s, 1676w, 1599w, 1493w, 1248s, 1150m; ¹H NMR (400 MHz) δ 5.72 (t, J = 9, 0.5H), 2.51 (q, J = 8, 1H), 1.68 (d, J = 8.5, 1H), 0.99 (t, J = 7.5, 1.5H), 0.09 (s, 4.5H) (*E*-isomer), 5.48 (t, J = 9, 0.5H), 2.35 (q, J = 7.5, 1H), , 1.39 (d, J = 8.5, 1H), 0.98 (t, J = 7.5, 1.5H), -0.03 (s, 4.5H) (*Z*-isomer), 7.39–7.13 (m, 5H) (*E*- and *Z*-isomers); ¹³C NMR (100 MHz) δ 143.5, 141.9, 140.9, 139.1, 128.8, 128.1, 127.9, 126.2, 126.0, 126.0, 124.4, 121.4, 32.5, 22.3, 19.8, 19.0, 13.5, 13.3, -1.6, -1.7; MS m/z (CI) 219 ($M+H^+$, 100), 218 (61), 148 (12), 90 (80), 73 (20); HRMS calcd for C₁₄H₂₃Si ($M+H^+$) 219.1569, found 219.1573. The geometry of the double bond was determined by a NOSEY experiment.

When allylic alcohol **52** was observed it was identified as follows:

2-Dodecen-1-ol⁴¹ **52**

Colorless oil; *E:Z* = 87:13 (by ¹H NMR analysis of allylic protons in the δ 4.50–4.00 region); R_f 0.10 (10% Et₂O in petrol); IR (neat)/cm⁻¹ 3338m, 2924s, 2843s, 1670w; ¹H NMR (400 MHz) δ 4.09 (d, J = 5.5, 1.6H) (*E*-isomer), 4.19 (d, J = 6, 0.4H) (*Z*-isomer), 5.74–5.54 (m, 2 H), 2.04 (dt, J = 7.5, 7.5, 2H), 1.44–1.22 (m, 15H), 0.89 (t, J = 7, 3H) (*E*- and *Z*-isomer); ¹³C NMR (100 MHz) δ 133.6, 133.3, 128.8, 128.3, 63.8, 58.6, 32.2, 31.9, 29.6, 29.5, 29.3, 29.2, 29.1, 22.7, 14.1 (*E*- and *Z*-isomers); MS m/z (CI) 202 ($M+NH_4^+$, 22), 200 (44), 184 (100), 167 (34), 166 (100), 96 (33), 95 (45), 81 (60); HRMS calcd for C₁₂H₂₈NO ($M+NH_4^+$) 202.2171, found 202.2163.

(41) Barrett, A. G. M.; Head, J.; Smith, M. L.; Stock, N. S.; White, A. J. P.; Williams, D. *J. Org. Chem.* **1999**, *64*, 6005–6018.

(III) ^1H and ^{13}C NMR spectra of all products

S35

14b

S36

S37

S39

S40

S41

(*E,E*)-14h

S42

(*E,E*)-**14i**

S43

14j

S44

(Z,E)-14h

S45

S46

S47

S48

S49

14n

S50

14o

S51

S52

S53

16b

S54

S57

S58

S59

(Z)-16h

S60

S61

S62

S63

S64

S65

S66

S67

S68

S70

S71

S73

S79

S80

S85