SUPPORTING INFORMATION

The Three Component Reaction of Dicarbomethoxycarbene, Aldehydes and β -Nitrostyrenes: A Stereoselective Synthesis of Substituted Tetrahydrofurans.

Vijay Nair *, Sindhu Mathai and R. Luxmi Varma

Organic Chemistry Division, Regional Research Laboratory (CSIR), Trivandrum, India

*Corresponding author: vijaynair_2001@yahoo.com

Table of Contents :

^1H NMR and ^{13}C NMR Spectra of compounds 3a-3j

Experimental Section

General: All reactions were carried out in oven-dried glasswares. Progress of reaction was monitored by Thin Layer Chromatography while purification was effected using silica gel column chromatography. NMR spectra were recorded at 300 (^1H) and 75 (^{13}C) MHz respectively on a Bruker DPX-300 MHz NMR spectrometer. IR spectra were recorded in Bomem MB Series FT-IR spectrophotometer. Melting points were recorded on a Buchi melting point apparatus and are uncorrected.

Typical experimental procedure

To the mixture of 4-chloro- β -nitrostyrene **1** (1.20 mmol), *p*-tolualdehyde (0.30 mmol) and dimethyl diazomalonate (0.30 mmol) in 5ml of dry benzene, 2mol% Rh(II) acetate was added and refluxed for about 16 h under argon atmosphere. The reaction mixture was then subjected to chromatography on a silica gel column using 90:10 hexane-ethyl acetate solvent mixtures to afford the product

Tetrahydrofuran **3a**.

Colorless crystalline solid. **mp** 158-159 °C; **IR** (KBr) ν_{max} : 3036, 2955, 2854, 1742, 1553, 1492, 1438, 1371, 1303, 1249, 1115, 1027, 811, 670, 609 cm^{-1} ; **^1H NMR** (CDCl_3) δ 2.33 (s, 3H), 3.36 (s, 3H), 3.93 (s, 3H), 5.18 (d, 1H, $J = 6.2$ Hz), 5.68 (overlapping doublets, 1H, $J_1 = 7.2$ Hz, $J_2 = 6.4$ Hz), 6.16 (d, 1H, $J = 7.3$ Hz), 7.35-7.14 (m, 8H); **^{13}C NMR** (CDCl_3) δ 21.3,

52.6, 53.4, 53.6, 84.4, 90.5, 93.7, 126.2, 129.0, 129.3, 129.9, 130.6, 131.9, 134.8, 139.3, 167.0, 168.1, Anal. Calcd for $C_{21}H_{20}ClNO_7$: C, 58.14; H, 4.65; N, 3.23. Found: C, 58.58; H, 4.70; N, 3.47

Tetrahydrofuran 3b.

Colorless crystalline solid **mp** 147 °C; **IR** (KBr) ν_{max} : 3079, 3032, 2955, 1760, 1553, 1501, 1465, 1444, 1382, 1310, 1253, 1134, 1093, 1062, 1025, 855, 808, 756, 710, 612 cm^{-1} ; **^1H NMR** (CDCl_3) δ 3.36 (s, 3H), 3.94 (s, 3H), 5.18 (d, 1H, $J = 6.1$ Hz), 5.7 (overlapping doublets, 1H, $J_1 = 7.1$ Hz, $J_2 = 6.3$ Hz) 6.19 (d, 1H, $J = 7.3$ Hz) 7.35-7.28 (m, 9H); **^{13}C NMR** (CDCl_3) δ 52.6, 53.5, 53.7, 84.4, 90.5, 93.7, 126.3, 128.6, 129.1, 129.5, 130.0, 131.9, 133.6, 134.9, 168.1. Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{ClNO}_7$: C, 57.22; H, 4.22; N, 3.34. Found: C, 57.42; H, 4.00; N, 3.55.

Tetrahydrofuran 3c

Colorless crystalline solid. **mp** 188-190 °C; **IR** (KBr) ν_{max} : 3036, 2955, 1755, 1634, 1539, 1445, 1357, 1297, 1236, 1108, 818 cm^{-1} ; **^1H NMR** (CDCl_3) δ 2.35 (s, 3H), 3.45 (s, 3H), 3.94 (s, 3H), 5.27 (d, 1H, $J = 7.4$ Hz), 5.81 (overlapping doublets, 1H, $J_1 = 7.7$ Hz, $J_2 = 7.5$ Hz), 6.2 (d, 1H, $J = 7.8$ Hz), 7.26-7.15 (m, 5H), 7.58 (overlapping doublets, 1H, $J_1 = 7.77$ Hz, $J_2 = 8.82$ Hz), 7.76(d, 1H, $J = 7.08$ Hz), 8.22 (s, 1H); **^{13}C NMR** (CDCl_3) δ 21.4, 52.8, 53.2, 53.77, 84.00, 90.0, 93.0, 123.2, , 123.8, 126.4, 129.4, 129.9, 130.6, 135.3, 135.3, 139.5, 148.5, 166.8, 168.0, Anal. Calcd for $\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}_9$: C, 56.76; H, 4.54; N, 6.30. Found: C, 56.70; H, 4.32; N, 6.72.

Tetrahydrofuran 3d

Colorless crystalline solid. **mp** 148 °C; **IR** (KBr) $\nu_{\max}$: 3022, 2955, 2854, 1762, 1613, 1566, 1519, 1445, 1283, 1236, 1135, 1054, 939 cm^{-1} ; **^1H NMR**: δ 2.33 (s, 3H), 3.26 (s, 3H), 3.93 (s, 3H), 5.22 (d, 1H, J = 5.46 Hz), 5.69 (overlapping doublets, 1H, J = 6.9 Hz, J_2 = 5.58 Hz), 6.18 (d, 1H, J = 7.0 Hz), 7.34-7.14(m, 9H); **^{13}C NMR** (CDCl_3) δ 21.3, 52.4, 53.6, 54.2, 84.8, 90.9, 94.3, 126.2, 128.57, 128.6, 128.9, 129.2, 130.7, 133.8, 139.1, 167.1, 168.1. Anal. Calcd for $\text{C}_{21}\text{H}_{21}\text{NO}_7$: C, 63.15; H, 5.30; N, 3.51. Found: C, 63.31; H, 4.81; N, 3.71

Tetrahydrofuran 3e

Colorless crystalline solid **mp** 130-131 °C; **IR** (KBr) $\nu_{\max}$: 2982, 2955, 1748, 1566, 1445, 1371, 1283, 1108 cm^{-1} ; **^1H NMR** (CDCl_3) δ 3.27 (s, 3H), 3.95 (s, 3H), 5.23 (d, 1H, J = 5.3 Hz), 5.71 (overlapping doublets, 1H, J = 6.7 Hz, J_2 = 5.6 Hz), 6.21 (d, 1H, J = 7.0 Hz), 7.36-7.35 (m 10H); **^{13}C NMR** (CDCl_3) δ 52.4, 53.6, 54.3, 84.8, 91.0, 94.3, 126.3, 128.56, 128.59, 128.7, 128.9, 129.4, 133.7, 133.8, 167.1, 168.1; Anal. Calcd for $\text{C}_{20}\text{H}_{19}\text{NO}_7$: C, 62.33; H, 4.97; N, 3.63. Found: C, 62.26; H, 5.06; N, 3.83.

Tetrahydrofuran 3f

Colorless crystalline solid. **mp** 148-149 °C; **IR** (KBr) $\nu_{\max}$: 3027, 2960, 2929, 2856, 1750, 1620, 1558, 1517, 1439, 1351, 1289, 1243, 1113, 1077, 1020, 974, 932, 860, 818, 756, 741, 632 cm^{-1} ; **^1H NMR** (CDCl_3) δ 2.3 (s, 3H), 3.36 (s, 3H), 3.93 (s, 3H), 5.19 (d, 1H, J = 6.2 Hz), 5.68 (overlapping doublets, 1H, J = 7.1 Hz, J_2 = 6.4 Hz), 6.16 (d, 1H, J = 7.3 Hz), 7.36-7.03 (m, 8H);

^{13}C **NMR** (CDCl_3) δ 21.3, 52.6, 53.3, 53.6, 84.4, 94.0, 115.8, 116.1, 126.3, 129.3, 130.4, 130.5, 130.7, 167.1, 168.2. Anal.

Calcd for $\text{C}_{21}\text{H}_{20}\text{FNO}_7$: C, 60.43; H, 4.83; N, 3.36. Found: C, 60.54; H, 4.83; N, 3.42.

Tetrahydrofuran 3g

Colorless crystalline solid, **mp** 140 °C; **IR** (KBr) ν_{max} : 3068, 3032, 2996, 2970, 2950, 1744, 1615, 1558, 1522, 1444, 1387, 1300, 1289, 1237, 1170, 1118, 1067, 1031, 963, 943, 922, 855, 751, 705, 643, 596, 539 cm^{-1} ; ^1H **NMR** (CDCl_3) δ 3.35 (s, 3H), 3.94 (s, 3H), 5.2 (d, 1H, J = 5.94 Hz), 5.69 (overlapping multiplets, 1H, J_1 = 6.4 Hz, J_2 = 6.8 Hz), 6.2 (d, 1H, J = 7.2 Hz), 7.09-7.03 (m, 2H), 7.35-7.25 (m, 7H); ^{13}C **NMR** (CDCl_3) δ 52.5, 53.3, 53.6, 84.3, 90.5, 93.9, 115.7, 116.0, 126.2, 128.5, 129.2, 129.36, 130.2, 130.4, 133.5, 164.3, 166.9, 168.1. Anal. Calcd for $\text{C}_{20}\text{H}_{18}\text{FNO}_7$: C, 59.55; H, 4.50; N, 3.47. Found: C, 59.72; H, 4.34; N, 3.70.

Tetrahydrofuran 3h

Colorless crystalline solid **mp** 115 °C; **IR** (KBr) ν_{max} : 3022, 2955, 2939, 1744, 1615, 1563, 1532, 1460, 1444, 1372, 1268, 1232, 1222, 1186, 1118, 1082, 1041, 927, 860, 803, 736, 684 cm^{-1} ; ^1H **NMR** (CDCl_3) δ 2.34 (s, 3H) 3.35(s, 3H), 3.79 (s, 3H), 3.93 (s, 3H), 5.14 (d, 1H, J = 5.9 Hz) 5.65 (overlapping doublets, 1H, J_1 = 6.1Hz, J_2 = 6.9 Hz), 6.15 (d, 1H, J = 7.17 Hz), 6.86 (d, 2H, J

= 8.7 Hz) 7.26-7.14 (m, 6H); ^{13}C NMR (CDCl_3) δ 21.2, 52.4, 53.39, 53.42, 55.1, 84.3, 90.6, 94.2, 114.1, 125.3, 126.1, 129.1, 129.6, 130.7, 139.0, 159.6, 167.1, 168.2, Anal. Calcd for $\text{C}_{22}\text{H}_{23}\text{NO}_8$: C, 61.53; H, 5.40; N, 3.26. Found: C, 61.84; H, 5.33; N,

3.46

Tetrahydrofuran 3i.

Colorless crystalline solid. **mp** 175 °C; **IR** (KBr) ν_{max} : 3025, 2984, 2921, 1741, 1562, 1481, 1464, 1455, 1293, 1284, 1222, 1126, 1037, 942, 895, 750, 617cm^{-1} ; **^1H NMR** (CDCl_3) δ 2.32 (s, 3H), 3.43 (s, 3H), 3.91 (s, 3H), 5.10 (d, 1H, J = 6.93 Hz), 5.66 (overlapping doublets, 1H, J_1 = 7.2 Hz, J_2 = 7.35 Hz), 6.12 (d, 1H, J = 7.62 Hz), 7.43-7.11(m, 7H).); ^{13}C NMR (CDCl_3) δ 21.4, 52.7, 52.9, 53.7, 84.1, 90.1, 93.2, 126.4, 128.0, 129.3, 130.58, 130.6, 130.8, 133.2, 133.4, 139.3, 166.8, 168.0. Anal. Calcd for $\text{C}_{21}\text{H}_{19}\text{Cl}_2\text{NO}_7$: C, 53.86; H, 4.09; N, 2.99. Found: C, 54.29; H, 3.63; N, 2.76.

Tetrahydrofuran 3j.

Colorless crystalline solid. **mp** 155 °C; **IR** (KBr) ν_{max} : 3027, 2996, 2965, 2924, 1739, 1563, 1482, 1460, 1455, 1294, 1284, 1222, 1129, 1077, 1041, 937, 891, 803 751 , 705, 617cm^{-1} ; **^1H NMR** (CDCl_3) δ 3.45 (s, 3H), 3.94 (s, 3H) 5.13 (d, 1H, J = 6.8 Hz), 5.70 (overlapping doublets, 1H, J_1 = 6.9 Hz, J_2 = 7.5 Hz) 6.18 (d, 1H, J = 7.5 Hz) 7.46-7.19 (m, 8H); ^{13}C NMR (CDCl_3) δ

52.7, 52.9, 53.6, 84.1, 90.2, 93.3, 126.3, 127.8, 128.5, 129.5, 130.5, 130.7, 133.1, 133.2, 133.2, 133.4, 166.6, 167.9 Anal.

Calcd for $C_{20}H_{17}Cl_2NO_7$: C, 52.88; H, 3.77; N, 3.08. Found: C, 52.46; H, 3.68; N, 3.29.

Figure S1. ^1H NMR of tetrahydrofuran **3a**

Figure S2. ¹³CNMR of tetrahydrofuran **3a**

NAME chandru13c
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030408
Time 13.07
INSTRUM dpx300
PROBHD 5 mm Multinu
PULPROG zgdc
TD 32768
SOLVENT CDCl3
NS 176
DS 0
SMH 18939.395 Hz
FIDRES 0.577984 Hz
AQ 0.8651252 sec
RG 1149.4
DM 26.400 usec
DE 9.43 usec
TE 300.0 K
D11 0.0300000 sec
PL12 17.20 dB
CPDPRG2 waltz16
PCPD2 80.00 usec
SF02 300.1312700 MHz
NUC2 ¹H
PL2 120.00 dB
D1 5.00000000 sec
P1 12.50 usec
SF01 75.4750150 MHz
NUC1 ¹³C
PL1 -6.00 dB

F2 - Processing parameters
SI 16384
SF 75.4677486 MHz
WDW EM
SSB 0
LB 0.20 Hz
GB 0
PC 1.00

1D NMR plot parameters
CX 20.00 cm
F1P 201.454 ppm
F1 15203.26 Hz
F2P -4.754 ppm
F2 -358.74 Hz
PPMCH 10.31036 ppm/cm
HZCM 778.09991 Hz/cm

Figure S3. ¹H NMR of tetrahydrofuran 3b

Figure S4. ¹³C NMR of tetrahydrofuran 3b

Figure S5. ¹H NMR of tetrahydrofuran **3c**

Figure S6. ¹³CNMR of tetrahydrofuran 3c

Figure S7. ¹H NMR of tetrahydrofuran **3d**

Figure S8. ^{13}C NMR of tetrahydrofuran **3d**

Figure S9. ¹H NMR of tetrahydrofuran **3e**

Figure S10. ¹³C NMR of tetrahydrofuran **3e**

Figure S11. ^1H NMR of tetrahydrofuran **3f**

Figure S12. ^{13}C NMR of tetrahydrofuran 3f

Figure S13. ¹H NMR of tetrahydrofuran **3g**

Figure S14. ¹³CNMR of tetrahydrofuran 3g

Figure S15. ^1H NMR of tetrahydrofuran **3h**

Figure S16. ¹³C NMR of tetrahydrofuran 3h

Current Data Parameters
NAME saum1
EXPNO 2
PROCNO 1

F2 - Acquisition Parameters
Date_ 20030724
Time 17.00
INSTRUM dpx300
PROBHD 5 mm Multinu
PULPROG zg30
TD 32768
SOLVENT CDC13
NS 4
DS 0
SWH 4816.956 Hz
FIDRES 0.147002 Hz
AQ 3.4013684 sec
RG 322.5
DW 103.800 usec
DE 169.96 usec
TE 300.0 K
D1 1.00000000 sec
P1 11.00 usec
SF01 300.1320709 MHz
NUC1 1H
PL1 0.00 dB

F2 - Processing parameters
SI 16384
SF 300.1300149 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 5.00

1D NMR plot parameters
CX 20.00 cm
F1P 8.282 ppm
F1 2485.72 Hz
F2p -0.453 ppm
F2 -136.09 Hz
PPMCM 0.43678 ppm/cm
HZCM 131.09045 Hz/cm

Figure S17. ¹H NMR of tetrahydrofuran 3i

Figure S18. ¹³CNMR of tetrahydrofuran **3i**

Figure S19. ¹H NMR of tetrahydrofuran 3j

Figure S20. ^{13}C NMR of tetrahydrofuran 3j