Supporting information

The Total synthesis of Spirotenuipesines A and B

Mingji Dai,[†] Samuel J. Danishefsky*,^{†,‡}

[†] *Department of Chemistry, Columbia University, Havemeyer Hall, New York, NY 10027, USA and*

[‡] *Laboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research Center, 1275 York Avenue,*

Room RRL 1301, New York, NY 10021, USA

General Methods

All non-aqueous reactions were carried out in oven-dried glassware under a slight positive pressure of argon unless otherwise noted. All reagents were commercially available and used without further purification from Sigma-Aldrich and TCI America, unless indicated otherwise. Solvents were reagent grade and purified by standard techniques: THF was distilled from Na-benzophenone or filtered through a dry-solvent system; CH₂Cl₂ was distilled from CaH₂ or filtered through a dry-solvent system; all other solvents were Aldrich “anhydrous” grade solvents, unless indicated otherwise. Reactions were magnetically stirred and monitored by thin layer chromatography on Merck silica gel 60-F₂₅₄ coated 0.25 mm plates. Preparative thin layer chromatography was performed with Merck silica gel 60-F₂₅₄ coated 0.50 mm plates. Flash chromatography was performed with Sorbent Technology silica gel 60 (particle size 32-63 μm), unless indicated otherwise. Yields reported are for isolated, spectroscopically pure compounds. Melting points are uncorrected. CDCl₃ was allowed to stand over K₂CO₃ and 4 Å MS to neutralize and dry prior to NMR sample preparation. NMR spectra were recorded on Bruker DRX 300 or 400 MHz and DMX 500MHz spectrometers. Proton and Carbon chemical shifts were referenced to residual solvent peaks. Abbreviations for ¹H NMR: s = singlet, d = doublet, t = triplet, q = quartlet, m = multiplet, or br = broad. IR spectra were recorded on a Perkin-Elmer Paragon 1000 FTIR

spectrometer. High resolution mass spectra were acquired in the Columbia University Mass Spectral Core facility on a JEOL HX110 spectrometer. Optical rotations were measured on a JASCO DIP-1000 spectrometer.

Experimental Procedures and Spectra Data

To a solution of **8** (620 mg, 4.4 mmol) in 20 mL of DMF were added imidazole (362 mg, 5.3 mmol) and TBSCl (1.0 M in THF, 5.3 mmol) at 0°C. The reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction was quenched with 100 mL water, extracted with ether (3 x 50 mL). The combined organic phase was washed with water and brine respectively, dried over MgSO₄, concentrated and purified by flash chromatography (Hexane / EA: 50 / 1) to give desired product (1.07 g, 4.20 mmol, 95%).

¹H NMR (300MHz, CDCl₃): δ 4.78 (s br, 2H), 4.71 (s br, 2H), 3.94 (m, 1H), 2.16 (m, 4H), 1.74 (s, 6H), 0.88 (s, 9H), 0.04 (s, 6H); ¹³C NMR (100MHz, CDCl₃): δ 142.7, 112.9, 69.6, 45.9, 26.1, 23.2, 18.3, -4.2; MS (EI⁺) calcd for C₁₅H₃₀OSi (M): 254.21; found 199.26 (losing 2-methylpropenyl group (C₄H₇)).

S.M. (178 mg, 0.70 mmol) in 2 mL of CH₂Cl₂ was added over 3 min to a stirred suspension of selenium dioxide supported on silica gel (5% SeO₂ on silica gel, 78 mg) in CH₂Cl₂ and *t*-butylhydroperoxide (5-6 M in nonane, 0.32 mL). The reaction mixture was stirred at room temperature for 2 days. The mixture was filtered through a plug of silica gel and washed with CH₂Cl₂. The combined organic phase was

dried over Na₂SO₄, concentrated and purified by flash chromatography (Hexane / EA: 50 / 1 to 20 / 1) to give desired product **9** in 40-50% yield.

¹H NMR (400MHz, CDCl₃): δ 5.11 (s, 1H), 4.90 (s, 1H), 4.80 (s, 1H), 4.73 (s, 1H), 4.05 (d, *J* = 6.40 Hz, 2H), 3.98 (m, 1H), 2.81 (t, *J* = 6.40 Hz, 1H), 2.26 (m, 4H), 1.73 (s, 3H), 0.90 (s, 9H), 0.08 (s, 6H); ¹³C NMR (100MHz, CDCl₃): δ 145.6, 142.2, 114.0, 113.3, 70.4, 66.4, 45.2, 41.1, 25.8, 22.8, 18.0, -4.66, -4.71; HRMS (FAB⁺) calcd for C₁₅H₃₁O₂Si: 271.2093; found 271.2079.

To a refluxing solution of **9** (1.0 eq) in benzene (0.02 M) was added Grubbs 2nd generation catalyst (0.05 eq.) under argon during 3-4 hours. The reaction was refluxed for 1-2 hours more (detected by TLC until the disappearance of s.m.). After evaporation of the solvent, the crude product was purified by flash chromatography (Hexane / EA: 6 / 1) to give desired product **10** in 82% yield.

¹H NMR (400MHz, CDCl₃): δ 4.47 (m, 1H), 4.15 (s, 2H), 2.74 (dd, *J* = 16.6, 9.9 Hz, 1H), 2.58 (dd, *J* = 16.6, 6.8 Hz, 1H), 2.38 (d, *J* = 16.1 Hz, 1H), 2.28 (d, *J* = 16.8 Hz, 1H), 1.67 (s, 3H), 1.54 (br, 1H), 0.88 (s, 9H), 0.06 (s, 6H); ¹³C NMR (100MHz, CDCl₃): δ 133.1, 131.6, 70.9, 59.0, 48.7, 44.3, 26.1, 18.5, 14.0, -4.49, -4.50. HRMS (FAB⁺) calcd for C₁₃H₂₅O₂Si: 241.1624; found 241.1628.

The mixture of **10** (18 mg, 0.074 mmol) and propionic acid (0.331 mg, 0.0045 mmol) in triethylorthoacetate (0.15 mL) was heated at 180°C for 2 days before cooling down to room temperature. After evaporation of the solvent, the crude product was purified by flash chromatography (Hexane / EA: 20 / 1) to give a 1/1 inseparable mixture of **11** and **12** (20 mg, 87%).

^1H NMR (400MHz, CDCl_3): δ 4.88-4.91 (m, 1H), 4.78-4.80 (m, 1H), 4.24-4.30 (m, 1H), 4.07-4.14 (m, 2H), 2.62-2.69 (m, 1H), 2.60 (d, $J = 14.56$ Hz, 0.5H), 2.45 (d, $J = 14.48$ Hz, 0.5H), 2.36-2.42 (m, 1H), 2.34 (d, $J = 2.24$ Hz, 1H), 2.19 (dd, $J = 13.28, 5.64$ Hz, 0.5H), 2.03 (dd, $J = 13.08, 5.04$ Hz, 0.5H), 1.77 (dd, $J = 13.08, 4.88$ Hz, 0.5H), 1.64 (ddd, $J = 13.28, 4.72, 1.40$ Hz, 0.5H), 1.19-1.30 (m, 3H), 0.88 (s, 9H), 0.06 (s, 6H); ^{13}C NMR (100MHz, CDCl_3): δ 171.9, 171.6, 157.9, 105.5, 105.4, 71.3, 60.1, 59.9, 48.2, 47.9, 46.5, 46.3, 43.7, 43.5, 43.1, 42.7, 28.9, 27.8, 26.0, 18.2, 14.5, -4.6 HRMS (FAB+) calcd for $\text{C}_{17}\text{H}_{31}\text{O}_3\text{Si}$: 311.2042; found 311.2015.

To the solution of **10** (670 mg, 2.77 mmol) in 15 mL of CH_2Cl_2 was added DMAP (67 mg, 0.55 mmol), TEA (1.2 mL, 8.31 mmol) and Ac_2O (0.53 mL, 5.54 mmol) at room temperature. The reaction mixture was stirred for 2 hours before quenched with saturated aqueous NH_4Cl and diluted with Et_2O . The separated organic layer was washed with HCl (1 N in H_2O), saturated aqueous NaHCO_3 and brine, then dried over anhydrous MgSO_4 , filtrated. After evaporation of the solvent, the crude acetate was used directly in the next step.

^1H NMR (400MHz, CDCl_3): δ 4.63 (d, $J = 12.4$ Hz, 1H), 4.58 (d, $J = 12.4$ Hz, 1H), 4.48 (m, 1H), 2.69-2.57 (m, 2H), 2.34-2.29 (m, 2H), 2.05 (s, 3H), 1.70 (s, 3H), 0.89 (s, 9H), 0.06 (s, 6H).

To a solution of the above crude acetate in 15 mL THF was added TBAF (1.0 M in THF, 3.3 mL, 3.3 mmol). The reaction mixture was stirred overnight before quenched with saturated aqueous NH_4Cl . The mixture was extracted with ethyl acetate (x 3) and the combined organic phases were washed with brine, dried over anhydrous MgSO_4 and concentrated. The residue was purified by flash chromatography (Hexane / EA: 2 / 1, then 1 / 1) to give 395 mg desired product in 84% yield over two steps.

^1H NMR (400MHz, CDCl_3): δ 4.63 (d, $J = 12.4$ Hz, 1H), 4.59 (d, $J = 12.8$ Hz, 1H), 4.42 (m, 1H), 2.73-2.65 (m, 2H), 2.32 (d, $J = 17.2$ Hz, 1H), 2.28 (d, $J = 18.0$ Hz, 1H), 2.03 (s, 3H), 1.72 (s, 3H); ^{13}C NMR

(100MHz, CDCl₃): δ 171.1, 136.1, 127.4, 69.9, 60.6, 48.4, 44.6, 20.8, 13.8. The product is unstable under all MS conditions and no meaningful peaks were detected.

To a solution of s.m. (59 mg, 0.35 mmol) in 3 mL CH₂Cl₂ was added glyoxylic acid chloride tosylhydrazone (169 mg, 0.65 mmol) and N, N-dimethylaniline (77 mg, 0.64 mmol) at 0 °C. The dark green solution was stirred for 15 min prior to injection of triethylamine (88 mg, 0.87 mmol). The resulting orange suspension was stirred 10 min at 0 °C then for 15 min at room temperature before water was introduced and the mixture was concentrated in vacuo. Saturated aqueous citric acid and 1/1 ethyl acetate-hexanes were added and the layers were separated. The organic layer was washed with citric acid and the combined aqueous layers were extracted with 1/1 ethyl acetate-hexanes. The extract was washed with equal volume of citric acid solution and the combined organic layers were dried over anhydrous MgSO₄ and concentrated. The residue was purified by flash chromatography (Hexane / EA: 6 / 1) to give diazoester **13** (72 mg, 88%) as a yellowish oil.

¹H NMR (400MHz, CDCl₃): δ 5.38 (m, 1H), 4.73 (s, 1H), 4.63 (s, 2H), 2.84 (dd, J = 18.4, 7.2 Hz, 1H), 2.79 (dd, J = 17.6, 7.2 Hz, 1H), 2.45 (d, J = 18.0 Hz, 1H), 2.41 (d, J = 18.8 Hz, 1H), 2.07 (s, 3H), 1.73 (s, 3H); ¹³C NMR (100MHz, CDCl₃): δ 171.0, 166.7, 135.8, 127.5, 73.2, 60.3, 46.4, 45.3, 41.5, 20.8, 13.6; MS (FAB⁺) calcd for C₁₁H₁₄N₂O₄: 238.09; found 179.17, 154.11.(fragment peaks).

A solution of diazoester **13** (60 mg, 0.25 mmol) in 15 mL toluene was added slowly to a stirred refluxing solution of bis-(N-*t*-butylsalicylaldiminato) copper (II) catalyst (5.2 mg, 0.013 mmol) in 25

mL toluene over 1 hour at a constant rate. The solution was held at reflux for 30 min after completion of addition, cooled, concentrated and purified by flash chromatography (Hexane / EA: 3 / 1, then 1 / 1) to give 48 mg of desired product **14** in 91% yield.

^1H NMR (400MHz, CDCl_3): δ 4.61 (m, 1H), 4.28 (d, $J=12.4$ Hz, 1H), 4.16 (d, $J=12.4$ Hz, 1H), 2.05 (s, 3H), 2.02 (dd, $J=13.2, 2.8$ Hz, 1H), 1.93-1.85 (m, 3H), 1.77 (s, 1H), 1.35 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 170.7, 170.2, 73.5, 63.1, 37.7, 34.4, 33.6, 32.7, 30.8, 20.7, 14.7; HRMS (FAB+) calcd for $\text{C}_{11}\text{H}_{15}\text{O}_4^+$: 211.0970; found 211.0970.

To a solution of **14** (32 mg, 0.15 mmol) in 5 mL MeOH of was added K_2CO_3 (25 mg, 0.18 mmol). The mixture was stirred at room temperature for 5 h. The reaction mixture was filtered through a plug of silica gel and washed with ethyl acetate. The collected solution was concentrated to give the desired product in quantitative yield, which was used in the next step without any further purification.

^1H NMR (400MHz, CDCl_3): δ 4.63 (m, 1H), 3.89 (d, $J=12.0$ Hz, 1H), 3.72 (d, $J=12.0$ Hz, 1H), 3.46 (s, 1H), 2.11 (dd, $J=13.2, 3.2$ Hz, 1H), 1.95-1.86 (m, 3H), 1.73 (s, 1H), 1.38 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 171.2, 73.8, 61.6, 38.0, 36.8, 34.0, 32.5, 30.9, 14.6; HRMS (FAB+) calcd for $\text{C}_9\text{H}_{13}\text{O}_3$: 169.0865; found 169.0855.

To a solution of the above crude alcohol in THF was added KHMDS (0.5 M in tol., 1.2 equiv.) at room temperature. After 2 h, carbon disulfide (2.0 equiv.) was added. The mixture was stirred for 2 h and then iodomethane (2.0 equiv.) was added. After a further 2 h, the reaction was quenched with acetic acid and diluted with a saturated aqueous ammonium chloride solution, before extracting with EtOAc (x 3). The

combined organic layers were dried over anhydrous MgSO_4 and concentrated to give the crude *bis*-thiocarbonate **15** which was used directly in the next step.

^1H NMR (400MHz, CDCl_3) of the crude product: δ 4.89 (d, $J=12.0$ Hz, 1H), 4.67 (d, $J=12.0$ Hz, 1H), 4.66 (m, 1H), 2.58 (s, 3H), 2.13 (dd, $J=13.2, 3.2$ Hz, 1H), 2.02-1.92 (m, 3H), 1.89 (s, 1H), 1.41 (s, 3H).

To a solution of the above *bis*-thiocarbonate **15** in 3 ml toluene was added AIBN (0.5 equiv) and tributyltinhydride (2.0 equiv.). The reaction mixture was heated at 110°C for 4h, then cooled, concentrated and purified by flash chromatography (Hexane / EA: 1 / 0, 6 / 1, then 4 / 1) to give 14 mg desired product **16** in 60% yield over 3 steps.

^1H NMR (400MHz, CDCl_3): δ 5.03-5.01 (m, 2H), 4.89 (m, 1H), 2.87 (d, $J=18.0$ Hz, 1H), 2.67 (dd, $J=18.0, 4.0$ Hz, 1H), 2.59 (d, $J=18.0$ Hz, 1H), 2.49 (dd, $J=18.0, 2.4$ Hz, 1H), 1.98 (dd, $J=12.8, 3.2$ Hz, 1H), 1.77 (td, $J=12.4, 2.8$ Hz, 1H), 1.26 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 170.3, 153.1, 107.5, 78.4, 48.2, 42.7, 41.2, 29.7, 22.4; MS (APCI+) calcd for $\text{C}_9\text{H}_{13}\text{O}_2$: 153.08; found 153.13.

To a solution of **16** (8 mg, 0.053 mmol) in 2 mL MeOH was added 8 drops of KOH (2 N in water). The reaction mixture was heated at 60°C for 4 h. After the reaction was complete, MeOH was removed in vacuo. The left residue was neutralized with HCl (6 N in water) to pH 1-2, then treated with NaHCO_3 (solid) until no bubbles came out of the solution.

To the above solution (about 2 mL) was added 2 mL THF, followed by KI (17.4 mg, 0.11 mmol) and iodine (27 mg, 0.11 mmol). The reaction mixture was stirred overnight, then quenched with 10% $\text{Na}_2\text{S}_2\text{O}_3$, extracted with ethyl acetate (x 3). The combined organic layers were washed with brine, dried

over anhydrous MgSO_4 and concentrated to give the crude iodolactone which was used directly in the next step.

^1H NMR (400MHz, CDCl_3) of the crude iodolactone: δ 4.39 (m, 1H), 3.46 (d, $J=11.2$ Hz, 1H), 3.36 (d, $J=11.2$ Hz, 1H), 2.82 (d, $J=18.4$ Hz, 1H), 2.64 (d, $J=18.0$ Hz, 1H), 2.47 (td, $J=15.2, 2.4$ Hz, 1H), 2.15 (td, $J=14.4, 2.4$ Hz, 1H), 2.08 (dd, $J=15.2, 5.2$ Hz, 1H), 2.00 (dd, $J=14.4, 4.8$ Hz, 1H), 1.30(s, 3H).

To a solution of the above crude iodolactone and 2,6-lutidine (8 equiv.) in 5 ml CH_2Cl_2 was added TBSOTf (2.0 equiv.) slowly at 0°C . The reaction mixture was allowed to warm up to room temperature and stirred for 5 h. The reaction was quenched with saturated aqueous NH_4Cl and extracted with ether (x 3). The combined organic layers were washed with 1N HCl, saturated aqueous sodium bicarbonate, water and brine respectively, then dried over anhydrous MgSO_4 , concentrated and purified by flash chromatography (Hexane / EA: 8 / 1) to give **17** (17 mg, 77% over three steps).

^1H NMR (400MHz, CDCl_3): δ 4.29 (m, 1H), 3.42 (d, $J=10.8$ Hz, 1H), 3.37 (d, $J=10.8$ Hz, 1H), 2.86 (d, $J=18.0$ Hz, 1H), 2.58 (d, $J=18.0$ Hz, 1H), 2.37 (d, $J=15.2$ Hz, 1H), 2.05 (d, $J=14.0$ Hz, 1H), 1.99 (dd, $J=14.8, 4.8$ Hz, 1H), 1.88 (dd, $J=14.0, 4.8$ Hz, 1H), 1.29 (s, 3H), 0.86 (s, 9H), 0.05 (s, 3H), 0.04 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 174.4, 94.0, 70.9, 52.5, 49.0, 46.7, 45.8, 25.7, 23.7, 18.0, 10.2, -4.9, -5.1; HRMS (FAB+) calcd for $\text{C}_{15}\text{H}_{28}\text{O}_3\text{SiI}$: 411.0853; found 411.0861.

To a solution of iodolactone **17** (175 mg, 0.43 mmol) in 30 ml toluene was added AIBN (70 mg, 0.43 mmol) and tributyltinhydride (373 mg, 0.34 ml, 1.28 mmol). The reaction mixture was heated to 60°C and bubbled with dry air for 2 days (more AIBN and Bu_3SnH if necessary). The reaction was cooled to 0°C , treated with 30 mg NaBH_4 in 8 ml ethanol and stirred for 2 hours before quenching with saturated

aqueous NH_4Cl . The organic layer was separated. The aqueous layer was washed with ether (x 3). The combined organic layers were dried over anhydrous MgSO_4 , filtered, concentrated and purified by flash chromatography (Hexane / EA: 4 / 1, then 2 / 1) to give 101 mg desired product in 79% yield.

^1H NMR (400MHz, CDCl_3): δ 4.34 (m, 1H), 3.77 (d, $J=12.4$ Hz, 1H), 3.70 (d, $J=12.4$ Hz, 1H), 2.78 (d, $J=18.0$ Hz, 1H), 2.57 (d, $J=18.0$ Hz, 1H), 2.09 (dt, $J=14.8, 2.4$ Hz, 1H), 1.95 (dt, $J=13.6, 2.4$ Hz, 1H), 1.86 (dd, $J=15.6, 5.6$ Hz, 1H), 1.83 (dd, $J=14.0, 4.8$ Hz, 1H), 1.29 (s, 3H), 0.87 (s, 9H), 0.05 (s, 3H), 0.04 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 176.7, 97.2, 71.9, 64.3, 51.4, 46.2, 45.5, 43.4, 25.6, 22.9, 17.8, -5.0, -5.2; HRMS (FAB+) calcd for $\text{C}_{15}\text{H}_{29}\text{O}_4\text{Si}$: 301.1835; found 301.1833.

To a solution of s.m. (100 mg, 0.33mmol) and PMB trichloroacetimidate (188 mg, 0.66 mmol) in 10 ml CH_2Cl_2 was added 10-camphorsulfonic acid (8 mg, 0.033 mmol). The reaction mixture was stirred overnight before the solvent was removed. The residue was purified by flash chromatography (Hexane / EA: 20/ 1) to give 132 mg desired product **18** in 95% yield.

^1H NMR (400MHz, CDCl_3): δ 7.21 (d, $J=8.4$ Hz, 2H), 6.87 (d, $J=8.8$ Hz, 2H), 4.49 (d, $J=11.6$ Hz, 1H), 4.42 (d, $J=11.6$ Hz, 1H), 4.32 (m, $J=5.2$ Hz, 1H), 3.80 (s, 3H), 3.56 (d, $J=10.4$ Hz, 1H), 3.49 (d, $J=10.8$ Hz, 1H), 2.67 (d, $J=17.6$ Hz, 1H), 2.55 (d, $J=17.6$ Hz, 1H), 2.04 (dd, $J=15.2, 4.0$ Hz, 1H), 1.98 (dd, $J=15.2, 6.0$ Hz, 1H), 1.89 (dd, $J=13.6, 5.2$ Hz, 1H), 1.83 (dd, $J=13.2, 5.2$ Hz, 1H), 1.26 (s, 3H), 0.86 (s, 9H), 0.03 (s, 3H), 0.02 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 176.1, 159.2, 129.8, 129.2, 113.7, 95.6, 73.3, 71.8, 71.3, 55.2, 50.9, 46.6, 45.0, 43.5, 25.7, 22.5, 17.9, -4.9, -5.0; HRMS (FAB+) calcd for $\text{C}_{23}\text{H}_{35}\text{O}_5\text{Si}$: 419.2254; found 419.2254.

To a solution of LDA (0.58 mmol) in 2 mL anhydrous THF was added **18** (165 mg, 0.393 mmol) in 3 x 2 mL anhydrous THF at -78 °C. The reaction mixture was stirred at -78 °C for half an hour then warmed up to 0 °C. CH₂O in argon stream was carried over the reaction surface for 5 minutes (CH₂O was generated by cracking (CH₂O)_n with heat gun.) The reaction was diluted with THF, allowed to warm up to room temperature and stirred overnight before quenched with saturated aqueous NH₄Cl. After a filtration, the reaction mixture was diluted with ether, washed with brine, dried over anhydrous MgSO₄, filtered, concentrated and purified by flash chromatography (Hexane / EA: 40/ 1, then 4 / 1) to give α-methylenelactone **20** (46.7 mg, 28%) and β-hydroxylactone **19** (114 mg, 65%).

One diastereomer of **19**: ¹H NMR (400MHz, CDCl₃): δ 7.20 (d, *J* = 8.4 Hz, 2H), 6.87 (d, *J* = 8.4 Hz, 2H), 4.49 (d, *J* = 11.6 Hz, 1H), 4.39 (d, *J* = 11.6 Hz, 1H), 4.32 (m, *J* = 6.0 Hz, 1H), 3.93 (t, *J* = 8.4 Hz, 1H), 3.80 (s, 3H), 3.71 (td, *J* = 10.0, 6.0 Hz, 1H), 3.57 (d, *J* = 10.8 Hz, 1H), 3.49 (d, *J* = 10.8 Hz, 1H), 2.95 (dd, *J* = 8.4, 6.0 Hz, 1H), 2.84 (d, *J* = 6.8 Hz, 1H), 2.02-1.90 (m, 3H), 1.66 (dd, *J* = 12.4, 5.6 Hz, 1H), 1.23 (s, 3H), 0.86 (s, 9H), 0.04 (s, 3H), 0.02 (s, 3H); ¹³C NMR (100MHz, CDCl₃): δ 178.2, 159.3, 129.5, 129.3, 113.8, 95.1, 73.4, 71.8, 71.1, 59.8, 55.2, 53.1, 49.0, 44.7, 42.7, 25.7, 21.7, 18.0, -4.8, -4.9; HRMS (FAB+) calcd for C₂₄H₃₇O₆Si: 449.2359; found 449.2358. The other diastereomer of **19**: ¹H NMR (400MHz, CDCl₃): δ 7.22 (d, *J* = 8.4 Hz, 2H), 6.87 (d, *J* = 8.4 Hz, 2H), 4.49 (d, *J* = 11.6 Hz, 1H), 4.43 (d, *J* = 11.6 Hz, 1H), 4.32 (m, 1H), 3.84 (m, 2H), 3.80 (s, 3H), 3.55 (d, *J* = 10.0 Hz, 1H), 3.48 (d, *J* = 10.0 Hz, 1H), 3.09 (t, *J* = 5.6 Hz, 1H), 2.86 (d, *J* = 6.8 Hz, 1H), 2.06 (d, *J* = 4.0 Hz, 1H), 1.98 (d, *J* = 14.0 Hz, 1H), 1.80 (dd, *J* = 13.6, 4.8 Hz, 1H), 1.24 (d, *J* = 9.2 Hz, 1H), 1.15 (s, 3H), 0.86 (s, 9H), 0.04 (s, 3H), 0.03 (s, 3H); ¹³C NMR (100MHz, CDCl₃): δ 178.5, 159.3, 129.4, 129.3, 113.8, 95.5, 73.4, 72.0, 70.9, 60.1, 55.2, 54.0, 53.0, 48.5, 44.2, 25.7, 17.9, 17.7, -4.9, -5.1; HRMS (FAB+) calcd for C₂₄H₃₇O₆Si: 449.2359; found 449.2358.

To a solution of β -hydroxylactone **19** (110 mg, 0.24 mmol) and triethylamine (0.14 mL, 0.98 mmol) in 5 mL of dry dichloromethane was added dropwise a solution of MsCl (56 mg, 0.49 mmol) in 2 mL CH_2Cl_2 at 0 °C. The reaction mixture was warmed to r.t. and stirred for 4 h. The dichloromethane was evaporated and the residue was dissolved in ether. The ether solution was washed with water, dried over anhydrous MgSO_4 , filtered and concentrated. The residue was dissolved in 6 mL of dichloromethane, and DBU (112mg, 0.733 mmol) was added. The mixture was stirred for 3 h, diluted with ether and washed with 10% HCl, saturated NaHCO_3 and brine, dried over anhydrous MgSO_4 , filtered and concentrated at reduced pressure. Chromatography of the residue (hexane / EA = 8 / 1) afforded 111 mg α -methylenelactone **20** in 95% yield.

^1H NMR (400MHz, CDCl_3): δ 7.21 (d, J = 8.4 Hz, 2H), 6.86 (d, J = 8.4 Hz, 2H), 6.10 (s, 1H), 5.39 (s, 1H), 4.49 (d, J = 12.0 Hz, 1H), 4.45 (d, J = 12.0 Hz, 1H), 4.29 (m, 1H), 3.80 (s, 3H), 3.53 (s, 2H), 2.05 (s, 2H), 1.99 (d, J = 13.6 Hz, 1H), 1.90 (dd, J = 13.2, 4.0 Hz, 1H), 1.32 (s, 3H), 0.82 (s, 9H), 0.01 (s, 3H), -0.03 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 169.8, 159.2, 147.7, 129.9, 129.2, 118.7, 113.7, 93.9, 73.2, 72.2, 71.0, 55.2, 53.7, 49.7, 43.7, 25.6, 22.8, 17.9, -5.0, -5.2 ; HRMS (FAB+) calcd for $\text{C}_{24}\text{H}_{35}\text{O}_5\text{Si}$: 431.2254; found 431.2263.

A toluene solution of dienophile **20** (112 mg, 0.26 mmol) and Danishefsky's diene **21** (447 mg, 2.59 mmol) was heated in a sealed tube at 180 °C for 48 hours at the presence of 0.01 eq methylene blue as stablizer. After the reaction was cooled down to room temperature, the solvent was removed. The

residue was dissolved in CH₂Cl₂ and treated with amberlite I-120 (acidic, 2 g) for 1 hour. After filtration and concentration, the crude product was purified by preparative TLC (Hexane / EA: 3/ 1) to give 115 mg desired product in 90% yield as an 8 / 1 mixture of diastereomers with the desired product **22** as major.

¹H NMR (400MHz, CDCl₃): δ 7.21 (d, *J* = 8.8 Hz, 2H), 7.15 (d, *J* = 10.4 Hz, 1H), 6.86 (d, *J* = 8.4 Hz, 2H), 5.92 (d, *J* = 10.0 Hz, 1H), 4.53 (d, *J* = 11.2 Hz, 1H), 4.43 (d, *J* = 11.2 Hz, 1H), 4.29 (quint, *J* = 5.2 Hz, 1H), 3.79 (s, 3H), 3.64 (d, *J* = 11.2 Hz, 1H), 3.50 (d, *J* = 11.2 Hz, 1H), 2.96 (m, 1H), 2.48 (m, 1H), 2.35 (m, 1H), 2.25 (dd, *J* = 13.2, 4.8 Hz, 1H), 2.12 (m, 1H), 2.05 (dd, *J* = 14.0, 3.6 Hz, 1H), 1.90 (dd, *J* = 14.4, 5.6 Hz, 1H), 1.70 (dd, *J* = 13.6, 4.4 Hz, 1H), 1.13 (s, 3H), 0.86 (s, 9H), 0.41 (s, 3H), 0.33 (s, 3H); ¹³C NMR (100MHz, CDCl₃): δ 198.1, 176.1, 159.4, 148.9, 129.6 (2C), 129.0, 113.8, 94.2, 73.6, 71.3, 70.6, 55.2, 52.7, 51.7, 47.8, 44.5, 33.4, 28.2, 25.7, 21.1, 18.0, -4.9, -5.1; HRMS (FAB+) calcd for C₂₈H₃₉O₆Si: 499.2516; found 499.2512.

To a stirring suspension of Ph₃P⁺CH₃I⁻ (186 mg, 0.46 mmol) in 5 mL THF was added KHMDS (0.5 M in toluene, 0.74 mL, 0.37 mmol) at -78 °C. The solution was warmed up to 0 °C and stirred for 30 min before cooled down to -78 °C. Enone **22** in 5 mL THF was added via cannula and the solution was warmed up to r.t. overnight (10 h) before quenched with saturated aqueous NH₄Cl solution, extracted with Et₂O (x 3). The organic layers were combined, washed with water, brine and dried over anhydrous MgSO₄, filtered, and concentrated under reduced pressure. The residue was purified by preparative TLC with Hexane / EtOAc (4 / 1) as the eluant to afford 88 mg colorless oil in 96% yield.

¹H NMR (400MHz, CDCl₃): δ 7.24 (d, *J* = 8.8 Hz, 2H), 6.87 (d, *J* = 8.4 Hz, 2H), 6.20 (d, *J* = 10.0 Hz, 1H), 5.90 (d, *J* = 10.0 Hz, 1H), 4.88 (s, 1H), 4.87 (s, 1H), 4.53 (d, *J* = 11.2 Hz, 1H), 4.46 (d, *J* = 11.6 Hz,

1H), 4.26 (quint, $J = 6.0$ Hz, 1H), 3.80 (s, 3H), 3.58 (d, $J = 10.8$ Hz, 1H), 3.50 (d, $J = 11.2$ Hz, 1H), 2.82 (m, 1H), 2.31 (m, 1H), 2.18-2.02 (m, 4H), 1.82 (m, 1H), 1.68 (dd, $J = 11.6, 6.4$ Hz, 1H), 1.06 (s, 3H), 0.86 (s, 9H), 0.04 (s, 3H), 0.03 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 178.2, 159.3, 140.9, 131.1, 129.5, 129.4, 128.8, 113.7, 112.6, 93.2, 73.4, 71.3, 71.2, 55.2, 52.4, 51.2, 47.2, 44.8, 28.1, 26.0, 25.7, 20.5, 18.0, -4.8, -5.0; HRMS (FAB+) calcd for $\text{C}_{29}\text{H}_{41}\text{O}_5\text{Si}$: 497.2723; found 497.2745.

To a solution of s.m. (87mg, 0.175mmol) in 10 ml THF was added TBAF (1.0 M in THF, 0.35 mL, 0.35 mmol). The reaction mixture was stirred for 1h before it was quenched with saturated aqueous NH_4Cl . The mixture was extracted with ethyl acetate (x 3) and the combined organic phases were washed with brine, dried over MgSO_4 and concentrated. The residue was purified by preparative TLC (Hexane / EA: 1/ 2) to give 67 mg desired product **23** in 100% yield.

^1H NMR (400MHz, CDCl_3): δ 7.24 (d, $J = 8.4$ Hz, 2H), 6.88 (d, $J = 8.8$ Hz, 2H), 6.22 (d, $J = 10.4$ Hz, 1H), 5.89 (d, $J = 10.0$ Hz, 1H), 4.90 (s, 1H), 4.89 (s, 1H), 4.53 (d, $J = 11.6$ Hz, 1H), 4.47 (d, $J = 11.6$ Hz, 1H), 4.36 (quint, $J = 5.6$ Hz, 1H), 3.81 (s, 3H), 3.61 (d, $J = 10.8$ Hz, 1H), 3.54 (d, $J = 10.8$ Hz, 1H), 2.85 (m, 1H), 2.34 (m, 1H), 2.29-2.11 (m, 3H), 2.05 (ddd, $J = 14.8, 5.2, 0.8$ Hz, 1H), 1.80 (m, 1H), 1.74 (ddd, $J = 13.2, 4.8, 0.8$ Hz, 1H), 1.08 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 178.7, 159.3, 140.8, 131.3, 129.4 (2C), 128.4, 113.7, 112.8, 93.8, 73.4, 71.1, 70.7, 70.6, 52.9, 51.3, 46.6, 44.5, 28.1, 26.0, 20.5, 20.4; HRMS (FAB+) calcd for $\text{C}_{23}\text{H}_{28}\text{O}_5$: 384.1937; found 384.1942.

To the solution of **23** (8.5 mg, 0.022 mmol) in 2 mL anhydrous CH₂Cl₂ was added DIBAL-H (0.05 mL, 0.049 mmol, 1.0 M in toluene) at -78 °C during 2 minutes. The reaction mixture was stirred at -78 °C for 20 minutes more before quenched with 0.05 mL saturated aqueous NH₄Cl and warmed up to room temperature. The resulting mixture was diluted with 8 mL Et₂O, stirred for 1.5 hours and treated with anhydrous MgSO₄ then filtrated through a plug of silica gel. After evaporation of the solvent, the crude hemilactal (d.r. 2/1) was used directly in the next step.

To a solution of the above hemilactal in 5 mL CH₂Cl₂ under 4 Å molecular sieves was added CSA (2.4 mg, 0.01 mmol). The reaction mixture was stirred at r.t. for 75 min before filtration. After evaporation of the solvent, the crude residue was purified by preparative TLC (Hexane / EA: 2 / 1) to afford 7.2 mg the desired product **24** in 90% yield.

¹H NMR (400MHz, CDCl₃): δ 7.28 (d, *J* = 9.2 Hz, 2H), 6.88 (d, *J* = 8.8 Hz, 2H), 6.16 (d, *J* = 10.4 Hz, 1H), 5.73 (d, *J* = 10.4 Hz, 1H), 5.05 (s, 1H), 4.81 (s, 1H), 4.80 (s, 1H), 4.55 (s, 2H), 4.46 (s, 1H), 3.81 (s, 3H), 3.65 (d, *J* = 10.8 Hz, 1H), 3.62 (d, *J* = 10.4 Hz, 1H), 2.41 (m, 3H), 2.21 (dd, *J* = 11.2, 2.8 Hz, 1H), 2.15 (d, *J* = 13.2 Hz, 1H), 1.53 (m, 2H), 1.19 (d, *J* = 12.8 Hz, 1H), 1.05 (s, 3H); ¹³C NMR (100MHz, CDCl₃): δ 142.3, 133.8, 130.1, 129.3, 113.7, 110.9, 102.9, 89.3, 76.3, 73.3, 71.3, 55.3, 52.9, 49.5, 43.5, 41.0, 29.7, 28.4, 25.7, 16.1, 14.1; HRMS (FAB⁺) calcd for C₂₃H₂₈O₄: 368.1988; found 368.2010.

To a solution of **24** (5 mg, 0.0136 mmol) in 1 mL dry pyridine was added a solution of OsO₄ in toluene (0.05 M, 0.3 mL, 0.015 mmol). The dark reaction mixture was stirred overnight. 2 mL 1M aq NaHSO₃ was added. After a further 90 min the mixture was diluted with 2 mL water and extracted with CHCl₃ (3 x 4 mL) and then with AcOEt (3 x 4 mL). Drying over MgSO₄ and evaporation of the solvent from the organic phase gave an oily residue **25** (d.r. 2.5/1 with the undesired diastereomer as major) which was used directly to the next step.

To a solution of the above crude diol **25** in 1.6 mL THF / H₂O (1 / 1) was added NaIO₄ (3.5 mg, 0.0162 mmol) at 0 °C. The solution was warmed to r.t. and stirred for 2 h, and then diluted with ethyl acetate before it was quenched with aqueous 20% Na₂S₂O₃ solution. The organic layer was separated, washed with aqueous 20% Na₂S₂O₃ solution, water and brine, dried over anhydrous MgSO₄, filtered and concentrated. The residue was purified by preparative TLC (Hexane/EA = 1/1) to afford the desired enone **26** 4.5 mg in 90% yield over two steps.

¹H NMR (400MHz, CDCl₃): δ 7.26 (d, *J* = 8.8 Hz, 2H), 7.05 (dd, *J* = 10.4, 1.6 Hz, 1H), 6.88 (d, *J* = 8.4 Hz, 2H), 5.98 (d, *J* = 10.8 Hz, 1H), 5.19 (s, 1H), 4.58 (d, *J* = 11.6 Hz, 1H), 4.51 (d, *J* = 11.6 Hz, 1H), 4.50(s, 1H), 3.81 (s, 3H), 3.71 (d, *J* = 11.2 Hz, 1H), 3.60 (d, *J* = 10.8 Hz, 1H), 2.49 (d, *J* = 9.6 Hz, 1H), 2.48 (d, *J* = 8.8 Hz, 1H), 2.37 (d, *J* = 13.2 Hz, 2H), 2.19 (dd, *J* = 13.2, 9.2 Hz, 1H), 1.92 (dt, *J* = 13.2, 9.2 Hz, 1H), 1.53 (d, *J* = 12.0 Hz, 1H), 1.25 (d, *J* = 13.2 Hz, 1H), 1.14 (s, 3H); ¹³C NMR (100MHz, CDCl₃): δ 198.8, 159.3, 154.2, 129.8, 129.5, 129.3, 113.8, 101.5, 89.6, 76.4, 73.5, 70.5, 55.2, 53.5, 49.8, 43.3, 40.9, 35.2, 25.2, 16.4; HRMS (FAB+) calcd for C₂₂H₂₇O₅: 371.1858; found 371.1870.

Table 1.

Nu ⁻ M ⁺	s.m.	dp	udp	conv. (dp)
MeLi	1	5.5	1	87% (73%)
MeMgBr	6.4	16	1	73% (68%)
MeLi/CeCl ₃	0.35	6	1	95% (82%)
MeMgBr/CeCl ₃	3	10	1	79% (71%)

A suspension of anhydrous CeCl₃ (107 mg, 0.43 mmol) in THF (2 mL) was stirred at 0 °C for 2 h, cooled to -78 °C, and treated with CH₃Li (0.27 mL, 0.43 mmol, 1.6 M in Et₂O). After 30 min at -78 °C, a solution of enone **26** (16 mg, 0.043 mmol) in THF (2 x 1 mL) was added and the reaction mixture was allowed to warm to 0 °C over 30 min. The mixture was stirred at 0 °C for 3 h and then quenched at 0 °C by slow addition of excess saturated NaHCO₃. Dilution with H₂O, extraction with EtOAc, brine wash, and workup afforded crude material. Flash chromatography (EtOAc / Hexane = 1 / 2) afforded 14 mg pure desired alcohol **27** in 84% yield with 13% undesired **C9-epi-27** which was contaminated with some unknown impurities. The ratio of the desired **27** and **C9-epi-27** was determined by ¹H NMR of the crude mixture before column purification. More condition evaluations were tabulated (table 1).

¹H NMR (400MHz, CDCl₃): δ 7.26 (d, *J* = 8.4 Hz, 2H), 6.88 (d, *J* = 8.8 Hz, 2H), 5.68 (d, *J* = 10.4 Hz, 1H), 5.56 (d, *J* = 10.8 Hz, 1H), 4.99 (s, 1H), 4.55 (d, *J* = 12.0 Hz, 1H), 4.52 (d, *J* = 12.4 Hz, 1H), 4.44(s, 1H), 3.81 (s, 3H), 3.62 (d, *J* = 12.8 Hz, 1H), 3.59 (d, *J* = 13.2 Hz, 1H), 2.33 (d, *J* = 12.8 Hz, 1H), 2.20 (dd, *J* = 11.6, 3.2 Hz, 1H), 2.10 (d, *J* = 13.6 Hz, 1H), 1.88 (d, *J* = 12.8 Hz, 1H), 1.68 (td, *J* = 14.0, 3.2 Hz, 1H), 1.54 (d, *J* = 10.0 Hz, 1H), 1.50 (td, *J* = 14.0, 3.2 Hz, 1H), 1.27 (s, 3H), 1.17 (d, *J* = 12.8 Hz, 1H), 1.06 (s, 3H); ¹³C NMR (100MHz, CDCl₃): δ 159.1, 136.5, 130.9, 130.2, 129.2, 113.7, 102.7, 89.3, 76.1, 73.3, 71.1, 69.3, 55.2, 52.9, 49.4, 43.4, 40.7, 37.1, 27.4, 24.3, 15.9; HRMS (FAB⁺) calcd for C₂₃H₂₉O₅: 385.2015; found 385.2009.

To a solution of PMB ether **27** (8 mg, 0.0207 mmol) in 2 mL CH_2Cl_2 / buffer solution pH = 7.00 (18 / 1) was added DDQ (9.4 mg, 0.0414 mmol). The reaction was stirred at room temperature for 5 h, then quenched with 3 mL saturated aqueous NaHCO_3 , extracted with AcOEt (5 x 3 mL), dried over anhydrous MgSO_4 , filtered, concentrated under reduced pressure and purified on a silica gel column using Hexane/ EtOAc (1 / 2 to 1 / 4) as the eluant to afford spirotenuipesine A (**1a**) 4.4 mg in 80% yield.

^1H NMR (400MHz, CDCl_3): δ 5.72 (dd, J = 10.4, 1.6 Hz, 1H), 5.48 (dd, J = 10.4, 1.6 Hz, 1H), 5.02 (s, 1H), 4.47 (quint, J = 1.6 Hz, 1H), 3.85 (dd, J = 12.0, 5.6 Hz, 1H), 3.74 (dd, J = 12.0, 7.2 Hz, 1H), 2.38 (d, J = 12.8 Hz, 1H), 2.24 (dd, J = 11.6, 3.6 Hz, 1H), 2.12 (dddd, J = 13.6, 4.8, 3.2, 1.6 Hz, 1H), 1.90 (dddd, J = 12.8, 5.2, 3.6, 1.6 Hz, 1H), 1.83 (dd, J = 7.2, 5.6 Hz, 1H), 1.70 (td, J = 13.2, 3.2 Hz, 1H), 1.52 (td, J = 14.0, 3.2 Hz, 1H), 1.50 (dq, J = 12.8, 0.8 Hz, 1H), 1.28 (s, 3H), 1.21 (br d, J = 12.8 Hz, 1H), 1.03 (s, 3H); ^{13}C NMR (100MHz, CDCl_3): δ 136.9, 130.5, 102.9, 90.0, 76.1, 69.2, 64.8, 52.9, 49.1, 43.1, 40.8, 37.0, 27.5, 24.4, 15.7; HRMS (FAB+) calcd for $\text{C}_{15}\text{H}_{23}\text{O}_4$: 267.1596; found 267.1592.

To a solution of **1a** (2.0 mg, 0.0075 mmol) in dichloromethane-methanol-phosphate buffer (pH 9.2) (1:4:2) (1.5 mL) was added acetone (0.03 mL) and 0.25 M Oxone solution in water (0.5 mL) at 0 °C. After being stirred for 6 h, this mixture was poured into water (5 mL) and extracted with 1-butanol (10

mL) three times. The organic layer was evaporated. The residue was chromatographed over silica gel eluted by chloroform-methanol (9:1) to afford **1b** (1.9 mg, 90% yield).

^1H NMR (400MHz, CDCl_3): δ 5.23 (s, 1H), 4.44 (m, 1H), 3.86 (d, $J = 12.8$ Hz, 1H), 3.78 (d, $J = 12.8$ Hz, 1H), 3.44 (dd, $J = 4.0, 1.6$ Hz, 1H), 3.04 (dd, $J = 4.0, 1.2$ Hz, 1H), 2.29 (d, $J = 12.8$ Hz, 1H), 2.20 (dd, $J = 11.6, 3.6$ Hz, 1H), 2.01 (br, 1H), 1.86 (dt, $J = 14.4, 4.0$ Hz, 1H), 1.60 (td, $J = 13.6, 3.2$ Hz, 1H), 1.46-1.52 (m, 2H), 1.30 (s, 3H), 1.26 (m, 1H), 1.22 (s, 3H), 1.17 (br d, $J = 12.4$ Hz, 1H); ^{13}C NMR (100MHz, CDCl_3): δ 101.3, 89.6, 75.7, 69.8, 64.2, 60.7, 59.5, 49.5, 48.4, 43.2, 41.2, 32.0, 23.8, 22.3, 15.2; HRMS (FAB $^+$) calcd for $\text{C}_{15}\text{H}_{23}\text{O}_5$: 283.1545; found 283.1533.