This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

The first study about the relationship between the extractability of thiacalix[4]arene derivatives and the position of the coordination binding sites

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1039/C4OB02393E>

PUBLISHER

© The Royal Society of Chemistry

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Zhao, Jiang-Lin, Hirotsugu Tomiyasu, Xin-Long Ni, Xi Zeng, Mark R.J. Elsegood, Carl Redshaw, Shofiur Rahman, P.E. Georgiou, Simon J. Teat, and Takehiko Yamato. 2019. "The First Study About the Relationship Between the Extractability of Thiacalix[4]arene Derivatives and the Position of the Coordination Binding Sites". figshare. <https://hdl.handle.net/2134/18137>.

1

The first study about the relationship between the extractability of thiacalix[4]arene derivatives and the position of the coordination binding sites

Jiang-Lin Zhao, Hirotugu Tomiyasu, Xin-Long Ni, Xi Zeng, Mark R. J. Elsegood, Carl Redshaw, Shofiur Rahman, Paris E. Georghiou and Takehiko Yamato*

The extractability of thiacalix[4]arene derivatives **2–4** are largely dependent on the position of the binding sites.

Please check this proof carefully. **Our staff will not read it in detail after you have returned it.**

Translation errors between word-processor files and typesetting systems can occur so the whole proof needs to be read. Please pay particular attention to: tabulated material; equations; numerical data; figures and graphics; and references. If you have not already indicated the corresponding author(s) please mark their name(s) with an asterisk. Please e-mail a list of corrections or the PDF with electronic notes attached – do not change the text within the PDF file or send a revised manuscript. Corrections at this stage should be minor and not involve extensive changes. All corrections must be sent at the same time.

Please bear in mind that minor layout improvements, e.g. in line breaking, table widths and graphic placement, are routinely applied to the final version.

We will publish articles on the web as soon as possible after receiving your corrections; **no late corrections will be made.**

Please return your **final** corrections, where possible within **48 hours** of receipt, by e-mail to: obc@rsc.org

Queries for the attention of the authors

Journal: **Organic & Biomolecular Chemistry**

Paper: **c4ob02393e**

Title: **The first study about the relationship between the extractability of thiacalix[4]arene derivatives and the position of the coordination binding sites**

Editor's queries are marked like this [Q1, Q2, ...], and for your convenience line numbers are indicated like this [5, 10, 15, ...].

Please ensure that all queries are answered when returning your proof corrections so that publication of your article is not delayed.

Query Reference	Query	Remarks
Q1	For your information: You can cite this article before you receive notification of the page numbers by using the following format: (authors), Org. Biomol. Chem., (year), DOI: 10.1039/c4ob02393e.	
Q2	Please check that the inserted CCDC number is correct.	
Q3	Please carefully check the spelling of all author names. This is important for the correct indexing and future citation of your article. No late corrections can be made.	
Q4	In Scheme 1, is it correct that the solvent is shown as "DMF&THF"? Or should it perhaps be shown as "DMF-THF" as it appears in the text? If the latter, please resupply the scheme, either as ChemDraw file or as TIF at 600 dpi.	
Q5	Please check that the changes to the English in the sentence beginning "This maybe attribute that when 3 complexes with Ag+, ..." have not affected the meaning.	

PAPER

The first study about the relationship between the extractability of thiacalix[4]arene derivatives and the position of the coordination binding sites†

Cite this: DOI: 10.1039/c4ob02393e

Jiang-Lin Zhao,^a Hirotugu Tomiyasu,^a Xin-Long Ni,^b Xi Zeng,^b Mark R. J. Elsegood,^c Carl Redshaw,^d Shofur Rahman,^e Paris E. Georghiou^e and Takehiko Yamato^{*a}

Three organic ionophores (**2–4**) based on the *p*-tert-butylthiacalix[4]arene backbone, blocked in the 1,3-*alternate* conformation, bearing two pyridyl coordinating moieties (*ortho* for **2**, *meta* for **3** and *para* for **4**), have been synthesized and characterized in the solid state. The solvent extraction experiments with the metal ions showed that the ability of these derivatives to complex with Ag⁺ appeared to be largely dependent on the position of the nitrogen atoms of the pyridyl ring. Two different complexation modes have been confirmed by ¹H NMR titration, ionophore **2** armed with two pyridyl moieties, complexed with Ag⁺ cation through N...Ag⁺...S interactions; however, ionophore **3** and ionophore **4**, complexed with Ag⁺ through metal–nitrogen (N...Ag⁺) interactions. The DFT computational studies were consistent with the experimental findings. These findings will provide us an important rule to design an appropriate thiacalix-arene ionophore in the future. Another study on the possibility for application of ionophores **2–4** for the treatment of waste water containing Cr(vi) and Cr(III), showed that ionophore **3** was useful in the application of the solvent extraction method in selective treatment of waste water containing Cr(vi) and Cr(III) prior to discharge.

Received 13th November 2014,
Accepted 21st January 2015

DOI: 10.1039/c4ob02393e

www.rsc.org/obc

Introduction

Thiacalix[4]arene is widely used as a macrocyclic platform for designing and building synthetic receptors toward metal cations.¹ The complexation properties of these molecules appear to be highly dependent upon the nature and number of donor atoms and also the conformation of the calix[*n*]arene moiety.² It is found that thiacalix[4]arene has a very high ability to bind transition metal ions,³ which has been quite unexpected considering the poor binding ability of calix[4]-arene. The 1,3-*alternate* stereoisomer, which shows an allosteric effect in metal cation binding, or offers divergently

oriented binding sites, is of special interest.^{1,4} For the synthesis of macrocycles with controlled (switchable) binding sites of metal cations,⁵ there is a need for the development of novel approaches to the design of tetrasubstituted thiacalix[4]-arenes with various groups with specific conformations. Recently, we reported the regioselective synthesis of *distal*-bis-[(2-pyridylmethyl)oxy]tetrathiacalix[4]arene in the 1,3-*alternate* conformation by a protection–deprotection method using benzyl groups as protecting groups.⁶ Pyridine derivatives of thiacalix[4]arene can exist as positional isomers which differ by the positions of the nitrogen (N) atom on the pyridyl unit which can be *ortho*, *meta* and *para* to the phenolic oxygen attachment position. The N-hetero atoms can serve as additional coordination sites due to their electron lone pairs and can also undergo facile further modification. Given that the position of the nitrogen atoms of the pyridyl ring can differ in thiacalix[4]arene derivatives, it is interesting to assess what kind of ability these derivatives will provide to interact with metal cations (hard or soft).

Chromium(III) has been reported to be biologically essential to mammals as it maintains effective glucose, lipid, and protein metabolisms. However, chromium(VI) can be toxic, as it can diffuse as Cr₂O₇^{2−} or HCr₂O₇[−] through cell membranes and oxidize biological molecules.⁷ Therefore, selective treatment of waste water containing Cr(vi) and Cr(III) prior to dis-

^aDepartment of Applied Chemistry, Faculty of Science and Engineering, Saga University, Honjo-machi 1, Saga 840-8502, Japan. E-mail: yamatot@cc.saga-u.ac.jp

^bDepartment Key Laboratory of Macrocyclic and Supramolecular Chemistry of Guizhou Province, Guizhou University, Guiyang, Guizhou 550025, China

^cChemistry Department, Loughborough University, Loughborough, LE11 3TU, UK

^dDepartment of Chemistry, The University of Hull, Cottingham Road, Hull, Yorkshire HU6 7RX, UK

^eDepartment of Chemistry, Memorial University of Newfoundland, St. John's, Newfoundland and Labrador, Canada A1B3X7

†Electronic supplementary information (ESI) available: Details of single-crystal X-ray crystallographic data. ¹H, ¹³C NMR, MS and IR spectra of **3**, computational study of **2–4** with Ag⁺. CCDC 1021161. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4ob02393e

charge is essential. Solvent extraction is one of the most commonly used treatment methods and employs a selective complexant especially for ions in aqueous solution. Thus, the development of efficient extractants for anions has received considerable attention in recent years.⁸ The dichromate ($\text{Cr}_2\text{O}_4^{2-}$ and HCr_2O_7^-) ions are anions with oxide functionalities at their periphery. These oxide moieties are potential sites for hydrogen bonding to the complexant or host molecule(s). Thiacalix[4]arene derivatives with nitrogen functionalities such as pyridine, amino, or imino groups on their lower rim have been shown to be capable of interacting with anions by hydrogen bonds as efficient extractants for oxoanions.⁹ Thus, the introduction of a pyridyl moiety to thiacalix[4]arene would potentially lead to an effective extractant for dichromate anions.

In this study, a series of 1,3-*alternate* thiacalix[4]arenes bearing pyridyl moieties (*ortho*, *meta* and *para*) at the lower rim which should have the appropriate encapsulating ionophilic cavity were targeted for synthesis. The relationship between the position of the nitrogen atoms of the pyridyl ring and the ability of these derivatives to interact with various ionic species were evaluated.

Results and discussion

The synthesis of the new thiacalix[4]arene derivatives is given in Scheme 1. For the synthesis of thiacalix[4]arene derivatives based on different functional units (1,3-*alternate*-2, 1,3-*alternate*-3 and 1,3-*alternate*-4), the parent compound (*distal*-1) was prepared according to published literature procedures.⁶ The reaction of bisbenzylated compound *distal*-1 with 3-(chloromethyl)pyridine in THF-DMF in the presence of Cs_2CO_3 as base yielded 1,3-*alternate*-3 in 59% yield. 1,3-*alternate*-2 and 1,3-*alternate*-4 were prepared as following a published procedure.^{6,10} All of the structures were confirmed by their ^1H and ^{13}C NMR and IR spectra, MS, elemental analyses and by X-ray crystallography.

Scheme 1 O-Alkylation of *distal*-1 with (chloromethyl)pyridine in the presence of Cs_2CO_3 .

The ^1H NMR spectrum of 1,3-*alternate*-3 shows two singlets for *tert*-butyl protons, in which both *tert*-butyl protons were observed at higher field, at δ 0.85 and 0.86 ppm due to the ring current effect arising from the two benzyl benzene rings and the two pyridine rings introduced; two singlets for the methylene protons at δ 5.06 ppm ($\text{OCH}_2\text{Benzyl}$) and 5.19 ppm ($\text{OCH}_2\text{Pyridyl}$), respectively, indicating a C_2 -symmetric structure for the 1,3-*alternate*-3 (Fig. S1†).

X-ray quality colourless crystals of 1,3-*alternate*-2, and 1,3-*alternate*-3 were obtained by recrystallizations from mixed MeOH and CHCl_3 solutions. The single crystal X-ray diffraction Ortep (Pluto) representations of 2 and 3 are shown in Fig. 1. It is clear that these compounds adopt 1,3-*alternate* conformations. Interestingly, both of the pyridine nitrogen atoms in 2 are orientated outwards, the distance between them being 9.079 Å. However, the pyridine nitrogen atoms in 3 are orientated inwards, the distance between them being only 3.883 Å.

Fig. 1 X-ray structures of (a) 2⁶ and (b) 3. Hydrogen atoms have been omitted for clarity.

This may be attributed to the distances between the pyridine nitrogen atoms and the oxygen atoms (N1...O1 and N2...O2). In the case of compound **2**, the distances between N1...O1 and N2...O2 are shorter; but for **3** the corresponding N1...O1 and N2...O2 distances are longer.

The shorter distances and hence the stronger electron repulsion could therefore be factors which control the different orientations of the nitrogen atoms toward each other.

Recently, the synthesis of calix[4]arenes bearing pendant pyridine groups at the lower rim as potential ligands for transition metal cations have been reported.¹¹ A similar investigation has also been conducted using hexahomotrioxacalix[3]-arene and homocalix[3]arene-based derivatives.¹² It is well known that the metal selectivity and extractability of these types of receptors are dependent on the ring size and the nature of the *O*-alkyl substituents. However, it is still unknown whether the metal extractability can be affected by the position of the coordination binding sites of the substituents themselves. Therefore, it is of importance to assess the relationship between the position of the nitrogen atoms of the pyridyl ring and the ability of these derivatives (**2–4**) to interact with ions. Experiments for solvent extraction of aqueous phase metal ions to the organic phase were therefore undertaken with **2–4**. The results showed that the extraction of transition metals by all three receptors **2–4** was higher than for the extraction of alkali metals, especially for Cr³⁺ and Ag⁺ (Fig. 2). The *E*% values of Cr³⁺ i.e. 43%, 61% and 52% for **2–4**, respectively, showed that a higher Cr³⁺ affinity exists for these molecules. However, what is surprising is that the extractability for Ag⁺, the *E*% values of 95%, 52% and 36% for **2–4**, respectively, showed that the extractability of Ag⁺ by **2** to **4** decreased gradually. These compounds are positional isomers differing only by the position of the nitrogen atom on the pyridyl ring. The position of the N atoms on the pyridyl rings (*ortho* for **2**, *meta* for **3** and *para* for **4**), which determines the distances between the nitrogen and the diaryl thiaether linkages were also reduced gradually. Recently, Ferlay reported a 1,3-*alternate* conformation thiocalix[4]arene armed with four pyridyl (*ortho*), complexed with Ag⁺ cation through N...Ag⁺...S interactions.¹³

Fig. 2 Extraction percentages of metal picrates with ionophores **2–4** ([host] = 4.0×10^{-5} M in CH₂Cl₂, [guest] = 4.0×10^{-5} M in water at 25 °C).

Thus, the extractability (*E*%) of **2–4** which followed the order of **2** > **3** > **4**, may be attributed to the shorter distance, the stronger N...Ag⁺...S interactions, the higher extractability (*E*%). This hypothesis is supported by the stability constants, which follow the same order of **2** > **3** > **4**. The binding constant (*K*_a) values for the complexation with Ag⁺ ion was determined to be $2.05 \times 10^4 \pm 875$ M⁻¹ (**2**), $3.86 \times 10^3 \pm 572$ M⁻¹ (**3**), $2.25 \times 10^3 \pm 365$ M⁻¹ (**4**) based on the Benesi-Hildebrand equation,²³ respectively (Fig. S13–18†).

Due to the existence of the two potential metal-binding sites, namely, the pyridine moieties and two benzyl moieties, there are several possibilities for metal complexation for compounds **2–4**. Both 1 : 1 and 1 : 2 metal complexation might be possible, attributable to electrostatic interactions as well as cation- π interactions. Job plots of **3** and **4** were carried out in the H₂O-CH₂Cl₂ phases. The *E*% values reach maxima at 0.5 mol fraction when **3** or **4** with Ag⁺ are changed systematically (Fig. S12†). Similar 1 : 1 coordination of **2** with Ag⁺ was shown by Job plots in our previous study (Fig. S12†).⁶ Thus, it can be concluded that Ag⁺ forms 1 : 1 complexes with **2–4**. These results suggest the major contribution of receptors **2–4** to Ag⁺ binding are from the nitrogens of the pyridine rings, and not from the alternative cation- π -interactions.

Furthermore, in order to look further into the binding properties of receptors **2–4** with Ag⁺, ¹H NMR titration experiments were carried out in CD₃Cl-CD₃CN = 10 : 1 solution. The chemical shift changes for compound **2–4** on complexation with Ag⁺ are illustrated in Fig. 3 and are summarized in Fig. 4.

Significant changes were observed for the pyridine ring protons after the complexation of each of **2–4** with

Fig. 3 ¹H NMR spectral changes of ionophores **2–4** (5×10^{-3} M) on addition of AgClO₄ (300 MHz, CDCl₃-CD₃CN = 10 : 1, [ionophores **2–4**] = 5×10^{-3} M). (a) Free **2**; (b) **2** in the presence of 1.0 equiv. of AgClO₄; (c) free **3**; (d) **3** in the presence of 1.0 equiv. of AgClO₄; (e) free **4**; (f) **4** in the presence of 1.0 equiv. of AgClO₄.

Fig. 4 Chemical shift changes of **2**, **3** and **4** induced in the presence of AgClO_4 ; + denotes the downfield and – denotes the upfield shift.

1.0 equiv. Ag^+ . In the case of **2**, the protons in the pyridine rings were shifted to lower field with $\Delta\delta = +0.27$, $+0.21$, $+0.52$ and $+0.83$ ppm for H_6 , H_5 , H_4 , and H_3 protons, respectively. In contrast, the OCH_2Py methylene protons were shifted dramatically up field, with $\Delta\delta = -0.53$. This may be due to both pyridine nitrogens of **2** close to the diaryl thiaether linkages ($\text{N}2\cdots\text{S}1 = \text{N}2\cdots\text{S}2 = 5.333$ Å, Fig. 1a). Thus, when **2** complexes with Ag^+ , the Ag^+ is easily captured through $\text{N}\cdots\text{Ag}^+\cdots\text{S}$ interactions.¹³ As a result, since the pyridine moieties are orientated inwards, the ring current shielding effect¹⁴ operating in the two thiacalixarene benzene rings is destroyed, forcing the steric conformation change. This affects the protons H_6 , H_5 , H_4 and H_3 of the pyridine rings which shift to lower field, due to the deshielding effect. Also, the OCH_2Py methylene protons become folded into the thiacalix[4]arene cavity and are thus shifted strongly upfield (-0.53 ppm), due to the steric conformation changes.

However, a different phenomenon was observed in the complexation of **3** with Ag^+ . From the X-ray results, both pyridine nitrogen atoms in **3** were orientated inwards and far from the diaryl thiaether linkages ($\text{N}2\cdots\text{S}1 = 6.360$ Å and $\text{N}2\cdots\text{S}2 = 5.847$ Å, Fig. 1b), which is exactly opposite to what is seen with **2**. The ^1H NMR spectrum of the Ag^+ of **3** reveals that the protons in the pyridine rings were shifted to lower field with $\Delta\delta = +0.09$, $+0.42$ and $+0.43$ ppm for H_6 , H_5 and H_4 protons, respectively. In contrast, a remarkable shielding effect experienced by proton H_2 (-0.38 ppm) was observed. It may be that when **3** complexes with Ag^+ , the Ag^+ is trapped in the cavity formed by the nitrogen atoms in pyridine, and then the proton H_2 end up folded into the π -cavity formed by the two thiacalixarene benzene rings and are thus shifted strongly upfield (-0.38 ppm). Thus, **3** complexes Ag^+ through the metal-nitrogen interactions and therefore, due to the interaction of the nitrogens and the Ag^+ , the H_6 , H_5 and H_4 protons of the pyridine rings shift to lower fields.¹⁵

Similar phenomena were observed for the complexation of **4** with Ag^+ ; protons H_3 and H_5 in the pyridine rings of **4** shifted to lower field after complexation ($+0.05$ ppm), as they are deshielded due to the $\text{N}\cdots\text{Ag}^+$ interactions. Pyridine ring protons H_2 and H_6 in **4** shifted upfield after complexation

(-0.06 ppm), which may be attributed to the weaker repulsion between the nitrogen atoms in the pyridine rings.¹⁵

The chemical shift changes of the thiacalixarene benzene and benzyl protons may also be attributed to the conformational changes of **2–4** upon complexation. The chemical shift changes ($\Delta\delta$) of **2–4** upon complexation are in the order $2 > 3 > 4$, which corresponds with the extractability of Ag^+ which was found to be in the same order.

To better understand the binding properties of receptors **2–4** with Ag^+ , a computation study was carried out. The molecular geometry of the individual structures in the gas-phase were fully optimized using Gaussian09,²² with the B3LYP level of DFT and the lanl2dz basis set. Significant conformational changes were observed for the pyridine ring protons of **2–4** after the complexation with Ag^+ . The conformation changes for **2** on complexation with Ag^+ ion can be seen in Fig. 5 (see the ESI for details of the computational study, Fig. S19–24†). Fig. 5 shows the structure (right) of the $2 \supset \text{Ag}^+$ complex. The optimized molecular geometry suggests that the Ag^+ binds, in accord with the ^1H NMR complexation study, *via* a $\text{N}\cdots\text{Ag}^+\cdots\text{S}$ short contact distance bond, which results in the conformation change. The $\text{N}\cdots\text{N}$ distance between the pyridine ring nitrogens decreases from 8.001 to 3.761 (Å) since the nitrogen

Fig. 5 Geometry-optimized (ball and stick) structures of: left: **2** and right: $2 \supset \text{Ag}^+$ complex. Color code for Ag^+ = magenta, pyridine nitrogen = blue, sulphur = yellow and oxygen atom = red. Hydrogen atoms have been omitted for clarity.

atoms move inwards after complexing with the Ag^+ . All four bridge sulphur atoms are roughly the same distance from the Ag^+ and presumably take an equal part in the coordination bonding.

However, a different phenomenon was observed in the complexation of 3–4 with Ag^+ . The N...N distance between the pyridine ring nitrogens decreases from 9.305 to 4.234 (Å) for 3 and 10.138 to 3.798 (Å) for 4 after complexing with the Ag^+ (Fig. S19–S24, Table S1†). The optimized molecular geometry suggests that complexation of 3–4 with Ag^+ occurs *via* N... Ag^+ interactions. The calculated complexation energies (ΔE kJ mol^{−1}) of the Ag^+ complexes of 2–4 are −488.096, −464.022 and −372.966 kJ mol^{−1} respectively (Table S2†), which is in agreement with the trend observed for the experimentally observed complexation data.

A preliminary evaluation of the anion binding efficiencies of 2–4 as potential extractants for the dichromate anion has been carried out by solvent extraction of aqueous solutions of $\text{K}_2\text{Cr}_2\text{O}_7$ into dichloromethane at different pH values according to a reported procedure.^{15,16} The extraction results, summarized in Fig. 6, indicate that 3 showed to be more effective for the extraction of dichromate anions at low pH (pH 1.5) than either 2 and 4. This is also consistent with the solvent extraction results seen with Cr^{3+} (Fig. 2). This could be attributed to the shorter (3.883 Å) distances (Fig. 1) between the pyridine nitrogen atoms in 3, which easily formed an efficient ion-pair (hydrogen bonded) complex in the two-phase extraction system following proton transfer to the nitrogen atoms. As the pH of the solution increased from 1.5 to 2.5 to 5.5 to 7.0, the *E*% for all three receptor ionophores decreased. This may directly be attributed to decreased proton concentrations in the solution.¹⁷ In other words, 3 showed a high extractability with dichromate anions only at lower pH, but another high extractability of Cr^{3+} at higher pH. Since, $\text{Cr}(\text{VI})$ is highly toxic, carcinogenic and harmful to human beings because it can diffuse as $\text{Cr}_2\text{O}_7^{2-}$ or HCr_2O_7^- through cell membranes and oxidize biological molecules,⁷ whereas $\text{Cr}(\text{III})$ is an essential

ion for mammals as it maintains effective glucose, lipid, and protein metabolisms.¹⁸ Thus, 3 could be a meaningful extractant when applying a solvent extraction method for the selective treatment of waste water containing $\text{Cr}(\text{VI})$ and $\text{Cr}(\text{III})$ prior to discharge.

Conclusion

Three 1,3-*alternate* thiacalix[4]arenes bearing pyridyl moieties (*ortho* for 2, *meta* for 3 and *para* 4) at the lower rim were regio-selectively synthesized. The solvent extraction experiments of the metal ions showed that the ability of these derivatives to complex with Ag^+ (95%, 52% and 36% for 2, 3 and 4, respectively) appeared to be largely dependent on the position of the pyridine nitrogen atoms. The mode of binding of the C_{2v} -symmetrical dipyrindyl-substituted thiacalix[4]arenes, 2–4 with Ag^+ was elucidated clearly using a ¹H NMR titration method. Two different complexation modes were observed: 2 armed with two *ortho* pyridyl groups, complexed with Ag^+ *via* N... Ag^+ ...S interactions, whereas 3 and 4 complexed with Ag^+ through N... Ag^+ interactions. The DFT computational studies were consistent with the experimental findings. These findings will provide us an important rule to design an appropriate thiacalix-arene ionophore in the future.

Further studies aimed at the potential for application of these extractants for the treatment of waste water containing $\text{Cr}(\text{VI})$ and $\text{Cr}(\text{III})$ were initiated. The combination of the two-phase solvent extraction data of Cr^{3+} and the results of the dichromate anion extraction by 3, suggests that 3 could be useful in the application of a solvent extraction method for the selective treatment of waste water containing $\text{Cr}(\text{VI})$ and $\text{Cr}(\text{III})$ ions prior to discharge.

Experimental section

General

All melting points were determined using a Yanagimoto MP-S1. ¹H NMR spectra were determined at 300 MHz with a Nippon Denshi JEOL FT-300 NMR spectrometer with SiMe_4 as an internal reference; *J* values are given in Hz. IR spectra were measured as KBr pellets or as liquid films on NaCl plates in a Nippon Denshi JIR-AQ20M spectrophotometer. UV spectra were measured by a Shimadzu 240 spectrophotometer. Mass spectra were obtained on a Nippon Denshi JMS-01SG-2 mass spectrometer at an ionization energy of 70 eV using a direct inlet system through GLC. Elemental analyses were performed by a Yanaco MT-5.

Materials

25,27-Dibenzoyloxy-5,11,17,23-tetra-*tert*-butyl-2,8,14,20-tetra-thiacalix[4]arene-26,28-diol (*distal*-1) was prepared from 5,11,17,23-tetra-*tert*-butyl-2,8,14,20-tetrathiacalix[4]arene-25,26,27,28-tetraol in one step according to a reported procedure.⁶

Fig. 6 *E*% values of dichromate anion with ionophores 2–4 (2.0 × 10^{−4} M, 2 h at 25 °C) at pH 1.5–7.0 (H_2O – CH_2Cl_2 : 10/10 (v/v); $\text{K}_2\text{Cr}_2\text{O}_7$ = 1 × 10^{−4} M).

1 O-Alkylation of *distal*-1 with 3-(chloromethyl)pyridine in the presence of Cs₂CO₃

5 A mixture of *distal*-1 (400 mg, 0.44 mmol) and Cs₂CO₃ (1.60 g, 4.92 mmol) in dry tetrahydrofuran (THF; 8 mL) was heated at reflux for 1 h under N₂. A solution of 3-(chloromethyl)pyridine (4.92 mmol) [prepared by neutralization of 3-(chloromethyl)pyridine hydrochloride (807 mg, 4.92 mmol) in DMF (8 mL) with a solution of triethylamine (0.68 mL, 4.92 mmol) in THF (8 mL) at room temperature] was then added and the mixture heated at reflux for an additional 24 h. After cooling the reaction mixture to room temperature, it was acidified with 1 M HCl (10 mL) and extracted with CH₂Cl₂ (100 mL × 2). The combined extracts were washed with water (50 mL × 2), and dried (MgSO₄) and concentrated under reduced pressure to give a yellow oil. The residue was washed with methanol to give a mixture of tetra-*O*-alkylated products as a colorless precipitate. The precipitate was washed with ether (5 mL) to give a colorless solid. Recrystallization from MeOH-CHCl₃ (1 : 3) gave **3** as a colorless prisms (280 mg, 59%).

25 **25,27-Dibenzyloxy-26,28-bis[(3-pyridylmethyl)oxy]-5,11,17,23-tetra-*tert*-butyl-2,8,14,20-tetrathiacalix[4]arene (3).** Colourless prisms [MeOH-CHCl₃ (1 : 3)], m.p. 285.4–286.6 °C. IR $\nu_{\max}$ (KBr)/cm⁻¹: 3058, 3030, 2958, 2902, 2868, 1575, 1546 and 1496. ¹H NMR (400 MHz, CDCl₃) δ = 0.85 (s, 18H, tBu), 0.86 (s, 18H, tBu), 5.06 (s, 4H, Ar-OCH₂Ph), 5.19 (s, 4H, Ar-OCH₂Py), 6.92 (d, *J* = 7.2 Hz, 4H, Ph-*H*), 7.02 (t, *J* = 7.6 Hz, 6H, Ph-*H*), 7.07 (s, 4H, Ar-*H*), 7.10 (s, 4H, Ar-*H*), 7.12 (t, *J* = 7.6 Hz, 2H, Py-*H*₅), 7.24 (d, *J* = 8.0 Hz, 2H, Py-*H*₄), 8.22 (s, 2H, Py-*H*₂) and 8.46 (d, *J* = 4.8 Hz, 2H, Py-*H*₆) ppm. ¹³C NMR (100 MHz, CDCl₃) δ = 30.77, 30.78, 33.89, 33.90, 67.74, 70.64, 122.99, 126.83, 127.18, 127.98, 128.38, 128.42, 128.48, 128.60, 133.20, 134.81, 137.47, 146.31, 146.58, 148.33, 148.92, 155.59 and 156.61 ppm. FABMS: *m/z*: 1083.30 (M⁺). Anal. calcd for C₆₆H₇₀N₂O₄S₄ (1083.53): C 73.16, H 6.51, N 2.59%. Found: C 71.85, H 6.56, N 2.38%.

40 Preparation of 25,27-dibenzyloxy-26,28-bis[(2-pyridylmethyl)oxy]-5,11,17,23-tetra-*tert*-butyl-2,8,14,20-tetrathiacalix[4]arene (**2**) was carried out following our previous report.⁶

45 **25,27-Dibenzyloxy-26,28-bis[(2-pyridylmethyl)oxy]-5,11,17,23-tetra-*tert*-butyl-2,8,14,20-tetrathiacalix[4]arene (2).** Colourless prisms [MeOH-CHCl₃ (1 : 3)], m.p. 274.8–275.5 °C. IR $\nu_{\max}$ (KBr)/cm⁻¹: 3058, 3029, 3008, 2955, 2901, 2866, 1571, 1588, 1546, 1496. ¹H NMR (400 MHz, CDCl₃) δ = 0.83 (s, 18 H, tBu), 0.85 (s, 18 H, tBu), 5.07 (s, 4 H, Ar-OCH₂Ph), 5.20 (s, 4 H, Ar-OCH₂Py), 6.66 (d, *J* = 7.2 Hz, 2 H, Py-*H*₃), 6.94 (d, *J* = 7.0 Hz, 4 H, Ph-*H*), 7.02 (t, *J* = 7.5 Hz, 6 H, Ph-*H*), 7.09 (4 H, s, Ar-*H*), 7.12 (4 H, s, Ar-*H*), 7.35 (t, *J* = 6.9 Hz, 4 H, Py-*H*_{4,5}) and 8.49 (d, *J* = 4.8 Hz, 2 H, Py-*H*₆) ppm. ¹³C NMR (100 MHz, CDCl₃) δ = 30.73, 30.82, 30.88, 33.88, 70.65, 71.46, 121.99, 126.80, 127.19, 127.99, 128.23, 128.60, 128.71, 137.52, 146.15, 146.35, 148.33, 156.01, 156.67 and 157.70 ppm.

55 Preparation of 25,27-dibenzyloxy-26,28-bis[(4-pyridylmethyl)oxy]-5,11,17,23-tetra-*tert*-butyl-2,8,14,20-tetrathiacalix[4]arene (**4**) was carried out following our previous report.¹⁰

1 **25,27-Dibenzyloxy-26,28-bis[(4-pyridylmethyl)oxy]-5,11,17,23-tetra-*tert*-butyl-2,8,14,20-tetrathiacalix[4]arene (4).** Colourless prisms [MeOH-CHCl₃ (1 : 3)], m.p. 283–285 °C. IR $\nu_{\max}$ (KBr)/cm⁻¹: 3055, 3029, 2952, 2921, 2853, 1604, 1572, 1562. ¹H NMR (400 MHz, CDCl₃) δ = 0.84 (s, 18 H, tBu), 0.86 (s, 18 H, tBu), 5.07 (s, 4 H, Ar-OCH₂Ph), 5.12 (s, 4 H, Ar-OCH₂Py), 6.90 (d, *J* = 5.5 Hz, 4 H, Py-*H*_{3,5}), 6.94 (d, *J* = 7.4 Hz, 4 H, Ph-*H*), 7.04 (t, *J* = 7.6 Hz, 4 H, Ph-*H*), 7.10 (s, 4 H, Ar-*H*), 7.12 (s, 4 H, Ar-*H*), 7.13–7.18 (m, 2 H, Ph-*H*) and 8.40 (d, *J* = 5.8 Hz, 4 H, Py-*H*_{2,6}) ppm. ¹³C NMR (100 MHz, CDCl₃) δ = 30.70, 30.73, 33.86, 33.89, 69.10, 70.80, 121.80, 126.90, 127.14, 128.03, 128.26, 128.75, 129.11, 137.35, 146.39, 146.50, 146.71, 149.38, 155.97 and 156.72 ppm. FABMS: *m/z*: 1083.45 (M⁺).

15 Extraction experiments and stoichiometry of metal complexation

20 Metal picrates (4.0 × 10⁻⁵ M) were prepared *in situ* by dissolving the metal hydroxide (0.02 mol) in 4.0 × 10⁻⁵ M picric acid (1000 mL); triply distilled water was used for all aqueous solutions. Two-phase solvent extraction was carried out between aqueous picrates (10 mL, [metal picrate] = 4.0 × 10⁻⁵ M) and host (10 mL, [host] = 4 × 10⁻⁵ M in CH₂Cl₂). The two-phase mixture in a stoppered flask was immersed in a thermostated water bath at 25 °C which was shaken at 300 strokes per min for 4 h and then kept at the same temperature for 1 h, allowing the complete separation of the two phases. This was repeated 3 times. The absorbance of each solution was determined by UV spectroscopy (λ = 356 nm). The method of continuous variation was employed to determine the stoichiometry in the complexes involving the host receptors **2**, **3** or **4**. The molar ratios of both the host and metal picrate were varied from 0 to 1, while the total concentration was kept at several constant levels. Job plots were generated by plotting the extracted [M⁺] versus the mole fraction of metal. We confirmed that this period was sufficient to attain the distribution equilibrium. The extractability was determined spectrophotometrically from the decrease in the absorbance of the picrate ion in the aqueous phase, as described by Pedersen.¹⁹

40 ¹H-NMR complexation experiments

45 To a CDCl₃-CH₃CN (10 : 1, v/v) solution (5 × 10⁻³ M) of **2**, **3** or **4** in an NMR tube was added a CD₃CN solution (5 × 10⁻³ M) of AgClO₄. The spectra were recorded after the additions. The temperature of the NMR probe was kept constant at 27 °C. The ¹H NMR data of the most-representative complexes are given below:

50 **2**: ¹H NMR (300 MHz, CHCl₃-CH₃CN, 10 : 1, v/v): δ = 0.83 (s, 18H, tBu), 0.85 (s, 18H, tBu), 5.07 (s, 4 H, CH₂-Ph), 5.19 (s, 4H, CH₂-Py), 6.67 (d, *J* = 7.8 Hz, 2H, Py-*H*₃), 6.94 (d, *J* = 7.6 Hz, 4H, Ph-*H*), 7.03 (t, *J* = 6.6 Hz, 6H, Ph-*H*), 7.10 (s, 4H, Ar-*H*), 7.13 (s, 4H, Ar-*H*), 7.36 (t, *J* = 7.8 Hz, 4H, Py-*H*_{4,5}) and 8.49 (d, *J* = 4.7 Hz, 2H, Py-*H*₆) ppm.

55 **2** ⊃ Ag⁺: ¹H NMR (300 MHz, CHCl₃-CH₃CN, 10 : 1, v/v): δ = 0.85 (s, 18H, tBu), 0.98 (s, 18H, tBu), 4.65 (s, 4H, CH₂-Py), 5.06 (s, 4H, CH₂-Ph), 6.98 (d, *J* = 7.5 Hz, 4H, Ph-*H*), 7.10–7.14 (m, 4H, Ph-*H*), 7.16 (s, 4H, Ar-*H*), 7.21 (t, *J* = 6.6 Hz, 2H, Ph-*H*),

7.26 (s, 4H, Ar-H), 7.50 (d, $J = 7.8$ Hz, 2H, Py-H₃), 7.57 (t, $J = 5.7$ Hz, 2H, Py-H₄), 7.82–7.92 (m, 2H, Py-H₅) and 8.77 (d, $J = 4.9$ Hz, 2H, Py-H₆) ppm.

3: ¹H NMR (300 MHz, CHCl₃–CH₃CN, 10 : 1, v/v): δ = 0.85 (s, 18H, tBu), 0.86 (s, 18H, tBu), 5.06 (s, 4H, CH₂–Ph), 5.19 (s, 4H, CH₂–Py), 6.93 (d, $J = 7.2$ Hz, 4H, Ph-H), 7.00–7.05 (m, 6H, Ph-H), 7.08 (s, 4H, Ar-H), 7.10 (s, 4H, Ar-H), 7.13 (m, 2H, Py-H₅), 7.27 (d, $J = 7.8$ Hz, 2H, Py-H₄), 8.20 (s, 2H, Py-H₂) and 8.46 (d, $J = 3.9$ Hz, 2H, Py-H₆) ppm.

3 $\supset$ Ag⁺: ¹H NMR (300 MHz, CHCl₃–CH₃CN, 10 : 1, v/v): δ = 0.84 (s, 18H, tBu), 0.91 (s, 18H, tBu), 5.07 (s, 4H, CH₂–Ph), 5.19 (s, 4H, CH₂–Py), 6.98 (d, $J = 7.5$ Hz, 4H, Ph-H), 7.07 (s, 4H, Ar-H), 7.08–7.12 (m, 4H, Ph-H), 7.17–7.20 (m, 2H, Ph-H), 7.21 (s, 4H, Ar-H), 7.52–7.57 (m, 2H, Py-H₅), 7.69 (d, $J = 7.9$ Hz, 2H, Py-H₄), 7.82 (s, 2H, Py-H₂) and 8.55 (d, $J = 5.1$ Hz, 2H, Py-H₆) ppm.

4: ¹H NMR (300 MHz, CHCl₃–CH₃CN, 10 : 1, v/v): δ = 0.84 (s, 18H, tBu), 0.85 (s, 18H, tBu), 5.07 (s, 4H, CH₂–Ph), 5.13 (s, 4H, CH₂–Py), 6.91 (d, $J = 5.5$ Hz, 4H, Py-H_{3,5}), 6.94 (d, $J = 7.4$ Hz, 4H, Ph-H), 7.02–7.07 (m, 4H, Ph-H), 7.11 (s, 4H, Ar-H), 7.13 (s, 4H, Ar-H), 7.14–7.18 (m, 2H, Ph-H) and 8.38 (d, $J = 5.9$ Hz, 4H, Py-H_{2,6}) ppm.

4 $\supset$ Ag⁺: ¹H NMR (300 MHz, CHCl₃–CH₃CN, 10 : 1, v/v): δ = 0.86 (s, 18H, tBu), 0.91 (s, 18H, tBu), 5.11 (s, 4H, CH₂–Ph), 5.15 (s, 4H, CH₂–Py), 6.94–6.99 (m, 4H, Py-H_{3,5}), 7.09 (d, $J = 7.3$ Hz, 2H, Ph-H), 7.13 (s, 4H, Ar-H), 7.16–7.21 (m, 6H, Ph-H), 7.21 (s, 4H, Ar-H) and 8.32 (d, $J = 6.0$ Hz, 4H, Py-H_{2,6}) ppm.

Crystallographic analyses of 3

Diffraction data were collected on a Bruker APEX 2 CCD diffractometer equipped with graphite-monochromated Mo-K α radiation at 150(2) K.²⁰ Data were corrected for Lorentz and polarisation effects and for absorption.²⁰ The structures were solved by direct methods and refined by full-matrix least-squares methods, on F^2 .²¹ H atoms were refined using a riding model except for those on hetero atoms in 3 which were freely refined.

Crystal data for 3. C₆₆H₇₀N₂O₄S₄, $M = 1083.48$. Orthorhombic, space group $Pmn2_1$, $a = 15.1668$ (6), $b = 14.7772$ (7), $c = 12.7612$ (6) Å, $V = 2860.1$ (2) Å³. $Z = 2$, $D_c = 1.258$ g cm⁻³, $F(000) = 1152$, $T = 100$ K, $\mu(\text{Mo-K}\alpha) = 0.17$ mm⁻¹, $\lambda(\text{Mo-K}\alpha) = 0.6525$ Å, colourless crystal of size $0.20 \times 0.20 \times 0.06$ mm³. The total number of reflections measured, to $\theta_{\text{max}} = 30.3^\circ$, was 345 676 of which 11 331 were unique ($R_{\text{int}} = 0.087$); 10 920 were ‘observed’ with $I > 2\sigma(I)$. For the ‘observed’ data only, $R_1 = 0.037$; $wR_2 = 0.101$ for all 11 331 reflections and 400 parameters. Residual electron density within ± 0.48 eÅ⁻³.

Crystallographic data (excluding structure factors) for the structures in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication numbers CCDC 245644 for 2⁶ and 1021161 for 3, respectively.

Acknowledgements

This work was performed under the Cooperative Research Program of “Network Joint Research Center for Materials and

Devices (Institute for Materials Chemistry and Engineering, Kyushu University)”. We would like to thank the OTEC at Saga University and the International Cooperation Projects of Guizhou Province (no. 20137002), The Royal Society of Chemistry for financial support and the EPSRC for an overseas travel grant to C.R.

Notes and references

- (a) N. Morohashi, F. Narumi, N. Iki, T. Hattori and S. Miyano, *Chem. Rev.*, 2006, **106**, 5291–5316; (b) R. Kumar, Y. O. Lee, V. Bhalla, M. Kumar and J. S. Kim, *Chem. Soc. Rev.*, 2014, **43**, 4824–4870.
- (a) F. Arnaud-Neu, G. Barrett, S. J. Harris, M. Owens, M. A. McKerverey, M.-J. Schwing-Weill and R. Schwinte, *Inorg. Chem.*, 1993, **32**, 2644–2650; (b) E. Ghidini, F. Ugozzoli, R. Ungaro, S. Harkema, A. Abu ElFadl and D. N. Reinhoudt, *J. Am. Chem. Soc.*, 1990, **112**, 6979–6985; (c) S. Shinkai, T. Otsuka, K. Fujimoto and T. Matsuda, *Chem. Lett.*, 1990, **19**, 835–838; (d) T. Tuntulani, P. Thavornnyutikarn, S. Poompradub, N. Jaiboon, V. Ruangpornvisuti, N. Chaichit, Z. Asfari and J. Vicens, *Tetrahedron*, 2002, **58**, 10277–10285.
- (a) M. Kumar, N. Kumar and V. Bhalla, *Org. Biomol. Chem.*, 2012, **10**, 1769–1774; (b) S. N. Podyachev, N. E. Burmakina, V. V. Syakaev, S. N. Sudakova, W. D. Habicher and A. I. Konovalov, *J. Inclusion Phenom. Macrocyclic Chem.*, 2011, **71**, 161–168; (c) T. Kajiwarra, N. Iki and M. Yamashita, *Coord. Chem. Rev.*, 2007, **251**, 1734–1746.
- (a) F. Botha, J. Budka, V. Eigner, O. Hudeček, L. Vrzal, I. Císařová and P. Lhoták, *Tetrahedron*, 2014, **70**, 477–483; (b) H. Zhao, J. Zhan, Z. Zou, F. Miao, H. Chen, L. Zhang, X. Cao, D. Tian and H. Li, *RSC Adv.*, 2013, **3**, 1029–1032; (c) F. Miao, J. Zhan, Z. Zou, D. Tian and H. Li, *Tetrahedron*, 2012, **68**, 2409–2413.
- (a) *Calixarenes in the Nanoworld*, ed. J. Vicens and J. Harrowfield, Springer, Berlin, 2007; (b) M. Kumar, A. Dhir and V. Bhalla, *Eur. J. Org. Chem.*, 2009, 4534–4540.
- T. Yamato, C. P. Casas, H. Yamamoto, M. R. J. Elsegood, S. H. Dale and C. Redshaw, *J. Inclusion Phenom. Macrocyclic Chem.*, 2005, **54**, 261–269.
- P. G. Krishna, J. M. Gladis, U. Rambabu, T. P. Rao and G. R. K. Naidu, *Talanta*, 2004, **63**, 541–546.
- (a) A. Sap, B. Tabakci and A. Yilmaz, *Tetrahedron*, 2012, **68**, 8739–8745; (b) S. Sayin, M. Yilmaz and M. Tavasli, *Tetrahedron*, 2011, **67**, 3743–3753; (c) F. Ozcan, M. Ersoz and M. Yilmaz, *Mater. Sci. Eng., C*, 2009, **29**, 2378–2383; (d) S. Bozkurt, E. Kocabas, M. Durmaz, M. Yilmaz and A. Sirit, *J. Hazard. Mater.*, 2009, **165**, 974–979.
- (a) A. Yilmaz, S. Memon and M. Yilmaz, *Tetrahedron*, 2002, **58**, 7735–7740; (b) M. Tabakci, S. Memon, M. Yilmaz and D. M. J. Roundhill, *J. Inclusion Phenom. Macrocyclic Chem.*, 2003, **45**, 265–270; (c) M. Bayrakci, S. Ertul, O. Sahin and M. Yilmaz, *J. Inclusion Phenom. Macrocyclic Chem.*, 2009, **63**, 241–247; (d) M. Tabakci, S. Memon and M. Yilmaz, *Tetra-*

- hedron, 2007, **63**, 6861–6886; (e) D. M. Roundhill and H. F. Koch, *Chem. Soc. Rev.*, 2002, **31**, 60–67.
- 10 C. Pérez-Casas, S. Rahman, N. Begum, Z. Xi and T. Yamato, *J. Inclusion Phenom. Macrocyclic Chem.*, 2007, **60**, 173–185.
- 5 11 (a) P. M. Marcos, F. A. Teixeira, M. A. P. Segurado, J. R. Ascensoc, R. J. Bernardino, P. J. Cragg, S. Michel, V. Hubscher-Bruder and F. Arnaud-Neuf, *J. Phys. Org. Chem.*, 2013, **26**, 295–305; (b) C. Bonaccorso, F. Nicoletta, V. Zito, G. Arena, D. Sciotto and C. Sgarlata, *Supramol. Chem.*, 2013, **25**, 615–625; (c) J. J. Colleran, B. S. Creaven, D. F. Donlon and J. McGinley, *Dalton Trans.*, 2010, **39**, 10928–10936; (d) M. Tabakci, S. Memon and M. Yilmaz, *Tetrahedron*, 2007, **63**, 6861–6865.
- 10 12 (a) T. Yamato, M. Haraguchi, J.-I. Nishikawa, S. Ide and H. Tsuzuki, *Can. J. Chem.*, 1998, **76**, 989–996; (b) T. Yamato, M. Haraguchi and S. Ide, *J. Chem. Soc., Perkin Trans. 1*, 1998, 609–614; (c) T. Yamato, *J. Inclusion Phenom. Macrocyclic Chem.*, 1998, **32**, 195–207.
- 15 13 A. Ovsyannikov, M. N. Lang, S. Ferlay, S. E. Solovieva, I. S. Antipin, A. I. Konovalov, N. Kyritsakas and M. W. Hosseini, *Dalton Trans.*, 2013, **42**, 116–126.
- 20 14 (a) F. Vögtle, *Cyclophane Chemistry*, John Wiley & Sons Ltd, 1993; (b) M. Tashiro and T. Yamato, *J. Org. Chem.*, 1981, **46**, 4556–4562; (c) M. Tashiro and T. Yamato, *J. Org. Chem.*, 1983, **48**, 1461–1468.
- 25 15 (a) X. L. Ni, C. C. Jin, X. K. Jiang, M. Takimoto, S. Rahman, X. Zeng, D. L. Hughes, C. Redshaw and T. Yamato, *Org. Biomol. Chem.*, 2013, **11**, 5435–5442; (b) J. L. Zhao, H. Tomiyasu, X. L. Ni, X. Zeng, M. R. J. Elsegood, C. Redshawd, S. Rahman, P. E. Georgiou and T. Yamato, *New J. Chem.*, 2014, **38**, 6041–6049.
- 30 16 S. Sayin, F. Ozcan and M. Yilmaz, *Mater. Sci. Eng., C*, 2013, **33**, 2433–2439.
- 35 17 A. Sap, B. Tabakci and A. Yilmaz, *Tetrahedron*, 2012, **68**, 8739–8745.
- 18 M. E. Losi, C. Amrhein and W. T. Frankenberger, *Rev. Environ. Contam. Toxicol.*, 1994, **136**, 91–121.
- 5 19 (a) C. J. Pedersen, *J. Am. Chem. Soc.*, 1967, **89**, 7017–7036; (b) C. J. Pedersen, *J. Am. Chem. Soc.*, 1967, **89**, 2495–2496; (c) C. J. Pedersen, *J. Am. Chem. Soc.*, 1970, **92**, 391–394.
- 10 20 SAINT and APEX 2 software for CCD diffractometers, Bruker AXS Inc., Madison, USA, 2008.
- 21 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Fundam. Crystallogr.*, 2008, **64**, 112–122.
- 22 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven Jr., J. A. Montgomery, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski and D. J. Fox, *Gaussian 09, Revision D.01*, Gaussian, Inc., Wallingford CT, 2013.
- 23 H. A. Benesi and J. H. Hildebrand, *J. Am. Chem. Soc.*, 1949, **71**, 2703–2707.
- 35 35
- 40 40
- 45 45
- 50 50
- 55 55