This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

The influence of local skin temperature on the sweat glands maximum ion reabsorption rate

PLEASE CITE THE PUBLISHED VERSION

<https://doi.org/10.1007/s00421-018-04059-5>

PUBLISHER

© Springer

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This is a post-peer-review, pre-copyedit version of an article published in European Journal of Applied Physiology. The final authenticated version is available online at: <https://doi.org/10.1007/s00421-018-04059-5>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Gerrett, Nicola, Tatsuro Amano, George Havenith, Yoshimitsu Inoue, and N. Kondo. 2019. "The Influence of Local Skin Temperature on the Sweat Glands Maximum Ion Reabsorption Rate". Loughborough University. <https://hdl.handle.net/2134/36918>.

The influence of local skin temperature on the sweat glands maximum ion reabsorption rate

Gerrett, N¹. Amano, T². Havenith, G³. Inoue, Y⁴. & Kondo, N¹.

¹ Laboratory for Applied Human Physiology, Graduate School of Human Development and Environment, Kobe University, Kobe, Japan

² Laboratory for Exercise and Environmental Physiology, Faculty of Education, Niigata University, Niigata, Japan

³ Environmental Ergonomics Research Centre, Loughborough Design School, Loughborough University, Loughborough, United Kingdom.

⁴ Laboratory for Human Performance Research, Osaka International University, Osaka, Japan

Address for correspondence: Narihiko Kondo, Ph.D., Laboratory for Applied Human Physiology, Graduate School of Human Development and Environment, Kobe University, 3-11 Tsurukabuto, Nada-Ku, Kobe 657-8501, Japan, Tel and Fax: +81- 78-803-7816, Email: kondo@kobe-u.ac.jp

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

Acknowledgements

We thank our participants for volunteering their time. We also thank Drs S. Koga and D. Okushima for their insightful comments during the preparation of this manuscript. Finally, we thank Dr Koji Sato for his assistance with aldosterone analysis.

Grants

This study was supported by a Grant-in-Aid for Scientific Research (16H04851 and 17H0253) from the Japan Society for the Promotion of Science from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

Disclosures

The authors declare no conflicts of interest, financial or otherwise.

Abstract

Purpose

Changes in mean skin temperature (T_{sk}) have been shown to modify the maximum rate of sweat ions reabsorption. This study aims to extend this knowledge by investigating if modifications could also be caused by local T_{sk} .

Methods

The influence of local T_{sk} on the sweat glands maximum ion reabsorption rates was investigated in 10 healthy volunteers (3 female and 7 male; 20.8 ± 1.2 yrs, 60.4 ± 7.7 kg, 169.4 ± 10.4 cm) during passive heating (water perfused suit and lower leg water immersion). In two separate trials, in a randomised order, one forearm was always manipulated to 33°C (Neutral), whilst the other was manipulated to either 30°C (Cool) or 36°C (Warm) using water perfused patches. Oesophageal temperature (T_{es}), forearm T_{sk} , sweat rate (SR), galvanic skin conductance (GSC) and salivary aldosterone concentrations were measured. The sweat glands maximum ion reabsorption rates were identified using the ΔSR threshold for an increasing ΔGSC .

Results

Thermal (T_{es} and body temperature (T_b)) and non-thermal responses (aldosterone) were similar across all conditions ($p>0.05$). A temperature dependent response for the sweat glands maximum ion reabsorption rates was evident between 30°C (0.18 ± 0.10 $\text{mg}/\text{cm}^2/\text{min}$) and 36°C (0.28 ± 0.14 $\text{mg}/\text{cm}^2/\text{min}$, $d = 0.88$, $p<0.05$), but not for 33°C (0.22 ± 0.12 $\text{mg}/\text{cm}^2/\text{min}$), $d = 0.44$ and $d = 0.36$, $p>0.05$).

Conclusion

The data indicates that small variations in local T_{sk} may not affect the sweat glands maximum ion reabsorption rates but when the local T_{sk} increases by $>6^{\circ}\text{C}$, ion reabsorption rates also increase.

79 **Keywords:** sweat ion regulation, sweat glands, skin temperature, aldosterone

80

81 **Abbreviations**

82 ANOVA analysis of variance

83 CFTR cystic fibrosis transmembrane channels

84 Cl^- Chloride

85 CVC cutaneous vascular conductance (%)

86 ENaC epithelial sodium channel

87 GSC galvanic skin conductance (μS)

88 HASG heat-activated sweat glands

89 HR heart rate (bpm)

90 K^+ potassium

91 MAP mean arterial pressure (mmHG)

92 Na^+ sodium

93 SGO sweat gland output

94 SR sweat rate ($\text{mg}/\text{cm}^2/\text{min}$)

95 T_b body temperature ($^{\circ}\text{C}$)

96 T_{es} Oesophageal temperature ($^{\circ}\text{C}$)

97 T_{sk} skin temperature ($^{\circ}\text{C}$)

98 $\dot{V}\text{O}_{2\text{max}}$ maximum oxygen uptake ($\text{ml}/\text{kg}/\text{min}$)

99

100 **Introduction**

101 Sweating is an essential physiological function for maintaining thermal homeostasis

102 and the content of sweated ions important for maintaining epidermal barrier homeostasis

103 and antimicrobial function of the skin. Despite rapid developments in biosensor technology

that aim to determine sweat rate and its ion concentration (Gao et al. 2018), our knowledge of sweat ion regulation is limited. It is well established that when eccrine sweat glands are stimulated, the secretory coil, located at the base of the sweat gland, forms an isotonic fluid and as sweat traverses the straight duct ion reabsorption occurs; resulting in a hypotonic sweat released on the skin surface (Sato 1977; Shamsuddin et al. 2005b). It has been suggested that with an increasing sweat rate (SR) the secretion rate increases proportionally faster than the reabsorption rate, resulting in a linear relationship between SR and sweat Na^+ concentration (Buono et al. 2008). Whilst this may be the case there are numerous studies providing evidence that the reabsorption rate can be enhanced, most notably following heat acclimation whereby a reduction in sweated ions occurs despite an increased sweat production (Buono et al. 2007; Amano et al. 2016). Furthermore, the maximum rate of ion reabsorption has been shown to differ between habitually trained and sedentary individuals and is also known to vary across the body (Amano et al. 2017). What regulates eccrine sweat glands ion reabsorption remains unknown.

A regulatory mechanism that has been previously reported to affect the maximum ion reabsorption rate is T_{sk} . The influence of T_{sk} on sweat production is well established (Nadel et al. 1971) but its impact on sweated ions is somewhat contradictory with studies reporting an influential role (Johnson et al. 1944; Robinson et al. 1985) and others reporting little or no effect (Bulmer and Forwell 1956). These earlier studies were primarily focused on the concentration of sweated ions but more recently Shamsuddin *et al.* (2005a) investigated the reabsorption of these ions, that ultimately affects the sweated ion concentration. By changing the ambient conditions (15°C or 25°C), Shamsuddin et al. (2005a) manipulated mean T_{sk} by approximately 3°C during an exercising protocol. The result of which was a significantly lower ion reabsorption rate at a lower compared to higher mean T_{sk} (0.21 ± 0.04 vs. 0.52 ± 0.06 mg/cm²/min, respectively). The inhibition of the

sweat glands ion reabsorption capacity was associated with the temperature dependency of epithelial sodium channel (ENaC) excitability and open probability (Chraïbi and Horisberger 2003). However, they did not report the local T_{sk} where the ion reabsorption rates were measured and also only focused on the effects of low T_{sk} ($<31^{\circ}\text{C}$) on the sweat glands maximum ion reabsorption rates. Therefore, it is still unknown whether the maximum ion reabsorption rates are proportional to local changes in T_{sk} systematically.

It is also necessary to determine the influence of local T_{sk} on the sweat glands maximum ion reabsorption rates as there is cumulative evidence for regional differences in ion reabsorption rates, with the torso generally having a higher reabsorption rate than the extremities (Amano et al. 2017; Gerrett et al. 2018a). However these regional differences were also mirrored by regional differences in local T_{sk} . It would be advantageous then to investigate the influence of local T_{sk} on the sweat glands ions reabsorption rate within temperatures ranges expected whilst sweating.

Shamsuddin et al. (2005a) speculated that higher T_{sk} might enhance the responsive ability of the sweat glands to a given concentration of circulating aldosterone; a hormone considered to be a primary effector of sodium exchange at the sweat gland (Hegarty and Harvey 1998; Harvey and Higgins 2000). Aldosterone concentrations and other water regulatory hormones such as atrial natriuretic peptide, vasopressin and renin are higher during exercise and are considered to be exercise intensity dependent (Convertino et al. 1983; Freund et al. 1991; Yoshida et al. 2006). Shamsuddin et al. (2005a) employed an exercise protocol (cycling at $60\% \dot{V}O_{2\text{max}}$) sufficient enough for the release of these hormones; although their influence on sweat glands ion reabsorption is inconclusive (Yoshida et al. 2006; Hew-Butler et al. 2010, 2014). As such it is important to investigate the role of T_{sk} on the sweat glands maximum ion reabsorption rates when hormonal responses associated with exercise are isolated.

In order to determine the role of local T_{sk} on the sweat glands ion reabsorption rates we aimed to manipulate local forearm T_{sk} to elicit a cool (30°C), neutral (33°C) and warm (36°C) T_{sk} , with the latter reflecting typical T_{sk} experienced during heat stress on the sweat glands ion reabsorption. In order to minimise any non-thermoregulatory mechanisms affecting sweat glands ion reabsorption rates the influence of local T_{sk} on ion reabsorption during a passive heating protocol was selected. We hypothesized that eccrine sweat glands maximum ion reabsorption rates are temperature dependent, with maximum ion reabsorption occurring at a higher sweat rate (SR) with a higher local T_{sk} (36°C) compared to neutral (33°C) and cool T_{sk} (30°C).

Methods

Participants were informed about the study purpose and procedures prior to providing verbal and written consent. The Human Subjects Committee of the Graduate School of Human Development and Environment at Kobe University (Japan) approved the study (report no. 259), which conforms to the standards set out by the Declaration of Helsinki (except for registration in a database).

Participants

Ten young healthy recreationally active participants (3 female and 7 male; 20.8 ± 1.2 yrs, 60.4 ± 7.7 kg, 169.4 ± 10.4 cm) were recruited for this study during winter in Japan and thus were considered unacclimated. Both males and females were recruited, as our previous research indicated no sex related differences in ion reabsorption rates (Amano et al. 2017). Participants were asked to refrain from consuming high sodium foods, caffeine or alcohol and to avoid any strenuous exercise 24 hours preceding the trials. For the experimental trials they were instructed to record their food and beverage intake during the

preceding 24 hours and asked to replicate this for each trial. To promote euhydration, participants were instructed to consume 500 ml of water 1-2 hours prior to the experiments. Upon arrival to the laboratory, participants provided a urine sample and hydration status was checked by a handheld refractometer (Atago Co.Ltd, Tokyo, Japan). All participants met the criteria for adequate hydration (USG <1.025) except two participants both on one occasion each. In these cases the participants were required to consume a further 300ml prior to the start of the experiments. All participants were non-smokers and were not taking any medications. Menstrual cycle phase per se was not controlled for in the female participants but each female was tested within the same stage of their cycle.

Experimental trials

Participants visited the laboratory twice for the manipulation of local forearm T_{sk} . Each trial was separated by at least 48 hours. All experimental trials were conducted in a climatic chamber (SR-3000; Nagano Science, Osaka, Japan) controlled at 27°C, 50% relative humidity, with minimal air movement. All tests were completed at the same time of day (± 1 hr), at least 2 hours after their last meal. During all experimental trials, participants wore standardized shorts and females also wore a sports bra underneath a water-perfused suit (Allen-Vanguard, Ottawa, Canada) to help elevate the thermal load and in an attempt to maintain a similar T_{sk} across the body.

Prior to each experiment participants provided a urine sample for the analysis of aldosterone. Prior to entering the experimental chamber participant self-inserted an oesophageal thermometer, as an index of core temperature (T_{es}), to the distance of one-fourth of standing height from the external nares (Mekjavic and Rempel 1990). After entering the chamber participants donned a water-perfused suit that covered the body except face, forearms, hands and lower legs (from above the knee). Participants then rested

in a semi supine position for approximately 60mins whilst the measuring instruments were attached (detailed below) and local T_{sk} of the forearm was manipulated and remained stable at the desired local T_{sk} (detailed below). During the instrumentation preparation phase, water at 34°C was passed through the suit at a flow rate of 750 ml/min to maintain a stable resting mean T_{sk} . Following instrumentation, baseline data were recorded for 5 minutes and a salivary aldosterone sample collected. Participants were then heated exogenously for 45 minutes separated into two phases. The first phase involved submerging the lower legs into a water bath set at 42°C and increasing the water temperature inside the suit to 40°C for 20 minutes. For the remaining 25 minutes, the lower leg water bath temperature was increased to 43°C and the temperature of the water inside the suit was increased to 45°C.

During all experiments, one forearm was controlled to elicit a neutral T_{sk} of 33°C (Neutral), whilst the other forearm was either manipulated to elicit a cool T_{sk} of 30°C (Cool) or a warm temperature of 36°C (Warm) in a balanced order. For clarity, in the rest of the document the phrase *Neutral-Cool* will be used to indicate when one forearm was manipulated to be Neutral and the other Cool, and *Neutral-Warm* when one forearm was manipulated to be Neutral and the other Warm. Sweat rate on bilateral forearms are comparable (Buono et al. 2011; Kenefick et al. 2012; Smith and Havenith 2012) and thus we assume asymmetry between the left and right arm and use the mean of the two neutral conditions. To manipulate local forearm T_{sk} , custom-made water perfused patches were wrapped around the left and right ventral forearm during each experiment. The water perfused patches consisted of tubes (inside diameter=2mm, outside diameter =4mm) running in a circular motion with no gaps between tubes. A small hole positioned in the centre of the patch allowed for entrance and exit of the sweat capsule tubes. Both patches were of equal surface area (16x13cm) and water was supplied to each patch at a flow rate of 250 ml/min (Variable-flow Chemical Transfer Pump, Fisher Scientific™). The patches

were connected to two separate water baths (Taitec Therminder DX-10 and SP12, Taitec Corporation, Saitama, Japan) where the water temperature was altered to achieve the desired local T_{sk} . The water temperature used varied for each participant but was approximately $29.4 \pm 3.2^{\circ}\text{C}$ for Neutral, $15.8 \pm 5.8^{\circ}\text{C}$ for Cool and $40.2 \pm 2.2^{\circ}\text{C}$ for Warm conditions. To control the cold-water temperature an immersion cooler (Taitec Cool pipe 300LF, Taitec Corporation, Saitama, Japan) was used concomitantly with the water bath.

The water patches were positioned onto the forearms for a minimum of 30 minutes prior to the start of the test, whereby local T_{sk} was altered until the desired temperatures were reached. Local T_{sk} was maintained at the desired temperature for a minimum of 5 minutes prior to the start of the experiment. If the local T_{sk} of the forearms during passive heating increased or decreased by more than 1°C of the target local T_{sk} , the temperature of the water bath was altered.

During the final minute of each experiment, a salivary aldosterone sample was collected. This was then followed by the removal of the forearm skin patches and all instruments so that the number of heat-activated sweat glands (HASG) using the starch-iodine technique (Inoue 1996) could be measured. Once these samples were collected the passive heating protocol was completed.

Measurements

For the determination of mean skin temperature (mean T_{sk}), 4 sites (chest, forearm, thigh and calf) were measured. In addition, local T_{sk} underneath the skin patch at the left and right forearm was measured at two locations adjacent to the sweat capsule and averaged. Both T_{es} and T_{sk} were measured using copper-constantan thermocouples (Inui Engineering, Higashi Osaka, Japan). The tip of the oesophageal thermometer was covered with silicon and the skin thermocouples were uncovered and attached to the skin using

Medipore tape. Mean T_{sk} and mean body temperature (T_b) were calculated using the following respective formula (Stolwijk and Hardy 1966; Gagge and Nishi 2011);

$$T_b = (0.8 \times T_{es}) + (0.2 \times \text{mean } T_{sk})$$

$$\text{Mean } T_{sk} = (\text{chest} \times 0.34) + (\text{calf} \times 0.18) + (\text{thigh} \times 0.33) + (\text{forearm} \times 0.15)$$

For the calculation of mean T_{sk} the forearm T_{sk} , was based on the average of both the left and right forearm T_{sk} .

SR was measured using the ventilated capsule method on the right and left mid-ventral forearm. Dry nitrogen gas was flushed (500ml/min) through the apparatus approximately 1 hour prior to each experiment to ensure stable readings. Each capsule (3.14 cm²) was affixed to the skin using double-sided tape at least 30 minutes prior to data collection. The temperature and humidity of the air flowing out of the capsule was measured using a capacitance hygrometer (HMP50; Vaisala, Helsinki, Finland). Two Ag/AgCl electrodes (Vitrode J, Nihon Kohden, Tokyo, Japan) for measuring GSC were attached either side of the sweat capsules, approximately 3cm apart (MP100 and GSC100C; Biopac, Goleta CA, USA). GSC is expressed as a change from baseline (Δ GSC), recorded during the 5 minute resting phase prior to heating. Cutaneous vascular conductance (CVC) was estimated by measuring skin blood flow on each forearm using laser-Doppler velocimetry (ALF21; Advanced, Tokyo, Japan). CVC was calculated as a percentage of the baseline value recorded during the resting phase prior to heating. T_{es} , local T_{sk} , $SkBf$, SR and GSC were recorded every second by a data logger (MX100; Yokogawa, Tokyo, Japan) and 1 min averaged calculated. Heart rate and arterial blood pressure were continuously measured on the left middle finger using a Finometer (Finometer; Finapres Medical Systems, Amsterdam, The Netherlands); mean arterial pressure (MAP) was subsequently calculated and averaged over 5-min periods.

The number of HASG was determined using the starch-iodine technique (Inoue 1996). Briefly, the residual sweat on the skin was first wiped away, followed by placing a small amount of iodine onto the skin with a cotton guaze. Excess iodine was removed by blotting the area with tissue paper. Starch paper attached to a small wooden block was then held in place over the measurement area for approximately 3 sec. The iodine was transferred from the HASG to the paper as indicated by a small dot. The same investigator counted the dots within a defined area (1cm^2). The sweat gland output per gland (SGO) at the respective site was calculated by dividing the SR (averaged from the last 5 minutes of the experiment) by the number of HASG.

Salivary and urine aldosterone was collected using SalivettesTM (Sarstedt, Newton, NC, USA) whereby a plain cotton swab was inserted into the mouth and chewed for 60 sec. The cotton swab was then returned into the SalivetteTM tube and spun at 4000RPM for 10 minutes. Samples were then frozen at -30°C until analysis. After thawing, salivary aldosterone (pg/ml) levels were quantified by competitive ELISA (LDN, GmbH & Co.KG, Germany). The sensitivity of the assay for aldosterone was 14pg/ml and the inter- and intra-assay coefficient of variation were between 3.9-7.5% and 9.4-9.7%, respectively.

Data analysis

As described in our previous experiments (Amano et al. 2016, 2017; Gerrett et al. 2018a), the maximum reabsorption rate of the sweat glands was obtained by plotting ΔGSC against ΔSR . By plotting this relationship, it is possible to identify three distinct phases; representing different stages of sweat production. In the first phase, there is an increase in ΔGSC but no change in ΔSR , which represents the isosmotic precursor sweat production in the proximal secretory coil. Such changes in ΔGSC and no changes in ΔSR are frequently utilised to identify pre-secretory sweat gland activity (Thomas and Korr 1957; Darrow

1964; Machado-Moreira et al. 2009; Gerrett et al. 2018b). In the second phase, an increased ΔSR without an increase in ΔGSC can be observed. As ΔGSC is influenced by both the amount of sweat produced as well as the electrolyte concentration the fact that ΔSR increases but there is no change in ΔGSC represents reabsorption of sweated ions in the sweat duct. Once the rate of sweat ion secretion exceeds its reabsorption limit in the duct then the third phase occurs where there is a proportional increase in ΔGSC with increasing ΔSR . The point at which the 2nd and 3rd phase intersect is used to identify the maximum rate of sweat glands ion reabsorption. In the present study, the thresholds were determined using segmented regression analysis on GraphPad Prism (version 7) software.

To investigate the effect of local T_{sk} on local maximum ion reabsorption, local SR, local ΔGSC and local CVC, a one-way repeated measure ANOVA with condition (Warm, Neutral and Cool) was carried out. The relation between temperature and ion reabsorption rates were analysed with correlation analysis; using normalized data to reduced individual effects. Correlation coefficients were considered as strong (≥ 0.60), moderate (0.40–0.59), and weak (0.20–0.39) (Cohen 1977).

A two-way ANOVA was used to assess the remaining thermophysiological parameters (T_{es} , T_b and mean and local T_{sk}) and cardiovascular responses (CVC, a one-way repeated measure ANOVA) to determine any differences between conditions (Neutral-Cool vs. Neutral-Warm) and over time. When any significant effects were observed, post hoc comparison using the Bonferroni test were carried out. All data were checked for sphericity and normality (Shapiro-Wilk test). As ANOVA's are fairly robust to violations of normality, if the data were approximately normal then the data was assessed with parametric data analysis. If the data violated this assumption substantially then a Friedman's test was performed (comparing Neutral, Cool and Warm data); this was the case for the following sets of data: local forearm, chest and thigh T_{sk} and SR. All data were

analysed using GraphPad Prism (version 7). Effect sizes (Cohen's d) were calculated for the local maximum ion reabsorption rates with the following criteria; an effect size of <0.20 is classified as 'trivial', $0.21-0.49$ as 'small', $0.50-0.79$ as 'moderate' and >0.80 as a 'large' effect. Values are means and standard deviations ($\pm$ SD) and statistical significance was set at $p<0.05$.

Results

Local skin temperature

The local forearms T_{sk} are illustrated in Figure 1. Forearm T_{sk} during Warm ($36.4 \pm 0.4^{\circ}\text{C}$) were higher ($p<0.05$) than Neutral ($33.5 \pm 0.7^{\circ}\text{C}$) and the Cool ($30.5 \pm 1.0^{\circ}\text{C}$) conditions ($p<0.05$). The latter was also lower than both the Neutral conditions ($p<0.05$). The local T_{sk} on the Neutral arm was not affected by the temperature on the experimental arm (Neutral-Cool; $33.6 \pm 0.7^{\circ}\text{C}$ and Neutral-Warm; $33.7 \pm 0.7^{\circ}\text{C}$, $p>0.05$). Forearm T_{sk} increased over time in all conditions ($p<0.05$) and was significantly higher than baseline after 40, 25 and 30 mins until the end of the experiment for Warm, both Neutral and Cool conditions, respectively. The increases from baseline to the end of passive heating were however similar between conditions (Warm; $+0.9 \pm 0.4^{\circ}\text{C}$, Neutral $+1.2 \pm 0.5^{\circ}\text{C}$, Cool; $+1.7 \pm 0.8^{\circ}\text{C}$, no interaction effect, $p>0.05$).

Maximum ion reabsorption

The maximum ion reabsorption thresholds, as indicated by the Δ SR threshold for an increasing Δ GSC, are illustrated in Figure 2. There is a significant pattern for a temperature dependent response for maximum ion reabsorption threshold (Warm; $0.28 \pm 0.14 \text{ mg/cm}^2/\text{min}$, Neutral; $0.22 \pm 0.12 \text{ mg/cm}^2/\text{min}$, Cool; $0.18 \pm 0.10 \text{ mg/cm}^2/\text{min}$, $p<0.05$), with differences noted between Warm and Cool only ($d = 0.88$, $p=0.024$). There

were small effect sizes for the maximum ion reabsorptions between Neutral and Warm ($d = 0.44, p > 0.05$) and Neutral and Cool ($d = 0.36, p > 0.05$). A weak, significant relation exists between local T_{sk} and the ΔSR threshold for maximum ion reabsorption ($r^2 = 0.36, p < 0.05$)

Thermo-physiological measurements

T_{es} , T_b and mean T_{sk} are illustrated in Figure 3A-C. T_{es} were similar between conditions (Neutral-Warm; $37.1 \pm 0.3^\circ\text{C}$ and Neutral-Cool $37.2 \pm 0.3^\circ\text{C}$, $p > 0.05$) and both gradually increased over time ($p < 0.05$), although the increase was similar between conditions (ΔT_{es} : Neutral-Warm; $0.92 \pm 0.2^\circ\text{C}$ and Neutral-Cool $0.77 \pm 0.1^\circ\text{C}$ $p > 0.05$). T_b were similar between conditions (Neutral-Warm; $36.9 \pm 0.3^\circ\text{C}$ and Neutral-Cool $36.8 \pm 0.3^\circ\text{C}$, $p > 0.05$) and both gradually increased over time ($p < 0.05$), although the increases were similar between conditions (ΔT_{es} : Neutral-Warm; $1.4 \pm 0.03^\circ\text{C}$ and Neutral-Cool $1.3 \pm 0.02^\circ\text{C}$ $p > 0.05$).

Mean T_{sk} was higher during the Neutral-Warm ($35.7 \pm 0.6^\circ\text{C}$) compared to Neutral-Cool ($34.9 \pm 1.0^\circ\text{C}$) conditions ($p < 0.05$) and did increase over time ($p < 0.05$), although the increase was similar between conditions ($\Delta \text{mean } T_{sk}$: Neutral-Warm; $3.3 \pm 0.5^\circ\text{C}$ and Neutral-Cool $3.2 \pm 0.3^\circ\text{C}$; no interaction effect, $p > 0.05$). Local T_{sk} of the chest, thigh and calf were similar between conditions (Neutral-Warm; $35.4 \pm 5.6^\circ\text{C}$, $33.4 \pm 5.2^\circ\text{C}$, $42.0 \pm 7.0^\circ\text{C}$ and Neutral-Cool: $35.3 \pm 5.5^\circ\text{C}$, $33.2 \pm 5.3^\circ\text{C}$, $41.9 \pm 7.0^\circ\text{C}$, $p < 0.05$, respectively). They all increased over time ($p < 0.05$) although the increase was similar between conditions (no interaction effect, $p > 0.05$).

Sweating responses

SR and ΔGSC measured at the forearms whilst local forearm T_{sk} were maintained at a Warm (36°C), Neutral (33°C) or Cool (30°C) temperatures are illustrated in Figure 4.

SR was higher during both Warm (0.43 ± 0.26 mg/cm²/min) and the Neutral conditions (Neutral-Warm; 0.40 ± 0.3 mg/cm²/min, and Neutral-Cool 0.40 ± 0.2 mg/cm²/min) than Cool (0.35 ± 0.2 mg/cm²/min). SR increased over time in all conditions ($p < 0.05$) and was significantly higher than baseline from 30 mins until the end of passive heating. The increases from baseline to the end of passive heating were similar between conditions (Warm; $+0.62 \pm 0.24$ mg/cm²/min, Neutral $+0.61 \pm 0.23$ mg/cm²/min, Cool; $+0.54 \pm 0.15$ mg/cm²/min, no interaction effect, $p > 0.05$).

Δ GSC was not significantly different between Warm (10.3 ± 11.5 μ S), Neutral (10.1 ± 9.8 μ S), or Cool (8.6 ± 11.3 μ S) conditions ($p > 0.05$). Δ GSC increased from baseline after 10 mins for all conditions ($p < 0.05$). There was no interaction effect as the increases from baseline to the end of passive heating were similar between conditions (Warm; 18.1 ± 14.6 μ S, Neutral; 19.4 ± 12.8 μ S, Cool; 19.35 ± 14.1 μ S, $p > 0.05$).

The HASG was similar between conditions: Warm; 107 ± 20 gland/cm², Neutral; 120 ± 24 gland/cm², Cool; 115 ± 18 gland/cm² ($p > 0.05$). The SGO was similar between conditions: Warm; 6.9 ± 2.7 μ g/gland/min, Neutral; 5.6 ± 1.7 μ g/gland/min, Cool; 5.3 ± 1.1 μ g/gland/min, $p > 0.05$).

The T_{es} threshold for SR was not significantly different between conditions (Warm; $36.72 \pm 0.4^\circ\text{C}$, $p > 0.05$, Neutral; $36.92 \pm 0.4^\circ\text{C}$, Cool; $37.08 \pm 0.3^\circ\text{C}$). The slope was also not significantly different (Warm; 0.74 ± 0.4 , Neutral; 1.00 ± 0.3 , Cool; 1.13 ± 0.5 , $p > 0.05$).

Urine and Salivary Aldosterone

Urine aldosterone concentrations were similar between conditions (Neutral-Warm; 5.03 ± 2.6 μ g/L and Neutral-Cool; 5.37 ± 2.5 μ g/L, $p > 0.05$). The urine specific gravity was also similar between conditions (Neutral-Warm; 1.021 ± 0.01 and Neutral-Cool; 1.021 ± 0.01 , $p > 0.05$). Salivary Aldosterone samples were similar between conditions and over

time (pre vs. post); there was also no interaction effect (Neutral-Warm-pre; 122.2 ± 40.5 pg/ml, Neutral-Warm-post; 116.1 ± 43.0 pg/ml, and Neutral-Cool-pre; 113.7 ± 28.9 pg/ml and Neutral-Cool post; 115.8 ± 41.7 pg/ml, $p > 0.05$).

Cardiovascular measurements

HR, MAP, CVC are presented in Table 1. HR's were similar between conditions (Neutral-Warm; 78.0 ± 8 bpm and Neutral-Cool 79.0 ± 9 bpm, $p > 0.05$) and both gradually increased over time ($p < 0.05$) and the increase was similar between conditions (Δ HR: Neutral-Warm; 30.0 ± 2 bpm and Neutral-Cool 31.2 ± 2 bpm, $p > 0.05$). MAP were similar between conditions (Neutral-Warm; 99.2 ± 2 mmHg and Neutral-Cool 99.1 ± 2 mmHg, $p > 0.05$) and did not increase over time in either condition ($p > 0.05$).

Discussion

The aim of the present study was to investigate the influence of local T_{sk} on the eccrine sweat glands maximum ion reabsorption rates during passive heating. We hypothesised that the sweat glands maximum ion reabsorption rates would occur at a higher sweat rate (SR) with a higher local T_{sk} (36°C) compared to neutral (33°C) and cool T_{sk} (30°C). The data indicates that there is a temperature dependent response, which was most prominent when local T_{sk} differed by 6°C (30°C vs. 36°C) whilst not when local T_{sk} differed by only 3°C (30°C vs. 33°C and 33°C vs. 36°C). In practical terms, a higher Δ SR for maximum ion reabsorption means that a more dilute sweat would be secreted onto the skin surface for a given SR, which may be advantageous when sweat rate is high to prevent the excess loss of ions. By selecting a passive heating protocol, we ensured that the values were obtained in controlled conditions to eliminate any potential effect from non-thermal control mechanisms, such as aldosterone, and other potential thermal controllers such as core and

body temperature. Mean T_{sk} was significantly different by only approximately 0.6°C between conditions, which is likely due to differences in local T_{sk} manipulations as all other measured local skin site temperatures were similar between conditions.

Thermal mechanism

We hypothesized that the maximum sweat gland ion reabsorption rate would be affected by local T_{sk} as previous studies, in vivo and in vitro, have reported a temperature dependency of Na^{+} channels excitability and ion reabsorption regulation (Ruff 1999; Chraïbi and Horisberger 2003; Shamsuddin et al. 2005a). The reabsorption of NaCl is primarily driven by the movement of Na^{+} down a steep concentration gradient that is generated by the $\text{Na}^{+}/\text{K}^{+}$ pump via ENaC (Bovell 2015). In vitro studies utilising a wide temperature range (19°C and 37°C) have demonstrated the temperature dependency of Na^{+} channels excitability; where 30% and 93% of the channels were excitable at these respective temperatures (Ruff 1999). However, the percentages of excitable channels became less prominent when comparing 31°C and 37°C (85% vs. 93%, respectively), which are closer to the physiological temperatures experienced whilst sweating. Chraïbi and Horisberger (2003) more recently showed that ENaC open probability was greater at lower (15°C) compared to higher temperatures ($24\text{-}30^{\circ}\text{C}$). Indeed other studies showing temperature dependency of ENaC , and cystic fibrosis transmembrane channels (CFTR) that are responsible for the reabsorption of Cl^{-} and interacts with ENaC (Reddy and Quinton 2003), show the effects of temperature are more prominent outside the physiological ranges examined in the present study. Our data indicates that when T_{sk} is within appropriate physiological ranges of $30\text{-}33\text{-}36^{\circ}\text{C}$, a T_{sk} difference of $\sim 3^{\circ}\text{C}$ is an insufficient temperature stimulus to affect the maximum rate of ion reabsorption but differences of $\sim 6^{\circ}\text{C}$ may elicit changes of approximately $0.1\text{mg}/\text{cm}^2/\text{min}$. Whilst the inter and intra-individual variability

is high for sweating responses, the ventilated sweat capsule technique is highly reliable and accurate and thus we are confident that our method is sensitive to detect these differences. Whilst GSC is more variable, the values themselves are less meaningful but the pattern of the response in relation to SR is important for detecting the sweat glands ion reabsorption rates.

Contrasting to our findings, Shamsuddin et al. (2005a) reported differences in ion reabsorption rates when mean T_{sk} was clamped at $\sim 31^{\circ}\text{C}$ and $\sim 28^{\circ}\text{C}$, a differences of 3°C , during dynamic exercise. These findings, alongside the in vitro studies mentioned earlier may indicate that cooler temperatures ($<30^{\circ}\text{C}$) have stronger effects on ion regulation. Alternatively, smaller differences in T_{sk} may affect ion reabsorption rates during dynamic exercise when hormonal mediators are likely in effect.

Core temperature (T_c), as indicated by T_{es} , in our present study and by Shamsuddin et al. (2005a) were not significantly different between the two conditions; thereby eliminating the role of T_{es} as a controlling mechanism. However, that is not to say that T_c per se will not influence the sweat glands maximum ion reabsorption rates in other conditions outside the realms of the studies under discussion. Temperature stimulation has been deemed an important regulatory mechanism for sudomotor activity. We reported no significant differences between conditions in the T_{es} threshold for the onset of neither sweating, nor the slope, or the SGO in this study. Indeed, the relative importance of T_c on SR compared to mean T_{sk} is well known (Nadel et al. 1971) and both Na^+ secretion and Na^+ reabsorption increase linearly with increasing SR (Buono et al. 2008). As SR increases it has been suggested that there becomes insufficient time for sweat ion reabsorption to occur and hence a maximum reabsorption rate is reached. The SR for maximum ion reabsorption occurred between 0.18 and 0.28 $\text{mg}/\text{cm}^2/\text{min}$ across the 3 temperature ranges,

which typically occurred within 15 minutes of passive heating. As can be seen in left panel of Figure 4, the SR response over time is similar between conditions, as also confirmed by a non-significant interaction effect (condition x time). It seems therefore that if any thermal controls exist over the sweat glands maximum ion reabsorption rates, it requires a stronger thermal input (e.g. from mean T_{sk} and/or T_c) than from small changes ($\leq 3^\circ\text{C}$) in local T_{sk} . It would be interesting to determine the minimum change in local T_{sk} that would affect the sweat glands maximum ion reabsorption rate. The role of T_c on SR and ion reabsorption seems likely, but certainly warrants clarification, as to does the role of mean and local T_{sk} under varying T_c responses.

Previous studies have reported higher reabsorption rates on the torso compared to the extremities but have been unable to determine any contributing mechanism as regional differences in local T_{sk} existed (Inoue et al. 1998; Amano et al. 2017). The findings of the present study provide important information to help elucidate why these regional differences occur and variations in local T_{sk} can be ruled out, as regional differences were less than $\sim 2^\circ\text{C}$. Instead, we hypothesize that structural differences in the sweat glands across the body may account for the regional differences reported in aforementioned studies.

Non-thermal mechanism

Regulation of ion loss is predominated in the literature by renal function despite potential large fluid and ion losses from eccrine sweat glands during exercise and/or heat exposure. Although structurally and functionally different to the kidneys, the influence of various water regulating hormones (arginine vasopressin, aldosterone and atrial natriuretic peptide) on sweat output have been investigated (Kirby and Convertino 1986; Hew-Butler et al. 2010, 2014). The precise mechanism for the regulation of the ion reabsorption is unknown but it is hypothesized that aldosterone, an important hormone in renal sodium

regulation, plays a role. Acting on mineralocorticoid receptors, aldosterone increases intracellular calcium, which regulates epithelial Na^+ and K^+ channels, reportedly by increasing either the permeability of the membrane to Na^+ , increasing active transport of Na^+ out of the cell and/or increasing the energy available to the Na^+/K^+ pump (Hegarty and Harvey 1998; Harvey and Higgins 2000).

By using a passive heating protocol, our study aimed to reduce non-thermal mediators, such as water-regulatory hormones (e.g. aldosterone, vasopressin and plasma renin activity) that are released during exercise (Convertino et al. 1983; Freund et al. 1991; Yoshida et al. 2006). We confirmed similar aldosterone concentrations between our two conditions, both before and after passive heating. In addition, hydration statuses were similar between both conditions so to rule out any potential effects of hydration on circulating hormones. Previous research by Shamsuddin et al. (2005a) reported a role of mean T_{sk} on ion reabsorption, but this occurred during dynamic exercise (cycling at 60% $\dot{V}\text{O}_{2\text{max}}$) where aldosterone concentrations would have been elevated, as demonstrated in our previous research comparing passive heating and cycling at 60% $\dot{V}\text{O}_{2\text{max}}$ (Gerrett et al. 2018a). Shamsuddin et al. (2005a), speculated that higher T_{sk} might enhance the responsiveness of the sweat glands to a given aldosterone concentration. In the present study however, aldosterone concentrations remained unchanged but local T_{sk} differed and we hypothesize that small differences ($\leq 3^\circ\text{C}$) in T_{sk} may only affect ion reabsorption rates when aldosterone (or other water regulatory hormones) are elevated. Further support is provided from our previous research where higher maximum ion reabsorption rates were reported during moderate intensity exercise (supine cycling at 60% $\dot{V}\text{O}_{2\text{max}}$) compared to passive heating (lower leg submersion 43°C water), despite a lower mean T_{sk} . We attributed those differences partially to the elevated salivary aldosterone concentrations during exercise compared to passive heating. To date no studies have determined the effect of

527 varying T_{sk} and/or T_c on aldosterone concentrations during exercise but it certainly warrants
528 investigation.

529 *Perspective*

530 Sodium chloride plays an important role in the formation of sweat within the
531 secretory coil. Why we do not, or cannot, reabsorb all the sodium chloride ions in the
532 reabsorptive duct is not clear; it may play a role in the evaporative potential of sweat on the
533 skin surface, it may aid skin barrier function and protection. Yet there is an adaptive
534 response as seen with heat acclimation (Buono et al. 2008; Amano et al. 2016). We know
535 from cases of cystic fibrosis that preventing an excess loss of these ions is an important
536 regulatory mechanism yet, it is poorly understood and often overlooked in favour of its
537 thermoregulatory role. The regulation of ion reabsorption is a fundamental research
538 question that requires further consideration.

539 Furthermore, in recent years there has been a drive to produce non-invasive
540 techniques to inform us about the human condition, using sweat as the medium. As a result,
541 the development of sensors to continuously measure sweat content is a rapidly growing
542 field in biomedical engineering but our knowledge of sweat ion regulation is limited and
543 this fundamental study adds to our knowledge. We provide further insight into the
544 methodological considerations for future studies in this area. In particular, our research
545 group has been accumulating consistent evidence of regional differences in ion
546 reabsorption across the body and the current study suggest these regional differences may
547 not have been due to differences in T_{sk} but rather structural or regulatory differences at the
548 level of the sweat gland across the body. If T_{sk} is expected to vary considerably then
549 knowing the T_{sk} is important. Consistent sensor placement in the same locations will allow
550 for better comparisons between conditions and from different studies.

Conclusion

Local T_{sk} within physiological ranges of 30-33-36°C only influenced the sweat glands maximum ion reabsorption rates during a passive heating protocol when local T_{sk} differed by 6°C (30°C vs. 36°C), whilst no differences were observed when local T_{sk} differed by 3°C (30°C vs. 33°C and 33°C vs. 36°C). These findings were observed when all other potential thermal and non-thermal controlling mechanisms were similar between conditions. Information from the literature and data from our study indicate that thermal controllers may exist but most probably from stronger thermal stimulus, such as mean T_{sk} and/or T_c , compared to smaller changes ($\leq 3^\circ\text{C}$) in local T_{sk} . We provide important insights for previous studies that have reported regional differences in maximum ion reabsorption rates but have been unable to confirm such differences in the presence of differing local T_{sk} . The data provides useful information for furthering our understanding of sweat gland ion reabsorption and potential controlling mechanisms. The application of which may be useful in the fields of thermoregulation, hypo/hyponatremia, dermatology and biosensor technology research.

References

- Amano T, Gerrett N, Inoue Y, et al (2016) Determination of the maximum rate of eccrine sweat glands' ion reabsorption using the galvanic skin conductance to local sweat rate relationship. *Eur J Appl Physiol* 116:281–290
- Amano T, Hirose M, Konishi K, et al (2017) Maximum rate of sweat ions reabsorption during exercise with regional differences, sex, and exercise training. *Eur J Appl Physiol* 117:. doi: 10.1007/s00421-017-3619-8
- Bovell D (2015) The human eccrine sweat gland: Structure, function and disorders. *J Local Glob Heal Sci* 2015:5. doi: 10.5339/jlghs.2015.5

576 Bulmer MG, Forwell GD (1956) The concentration of sodium in thermal sweat. *J Physiol*
 577 32:5–122

578 Buono MJ, Claros R, Deboer T, Wong J (2008) Na⁺ secretion rate increases
 579 proportionally more than the Na⁺ reabsorption rate with increases in sweat rate. *J*
 580 *Appl Physiol* 105:1044–1048. doi: 10.1152/jappphysiol.90503.2008.

581 Buono MJ, Jechort A, Marques R, et al (2007) Comparison of infrared versus contact
 582 thermometry for measuring skin temperature during exercise in the heat. *Physiol*
 583 *Meas* 28:855–859. doi: 10.1088/0967-3334/28/8/008

584 Buono MJ, Tabor B, White A (2011) Localized β -adrenergic receptor blockade does not
 585 affect sweating during exercise. *Am J Physiol Integr Comp Physiol* 300:R1148–
 586 R1151. doi: 10.1152/ajpregu.00228.2010

587 Chraïbi A, Horisberger J-D (2003) Dual effect of temperature on the human epithelial Na
 588 ⁺ channel. *Pflugers Arch -Eur J Physiol* 447:316–320. doi: 10.1007/s00424-003-
 589 1178-9

590 Cohen J (1977) *Statistical Power Analysis for the Behavioral Sciences*. Lawrence
 591 Erlbaum Associates, Hillsdale, New Jersey

592 Convertino V a, Keil LC, Greenleaf JE (1983) Plasma volume, renin, and vasopressin
 593 responses to graded exercise after training. *J Appl Physiol* 54:508–514

594 Darrow CW (1964) The rationale for treating the change in galvanic skin response as a
 595 change in conductance. *Psychophysiology* 1:31–38

596 Freund BJ, Shizuru EM, Hashiro GM, et al (1991) Hormonal, electrolyte, and renal
 597 responses to exercise are intensity dependent. *J Appl Physiol* 70:900–906

598 Gagge AP, Nishi Y (2011) Heat exchange between human skin surface and thermal
 599 environment. In: *Comprehensive Physiology*. John Wiley & Sons, Inc., Hoboken,
 600 NJ, USA, pp 69–92

601 Gao W, Brooks GA, Klonoff DC (2018) Wearable physiological systems and
602 technologies for metabolic monitoring. *J Appl Physiol* 1:548–556

603 Gerrett N, Amano T, Inoue Y, et al (2018a) The effects of exercise and passive heating on
604 the sweat glands ion reabsorption rates. *Physiol Rep* 6:. doi: 10.14814/phy2.13619

605 Gerrett NM, Griggs KE, Redortier B, et al (2018b) Sweat-from gland to skin surface -
606 production, transport and skin absorption. *J Appl Physiol* japplphysiol.00872.2017.
607 doi: 10.1152/japplphysiol.00872.2017

608 Harvey BJ, Higgins M (2000) Nongenomic effects of aldosterone on Ca²⁺ in M-1
609 cortical collecting duct cells. *Kidney Int* 57:1395–1403. doi: 10.1046/j.1523-
610 1755.2000.00981.x

611 Hegarty J, Harvey BJ (1998) Aldosterone increases intracellular calcium in cultured
612 human sweat gland epithelial cells by a non-genomic mechanism of action. *J Physiol*
613 511P

614 Hew-Butler T, Hummel J, Rider BC, Verbalis JG (2014) Characterization of the effects
615 of the vasopressin V2 receptor on sweating, fluid balance, and performance during
616 exercise. *Am J Physiol Regul Integr Comp Physiol* 307:R366-75. doi:
617 10.1152/ajpregu.00120.2014

618 Hew-Butler T, Noakes TD, Soldin SJ, Verbalis JG (2010) Acute changes in arginine
619 vasopressin, sweat, urine and serum sodium concentrations in exercising humans:
620 does a coordinated homeostatic relationship exist? *Br J Sports Med* 44:1710–715.
621 doi: 10.1136/bjsm.2008.051771

622 Inoue Y (1996) Longitudinal effects of age on heat-activated sweat gland density and
623 output in healthy active older men. *Eur J Appl Physiol Occup Physiol* 74:72–77. doi:
624 10.1007/BF00376497

625 Inoue Y, Nakao M, Ishizashi H, et al (1998) Regional differences in the Na +

626 reabsorption of sweat glands. *Appl Hum Sci* 17:219–221

627 Johnson RE, Pitts GC, Consolazio FC (1944) Factors influencing chloride concentration
 628 in human sweat 1. *Am J Physiol* 141:575–589. doi: 10.220.33.3

629 Kenefick RW, Cheuvront SN, Elliott LD, et al (2012) Biological and analytical variation
 630 of the human sweating response: implications for study design and analysis. *Am J*
 631 *Physiol Regul Integr Comp Physiol* 302:252–258. doi:
 632 10.1152/ajpregu.00456.2011.-Appropriate

633 Kirby CR, Convertino VA (1986) Plasma aldosterone and sweat sodium concentrations
 634 after exercise and heat acclimation. *J Appl Physiol* 61:967–70

635 Machado-Moreira CA, Edkins E, Iabushita AS, et al (2009) Sweat gland recruitment
 636 following thermal and psychological stimuli. In: Castellini JW (ed) 13th
 637 International Conference of Environmental Ergonomics

638 Mekjavic IB, Rempel ME (1990) Determination of esophageal probe insertion length
 639 based on standing and sitting height. *J Appl Physiol* 69:376–379. doi:
 640 10.1152/jappl.1990.69.1.376

641 Nadel ER, Bullard RW, Stolwijk JA (1971) Importance of skin temperature in the
 642 regulation of sweating. *J Appl Physiol* 31:80–87

643 Reddy M, Quinton P (2003) Functional interaction of CFTR and ENaC in sweat glands.
 644 *Pflügers Arch - Eur J Physiol* 445:499–503. doi: 10.1007/s00424-002-0959-x

645 Robinson S, Gerking SD, Tuerell ES, Kincaid RK (1985) Effects of skin temperature of
 646 salt concentration of sweat. *J Appl Physiol* 2:654–62

647 Ruff RL (1999) Effects of temperature on slow and fast inactivation of rat skeletal muscle
 648 Na(+) channels. *Am J Physiol* 277:C937-47

649 Sato K (1977) The physiology, pharmacology, and biochemistry of the eccrine sweat
 650 gland. Springer Berlin Heidelberg, pp 51–131

651 Shamsuddin AKM, Kuwahara T, Oue A, et al (2005a) Effect of skin temperature on the
652 ion reabsorption capacity of sweat glands during exercise in humans. *Eur J Appl*
653 *Physiol* 94:442–447. doi: 10.1007/s00421-005-1354-z

654 Shamsuddin AKM, Yanagimoto S, Kuwahara T, et al (2005b) Changes in the index of
655 sweat ion concentration with increasing sweat during passive heat stress in humans.
656 *Eur J Appl Physiol* 94:292–297. doi: 10.1007/s00421-005-1314-7

657 Smith CJ, Havenith G (2012) Body mapping of sweating patterns in athletes: A sex
658 comparison. *Med Sci Sports Exerc* 44:2350–2361. doi:
659 10.1249/MSS.0b013e318267b0c4

660 Stolwijk JA, Hardy JD (1966) Partitional calorimetric studies of responses of man to
661 thermal transients. *J Appl Physiol* 21:967–977

662 Thomas PE, Korr IM (1957) Relationship between sweat gland activity and electrical
663 resistance of the skin. *J Appl Physiol* 10:505–510

664 Yoshida T, Shin-ya H, Nakai S, et al (2006) Genomic and non-genomic effects of
665 aldosterone on the individual variation of the sweat Na⁺ concentration during
666 exercise in trained athletes. *Eur J Appl Physiol* 98:466–471. doi: 10.1007/s00421-
667 006-0295-5

668

669

670 **Figures and Tables**

671 **Table 1:** Cardiovascular responses measured continuously and averaged over 5 minutes during the passive heating protocol. There were no
672 differences in heart rate (HR) or mean arterial pressure (MAP) between the Neutral-Warm and Neutral-Cool conditions ($p>0.05$). There were no
673 differences in CVC between both Neutral (33°C) conditions (Neutral-Warm and Neutral-Cool), Cool (30°C), or Warm (36°C) conditions ($p>0.05$).
674 All variables increased over time but there was no interaction effect ($p>0.05$). Values are based on 5-min averages and expressed as mean $\pm$ SD
675 for 10 participants.
676

		BL	5	10	15	20	25	30	35	40	45
HR (bpm)	Neutral-Warm	61.3 ± 8.7	65.5 ± 6.9	67.9 ± 8.4	72.7 ± 8.8	77.0 ± 7.1	78.2 ± 9.4	79.2 ± 10.8	82.5 ± 11.3	86.9 ± 10.7	89.9 ± 10.6
	Neutral-Cool	61.9 ± 8.1	65.6 ± 8.7	67.0 ± 9.9	72.2 ± 7.1	78.4 ± 7.0	79.7 ± 7.7	81.1 ± 8.1	84.3 ± 8.0	88.6 ± 8.1	91.4 ± 8.9
MAP (mmHg)	Neutral-Warm	93.9 ± 6.8	98.9 ± 10.6	99.1 ± 8.8	99.8 ± 9.6	101.5 ± 9.6	100.4 ± 9.9	97.4 ± 6.8	99.7 ± 8.5	99.4 ± 6.4	97.3 ± 6.8
	Neutral-Cool	95.1 ± 7.1	98.3 ± 7.6	99.7 ± 9.7	99.2 ± 8.5	96.8 ± 8.7	97.8 ± 8.7	98.9 ± 8.9	99.9 ± 8.5	99.5 ± 8.5	101.5 ± 7.4
CVC (%)	Warm	100.0 ± 0.0	143.0 ± 97.2	205.0 ± 158.2	351.5 ± 291.0	480.2 ± 310.8	578.0 ± 304.9	620.5 ± 301.7	643.4 ± 288.7	648.8 ± 298.5	655.0 ± 310.1
	Neutral	100.0 ± 0.0	120.5 ± 42.7	159.8 ± 59.9	252.4 ± 129.5	384.7 $\pm 163.$	460.8 ± 173.0	524.7 ± 170.3	601.7 ± 200.7	645.6 ± 212.3	663.6 ± 232.8
	Cool	100.0 ± 0.0	101.4 ± 19.1	120.4 ± 35.0	165.4 ± 51.9	224.9 ± 79.4	281.6 ± 74.5	357.1 ± 105.6	464.5 ± 114.0	590.9 ± 180.9	645.5 ± 215.7

677

Warm
 Neutral
 Cool

Figure 1: Local forearm skin temperature (T_{sk}) measured whilst manipulating local T_{sk} to 36°C (Warm), 33°C (Neutral) and 30°C (Cool) during a 45-minute passive heating protocol. Forearm T_{sk} was higher during Warm compared to both Neutral and Cool condition and Neutral was also higher than the Cool condition ($p < 0.05$). Forearm T_{sk} increased over time in all conditions ($p < 0.05$) and * indicates when forearm T_{sk} was higher than baseline from each condition. The increases were similar between conditions; hence there was no interaction effect between condition and time ($p > 0.05$). Values are based on 1-min averages and expressed as mean $\pm$ SD for 10 participants. To aid clarity error bars are provided at 5-min intervals.

Figure 2: The relation between local T_{sk} and the ΔSR threshold for an increasing ΔGSC . The solid line and black cycle represent the mean data (n=10), whilst the dashed line and empty circles are the individual responses. Warm was significantly higher than Cool (* indicates $p < 0.05$)

Figure 3: Esophageal temperature (T_{es}) (A), body temperature (T_b) (B) and mean skin temperature (T_{sk}) (C) during the 45-minute passive heating protocol when local forearms skin temperature were stimulated to be Neutral and Warm or Neutral and Cool. There were

no significant differences found between conditions for T_{es} or T_b and whilst they both increased over time there was no interaction effect ($p>0.05$). Mean T_{sk} was higher during the Neutral-Warm compared to Neutral-Cool conditions ($*p<0.05$) likely due to differences in local forearm T_{sk} as all other skin sites were not different ($p>0.05$). Mean T_{sk} increased over time ($p<0.05$) but this was similar between conditions ($p>0.05$). Values based on 1-min averages and are expressed as mean $\pm$ SD for 10 participants. To aid clarity error bars are provided at 5-min intervals.

Figure 4: Local sweat rate (SR) (left) and ΔGSC (right) measured whilst stimulating local T_{sk} at 36°C (Warm), 33°C (Neutral) and 30°C (Cool) during a 45-minute passive heating protocol. SR was significantly different between all conditions ($p<0.05$) and whilst SR increases over time the increase was similar between conditions (no interaction effect, $p>0.05$). There were no differences in ΔGSC between conditions ($p>0.05$). Whilst ΔGSC increased over time the increase was similar between conditions (no interaction effect, $p>0.05$). Values are based on 1-min averages and expressed as mean $\pm$ SD for 10 participants. To aid clarity error bars are provided at 5-min intervals.