This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

The interactive effect of cooling and hypoxia on forearm fatigue development

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1007/s00421-015-3181-1>

PUBLISHER

© Springer-Verlag

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Lloyd, Alex, S.G. Hodder, and George Havenith. 2019. "The Interactive Effect of Cooling and Hypoxia on Forearm Fatigue Development". figshare. <https://hdl.handle.net/2134/17889>.

Title:

The interactive effect of cooling and hypoxia on forearm fatigue development

(Accepted for publication in European Journal of Applied Physiology 30/04/15)

Authors:

Alex Lloyd

Simon Hodder

George Havenith

Affiliation:

Environmental Ergonomics Research Centre, Loughborough University, Loughborough, UK

Running Head:

Fatigue in combined hypoxia and cold

Corresponding Author:

George Havenith

Environmental Ergonomics Research Centre,

James France Bldg,

Loughborough,

Leicestershire

UK

LE11 2TU

Email: G.Havenith@lboro.ac.uk

ABSTRACT

PURPOSE: To examine the effect of separate and combined exposure to hypoxia (normoxia ($F_{I}O_2 = 0.21$) vs. moderate altitude ($F_{I}O_2 = 0.13$)) and temperature (thermoneutral ($22^{\circ}C$) vs. cold ($5^{\circ}C$)) on muscle fatigue development in the forearm, after repeated low resistance contractions. **METHODS:** Eight males were exposed for 70-mins to four separate conditions in a balanced order. Conditions were normoxic-thermoneutral (N), hypoxic-thermoneutral, normoxic-cold and hypoxic-cold. After 15-min seated rest, participants carried out intermittent dynamic forearm exercise at 15% maximal isometric voluntary contraction (MVC) for eight consecutive, 5-min work bouts. Each bout was separated by 110-s rest during which MVC force was collected. **RESULTS:** When exposed to hypoxia and cold independently, the exercise protocol decreased MVC force of the finger flexors by 8.1 and 13.9% respectively, compared to thermoneutral normoxia. When hypoxia and cold were combined, the decrease in MVC force was 21.4% more than thermoneutral normoxia, reflecting an additive effect and no interaction. EMG relative to force produced during MVC, increased by 2 and 1.2 μV per kg (36 & 23% of N) for cold and hypoxia respectively. When the stressors were combined the effect was additive, increasing to 3.1 μV per kg (56% of N). **CONCLUSION:** When compared to exercise in thermoneutral normoxic conditions, both cold and hypoxia significantly reduce brief MVC force output. This effect appears to be of mechanical origin, not a failure in muscle fibre recruitment per se. Additionally, the reduction in force is greater when the stressors are combined, showing an additive effect.

INTRODUCTION

Passive cold exposure can reduce a muscles mechanical response (e.g. power) to a given electrophysiological excitation or descending voluntary drive (Feretti 1992; Oksa et al., 2002). This is widely attributed to reductions in muscle temperature (Bergh & Ekblom 1979a) which reduces contractile function due to slowed intramuscular energetics and peripheral nerve conduction velocities (Kossler et al. 1987; Bigland Ritchie et al. 1981; Bergh, 1980; Faulkner et al. 1990; Sweitzer & Moss 1990; Feretti 1992; De Ruiter & De Haan 2000; Allen et al 2008; Racinais & Oksa, 2010; Cahill et al. 2011; Cè et al. 2012). Several studies report that action potential propagation, ATP hydrolysis, Ca^{2+} handling and sensitively as well as cross-bridge force kinetics are adversely affected by lower tissue

temperatures (Kossler et al. 1987; Sweitzer & Moss 1990; Mucke & Heuer 1989; Feretti 1992; Oksa et al. 2002; C et al. 2012). However, the slowing of mechanical processes, as well as efferent and afferent nerve conduction, occur independently of exercise (present during passive cold exposure), and thus may even serve to attenuate metabolite production, and/ or increase central drive, during prolonged isometric contractions (Ray et al. 1997; Segal et al. 1986; De Ruiter & De Haan 2000; Todd et al. 2005; Allen et al. 2008; Cahill et al. 2011; Lloyd et al. 2014). Nevertheless, during dynamic exercise in cooled muscle (i.e. active cold exposure), significant increases in skeletal muscle fatigue are reported (Bergh and Ekblom 1979b; Bergh, 1980; Faulkner et al. 1990; Racinais & Oksa, 2010). This is predominantly due to co-activation of the agonist- antagonist pair (Oksa et al. 1997) resulting in higher workload for the agonist muscle (Oksa et al. 2002) thereby reducing aerobic- mechanical efficiency (McArdle et al, 1976). Furthermore, dynamic exercise in cold muscle is likely affected by reductions in muscle blood flow (Yanagisawa et al, 2007; Gregson et al, 2011), which may hinder oxygen delivery (Amann & Calbet, 2007) and diminish the removal metabolic by-products (Blomstrand et al. 1984).

Contrary to tissue cooling, passive exposure to hypoxia does not appear to affect maximal force generating capacity or action potential propagation (Perrey & Rupp 2009). However, increases in muscle fatigue during prolonged exercise in hypoxia have been observed during both whole-body (Amann & Calbet 2007) and repeated contractions of isolated muscle groups (Fulco 1994; 1996; Katayama et al. 2007; Perrey & Rupp 2009; Millet et al. 2008; 2012; Christian et al. 2014a). The rise in muscle fatigue during hypoxia can be largely attributed to a shift of the relative exercise intensity, higher muscle fibre recruitment, and thereby increased intramuscular metabolic disturbance (Edwards 1981; Fulco et al. 1996; Amann et al. 2006a; 2006b; 2007a; 2007b; Fulco et al. 1994, Katayama et al. 2007; Christian et al. 2014a). Specifically, the increase in inorganic phosphate, reactive oxygen species and hydrogen ion production and their interference with the contractile proteins and sarcoplasmic Ca^{2+} release mechanisms are thought to be a major factor behind the increase in muscle fatigue development (Haseler et al 1999; Hogan et al. 1999; Amann & Calbet 2007; Perrey & Rupp 2008). In hypoxia, evidence also suggests increased afferent feedback and decreased cerebral oxygenation can reduce voluntary drive to the muscle, exacerbating net fatigue (Amann & Dempsey 2007; Amann et al. 2006a; 2007b; Goodall et al. 2010; Millet, 2008; 2012). However, the relative contributions of afferent feedback and cerebral oxygenation, as well as the sense of effort, to changes in central drive during

fatiguing exercise in hypoxia remains subject to on-going investigations (Millet et al. 2008; 2012; Goodall et al. 2010; Amann et al., 2013; Christian et al. 2014a; Christian et al. 2014b).

While much research exists on these stressors separately, ascent to altitude often constitutes of exposure to both hypoxia and cold stress; however the interactive effects of these stressors in combination are not well understood (Tipton, 2012). Studies that have examined combined hypoxic-cold stress have focused largely on thermogenesis, skin blood flow and thermal sensitivity (Robinson & Haymes 1991; Johnston et al. 1996; Gautier et al. 1987, Wood 1991; Cipriano & Goldman 1975; Simmons et al. 2010; 2011), leaving fatigue development and human performance relatively unexamined (Tipton 2012). Given the potential for hypoxic-cold to severely compromise oxygen delivery to the active muscle - through simultaneous reductions in oxygen transport (muscle blood flow) and arterial oxygen content (hypoxemia) (Yanagisawa et al, 2007; Gregson et al, 2011; Amann & Calbet 2007) - as well as greatly increase metabolite production - through simultaneous rises in agonist-antagonist co-activation in the cold and type II recruitment in hypoxia (Edwards 1981; Fulco et al. 1996; Oksa et al., 2002; Amann et al. 2006a; 2006b; 2007a; 2007b; Fulco et al. 1994, Katayama et al. 2007; Christian et al. 2014a) - we sought to investigate the independent and combined (interactive) effects of hypoxia and cold on forearm fatigue.

To investigate the interaction between hypoxia and cold on fatigue development an additive effects model (standard ANOVA) was used. Using the additive model, stressor (in fact all variables) interactions are categorised as either synergistic or antagonistic (Folt et al. 1999). Significant interactions suggest the effect size of one variable has been reduced (antagonistic) or accentuated (synergistic) by the presence (or effect) of the other, whereas additive effects are seen during net stressor independence i.e. no interaction. Interactions are best illustrated using variable A and B:

Additive = A & B combined = A + B individually (1)

Synergistic = A & B combined > A + B individually (2)

Antagonistic = A & B combined < A + B individually (3)

Nullifying = A & B combined = A or B individually (4)

Multiplicative = A & B combined = A x B individually (5)

Importantly, nullifying interactions (Eq. 4) are strong antagonistic interactions in which the influence of one variable has been entirely abolished by the presence (or effect) of the other; while multiplicative effects describe strong synergistic interactions (Eq. 5).

To examine the interaction between hypoxia and cold on fatigue development, we quantified changes in the relationship between electromyogram (EMG) and maximal isometric voluntary contraction (MVC) force in response to low intensity, intermittent ‘gripping’ exercises. An isolated forearm model was used due to the importance of finger flexor function for climbing, mountaineering or those performing manual work at altitude, as well as due to the known exacerbation of fatigue during prolonged (> 4-min) low intensity (<30% MVC) isolated muscle exercise in both cold and hypoxic environments (e.g. Oksa et al. 2002; Perrey & Rupp 2009). It was hypothesised that 1) independent exposure to hypoxia or cold will induce a significant increase in post-exercise neuromuscular fatigue, compared to control conditions; and 2) during combined hypoxic-cold exposure, a synergistic interaction on fatigue will occur, with reductions in muscle blood flow (Yanagisawa et al, 2007; Gregson et al, 2011), and increased metabolite production (co-activation) during cold (Oksa et al., 2002) synergistically accentuated by the reductions in oxygen delivery and higher type II muscle fibre activation observed in hypoxia (e.g. Amann & Calbet 2007).

Keywords: cooling, hypothermia, high altitude, electromyography, combined stressors.

Abbreviations:

ANOVA: Analysis of Variance

AU: Arbitrary Units

C: Condition normoxic- cold

EMG: Electromyography

ED: extensor digitorum

FCR: flexor carpi radialis

FDS: flexor digitorum superficialis

FFT: Fast Fourier Transform

FI: Fatigue Index

F_IO₂: Fraction of Inspired Oxygen

H: Condition hypoxic- thermoneutrality

HC: Conditions hypoxic- cold

HR: Heart Rate

LDF: Laser Doppler flowmetry

MVC: Maximal Voluntary Contraction

N: Condition normoxic- thermoneutrality

RMS: Root Mean Square

RPE: Rate of Perceived Exertion

SpO₂: Peripheral Arterial Oxygen Saturation

T_{co}: Core temperature

T_a: Ambient temperature

T_{sk}: Skin Temperature

METHODS

Subjects

Eight healthy men volunteered as participants for this study. Their (mean $\pm$ SD) age was 21.9 ± 0.8 years and Body Mass Index was 23.5 ± 1.8 . Using the short International Physical Activity Questionnaire, a minimal activity level of 25 MET-hours per week was used when selecting volunteers. The average weekly exercise level was 41.5 ± 15.4 MET-hours per week. No participant was trained in a specific sport, but all participants were regularly participating in a range of physical activities, and thus appeared well accustomed to novel and strenuous exercise regimes. All participants were requested to abstain from caffeine, alcohol and exhaustive exercise 24 hours prior to the experiment.

The experimental protocol was approved by the Loughborough University Ethical Advisory Committee and was conducted in accordance with the World Medical Associations Declaration of Helsinki for medical research using human participants. All participants were given an information sheet that outlined the procedure, risks and requirements for the experiment. Participants provided written informed consent and completed a questionnaire based health screening.

Experimental Protocol

Systemic, 70-min exposures to four conditions were performed in T.I.S.S. Peak Performance (Series 2009) Chambers at Loughborough University Environmental Ergonomic Research Centre. Participants were exposed once to each of four conditions; control/ normoxic- thermoneutrality (N), hypoxic- thermoneutrality (H), normoxic- cold (C) and hypoxic- cold (HC). Thermoneutral conditions (N, H) were 22°C (50% RH) ambient temperature (T_a) and subjects were dressed in shorts, a t-shirt, socks and trainers. In cold conditions (C, HC) T_a was 5°C (50% RH) and participants wore the same clothing, minus any upper body insulation (t-shirt). 5°C T_a was selected in an attempt to reduce average skin temperature (T_{sk}) by approximately 5-10°C, which in turn was assumed to change forearm muscle temperature and cause an increase in fatigue (Oksa et al. 2002) when compared to thermoneutral conditions (N, H). Hypoxic exposures (H, HC) were 0.13 Fraction of Inspired Oxygen (F_{IO_2} : equivalent altitude = ~4000m) aiming to reduce peripheral arterial oxygen saturation (SpO_2) to approximately 85%, a moderate level assumed high enough to influence fatigue during isolated muscle exercise (Millet et al. 2008; 2012; Perrey & Rupp 2009; Christian et al. 2014a). The selection of temperature and F_{IO_2} also aimed to balance severity with ecological validity, in order to maintain relevance for those working or exercising at altitude. Normobaric hypoxia was achieved using an inbuilt chamber hypoxic air generator. Hypoxia was continuously monitored for consistency using a Servomex (570A, Sussex, UK) oxygen analyser as well as the inbuilt analyser on the T.I.S.S. Peak Performance (Series 2009) Chambers. Condition order was balanced and exposures were separated by at least 4 days to allow full recovery from the fatigue protocol. Participants were blinded to conditions prior to exposure.

Measurements of Temperature, Arterial O₂ Saturation, Skin Blood Flow, Heart Rate and Perceived Exertion

During all conditions aural (T_{co}) and local skin temperature (T_{sk}) from 4 different sites on the exercising arm (bicep, midline of the posterior forearm, midline of the anterior forearm & posterior of hand), were collected. Grant International skin and aural thermistors were secured using Transpore 3M medical grade tape. Aural thermistors were also insulated using cotton wool and earmuffs. Data was recorded at 1-min intervals from 1-min pre-exposure using a Squirrel Data Logger (1000 series, Grant Instruments, UK).

Immediately after each MVC was performed, percentage saturation of peripheral arterial blood (SpO_2) and heart rate (HR) were measured using a pulse oximeter attached to the middle finger of the non-exercising arm (Model 8500, Nonin Medical, Netherlands). HR and SpO_2 were collected once every 5-s over the first 20-s of each 110-s rest period. Baseline measures of resting HR and SpO_2 were collected pre-exposure to condition N, post 5-mins supine rest.

Laser Doppler flowmetry (LDF; arbitrary perfusion units [AU; Flux]) of the index finger on the non-exercising hand (Server, Satellite and Optic Probe, MoorLAB, Moor Instruments, UK) and Borg's Rate of Perceived Exertion (RPE) scores were also recorded immediately after each MVC was performed. LDF was calibrated prior to exposure using the Brownian motion of polystyrene microspheres diluted in water. This LDF device has been used in previous physiological studies to assess skin blood flow (Thompson, Holowatz & Kenney 2005) and is precise to $\pm 3\%$ and accurate to $\pm 10AU$ as determined by the manufacturer (Moor Instruments, 2004).

Fatigue Protocol & Force Measurement

Upon exposure to the test conditions, each participant was secured into a restraint system that maintained 90° flexion of the elbow, with the palm and anterior forearm facing vertically, while restricting any movement of the wrist. The system was used to isolate the working forearm muscles and maintain consistent muscle dynamics during exercise. Participants remained secured throughout the experiment. Once in the environmental conditions, participants undertook a 15-min rest period, allowing time for body heat to decline in cold (not reaching steady state) and arterial oxygenation to stabilise in hypoxia. After the rest period, participants performed a fatigue protocol that consisted of

eight 5-min work bouts, each separated by 110-s rest/ data collection periods, timed using a standard digital stopwatch. During each 5-min exercise bout, dynamic grip clenches (Fit66 Adjustable Grip Exerciser) were performed every 2-s (timed using an audio/visual metronome), at a workload of 15% of the MVC recorded at the start of the exposure ($MVC_{baseline}$) on their first experimental day. Given all subjects were healthy, regularly active, and accustomed to performing a wide range of physical activities, no separate familiarisation was deemed necessary; however to ensure an accurate prediction of workload was made, practise attempts were available during the MVC used to predict the workload. In all other circumstances, only one MVC was performed. Based on pilot studies, the fatigue protocol was designed to induce an estimated workload that was approximately equal to or greater than, 'hard' or 15 on Borg's RPE Scale, when conducted in thermoneutral normoxic conditions.

MVC force (3-s contraction) using a Grip Dynamometer (Takei™ No.1857) was collected at the start of each exposure ($MVC_{baseline}$) and after every second work bout (MVC_1 , MVC_2 , MVC_3 , MVC_4). A change in MVC, in conjunction with the corresponding EMG data was used to quantify fatigue. Baseline measures of EMG ($EMG_{baseline}$) and MVC were collected on immediate exposure to the experimental condition. A schematic overview of the experimental protocol and interventions is shown in Figure 1.

Electromyography & Fatigue Index

To evaluate the myoelectrical activity of the working muscles, surface EMG (Biometrics Ltd, UK) was measured on flexor carpi radialis (FCR), flexor digitorum superficialis (FDS) and extensor digitorum (ED). Other studies analysing forearm muscle fatigue have utilised similar muscle groups (West et al. 1995; Oksa et al. 2002). Flexor group (FCR & FDS) contractions were dynamic during grip clench exercises, and isometric during MVC. The extensor digitorum acts as a fixator and was isometric during both contraction manoeuvres.

The placement of each electrode sensor followed the recommendations outlined by SENIAM (Hermens et al., 1999). The skin was cleaned with alcohol and shaved when necessary. To ensure accurate placement, participants were positioned with the hand supinated and elbow flexed (FDS and FCR) or with the hand pronated and the elbow extended (ED). Each electrode (Biometric Ltd EMG pre

amplifier type no.SX230) was placed over the belly of the muscle, identified using palpation over the centre line between the origin and insertion of the muscle. Finger and grip movements were used to tense the appropriate muscle, inducing visible muscle tone, and aiding the accuracy and consistency of EMG placement. The placement of the FDS (1), the FCR (2) and ED (3) electrodes anatomically corresponded to: 1) 2/3 medial of the lateral border of the ventral forearm (in line with the 2nd middle phalanx) and 2/3 distal from antecubital fossa to the wrist- palm intersection; 2) medial of the lateral border of the ventral forearm (in line with the 1st middle phalanx) and 1/3 distal from the antecubital fossa to wrist- palm intersection; and finally 3) one half medial of the lateral border of the dorsal forearm (in line with the 2nd middle phalanx) and 1/3 distal from the line of the antecubital fossa to the ulna styloid process. All sensors and wires were secured using double sided tape with the two 2-cm spaced probe contacts running in parallel with the muscle fibres. The reference/ ground electrode (Biometrics Ltd earthing strap type no.R200) was placed above the ulna styloid process (inactive tissue) on the opposite wrist. All signals were zeroed prior to measurement and MVC data was logged on a DataLog P3X8 (Biometrics Ltd, UK).

EMG was recorded during all MVC measurements. The EMG sample rate used was 1000Hz, and was amplified 1000 times to minimise noise on the connecting cable. A signal band of 15Hz to 450Hz, minus the unwanted line frequency of 50 or 60Hz, including the harmonics of this frequency, was measured. Root Mean Square (RMS; μV) amplitude values were averaged over a 400-ms running average for the duration of the MVC. Analysis was carried out on the corresponding DataLog Software (Biometrics Ltd, UK) then confirmed and stored using Microsoft Excel 2007. For EMG analysis, a 1-s manually selected sample to include the steadiest values either side of the highest point in the EMG signal was used. Each muscle EMG_{rms} amplitude was calculated using the mean over each 1-s trace. To calculate the average combined muscles EMG_{rms} amplitude, FDS FCR and ED were weighted as 1:1:2 representing a value equal in its contributions from flexors and extensors compartments.

The variable used in this study to define the level of mechanical fatigue was the Fatigue Index (FI; Oksa et al. 2002). The FI quantifies fatigue using changes in MVC force and EMG amplitude of both the flexor and extensor muscles (mean flexor and extensor weighted 1:1; see above), relative to start and finish of the exercise protocol. FI is calculated as:

$$FI = (MVC_{\text{baseline}} / EMG_{\text{baseline}}) / (MVC_4 / EMG_4)$$

If FI = 1.0 then fatigue is not apparent, however the higher the value above 1.0, the greater the mechanical failure (force independent of excitation). FI differed in the present study from Oksa et al. (2002), because EMG data was collected during the MVC manoeuvre, not during submaximal (fatiguing) exercise; although FI in this case still represents electromechanical transmission failure as well as fatigue through intramuscular metabolic disturbance of the mechanical contraction. To represent the electromechanical ratio over time, EMG_{rms} relative to force produced during MVC (μV per kg) was used; this variable is analogous to FI.

Data analysis

Independent variables for this study were 1) $F_{I\text{O}_2}$; 2) T_a ; and 3) experimental time. Dependent Variables were MVC, EMG_{rms} amplitude, RPE, HR, LDF flux, T_{co} , T_{sk} , and FI. All LDF data were corrected for individuals occluded baseline value using a 2- min-vascular occlusion (rubber band applied to the finger base) during the 15-min rest-period in condition N. MVC and EMG were calculated to represent a change from baseline for each condition. In the figures, EMG data and FI were normalised to the control condition. The following calculation was used to express the variables in condition H, C, and HC, as a percentage change from condition N:

$$\text{Normalised change (\%)} = (\text{Treatment [H, C, HC]} / \text{Control [N]}) - 1 \times 100$$

Significance between conditions across the whole exposure was tested using a three-way [$2 \times 2 \times 5$; $F_{I\text{O}_2} \times T_a \times \text{time (fatigue)}$], repeated measures ANOVA. Significance was tested at a 95% confidence level (p , <0.05). To test FI and specific time point data for significance between $F_{I\text{O}_2}$ and T_a , two-way (2×2) repeated measures ANOVA was used. When no significant interaction was observed, the effect of cold and hypoxia were reported as C & HC collapsed or H & HC collapsed respectively. All results are displayed as mean $\pm$ SEM.

RESULTS

Temperature, Arterial O₂ Saturation, Skin Blood Flow, Rate of Perceived Exertion & Heart Rate

Table 1 shows body temperature, SpO₂, Skin Blood Flow, RPE & HR across conditions at each MVC intervention. A significant ($p = 0.001$) reduction in pre-exposure to MVC₄ ΔT_{co} was observed during cold conditions ($-0.46 \pm 0.18^{\circ}\text{C}$) whereas a significant increase ($p = 0.007$) in ΔT_{co} was observed in thermoneutral conditions ($0.19 \pm 0.05^{\circ}\text{C}$). The results showed a significant effect of cold T_a ($p = 0.003$) but no significant effect of hypoxia ($p = 0.6$) and no interaction ($p = 0.5$) between stressors. At MVC₄ T_{sk} of the posterior and anterior forearm was also significantly ($p < 0.05$) lower during cold exposures ($25.6 \pm 0.5^{\circ}\text{C}$) than thermoneutral ($32.7 \pm 0.3^{\circ}\text{C}$), therefore achieving the aimed reduction in local T_{sk} of 5-10°C. Forearm T_{sk} increased slightly at exercise commencement ($t = 15\text{-mins}$) in all conditions but remained significantly ($p < 0.05$) lower in cold. Mean skin blood flow (LDF flux) over all time points was also significantly ($p = 0.001$) lower in the cold ($-178 \pm 63 \text{ AU}$). HR varied similarly with temperature ($p = 0.03$), reducing from $85 \pm 2 \text{ bpm}$ in thermoneutral conditions to $78 \pm 2 \text{ bpm}$ in cold. We observed no significant effects of hypoxia on forearm T_{sk} ($p = 0.8$), HR ($p = 0.2$) or LDF flux ($p = 0.3$). Mean SpO₂ measurements across the four interventions showed a significant ($p = 0.001$) reduction from $97.5 \pm 0.3\%$ SpO₂ in normoxic conditions to $85.5 \pm 0.6\%$ SpO₂ in hypoxic conditions. While the effect of temperature on SpO₂ was significant ($p = 0.01$) the mean change was minimal ($\sim 1\%$).

RPE increased significantly ($p = 0.03$) over time (RPE first set 12.6 ± 0.7 ; RPE last set 14.0 ± 0.7) but no participants reached task failure during the fatigue protocol in any conditions. There was no main effect over time for temperature ($p = 0.4$) and hypoxia ($p = 0.1$). However, a significant effect of temperature was observed at MVC₃ ($p = 0.021$) and MVC₄ ($p = 0.014$) and a trend for the effect of hypoxia ($p = 0.074$) was observed at MVC₁ (Table 1).

Maximal Isometric Voluntary Contraction

Figure 2 shows MVC force produced at each intervention in each condition. MVC decreased significantly ($p < 0.05$) over time and varied significantly ($p < 0.05$) between conditions at various MVC time points (see Figure 2). However, when combined there was no significant statistical interaction

between cold and hypoxia (independent contributions) on the total decrease at MVC₄ ($p = 0.8$) or at the MVC time points i.e. the combined effect of cold and hypoxia was additive. Figure 2, Panel B also shows MVC plotted as a percentage decline from the baseline MVC.

Electromyography & Fatigue Index.

EMG_{rms} amplitude during MVC for the combined forearm muscles decreased incrementally with time ($p = 0.03$), however the effects of temperature ($p > 0.2$) and hypoxia ($p > 0.5$) were not significant between interventions (Figure 3, Panel A), except during MVC₁ where the normalised combined EMG_{rms} was higher ($p = 0.05$) during cold conditions (C + HC) by $32 \pm 13 \mu\text{V}$ (Figure 3, Panel B).

The FI of the forearm muscles increased to 1.25 in cold and 1.10 in hypoxia. Variance was significant for effects of temperature ($p = 0.003$) and hypoxia ($p = 0.01$) however there was no stressor interaction ($p = 0.9$): FI was equal 1.45 in combined conditions (Figure 4, Panel A). Expressed as a percentage of condition N (FI_%), exercise resulted in $24 \pm 7\%$ and $39 \pm 9\%$ higher fatigue in hypoxia and cold respectively. The combined effect was additive resulting in $62 \pm 11\%$ increase in FI i.e. there was no interaction (Figure 4, Panel B; Table 1). EMG_{rms} relative to force produced during MVC ($\mu\text{V per kg}$) was also significantly increased, with main effects for cold ($p = 0.02$) over time and a strong trend for hypoxia ($p = 0.06$) over time. By MVC₄ the effect was significant for both cold ($p = 0.003$) and hypoxia ($p = 0.008$), increasing by 2.0 and 1.2 $\mu\text{V per kg}$ (36 and 23% of condition N) respectively. The combined effect was additive (3.1 $\mu\text{V per kg}$ and 56% of condition N) showing no interaction ($p = 0.9$) between stressors (Figure. 5, Panel A).

DISCUSSION

Summary of main findings

This study quantified forearm muscle fatigue development during independent and combined reductions in ambient temperature and inspired oxygen concentration, during repeated low resistance exercise. The results demonstrate an additive, not interactive, effect on fatigue when humans are exposed to combined hypoxia and cold. Additive effects were observed for both MVC force output and electromechanical ratio (FI and $\mu\text{V per kg}$) at various time points.

Maximal voluntary contraction in hypoxic -cold

The decline in maximal voluntary force over time can be used to quantify net (overall) neuromuscular fatigue (Gandevia, 2001).

In this study, independent exposure to hypoxia and cold significantly reduced MVC force at various time points (Figure 2). In hypoxia, the decline in MVC force (Figure 2, Panel B) occurred early in the fatigue protocol, before plateauing at a lower level than thermoneutral normoxic conditions (Figure 2, Panel B). This differed from cold stress, which continuously reduced MVC over time, resulting in a greater impact of cooling [compared to hypoxia] in the later stages of the exercise protocol (Figure 2, Panel B). The differences in the temporal decline in MVC may be explained by the stabilisation of arterial oxygenation early in the hypoxic protocol (Table 1), contrary to a more progressive cold penetration through peripheral tissue with cold exposure.

When hypoxia and cold were combined, the temporal (shape of) decline in MVC force (Figure 2, Panel B) reflected equal (additive) contributions from both stressors, with neither hypoxia nor cold taking clear precedence in the combined condition (Figure 2, Panel B) i.e. during hypoxic-cold, MVC was subject to both an early (hypoxic) and progressive (cold) decrease over time. This is reflected in the final MVC (MVC_4), which reduced in force to a value equal to the summative effect of each stressor individually. Specifically during independent cold and hypoxic stress, MVC force declined by 8.1 and 13.9% more than thermoneutral normoxia, while MVC force decreased by 21.4% more during combined hypoxic-cold, closely matching the additive value of hypoxia and cold individually (22%).

Electromechanical mechanisms during fatigue

Force and EMG assessment has been widely used to understand central and peripheral contributions to fatigue (for reviews see: Gandevia, 2001 Amann & Calbet 2007). However EMG is perhaps better suited to subdividing electrophysiological excitation and mechanical fatigue further downstream from the neuromuscular junction – to a point within the muscle fibre itself (Allen et al. 2008). Importantly under this definition, mechanical fatigue still encompasses many of the intramuscular factors usually associated with peripheral fatigue (Enoka & Stuart 1992; Fitts 1994; Allen et al 2008), while the

changes in motor unit excitation remain partially representative of descending voluntary drive (Gandevia, 2001).

Mechanical fatigue in hypoxic-cold

In the present study, mechanical fatigue was disassociated from electrophysiological changes using the electromechanical ratio. These include EMG_{rms} relative to MVC force (in μV per kg) and the FI (Figure 4; Figure 5). EMG/MVC and the FI measure the direct mechanical response to net fibre excitation and can be attributed on an individual fibre basis to both electromechanical transmission failure and a reduced mechanical response (e.g. force or power) per unit of excitation (Allen et al. 2008).

In this study, independent cold exposure resulted in an increase in both $\Delta EMG/MVC$ and the FI (Figure 4; Figure 5). A similar yet smaller effect was observed during independent exposure to hypoxia. By MVC_4 (post-exercise), $\Delta EMG/MVC$ was significantly increased by 1.2 and 2 μV per kg in hypoxia and cold respectively, representing a 23 and 36% change from N (Figure 5). This corresponded to a 24 and 39% increase in FI respectively (Figure 4). FI and $\Delta EMG/MVC$ reflected the effect of hypoxia and cold on MVC force (Figure 2), suggesting fatigue was predominantly of mechanical origin in this study i.e. a failure distal of electrophysiological processes. This is supported by the time course of mechanical fatigue, which is similar to those observed in electrically stimulated muscle fibres in vitro; an early increase, a plateau, then a late increase (Allen et al 2008; Marcora & Staiano, 2010).

Mechanical failure independent of excitation during cold exposure can be attributed to number of factors, such as increases in the relative exercise intensity due to co-activation of the agonist-antagonist pair (Oksa et al. 1997; 2002; Racinais & Oksa 2010) and reduced muscle blood flow (Yanagisawa et al, 2007; Gregson et al, 2011). It may also result from progressive cold penetration through the muscle tissue (Oksa et al. 2002) gradually increasing the number of muscle fibres affected by slowed intramuscular energetics (Bergh, 1980; Faulkner et al. 1990; De Ruiter & De Haan 2000; Allen et al 2008; Racinais & Oksa, 2010; Cahill et al. 2011). Conversely, the mild effect of hypoxia on mechanical function (fatigue) has been widely attributed to increases in energetic metabolite

interference with Ca^{2+} handling and the contractile proteins (Edwards 1981; Fulco et al. 1996; Fitts 1994; Haseler et al. 1999; Hogan et al. 1999; Amann & Calbet 2007; Perrey & Rupp 2009; Christian et al., 2014a).

In this study, it was hypothesised that during combined hypoxic-cold stress, co-activation (Oksa et al. 2002) and muscle blood flow reductions (Yanagisawa et al, 2007; Gregson et al, 2011) in cold would accentuate the low oxygen delivery (Amann & Calbet 2007) during hypoxia, resulting in an interaction on net fatigue (MVC) due to altered function at the mechanical level (FI and $\Delta\text{EMG}/\text{MVC}$). However, contrary to this hypothesis the effect on mechanical fatigue was additive showing no interactions; $3.1 \pm 0.3 \mu\text{V}$ per kg (FI increased by 62% of N) (Figure 4; Figure 5). One explanation for this is that the inhibitory influence of increased energetic metabolites during hypoxia (Fitts 1994; Haseler et al. 1999; Hogan et al. 1999; Perrey & Rupp 2009), was not influenced by the direct slowing effect of cooling on ATP hydrolysis, Ca^{2+} handling and the contractile proteins during cold exposure (Kossler et al. 1987; Sweitzer & Moss 1990; Feretti 1992; Oksa et al. 2002). It suggests that despite each stressor hindering mechanical function, cold and hypoxia may influence fatigue through sufficiently independent cellular mechanisms, so as not to interact with one another during low resistance exercise.

The observed additive effect may also result from multiple inter-mechanism interactions and/or an interaction cancelation. A synergistic and antagonistic interaction of similar magnitude would result in net additive effects, and thus to investigate this further, interactive studies examining the individual mechanisms that contribute to hypoxic- cold fatigue are required.

Electrophysiological factors during fatigue in hypoxic- cold

In the present study, MVC sequentially decreased in peak force output and EMG_{rms} amplitude over time, across all conditions (Figure 3). This suggests changes in corticospinal drive (Gandevia, 2001) and/ or action potential propagation (Bigland Ritchie et al. 1981) were partially responsible for fatigue observed during the exercise protocol. However, the decline in combined forearm EMG_{rms} was not generally affected by condition, thus it is unlikely that electrophysiological factors are primarily responsible for the environmental influences on fatigue in the present study. Moreover, despite changes in cognitive function as a result of immediate exposure to environmental stress (Gaoua et al. 2011), we

observed no influence on MVC or EMG_{rms} at the start of the exposure, suggesting central drive and muscle fibre recruitment remained largely unaffected.

Perceptual responses to fatigue in hypoxic- cold

Despite no significant condition effect on motor unit recruitment (EMG_{rms}) during MVC (Figure 3), the rise in the relative work rate (recruitment/ voluntary drive) during submaximal repetitive exercise over time and across conditions did appear to mildly increase RPE (Table 1). RPE reflected the temporal decline in overall fatigue, showing a trend for decrease early in hypoxia (MVC₁) and a small effect on effort during the latter stages of the protocol with cooling (MVC₃ & MVC₄). Since participants received no specific detailing on the interpretation of RPE, the increase is likely in response to both a greater mental effort (Marcora & Staiano, 2010) and a higher peripheral discomfort (Christian et al. 2014b) with fatigue.

The effect on mechanical and perceived fatigue, yet not MVC recruitment could be because of the inherent limitations on conscious regulation during closed loop protocols. In this study, the only option was to a) stop exercise, or b) attenuate motor output during a brief MVC; each providing little or no relief from fatigue. As such, neither perceptual tolerance to fatigue, nor the volitional regulation of neural drive in response to high or maximal levels of peripheral fatigue were investigated. In fact, even given a greater scope for regulation, recovery between bouts may have resulted in maintained neural drive during brief maximal contractions, since afferent feedback is most relevant during a prolonged mental effort with no immediate recovery (Cahill et al., 2011; Amann et al. 2013; Christian et al. 2014a; Lloyd et al., 2014).

Additional perspectives and limitations

Previous studies have shown that hypoxia can cause vasodilation in non- acral skin during cold exposure (e.g. Cipriano & Goldman, 1975; Johnston et al. 1996; Simmons et al, 2010). However, the vasoconstrictor response of acral skin (finger pad) during cold exposure was not significantly affected by hypoxia in the present study. A possible explanation is that the large core to skin temperature gradient (~ -11°C) at the non- exercised finger was sufficient to abolish the hypoxic vasodilation effect (Simmons et al, 2011), contrary to observations of more proximal skin at milder temperatures for

shorter durations (Cipriano & Goldman, 1975; Simmons et al, 2010). Furthermore, to our knowledge no direct relationship has previously been shown between perfusion of skin microvasculature and local muscle blood flow. As such, substantiation of the link between skin, local muscle blood flow, and fatigue during both local cold, and combined hypoxic- cold stress, would be an interesting avenue for future studies.

In the present study aural temperature was used to illustrate changes in core temperature. The results showed a small but significant shift of -0.46°C during cold exposure, despite a probable increase in metabolic rate during exercise. It should be noted that some of this drop in aural temperature may have been caused by local tissue cooling of the aural canal in this study. However previous studies using a similar duration and severity of hypoxic- cold have reported similar changes after 75-mins rest measured by rectal temperature (-0.4°C ; Robinson & Haymes 1990) and a follow up study in our lab under similar conditions as the present study (using rectal temperature assessment) has also produced drops in core temperature, although smaller (-0.2°C after 40-min rest) and after an initial rise (0.15°C).

Surface EMG reflects not only descending drive to the muscle but also the electrode/ muscle interface. As such the effect of changes in local tissue temperature around electrode site cannot be ruled out as a contributing factor to the present observations (Racinais, 2013). Also, because flexor and extensor EMG was not measured during the submaximal exercise bouts, co-activation and the temporal rise in mechanical fatigue cannot be concluded from the present results. Additionally due to the use of small forearm muscles in this study, inter-muscle cross talk is a potential limitation. Finally, it should be recognised that the absence of a separate familiarisation for MVC trials is also a limitation. While this may be minimised by allowing practise attempts prior to the prediction of workloads, and by using young, regular exercisers who are well accustomed to physical activity, the importance of this on reproducibility, validity and reliability should be acknowledged in the context of this study.

Conclusion

In conclusion, the decrease in MVC force and increase in electromechanical ratio and FI (Figures 2, 4 & 5 respectively) support previous findings, suggesting that independent exposure to cold and hypoxia can significantly increase muscle fatigue compared to control conditions. The main finding of this

study is that when moderate hypoxia and cold exposure are combined, the decline in MVC force and rise in electromechanical ratio suggest the level of fatigue increases additively, with no interactions; however further research is warranted using alternative stressor severities and exercise modalities, as this may lead to different results.

REFERENCES

- Allen DG, Lamb GD & Westerblad H (2008) Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev* 88, 287 – 332.
- Amann M, Calbet JAL (2007) Convective oxygen transport and fatigue. *J Appl Physiol* 104: 861–870.
- Amann M, Dempsey JA (2007) Peripheral muscle fatigue from hyperoxia to moderate hypoxia – a carefully regulated variable? *Physiology News* 66: 28–29.
- Amann M, Eldridge MW, Lovering AT, Stickland MK, Pegelow DF, Dempsey JA (2006a) Arterial oxygenation influences central motor output and exercise performance via effects on peripheral locomotor muscle fatigue. *J Physiol* 575: 937–952.
- Amann M, Pegelow DF, Jacques AJ, Dempsey JA (2007a) Inspiratory muscle work in acute hypoxia influences locomotor muscle fatigue and exercise performance of healthy humans. *Am J Physiol Regul Integr Comp Physiol* 293: R2036–R2045.
- Amann M, Romer LM, Pegelow, DF, Jacques AJ, Hess, CJ, Dempsey JA (2006b) Effects of arterial oxygen content on peripheral locomotor muscle fatigue. *J Appl Physiol* 101: 119–127.
- Amann M, Romer LM, Subudhi AW, Pegelow DF, Dempsey JA (2007b) Severity of arterial hypoxaemia affects the relative contributions of peripheral muscle fatigue to exercise performance in healthy humans. *J Physiol* 581: 389–403.
- Amann M, Venturelli M, Ives SJ, McDaniel J, Layec G, Rossman MJ, Richardson RS (2013) Peripheral fatigue limits endurance exercise via a sensory feedback-mediated reduction in spinal motoneuronal output. *J Appl Physiol* 115: 355–364.
- Bergh U (1980) Human power at subnormal body temperatures. *Acta Physiol Scand* 478: 1–39.
- Bergh U, Ekblom B (1979a) Influence of muscle temperature on maximal muscle strength and power output in human skeletal muscles. *Acta Physiol Scand* 107:33–37.
- Bergh U, Ekblom B. (1979b) Physical performance and peak aerobic power at different body temperatures. *J Appl Physiol* 46: 885–889.

- Bigland-Ritchie B, Donovan EF, Roussos CS (1981) Conduction velocity and EMG power spectrum changes in fatigue of sustained maximal efforts. *J Appl Physiol* 51: 1300–1305.
- Blomstrand E, Bergh U, Essen-Gustavsson B, Ekblom B (1984) Influence of low muscle temperature on muscle metabolism during intense dynamic exercise. *Acta Physiol Scand* 120: 229–236.
- Cahill F, Kalmar JM, Pretorius T, Gardiner PF, Giesbrecht GG (2011) Whole-body hypothermia has central and peripheral influences on elbow flexor performance. *Exp Physiol* 96: 528–538
- Cè E, Rampichini S, Agnello L, Limonta E, Veicsteinas A, Esposito F. (2012) Combined effects of fatigue and temperature manipulation on skeletal muscle electrical and mechanical characteristics during isometric contraction. *J Electromyogr Kinesiol*. 22(3):348-355
- Christian RJ, Bishop DJ, Billaut F, Girard O (2014a) Peripheral fatigue is not critically regulated during maximal, intermittent, dynamic leg extensions. *J Appl Physiol* 117: 1063–1073.
- Christian RJ, Bishop DJ, Billaut F, Girard O. (2014b) The role of sense of effort on self-selected power output. *Frontiers in Physiology* 5: 115.
- Cipriano LF & Goldman RF (1975). Thermal responses of unclothed men exposed to both cold temperatures and high altitudes. *J Appl Physiol* 39: 796–800.
- De Ruiter CJ, De Haan A. (2000) Temperature effect on the force/velocity relationship of the fresh and fatigued human adductor pollicis muscle. *Pflügers Arch* 440: 163–170.
- Edwards R (1981) Human muscle function and fatigue. In: *Human Muscle Fatigue Physiological Mechanisms*, edited by Porter R and Whelan J. London: Pitman, p. 1–8.
- Enoka RM, Stuart DG (1992) Neurobiology of muscle fatigue. *J Appl Physiol* 72: 1631–1648.
- Faulkner JA, Zerba E, Brooks SV (1990) Muscle temperature of mammals: cooling impairs most functional properties. *Am J Physiol Regul Integr Comp Physiol* 28: R259–R265.
- Fitts R H (1994) Cellular mechanisms of muscle fatigue. *Physiol Rev* 74: 49–94.
- Ferretti G (1992) Cold and muscle performance. *Int J Sports Med* 13 Suppl 1: 185–187.
- Folt CL., Chen CY., Moore MV, Burnaford J (1999). Synergism and antagonism among multiple stressors. *Limnol. Oceanogr.*, 44, 864–877.
- Fulco CS, Cymerman A, Muza S R, Rock PB, Pandolf KB, Lewis SF (1994) Adductor pollicis muscle fatigue during acute and chronic altitude exposure and return to sea level. *J Appl Physiol* 77: 179–183.

- Fulco CS, Lewis SF, Frykman PN, Boushel R, Smith S, Harman EA, Cymerman A, Pandolf KB. (1996) Muscle fatigue and exhaustion during dynamic leg exercise in normoxia and hypobaric hypoxia. *J Appl Physiol* 81: 1891–1900.
- Gandevia SC (2001). Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev* 81: 1725-1789.
- Gaoua N, Grantham J, El Massioui F, Girard O, Racinais S (2011) Cognitive decrements do not follow neuromuscular alterations during passive heat exposure. *Int J Hyperthermia*: 27:10-19
- Gregson W, Black MA, Jones H, Milson J, Morton J, Dawson B, Atkinson G, Green D (2011) Influence of cold water immersion on limb and cutaneous blood flow at rest. *Am J Sports Med*: 39(6): 1316-1323
- Gautier H, Bonora M, Schultz SA, Remmers JE (1987) Hypoxia-induced changes in shivering and body temperature. *J Appl Physiol* 62: 2477–2484.
- Goodall S, Ross EZ, Romer L M (2010) Effect of graded hypoxia on supraspinal contributions to fatigue with unilateral knee-extensor contractions. *J Appl Physiol* 109: 1842-1851
- Haseler LJ, Hogan MC, Richardson RS (1999) Skeletal muscle phosphocreatine recovery in exercise-trained humans is dependent on O₂ availability. *J Appl Physiol* 86: 2013–2018.
- Hogan MC, Richardson RS, Haseler LJ (1999) Human muscle performance and PCr hydrolysis with varied inspired oxygen fractions: a ³¹P-MRS study. *J Appl Physiol* 86: 1367–1373.
- Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G (2000) Development of recommendations for SEMG sensors and sensor placement procedures. *J Electromyogr Kinesiol* 10:361–374.
- Johnston CE, White MD, Wu M, Bristow GK, Giesbrecht GG (1996) Eucapnic hypoxia lowers human cold thermoregulatory response thresholds and accelerates core cooling. *J Appl Physiol* 80: 422–429
- Katayama K, Amann M, Pegelow DF, Jacques AJ, Dempsey JA (2007) Effect of arterial oxygenation on quadriceps fatigability during isolated muscle exercise. *Am J Physiol Regul Integr Comp Physiol* 292: R1279–R1286.
- Kossler F, Lange F, and Kuchler (1987) Isometric twitch and tetanic contraction of frog skeletal muscles at temperatures between 0 to 30°C. *Biomed Biochim Acta* 46: 809–814.

- Lloyd A, Hodder S, Faulkner S, Fry A, Havenith G (2014) Muscle temperature limits isometric endurance via sensory feedback-mediated central fatigue. *Med Sci Sports Exerc* 46(5S):180-187.
- Marcora SM, Staiano W (2010) The limit to exercise tolerance in humans: mind over muscle? *Eur J Appl Physiol* 109(4):763-70.
- McArdle WD, Magel JR, Lesmes GR, Pechar GS (1976) Metabolic and cardiovascular adjustment to work in air and water at 18,25, and 33°C. *J Appl Physiol* 40(1): 85-90.
- Millet, GY, Aubert D, Favier, FB, Busso, T. and Benoît, H. (2008) Effect of acute hypoxia on central fatigue during repeated isometric leg contractions. *Scand J Med Sci Sports*, 19: 695–702.
- Millet GY, Muthalib M, Jubeau M, Laursen PB, Nosaka K. (2012) Severe hypoxia affects exercise performance independently of afferent feedback and peripheral fatigue. *J Appl Physiol* 112: 1335–1344
- Moor Instruments (2004) MoorLab laser Doppler blood flow user manual.
- Mucke R, Heuer D (1989) Behaviour of EMG parameters and conduction velocity in contractions with different muscle temperatures. *Biomed Bichim Acta* 5/6: 459–464
- Oksa J, Ducharme M, Rintamäki H (2002) Combined effect of repetitive work and cold on muscle function and fatigue. *J Appl Physiol* 92: 354–361.
- Oksa J, Rintamäki H, Rissanen S (1997) Muscle performance and electromyographic activity of the lower leg muscles with different levels of cold exposure. *Eur J Appl Physiol* 75: 484–490.
- Perrey S, Rupp T (2009) Altitude-induced changes in muscle contractile properties *High Altitude Medicine and Biology*. 10;175-182.
- Racinais S, Oksa J (2010) Temperature and neuromuscular function. *Scand J Med Sci Sports* 20: 1-18.
- Racinais S (2013) Hot ambient conditions shift the Force / EMG relationship. *Springerplus* 15: 2(1): 317.
- Ray CA, Hume KM, Gracey KH, Mahoney ET (1997) Muscle cooling delays activation of the muscle metaboreflex in humans. *Am J Physiol Heart Circ Physiol* 273: 2436-2441.
- Robinson KA, Haymes EM (1990) Metabolic effects of exposure to hypoxia plus cold at rest and during exercise in humans. *J Appl Physiol* 68: 720-725
- Segal SS, Faulkner JA, White TP (1986) Skeletal muscle fatigue in vitro is temperature dependent. *J Appl Physiol* 61: 660-665.

- Simmons GH, Fieger SM, Minson CT & Halliwill JR (2010). Hypoxic cutaneous vasodilation is sustained during brief cold stress and is not affected by changes in CO₂. *J Appl Physiol* 108, 788–792.
- Simmons GH, Barrett-O'Keefe Z, Minson CT, Halliwill JR (2011) Cutaneous vascular and core temperature responses to sustained cold exposure in hypoxia. *Exp. Physiol.*, 96 1062–1071.
- Sweitzer N and Moss R (1990) The effect of altered temperature on Ca²⁺-sensitive force in permeabilized myocardium and skeletal muscles. *J Gen Physiol* 96: 1221–1245.
- Tipton M (2012) A case for combined environmental stressor studies. *Extrem Physiol Med* 1:7-8
- Thompson, C S, Holowatz LA, Kenney WL (2005) Cutaneous vasoconstrictor responses to norepinephrine are attenuated in older humans. *Am J Physiol Regul Integr Comp Physiol* 288: R1108 –R1113.
- Todd G, Butler JE, Taylor JL, Gandevia SC (2005) Hyperthermia: a failure of the motor cortex and the muscle. *J Physiol* 563: 621-631.
- West W, Hicks A, Clements L, Downing J (1995) The relationship between voluntary electromyogram, endurance time and intensity of effort in isometric handgrip exercise. *Eur J Appl Physiol* 71: 301 30
- Wood SC (1991) Interactions between hypoxia and hypothermia. *Annu Rev Physiol* 53: 71–85.
- Yanagisawa O, Kudo H, Takahashi N, Yoshioka H (2004) Magnetic resonance imaging evaluation of cooling on blood flow and oedema in skeletal muscles after exercise. *Eur J Appl Physiol* 91:737–40

TABLE LEGENDS

Table 1: The effect of condition and time on aural temperature (T_{co}), whole arm and forearm skin temperature (T_{sk}), heart rate (HR), peripheral arterial oxygen saturation (SpO_2), index finger blood flow (LDF), and rate of perceived exertion (RPE). Temperature data are averaged over each 110-s rest period. HR, SpO_2 and LDF are averaged over 20-s after each MVC. N, thermoneutral (22°C T_a) normoxic (0.21 F_{IO_2}); H, thermoneutral (22°C T_a) hypoxic (0.13 F_{IO_2}); C, cold (5°C T_a) normoxic (0.21 F_{IO_2}); HC, cold (5°C T_a) hypoxic (0.13 F_{IO_2}). Symbols for effects within the same time point: @Trend for effects of hypoxia ($p < 0.08$). *Significant for effects of temperature ($p < 0.05$); #Significant for effects of F_{IO_2} ($p < 0.05$).

FIGURE LEGENDS

Fig 1 Schematic representation of the experimental protocol, measurements taken and interventions.

T_{co} , aural temperature; T_{sk} , skin temperature; MVC, maximal isometric voluntary contraction; EMG, electromyography; LDF, laser Doppler blood flow; SpO_2 , peripheral oxygen saturation percentage; HR, heart rate; RPE, rating of perceived exertion. Straight arrows indicate the timing of data collection interventions.

Fig 2 The effect of fatigue on maximal voluntary contraction (MVC) across conditions. Panel B shows maximal voluntary contraction force normalised to baseline for each intervention. N, thermoneutral ($22^{\circ}\text{C } T_a$) normoxic ($0.21 F_{I}O_2$); H, thermoneutral ($22^{\circ}\text{C } T_a$) hypoxic ($0.13 F_{I}O_2$); C, cold ($5^{\circ}\text{C } T_a$) normoxic ($0.21 F_{I}O_2$); HC, cold ($5^{\circ}\text{C } T_a$) hypoxic ($0.13 F_{I}O_2$). Symbols for effects within the same time point: *Significant for the effect of temperature to $p < 0.05$ level. #Significant for the effect of $F_{I}O_2$ to $p < 0.05$ level.

Fig 3 The effect of fatigue on root mean squared electromyogram amplitude (EMG) across conditions. Panel B shows EMG_{rms} normalised to baseline for each intervention. N, thermoneutral ($22^{\circ}\text{C } T_a$) normoxic ($0.21 F_{I}O_2$); H, thermoneutral ($22^{\circ}\text{C } T_a$) hypoxic ($0.13 F_{I}O_2$); C, cold ($5^{\circ}\text{C } T_a$) normoxic ($0.21 F_{I}O_2$); HC, cold ($5^{\circ}\text{C } T_a$) hypoxic ($0.13 F_{I}O_2$). Symbols for effects within the same time point: *Significant for the effect of temperature to $p < 0.05$ level (data significant when normalised percentage of baseline only).

Fig 4 The effect of condition on Fatigue Index (Panel A) and condition normalised (Δ) Fatigue index (Panel B); N, thermoneutral ($22^{\circ}\text{C } T_a$) normoxic ($0.21 F_{I}O_2$); H, thermoneutral ($22^{\circ}\text{C } T_a$) hypoxic ($0.13 F_{I}O_2$); C, cold ($5^{\circ}\text{C } T_a$) normoxic ($0.21 F_{I}O_2$); HC, cold ($5^{\circ}\text{C } T_a$) hypoxic ($0.13 F_{I}O_2$). *Significant for the effect of temperature to $p < 0.005$ level. #Significant for the effect of $F_{I}O_2$ to $p < 0.05$ level.

Fig 5 shows the EMG/MVC relative to baseline, and normalised to the thermoneutral condition for each time point in hypoxia (H), cold (C) and hypoxic- cold (HC). The right hand axis on both panels shows each variable as percentage change of thermoneutral normoxic condition. Symbols for effects within the same time point: @Trend for effects of temperature ($p < 0.08$). *Significant for effects of temperature ($p < 0.05$); **Significant for effects of temperature ($p < 0.01$). +Trend for effects of $F_{I}O_2$ ($p < 0.08$). #Significant for effects of $F_{I}O_2$ ($p < 0.05$); ##Significant for effects of $F_{I}O_2$ ($p < 0.01$).

Table1:

Variable	Time Point	N	H	C	HC
Aural T_{co} (°C)	Pre-exposure	36.45 ± 0.08	36.38 ± 0.07	36.60 ± 0.12	36.57 ± 0.07
	iMVC ₁ (ΔT_{co})	0.20 ± 0.06	0.21 ± 0.09	0.07 ± 0.20	-0.10 ± 0.11
	iMVC ₂ (ΔT_{co})	0.20 ± 0.06	0.21 ± 0.08	-0.08 ± 0.23*	-0.31 ± 0.14*
	iMVC ₃ (ΔT_{co})	0.18 ± 0.05	0.19 ± 0.08	-0.23 ± 0.24*	-0.45 ± 0.16*
	iMVC ₄ (ΔT_{co})	0.18 ± 0.05	0.21 ± 0.08	-0.33 ± 0.25*	-0.58 ± 0.17*
Whole Arm T_{sk} (°C)	Pre-exposure	30.66 ± 0.56	30.99 ± 0.36	32.03 ± 0.18	31.30 ± 0.24
	iMVC ₁	32.40 ± 0.39	32.29 ± 0.28	24.70 ± 0.56*	25.06 ± 0.81*
	iMVC ₂	32.76 ± 0.37	32.53 ± 0.26	24.90 ± 0.47*	25.13 ± 0.78*
	iMVC ₃	32.89 ± 0.35	32.53 ± 0.30	24.94 ± 0.53*	25.35 ± 0.72*
	iMVC ₄	32.86 ± 0.32	32.50 ± 0.25	24.86 ± 0.52*	25.08 ± 0.62*
Forearm T_{sk} (°C)	Pre-exposure	30.53 ± 0.46	30.62 ± 0.31	32.06 ± 0.22*	31.17 ± 0.38
	iMVC ₁	32.48 ± 0.36	32.20 ± 0.42	25.10 ± 0.73*	25.69 ± 0.93*
	iMVC ₂	33.06 ± 0.29	32.73 ± 0.39	25.68 ± 0.72*	26.27 ± 0.89*
	iMVC ₃	33.20 ± 0.26	32.80 ± 0.46	26.08 ± 0.79*	26.42 ± 0.88*
	iMVC ₄	33.17 ± 0.24	32.76 ± 0.36	26.08 ± 0.63*	26.35 ± 0.82*
HR (BPM)	Pre-exposure	61.5 ± 1.0	-	-	-
	iMVC _{baseline}	87.4 ± 4.7	85.3 ± 2.5	77.3 ± 3.6*	79.3 ± 4.4*
	iMVC ₁	81.6 ± 1.8	89.0 ± 3.2 [@]	77.0 ± 3.7*	81.4 ± 3.3* [@]
	iMVC ₂	81.6 ± 2.5	82.0 ± 4.0	78.1 ± 4.0	78.9 ± 2.4
	iMVC ₃	84.6 ± 3.1	90.4 ± 4.0 [#]	75.1 ± 4.0*	81.3 ± 3.4* [#]
	iMVC ₄	85.8 ± 2.9	87.8 ± 4.1	76.8 ± 3.1*	80.5 ± 4.3*
SpO ₂ (%)	Pre-exposure	98.0 ± 0.4	-	-	-
	iMVC _{baseline}	96.8 ± 0.3	87.4 ± 1.5 [#]	97.9 ± 0.5	88.6 ± 0.9 [#]
	iMVC ₁	97.4 ± 0.3	85.3 ± 0.6 [#]	98.1 ± 0.3	86.6 ± 1.2 [#]
	iMVC ₂	97.0 ± 0.4	85.0 ± 0.7 [#]	98.5 ± 0.2	87.0 ± 1.6 [#]
	iMVC ₃	96.6 ± 0.5	83.9 ± 1.3 [#]	98.5 ± 0.4*	84.8 ± 0.6* [#]
	iMVC ₄	97.5 ± 0.4	83.9 ± 0.9 [#]	98.3 ± 0.3*	86.0 ± 0.8* [#]
LDF (% of N)	iMVC _{baseline}	389 ± 14	408 ± 53	199 ± 25*	172 ± 27*
	iMVC ₁	345 ± 51	303 ± 47	76 ± 17*	65 ± 15*
	iMVC ₂	304 ± 46	256 ± 66	98 ± 19*	77 ± 20*
	iMVC ₃	317 ± 53	253 ± 68	84 ± 23*	96 ± 22*
	iMVC ₄	268 ± 44	272 ± 47	42 ± 12*	94 ± 25*
RPE	iMVC ₁	12.4 ± 0.8	13.4 ± 0.8 [@]	11.9 ± 1.0	12.8 ± 0.7 [@]
	iMVC ₂	12.5 ± 0.9	13.6 ± 0.6	13.0 ± 0.7	12.9 ± 0.8
	iMVC ₃	13.0 ± 0.8	14.0 ± 0.7	13.9 ± 0.6*	14.0 ± 0.8*
	iMVC ₄	13.1 ± 0.9	14.0 ± 0.7	14.0 ± 0.8*	14.8 ± 0.6*

Figure 1:

Figure 2:

Figure 3:

Figure 4:

Figure 5: