Total Enantioselective Synthesis of (-)-Cytisine

Bruno Danieli, Giordano Lesma,* Daniele Passarella, Alessandro Sacchetti, Alessandra Silvani,* Andrea Virdis

Dipartimento di Chimica Organica e Industriale e Centro Interdisciplinare Studi biomolecolari e applicazioni Industriali (CISI), Università degli Studi di Milano, via G. Venezian 21, 20133 Milano

Supporting Information Available

(3*S*,5*R*) (+)-3-Acetoxymethyl-5-formyl-piperidine-1-carboxylic acid benzyl ester 5 was prepared according to ref.12.

(3*S*,5*R*, 1'*S*) (-)-3-Acetoxymethyl-5-(1'-hydroxy-but-3'-enyl)-piperidine-1-carboxylic acid benzyl ester 6a

To a suspension of (-)-*B*-methoxydiisopinocampheylborane (1.2 g, 3.8 mmol) in ether (18 mL) under nitrogen at -78°C, a solution of allylmagnesium bromide (1.0 M in ether, 3.8 mL, 3.8 mmol) was added dropwise. The mixture was stirred for 15 min at -78°C and for 1 h at room temperature, and then it was recooled to -78°C. A solution of **5** (933 mg, 2.9 mmol) in ether (10 mL) was added, and the mixture was stirred for 2 h at -78°C and for 2 h at room temperature. A 3N sodium hydroxide solution (15 mL) was added, followed by 30% hydrogen peroxide solution (15 mL), and the mixture was heated for 3 h. After cooling, the mixture was diluted with ether (40 mL) and it was washed with water (30 mL) and brine (30 mL). The aqueous washings were extracted three times with diethyl ether (40 mL), and the combined ethereal extracts were dried over Na₂SO₄. Removal of the solvent under reduced pressure was followed by flash chromatography of the residue on silica gel (ethyl acetate/hexane 1:5), to yield 801 mg (76%) of **6a** as a colorless oil: *R_f* (ethyl acetate/hexane 1:5) 0.30; [α]_D²⁵ - 1.4 (*c* 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ_H 7.35 (5H, br, s), 5.88-5.70 (1H, m), 5.18-5.10 (2H, m), 5.12 (2H, s), 4.40 (1H, d, *J* = 13.3 Hz), 4.25 (1H, d, *J* = 13.3 Hz), 4.00 (1H, dd, *J* = 11.7, 6.7 Hz), 3.85 (1H, dd, *J* = 11.7, 7.3 Hz), 3.50-3.40 (1H, m), 2.55 (1H, t, *J* = 12.1 Hz), 2.40 (1H, t, *J* = 12.6 Hz), 2.30-2.10 (2H, m), 2.00 (3H, s), 1.90-1.80 (1H, m), 1.75 (1H, d, *J* = 12.2 Hz), 1.65-1.50 (1H, m), 1.08 (1H, q, *J* = 12.2 Hz); ¹³C NMR (75.4 MHz, CDCl₃) δ_C 170.8, 155.1, 136.8, 134.2, 127.1-130.2 (5C), 118.7, 71.7, 67.1, 66.6, 46.8, 40.8, 39.1, 35.4, 29.7, 20.8; HRMS calcd for C₂₀H₂₇NO₅; 361.4419. Found: 361.4431.

(3*S*,5*R*, 1'*R*) (-)-3-Acetoxymethyl-5-(1'-hydroxy-but-3'-enyl)-piperidine-1-carboxylic acid benzyl ester 6b

6b was prepared following the procedure described for **6a**, using (+)-*B*-methoxydiisopinocampheylborane in place of the (-)-antipode (67% yield). *R_f* (ethyl acetate/hexane 1:5) 0.30; [α]_D²⁵ - 1.6 (*c* 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ_H 7.35 (5H, br, s), 5.88-5.75 (1H, m), 5.20-5.12 (2H, m), 5.15 (2H, s), 4.30 (1H, d, *J* = 10.2 Hz), 4.20 (1H, d, *J* = 12.2 Hz), 4.04 (1H, dd, *J* = 11.2, 5.3 Hz), 3.91 (1H, dd, *J* = 11.2, 7.2 Hz), 3.58-3.50 (1H, m), 2.65 (1H, t, *J* = 12.3 Hz), 2.47 (1H, t, *J* = 12.3 Hz), 2.34 (1H, dt, *J* = 13.9, 4.8 Hz), 2.22 (1H, dt, *J* = 14.0, 8.3 Hz), 2.05 (3H, s), 2.00 (1H, d, *J* = 13.0 Hz), 1.95-1.83 (1H, m), 1.70-1.58 (1H, m), 1.10 (1H, q, *J* = 12.2 Hz); ¹³C NMR (75.4 MHz, CDCl₃) δ_C 171.0, 155.3,

136.8, 134.2, 128.5-127.8 (5C), 118.9, 71.6, 67.1, 66.5, 47.1, 46.8, 40.6, 39.3, 35.5, 28.3, 20.8; HRMS calcd for $C_{20}H_{27}NO_5$: 361.4419. Found: 361.4428.

(3S,5R,1'S) (+)-3-Acetoxymethyl-5-(1'-methanesulfonyloxy-but-3'-enyl)-piperidine-1-carboxylic acid benzyl ester
7

To a solution of **6a** (843 mg, 2.3 mmol) in anhydrous CH_2Cl_2 (20 mL), under nitrogen atmosphere at $0^\circ C$, Et_3N (1.6 mL, 11.7 mmol, d 0.7 g/mL), DMAP (28 mg, 0.23 mmol) were added and methanesulfonyl chloride (270 μL , 3.5 mmol, d 1.5 g/mL) was slowly dropped. The mixture was stirred for 1.5 h at room temperature, then washed with 5% aq. H_3PO_4 (2 x 20 mL). The organic layer was dried over Na_2SO_4 and concentrated to give 1 g (99%) of **7**, as an oil: R_f (ethyl acetate/hexane 1:1) 0.40; $[\alpha]_D^{25} + 3.2$ (c 1, $CHCl_3$); 1H NMR (300 MHz, $CDCl_3$) δ_H 7.35 (5H, br, s), 5.90-5.70 (1H, m), 5.18-5.10 (2H, m), 5.12 (2H, s), 4.50-4.30 (2H, m), 4.00 (1H, dd, $J = 15.4, 6.1$ Hz), 3.88 (1H, dd, $J = 15.4, 6.3$ Hz), 3.55-3.40 (1H, m), 3.00 (3H, s), 2.55 (1H, t, $J = 12.8$ Hz), 2.45 (1H, t, $J = 12.8$ Hz), 2.40-2.10 (2H, m), 2.05 (3H, s), 2.00-1.90 (1H, m), 1.90-1.80 (1H, m), 1.60-1.50 (1H, br), 1.18 (1H, q, $J = 12.8$ Hz); HRMS calcd for $C_{21}H_{29}NO_7S$: 439.5318. Found: 439.5328.

(3S,5R,1'R) (-)-3-Acetoxymethyl-5-(1'-azido-but-3'-enyl)-piperidine-1-carboxylic acid benzyl ester
8

To a solution of **7** (1g, 2.3 mmol) in anhydrous DMF (30 mL), NaN_3 (224 mg, 3.4 mmol) was added under nitrogen atmosphere and the mixture was stirred at $80^\circ C$ for 2 h. The suspension was diluted with diethyl ether (30 mL) and added to water (30 mL). The aqueous layer was extracted with diethyl ether (3 x 20 mL). The combined organic layers were washed with brine and dried over Na_2SO_4 . Removal of the solvent in *vacuum* gave an oily residue which was purified by flash chromatography on silica gel (ethyl acetate/hexane 2:5), affording 765 mg (87%) of **8**, as an oil: R_f (ethyl acetate/hexane 2:5) 0.23; $[\alpha]_D^{25} - 29.0$ (c 1, $CHCl_3$); 1H NMR (300 MHz, $CDCl_3$) δ_H 7.35 (5H, br, s), 5.80-5.55 (1H, m), 5.17-5.10 (2H, m), 5.12 (2H, s), 4.40-4.15 (2H, m), 4.00 (1H, dd, $J = 12.1, 6.1$ Hz), 3.90 (1H, dd, $J = 12.1, 7.0$ Hz), 3.34-3.25 (1H, m), 2.65 (1H, t, $J = 12.3$ Hz), 2.55 (1H, t, $J = 12.3$ Hz), 2.45-2.34 (2H, m), 2.00 (3H, s), 1.90-1.65 (2H, m), 1.60-1.46 (1H, m), 1.10 (1H, q, $J = 13.1$ Hz); ^{13}C NMR (75.4 MHz, $CDCl_3$) δ_C 170.8, 155.1, 133.2, 127.0-130.0 (5C), 118.7, 68.1, 66.2, 64.3, 47.1, 45.6, 39.2, 35.5, 28.8, 20.6; HRMS calcd for $C_{20}H_{26}N_4O_4$: 386.4546. Found: 386.4561.

(3S,5R,1'R) (+)-3-Acetoxymethyl-5-(1'-acryloylamino-but-3'-enyl)-piperidine-1-carboxylic acid benzyl ester
4

To a solution of **8** (141 mg, 0.36 mmol) in THF (5 mL), triphenylphosphine (96 mg, 0.36 mmol) was added. After the reaction was complete (controlled by TLC), H_2O (330 μL , 18.3 mmol) was added. The mixture was stirred overnight, then concentrated in *vacuum* and the water removed by azeotropic distillation with toluene. The crude amine was dissolved in anhydrous CH_2Cl_2 (6 mL) under nitrogen atmosphere and cooled to $0^\circ C$. To this stirred solution Et_3N (112 μL , 0.80 mmol, d 0.7 g/mL) and DMAP (4 mg, 0.04 mmol) were added and acryloyl chloride (39 μL , 0.47 mmol, d 1.1 g/mL) was slowly dropped. After stirring at $0^\circ C$ for 30 min and at room temperature for 4h, the reaction mixture was washed with 1N HCl and then with 5% aq. $NaHCO_3$. Removal of the solvent under reduced pressure was followed by flash chromatography on silica gel (ethyl acetate/hexane 3:2) to yield 89 mg (59%, over two steps) of **4**, as an oil: R_f

(ethyl acetate/hexane 3:2) 0.29; $[\alpha]_D^{25} + 8.0$ (c 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ_H 7.30 (5H, br, s), 6.24 (1H, dd, $J = 16.0, 5.0$ Hz), 6.06 (1H, dd, $J = 16.0, 10.0$ Hz), 5.81-5.65 (1H, m), 5.63 (1H, dd, $J = 10.3, 5.0$ Hz), 5.35 (1H, br, d, $J = 9.4$ Hz), 5.17-5.02 (2H, m), 5.10 (2H, s), 4.36-4.18 (2H, m), 4.06-3.95 (1H, m), 3.86 (1H, dd, $J = 16.1, 7.0$ Hz), 3.83 (1H, dd, $J = 11.1, 7.1$ Hz), 2.55-2.30 (2H, m), 2.26-2.13 (2H, m), 2.01 (3H, s), 1.92-1.78 (1H, m), 1.72-1.55 (2H, m), 0.92 (1H, q, $J = 12.8$ Hz); ¹³C NMR (75.4 MHz, CDCl₃) δ_C 170.8, 165.3, 155.1, 133.4, 130.6, 126.0-128.2 (5C), 118.5, 67.2, 66.3, 50.6, 47.1, 46.6, 39.0, 36.4, 35.5, 29.7, 20.8; HRMS calcd for C₂₃H₃₀N₂O₅: 414.5060. Found: 414.5072.

(2R,3'R,5'S) (+)-5'-Acetoxymethyl-6-oxo-1,2,3,6,3',4',5',6'-octahydro-2'H-[2,3']-bipyridinyl-1'- carboxylic acid benzyl ester 3

To a stirred solution of **4** (247 mg, 0.6 mmol) in CH₂Cl₂ (10 mL) Cl₂Ru(CHPh)(P(*c*-hexyl)₃)₂ (24 mg, 0.03 mmol, 5% mol) was added under nitrogen atmosphere. The mixture was refluxed for 12h. DMSO (106 μ L, 1.5 mmol, d 1.1 g/mL) was added and the mixture was stirred overnight at room temperature. The solvent was removed in *vacuum* and the residue was purified by flash chromatography on silica gel (ethyl acetate), to yield 183 mg (79%) of **3**, as an oil: R_f (ethyl acetate) 0.15; $[\alpha]_D^{25} + 8.4$ (c 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ_H 7.35 (5H, br, s), 6.62 (1H, dt, $J = 11.6, 2.9$ Hz), 6.26 (1H, s), 5.91 (1H, d, $J = 9.9$ Hz), 5.12 (2H, s), 4.38-4.16 (2H, m), 4.04 (1H, dd, $J = 11.1, 5.8$ Hz), 3.87 (1H, dd, $J = 11.2, 6.9$ Hz), 3.56-3.41 (1H, m), 2.56-2.41 (2H, m), 2.40-2.23 (2H, m), 2.02 (3H, s), 1.85 (1H, d, $J = 11.6$ Hz), 1.81-1.68 (2H, m), 1.04 (1H, q, $J = 11.6$ Hz); ¹³C NMR (75.4 MHz, CDCl₃) δ_C 170.7, 166.3, 155.1, 140.2, 136.7, 127.1-128.2 (5C), 124.6, 67.2, 66.1, 53.3, 46.3, 45.9, 39.9, 35.6, 29.9, 26.9, 20.6; HRMS calcd for C₂₁H₂₆N₂O₅: 386.4518. Found: 386.4523.

(2R,3'R,5'S) (+)-5'-Hydroxymethyl-6-oxo-1,2,3,6,3',4',5',6'-octahydro-2'H-[2,3']-bipyridinyl-1'- carboxylic acid benzyl ester 9

To a solution of **3** (205 mg, 0.53 mmol) in THF (2 mL) NaOH 0.27 M (3 mL, 0.81 mmol) was added and the mixture was stirred for 2 h at room temperature. After removing THF in *vacuum*, water is added, and the aqueous phase was extracted with ethyl acetate. The combined organic layers were dried over Na₂SO₄ and the solvent was removed under reduced pressure to yield 179 mg (98 %) of **9**, as a foam: R_f (ethyl acetate) 0.08; $[\alpha]_D^{25} + 23.6$ (c 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ_H 7.32 (5H, br, s), 6.61-6.51 (1H, m), 6.28-6.12 (1H, br s), 5.89 (1H, d, $J = 7.8$ Hz), 5.12 (2H, s), 4.31-4.16 (2H, m), 3.55-3.37 (3H, m), 2.53-2.34 (2H, m), 2.35-2.19 (2H, m), 1.86 (1H, d, $J = 11.1$ Hz), 1.79-1.62 (2H, m), 1.03 (1H, q, $J = 11.1$ Hz); ¹³C NMR (75.4 MHz, CDCl₃) δ_C 166.7, 155.3, 140.7, 136.7, 127.8-128.5 (5C), 124.3, 67.2, 64.9, 53.2, 46.3, 45.9, 39.8, 38.5, 29.6, 26.9; HRMS calcd for C₁₉H₂₄N₂O₄: 344.4141. Found: 344.4120.

(2R,3'R,5'S) (+)-5'-Methanesulfonyloxymethyl-6-oxo-1,2,3,6,3',4',5',6'-octahydro-2'H-[2,3']bipyridinyl-1'- carboxylic acid benzyl ester 10

To a solution of **9** (160 mg, 0.46 mmol), Et₃N (323 μ L, 2.32 mmol, d 0.73 g/mL), DMAP (10 mg, 0.08 mmol) in anhydrous CH₂Cl₂ (10 mL), methanesulfonyl chloride (53 μ L, 0.69 mmol, d 1.5 g/mL) was added at 0°C under nitrogen atmosphere. The mixture was stirred for 1.5 h at room temperature, then washed with 5% aq. H₃PO₄ (2 x 5 mL). The

organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. Flash chromatography (ethyl acetate) gave 130 mg (67%) of **10**, as an oil: *R_f* (ethyl acetate) 0.40; [α]_D²⁵ + 3.2 (*c* 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ _H 7.35 (5H, br, s), 6.61 (1H, dt, *J* = 10.8, 5.4 Hz), 6.50 (1H, br, s), 6.12 (1H, br, s), 5.93 (1H, d, *J* = 10.8 Hz), 5.20 (2H, s), 4.31 (1H, d, *J* = 10.8 Hz), 4.23 (1H, d, *J* = 10.8 Hz), 4.18-4.01 (2H, m), 3.51-3.42 (1H, m), 3.01 (3H, s), 2.65-2.36 (2H, m), 2.36-2.25 (2H, m), 1.93 (1H, d, *J* = 12.4 Hz), 1.82-1.60 (2H, m), 1.10 (1H, q, *J* = 12.4 Hz); ¹³C NMR (75.4 MHz, CDCl₃) δ _C 169.8, 166.2, 155.1, 140.0, 136.7, 127.8-128.5 (5C), 123.9, 70.8, 67.2, 53.3, 45.9, 46.3, 39.7, 37.5, 35.9, 29.5, 26.9; HRMS calcd for C₂₀H₂₆N₂O₆S: 422.5040. Found: 422.5033.

(6*R*,7*R*,9*R*) (+)-*N*-Cbz-5,6-dihydro-cytisine **11**

To a suspension of sodium hydride (57 mg, 0.88 mmol, 80% dispersion in mineral oil) in anhydrous THF (4 mL) cooled to 0°C, a solution of **10** (186 mg, 0.44 mmol) in THF (4 mL) was added. After 15 minutes at 0°C, the reaction mixture was stirred at room temperature for 2 h. A pH 7 phosphate buffer solution was added, THF was evaporated and the aqueous layer was extracted with ethyl acetate. The combined organic extracts were dried over Na₂SO₄ and concentrated to give 128 mg (89%) of **11**, as an oil: *R_f* (ethyl acetate) 0.29; [α]_D + 17.4 (*c* 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ _H 7.30 (5H, br, s), 6.53-6.47 (1H, m), 5.95 (1H, dd, *J* = 10.1, 2.4 Hz), 5.20 (1H, d, *J* = 12.3 Hz), 5.10 (1H, d, *J* = 12.3 Hz), 4.39 (1H, dd, *J* = 15.2, 10.1 Hz), 4.19- 3.98 (2H, m), 3.63-3.52 (1H, m), 3.00-2.91 (2H, m), 2.88 (1H, d, *J* = 15.2 Hz), 2.38-2.22 (3H, m), 1.92 (1H, d, *J* = 13.5 Hz), 1.80 (1H, br, s), 1.54 (1H, d, *J* = 13.5 Hz); ¹³C NMR (75.4 MHz, CDCl₃) δ _C 170.7, 169.1, 155.1, 138.1, 137.5, 127.8-128.5 (5C), 125.7, 67.4, 55.9, 48.4, 49.9, 43.8, 34.5, 32.7, 25.7, 24.7; HRMS calcd for C₁₉H₂₂N₂O₃: 326.3988. Found: 326.3991.

(-)-*N*-Cbz-cytisine **12**

To a solution of DDQ (98 mg, 0.43 mmol) in dioxane (4 mL) a solution of **11** (140 mg, 0.43 mmol) in 4 mL dioxane was added, under nitrogen atmosphere. The mixture was stirred and refluxed at 110°C for 4h. Removal of the solvent under reduced pressure was followed by flash chromatography (ethyl acetate/hexane 3:1) to yield 41 mg (50%) of **12**, as an oil: *R_f* (ethyl acetate/hexane 3:1) 0.24; [α]_D²⁵ - 104.7 (*c* 1, CHCl₃); ¹H NMR (300 MHz, CDCl₃) δ _H 7.37-7.11 (6H, m), 6.44 (1H, d, *J* = 9.1 Hz), 6.01 (1H, d, *J* = 6.5 Hz), 5.02 (1H, d, *J* = 12.4 Hz), 4.97 (1H, d, *J* = 12.3 Hz), 4.38-4.22 (2H, m), 4.16 (1H, d, *J* = 15.8 Hz), 3.86 (1H, dd, *J* = 15.8, 7.2 Hz), 3.18-2.96 (3H, m), 2.48-2.38 (1H, br, s), 2.05-1.88 (2H, m); ¹³C NMR (75.4 MHz, CDCl₃) δ _C 163.4, 155.3, 148.8, 138.6, 136.3, 127.8-128.5 (5C), 117.4, 105.4, 67.4, 50.0, 51.1, 49.0, 34.5, 27.3, 25.8; HRMS calcd for C₁₉H₂₀N₂O₃: 324.3829. Found: 324.3837.

(-)-cytisine **1**

To a solution of **12** (110 mg, 0.34 mmol) in THF (6 mL), 6N HCl (2.2 mL) was added and the solution was refluxed for 3h. Then water (3 mL) was added and THF was removed in *vacuum*. The aqueous phase was basified with 1N NaOH (pH 9-10) and extracted with diethyl ether. The combined organic extracts were dried over Na₂SO₄ and the solvent was evaporated to yield 50 mg (78%) of **1**, as an oil: *R_f* (ethyl acetate) 0.02; [α]_D²⁵ - 114.2 (*c* 1, EtOH); ¹H NMR (300 MHz, CDCl₃) δ _H 7.26 (1H, dd, *J* = 9.1, 6.8 Hz), 6.41 (1H, dd, *J* = 9.1, 1.3 Hz), 5.88 (1H, dd, *J* = 6.8, 1.3 Hz), 4.10 (1H, d, *J* = 15.6 Hz), 3.86 (1H, ddd, *J* = 15.6, 6.6, 0.9 Hz), 3.07 (1H, dd, *J* = 12.4, 1.3 Hz), 3.04 (1H, dd, *J* = 12.1, 2.4 Hz), 2.99

(2H, br, d, $J = 12.3$ Hz), 2.86 (1H, br, s), 2.31 (1H, br, s), 1.94 (2H, t, $J = 3.2$ Hz), 1.51 (1H, s); ^{13}C NMR (75.4 MHz, CDCl_3) δ_{C} 163.7, 151.1, 138.8, 116.8, 104.9, 54.00, 53.00, 49.7, 35.6, 27.8, 26.3; HRMS calcd for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{O}$: 190.2470. Found: 190.2465.