Supporting Information

Total Synthesis of (–)-Callystatin A.

Neil F. Langille and James S. Panek*

*Department of Chemistry and Center for Chemical Methodology and Library
Development, Metcalf Center for Science and Engineering, Boston University,
590 Commonwealth Avenue, Boston, Massachusetts 02215*

Experimental section: ^1H and ^{13}C NMR spectra were taken in CDCl_3 at 400 MHz and 75 MHz, respectively. Chemical shifts are reported in parts per million using the solvent internal standard (chloroform, 7.24 and 77.0 ppm, respectively). Data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad, ABq = AB quartet), coupling constant. A dagger (†) denotes spectrum included at end of Supporting Information (p. 19-29). Diastereomeric ratios were determined by ^1H NMR (400 MHz) analysis of crude mixtures, operating at signal/noise ratio of $>200/1$. Infrared resonance spectra were recorded on a NexusTM 670 FTIR spectrometer. High resolution mass spectra were obtained on a FiniganTM MAT-90 spectrometer. Optical rotations were recorded on an AUTOPOL III digital polarimeter at 589 nm. CH_2Cl_2 , MeOH, PhMe, Et_2O , PhH, and THF were obtained from a dry solvent system (alumina) and used without further drying. *i*-PrOH was distilled from Mg containing a small amount of I_2 . Et_3N , pyridine, and 2,6-lutidine were distilled from and stored over KOH. TMSCl was distilled from quinoline prior to use. $(\text{COCl})_2$ was distilled immediately prior to use. DMSO and EtCN were distilled from CaH_2 . ZnCl_2 was fused by exposure to flame under reduced pressure prior to use. $\text{Pd}(\text{PPh}_3)_4$ was synthesized¹ by a known method. CrCl_2 was dried by brief exposure to flame under reduced pressure. TiCl_4 was distilled from Cu powder immediately prior to use. All other reagents were used as supplied, unless otherwise noted. All reactions were carried out in oven-dried glassware, under a blanket of argon unless otherwise noted. Analytical thin layer chromatography was performed on Sorbent Technologies 0.20 mm silica gel 60 Å plates. Normal phase flash chromatography² was performed on Sorbent Technologies 32-63 μm 60 Å silica gel. Reverse phase flash chromatography³ was performed on C18-functionalized silica gel (Aldrich). Melting points are uncorrected.

Experimental Procedures and Spectroscopic Characterization of Novel Compounds

(1) Hegedus, L. S. In *Organometallics in Synthesis: A Manual*; Schlosser, M., Ed.; John Wiley & Sons Ltd.: New York, 1994; Chapter 5, pp. 448.

(2) Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, 43, 2923-2925.

(3) Farina, V. *J. Org. Chem.* **1991**, 56, 4985-4987.

(4) Journet, M.; Cai, D.; DiMichele, L. M.; Larsen, R. D. *Tetrahedron Lett.* **1998**, 39, 6427-6428.

the layers were separated. The aqueous layer was extracted with Et₂O (3×). The combined organic layers were washed with NaCl, dried over MgSO₄, filtered, and concentrated to yield 10.9 g (>99% yield) desired alcohol as a pale yellow oil, which was sufficiently pure and used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 1.04-1.07 (21H, m), 1.82 (1H, d, J=6.3, exchangeable), 2.47 (2H, m), 4.43 (1H, dd, J=5.9, 11.9), 5.15 (1H, d, J=1.3), 5.19 (1H, dd, J=1.3, 8.2), 5.88 (1H, m); ¹³C NMR (75 MHz, CDCl₃) δ 11.1, 18.6, 42.4, 62.1, 86.0, 108.0, 118.9, 133.0; IR (neat) ν_{max} 3560, 2960, 2870, 1734, 1464, 1405, 1293, 1180, 1047, 985; HRMS(EI) *m/z* calc'd for C₁₅H₂₈O₁Si₁ [M]⁺ 252.1909, found: 252.1845.

(*R*)-1-(Triisopropylsilyl)hex-5-en-1-yn-3-yl acrylate (**5**) and (*S*)-1-(triisopropylsilyl)hex-5-en-1-yn-3-ol ((-)-**10**). To a flask charged with racemic alcohol **10** (10.9 g, 43.3 mmol) and hexanes (215 mL, 0.200 M) at rt were added vinyl acrylate (12.5 mL, 120 mmol, 2.77 equiv.) and lipase Amano AK (5.46 g, 0.500 wt. equiv.). The mixture was stirred vigorously at rt, and monitored by ¹H NMR analysis. After 7 days,⁵ the reaction mixture was filtered, rinsed with Et₂O, and concentrated.⁶ Flash chromatography (silica, 1% EtOAc/hexanes) provides 6.10 g (46.0%) desired acrylate (**5**) as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 1.04 (21H, m), 2.55 (2H, dd, J=6.3, 6.9), 5.13 (2H, m), 5.50 (1H, dd, J=6.3, 6.9), 5.83 (2H, m), 6.11 (1H, m), 6.41 (1H, dd, J=1.0, 17.1); ¹³C NMR (75 MHz, CDCl₃) δ 11.0, 18.5, 39.5, 63.9, 87.3, 103.9, 118.7, 128.2, 131.3, 132.2, 164.9; IR (neat) ν_{max} 2944, 2893, 2866, 1734, 1464, 1405, 1294, 1260, 1181, 1042, 988; HRMS(CI, NH₃) *m/z* calc'd for C₁₅H₂₈O₁Si₁ [M+H]⁺ 307.2093, found: 307.2127; [α]_D = + 70.1 (c=0.11, CHCl₃). Chiral HPLC of the derived benzyl ether⁷ (Chiracel OD, hexanes eluent, r.t.(*R*) = 18.3 min, r.t.(*S*) = 19.1 min): >95% ee. Continued column elution (2→4% EtOAc/hexanes) provides 4.81 g (44.1%) (-)-**10**.⁸

(5) Experimentally, the reaction time varies from 5 days to 10 days, depending on the activity of the lipase sample. Monitoring the reaction by ¹H NMR (δ 5.50 in **5** (-CH₂OAc-) vs. δ 4.43 in (-)-**10** (-CH₂OH-)) ensures complete reaction with no over-resolution. No variation in stereochemical outcome has been observed in this reaction.

(6) After filtration, lipase *Pseudomonas* AK (Amano) can be used repeatedly with minimal loss of enzymatic activity over several months when stored at 4 °C.

(7) Racemic **5**, although UV active, proved impossible to resolve by chiral HPLC. The benzyl ether was synthesized (a. K₂CO₃, MeOH; b. NaH, BnBr, Bu₄NI, Et₂O) and analyzed by HPLC as above. Although none of the opposite enantiomer was observed by HPLC analysis, we only feel confident reporting >95% ee, due to the transformations required for analysis.

(8) HPLC analysis as per ref. 7 indicates >95% ee of (*S*)-enantiomer.

Stereochemical assignment of (*R*)-acrylate **5 and (*S*)-alcohol (**-**)-**10**.** To a sample of (*R*)-acrylate **5** (61.0 mg, 0.200 mmol) in MeOH (2.00 mL, 0.100 M) at rt was added K₂CO₃ (127 mg, 1.20 mmol, 6.00 equiv.), and the reaction was stirred 18 h. The reaction was quenched with 10% HCl to a pH of 5, and CH₂Cl₂ was added. The layers were separated, and the aqueous layer was extracted into CH₂Cl₂ (3×). The organic layers were washed with NaCl, dried over MgSO₄, filtered and concentrated to yield 50.2 mg desired (*R*)-alcohol (99.6%). The alcohol was dissolved in 2.00 mL CH₂Cl₂ (0.100 M), and cooled to 0 °C. To the mixture was added (*R*)-O-acetyl mandelic acid (26.0 mg, 0.218 mmol, 1.10 equiv.), DCC (45.0 mg, 0.218 mmol, 1.10 equiv.) and DMAP (2.40 mg, 0.0198 mmol, 10.0 mol%). The mixture was allowed to warm to rt, and stirred 36 h. The mixture was cooled to 0 °C, filtered, and concentrated. Purification by flash chromatography (silica, 0→1% EtOAc/hexane) provides 63.0 mg (73.7%) desired (*R*)-OAc-mandelate ester of (*R*)-alcohol as a colorless oil. Similar treatment of 50.0 mg (*S*)-alcohol (**-**)-**10** provides 67.0 mg (78.2%) (*R*)-OAc-mandelate ester of (*S*)-alcohol. ¹H NMR analysis⁹ of each compound confirms stereochemical assignment (key resonances are illustrated above and underlined below, with critical Δδ(*R,R*-ester – *R,S*-ester) values in **bold**).

- (*R*)-OAc-Mandelate ester of (*R*)-alcohol (**+**)-**10**: ¹H NMR† (400 MHz, CDCl₃) δ 0.95 (21H, s, Δδ_{*R,S*} = **-0.08**), 2.17 (3H, s), 2.52 (2H, m, Δδ_{*R,S*} = **+0.11**), 5.13 (2H, m, Δδ_{*R,S*} = **+0.18**), 5.41 (1H, dd, *J*=5.6, 6.6), 5.78 (1H, m, Δδ_{*R,S*} = **+0.17**), 5.89 (1H, s), 7.33-7.46 (5H, m).

- (*R*)-OAc-Mandelate ester of (*S*)-alcohol (**-**)-**10**: ¹H NMR† (400 MHz, CDCl₃) δ 1.03 (21H, s), 2.16 (3H, s), 2.41 (2H, m), 4.95 (2H, m), 5.42 (1H, m), 5.61 (1H, m), 5.93 (1H, s), 7.34-7.46 (5H, m).

(*R*)-6-((Triisopropylsilyl)ethynyl)-5,6-dihydropyran-2-one (11**).**

A sample of diene-yne **5** (5.00 g, 16.3 mmol) was dissolved in CH₂Cl₂ (327 mL, 0.050 M) and heated to reflux. To the mixture was added Grubbs I catalyst (721 mg, 0.817 mmol, 5.00 mol %), and heating was continued. After 2 h, an additional portion of Grubbs I (721 mg, 5.00 mol %) was added. The reaction was heated at reflux for 36 h, and then cooled to rt. Air was bubbled into the solution for 1 h, then the mixture was concentrated. Purification by flash chromatography (silica, 10% EtOAc/hexanes) provides 3.77 g (83.1%) desired lactone **11** as a dark brown oil.¹⁰ ¹H NMR (400 MHz, CDCl₃) δ 1.02 (21H, m), 2.66 (2H, m), 5.18 (1H, dd, *J*=5.6, 7.2), 6.04 (1H, d, *J*=10.0), 6.85 (1H, m); ¹³C NMR (75 MHz, CDCl₃)

(9) Whitesell, J. K.; Reynolds, D. *J. Org. Chem.* **1983**, 48, 3548-3551.

(10) Ruthenium content in lactone **11** was not measured.

δ 10.9, 18.4, 30.6, 67.4, 88.6, 102.4, 121.5, 143.9, 162.7; IR (neat) ν_{max} 3045, 2970, 2867, 1733, 1410, 1294, 1180, 989; HRMS(Cl, NH_3) m/z calc'd for $\text{C}_{16}\text{H}_{27}\text{O}_2\text{Si}_1$ $[\text{M}+\text{H}]^+$ 279.1780, found: 279.1771; $[\alpha]_{\text{D}} = +68.5$ ($c=0.30$, CHCl_3).

((*(2R,6R)*-6-isopropoxy-3,6-dihydro-2H-pyran-2-yl)ethynyl)triisopropylsilane (**12**). To a sample of lactone **11** (5.00 g, 18.0 mmol) in CH_2Cl_2 (180 mL, 0.100 M) at -78°C was added 1.00 M DIBAL-H/hexanes (19.8 mL, 1.10 equiv.), dropwise over 1 min. The reaction was stirred at -78°C 15 min, then quenched with 20 mL H_2O . After warming to rt, the mixture was extracted into CH_2Cl_2 (5 $\times$), dried over MgSO_4 , and concentrated. The crude residue was dissolved in PhH (180 mL, 0.100 M), and then PPTS (904 mg, 3.60 mmol, 20.0 mol %) and *i*-PrOH (27.4 mL, 360 mmol, 20.0 equiv.) were added. The reaction mixture was heated to 70°C for 18 h, then concentrated *in vacuo*. Purification by flash chromatography (2 $\rightarrow$ 4% EtOAc/hexanes) provides 4.76 g (82.0%) desired lactol **12** as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 1.05 (21H, m), 1.16 (3H, d, $J=6.4$), 1.25 (3H, d, $J=6.4$), 2.28 (2H, m), 4.02 (1H, qq, $J=6.4$, 6.4), 4.69 (1H, dd, $J=3.8$, 11.2), 5.11 (1H, s), 5.67 (1H, d, $J=9.6$), 5.92 (1H, m); ^{13}C NMR (75 MHz, CDCl_3) δ 11.0, 18.4, 22.1, 23.6, 31.7, 57.6, 70.1, 93.3, 106.6, 126.1, 128.0; IR (neat) ν_{max} 3046, 2972, 2931, 2850, 1745, 1402, 1318, 1182, 1100, 1024, 946; HRMS(EI) m/z calc'd for $\text{C}_{19}\text{H}_{34}\text{O}_2\text{Si}_1$ $[\text{M}]^+$ 322.2328, found: 322.2341; $[\alpha]_{\text{D}} = +23.6$ ($c=0.35$, CHCl_3).

(*(2R,6R)*-2-ethynyl-6-isopropoxy-3,6-dihydro-2H-pyran (**2**). In a round-bottomed flask were combined TBAF (1.00 M solution in THF, 7.00 mL) and glacial acetic acid (0.51 mL, 9.10 mmol, 1.30 equiv. vs. TBAF). The resulting solution was stirred 15 min. Separately, lactol **12** (1.00 g, 3.10 mmol) was dissolved in THF (24.8 mL), and the solution was stirred at rt under argon. A portion of the AcOH/TBAF solution (6.20 mL, 2.00 equiv.; total reaction volume = 31.0 mL, 0.100 M) was added dropwise, and the reaction was stirred at rt. After 24 h, the reaction was quenched with H_2O , extracted into Et_2O (3 $\times$), washed with brine, dried over MgSO_4 , filtered and concentrated carefully *in vacuo* at 0°C . Flash chromatography (silica, 4 $\rightarrow$ 6% Et_2O /pentane) provides 470 mg (91.2%) desired **2** as a volatile yellow oil. ^1H NMR † (400 MHz, CDCl_3) δ 1.15 (3H, d, $J=6.0$), 1.24 (3H, d, $J=6.0$), 2.18 (1H, ddd, $J=1.2$, 3.6, 17.6), 2.36 (1H, dddd, $J=2.8$, 5.6, 10.8, 17.6), 2.43 (1H, d, $J=2.0$), 4.02 (1H, qq, $J=6.0$, 6.0), 4.66 (1H, ddd, $J=2.0$, 3.6, 10.8), 5.09 (1H, brs), 5.67 (1H, ddd, $J=1.2$, 2.8, 10.0), 5.92 (1H, dd, $J=5.6$, 10.0); ^{13}C NMR (75 MHz, CDCl_3) δ 21.8, 23.7, 31.1, 56.9, 69.9, 72.6, 82.7, 92.9, 126.0, 127.6; IR (neat) ν_{max} 3293, 3048, 2972, 2931, 2905, 1659, 1402, 1318, 1182, 1126, 1100, 1024, 1000, 946; HRMS(EI) m/z calc'd for $\text{C}_{10}\text{H}_{14}\text{O}_2$ $[\text{M}]^+$ 166.0994, found: 166.0991; $[\alpha]_{\text{D}} = +83.8$ ($c=0.14$, CHCl_3).

(*R*)-1-Benzyloxy-5,5-dibromo-3-methyl-4-pentene. To a flask charged with PPh_3 (34.5 g, 132 mmol, 4.00 equiv.) and CH_2Cl_2 (100 mL) at 0°C was added CBr_4 (21.8 g, 65.7 mmol, 2.00 equiv.), and the reaction was stirred 10 min at 0°C . To the reaction mixture was added, by cannula, a

solution of aldehyde **16**¹¹ (6.31 g, 32.9 mmol) in CH₂Cl₂ (100 mL), followed by addition of 2,6-lutidine (7.66 mL, 65.7 mmol, 2.00 equiv.). The reaction mixture was allowed to warm to rt, then stirred 2 h at rt. The reaction was quenched with sat. aq. NH₄Cl and EtOAc was added. The mixture was decanted, the layers were separated, and the aqueous layer was extracted into EtOAc (4×). The combined organic layers were washed with H₂O, brine, dried over MgSO₄, filtered, and concentrated. Purification by flash chromatography (silica, 2→4% EtOAc/hexanes) provides 11.0 g (96.1%) desired dibromide as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 1.02 (3H, d, J=6.9), 1.65 (2H, m), 2.67 (1H, m), 3.46 (2H, m), 4.48 (2H, ABq, J=11.5), 6.19 (1H, d, J=9.6), 7.24-7.34 (5H, m); ¹³C NMR (75 MHz, CDCl₃) δ 19.3, 35.6, 36.0, 68.1, 73.1, 87.8, 127.7, 128.4, 138.5, 143.8; IR (neat) ν_{max} 3036, 3030, 2962, 2927, 2867, 1453, 1363, 1105, 697; HRMS(EI) *m/z* calc'd for C₁₃H₁₆Br₂O₁ [M]⁺ 345.9568, found: 345.9583; [α]_D = -23.4 (c=0.23, CHCl₃).

(*R*)-(5-(Benzyloxy)-3-methylpent-1-ynyl)trimethylsilane (**6**). A sample of dibromide (11.3 g, 32.4 mmol) in THF (162 mL, 0.200 M) was cooled to -78 °C, and *n*-BuLi (2.50 M in hexanes, 28.6 mL, 71.4 mmol, 2.20 equiv.) was added dropwise over 2 min. To the reaction mixture was added TMSCl (10.3 mL, 81.2 mmol, 2.50 equiv.). After 5 min at -78 °C, the reaction was warmed to rt, stirred 30 min. After quenching the reaction with H₂O and extracting into EtOAc (3×), washing the organic layers with H₂O, brine, drying with MgSO₄, filtering and concentrating *in vacuo* afforded 8.27 g (98.0%) desired alkyne **6**, which was sufficiently pure and required no further purification. ¹H NMR (400 MHz, CDCl₃) δ 0.11 (9H, s), 1.16 (3H, d, J=6.9), 1.69 (2H, m), 2.67 (2H, m), 3.59 (2H, dd, J=5.9, 6.3), 4.50 (2H, s), 7.26-7.33 (5H, m); ¹³C NMR (75 MHz, CDCl₃) δ 0.2, 21.0, 23.8, 36.8, 68.3, 73.0, 84.4, 111.3, 127.5, 127.6, 128.4, 138.6; IR (neat) ν_{max} 3064, 3030, 2961, 2859, 2167, 1453, 1249, 1097, 841; HRMS(EI) *m/z* calc'd for C₁₆H₂₄O₁Si₁ [M]⁺ 260.1596, found: 260.1607; [α]_D = -11.0 (c=0.3, CH₂Cl₂).

(*R,E*)-(5-(Benzyloxy)-1-iodo-3-methylpent-1-enyl)trimethylsilane (**17**). A sample of Cp₂ZrHCl (11.9 g, 46.2 mmol, 2.50 equiv.) was placed in an argon-filled flask. A solution of alkyne **6** (4.80 g, 18.5 mmol) in THF (462 mL, 0.0400 M) was added by cannula, the mixture was stirred vigorously. After 5 min, the mixture was heated to 55 °C to yield a clear yellow solution. After 60 min, the reaction was cooled briefly to rt, and a solution of I₂ (6.09 g, 24.0 mmol, 1.30 equiv.) in THF (46.2 mL, 0.400 M) was added by cannula, and the reaction was protected from light. After consumption of the vinyl zirconium species (~30 min by TLC), the reaction was quenched with sat. aq. Na₂S₂O₃ and pentane was added to initiate crystallization of residual zirconium reagent. The mixture was decanted, then extracted into pentane (4×). The organic layers were combined, washed with H₂O, brine, dried with MgSO₄, filtered, and concentrated. ¹H NMR analysis of the

(11) Myers, A. G.; Yang, B. H.; Chen, H.; McKinstry, L.; Kopecky, D. J.; Gleason, J. L. *J. Am. Chem. Soc.* **1997**, *119*, 6496-6511.

crude residue indicated >20:1 crude regioselectivity. Flash chromatography (silica, 0→1% EtOAc/hexanes) affords the desired vinyl iodide **17** as a yellow oil (6.37 g, 88.9% yield). ¹H NMR (400 MHz, CDCl₃) δ 0.25 (9H, s), 0.96 (3H, d, J=6.9), 1.56 (2H, m), 2.60 (1H, m), 3.43 (2H, m), 4.46 (2H, m), 6.92 (1H, d, J=10.9), 7.31 (5H, m); ¹³C NMR (75 MHz, CDCl₃) δ 1.2, 20.5, 36.6, 36.8, 68.1, 73.0, 106.1, 127.5, 127.6, 128.3, 138.5, 161.6; IR (neat) ν_{max} 3029, 2956, 2865, 2792, 1453, 1362, 1248, 1103, 841, 697; HRMS(EI) *m/z* calc'd for C₁₆H₂₅I₁O₁Si₁ [M]⁺ 388.0719, found: 388.0674; [α]_D = +20.2 (c=0.20, CH₂Cl₂).

(*R,Z*)-7-(Benzyloxy)-5-methylhept-3-en-3-yl)trimethylsilane (**18**). A sample of fused ZnCl₂ (12.6 g, 92.3 mmol, 10.0 equiv.) was dissolved in THF (114 mL),¹² and cooled to

0 °C. To the reaction mixture was added EtMgBr (3.0 M solution in Et₂O, 31.8 mL, 92.3 mmol) dropwise over 30 min. The ice bath was removed, and the reaction was stirred 1 h at rt to generate 92.3 mmol EtZnX solution. Separately, iodide **17** (3.58 g, 9.23 mmol), THF (20.0 mL), and Pd(PPh₃)₄ (533 mg, 0.462 mmol, 5.00 mol %) were combined at rt. To the iodide/Pd mixture was added 1/2 decanted EtZnX solution (73.0 mL, 46.2 mmol, 5.00 equiv.) dropwise at rt. After consumption of the vinyl iodide by TLC (~30 min), the reaction was quenched with H₂O, and the layers were separated. The aqueous layer was extracted into EtOAc (3×). The combined organic layers were washed with H₂O, brine, dried over MgSO₄, filtered and concentrated. Purification by flash chromatography (silica, 2% EtOAc/hexanes) provides 2.57 g (95.8%) **18** as a pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 0.11 (9H, s), 0.90-0.94 (6H, m), 1.56 (2H, m), 2.01 (2H, m), 2.46 (1H, m), 3.42 (2H, dd, J=5.6, 7.9), 4.45 (2H, ABq, J=11.9), 5.64 (1H, d, J=10.2), 7.23-7.32 (5H, m); ¹³C NMR (75 MHz, CDCl₃) δ 0.5, 15.6, 21.4, 30.9, 33.0, 37.4, 68.8, 72.9, 127.4, 127.6, 128.3, 138.7, 139.5, 147.6; IR (neat) ν_{max} 3030, 2958, 2927, 2867, 2361, 1612, 1453, 1248, 1101, 836; HRMS(CI, NH₃) *m/z* calc'd for C₁₈H₃₁O₁Si₁ [M+H]⁺ 291.2144, found: 291.2110; [α]_D = +14.8 (c=0.2, CH₂Cl₂).

(*R,Z*)-3-Methyl-5-(trimethylsilyl)hept-4-en-1-ol. To a sample of benzyl ether **18** (3.64 g, 14.0 mmol) was added CH₂Cl₂ (46.7 mL) and H₂O (23.3 mL, 0.200 M total). To the mixture

was added DDQ (6.36 g, 28.0 mmol, 2.00 equiv.), and the mixture was stirred vigorously for 18 h. The reaction was quenched with 5% aq. NaHCO₃, and the mixture was extracted into CH₂Cl₂ (3×). The combined organic layers were washed with H₂O, brine, dried over MgSO₄, filtered and concentrated. Purification by flash chromatography (10→15% EtOAc/hexanes) yields 2.33 g (83.1%) desired alcohol as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 0.13 (9H, s), 0.91-0.98 (6H, m), 1.29 (1H, s), 1.55 (2H, m), 2.03 (2H, m), 2.48 (2H, m), 3.62 (2H, m), 5.70 (1H, d, J=10.5); ¹³C NMR (75 MHz, CDCl₃) δ 0.3, 15.4, 21.5, 30.9, 33.2, 40.2, 61.5, 139.9, 147.6; IR (neat) ν_{max} 3650, 2962, 2930, 2320, 1650, 1453, 1248, 1090; HRMS(CI, NH₃) *m/z* calc'd for C₁₁H₂₅O₁Si₁ [M+H]⁺ 201.1675, found: 201.1670; [α]_D = -23.1 (c=0.30, CHCl₃).

(12) On large scale (>3.00 g ZnCl₂), sonication is frequently required to facilitate dissolution.

(*R,Z*)-3-Methyl-5-(trimethylsilyl)hept-4-enal (**19**).

To a $-78\text{ }^{\circ}\text{C}$ flask charged with CH_2Cl_2 (74.1 mL) and $(\text{COCl})_2$ (1.00 mL, 11.5 mmol, 1.15 equiv.) was added DMSO (1.58 mL, 22.0 mmol, 2.20 equiv.) dropwise over 1 min. After stirring 20 min at $-78\text{ }^{\circ}\text{C}$, primary alcohol (2.00 g, 9.98 mmol) in CH_2Cl_2 (36.7 mL, 0.0900 M total) was added by cannula. After stirring 20 min at $-78\text{ }^{\circ}\text{C}$, Et_3N (6.97 mL, 49.9 mmol, 5.00 equiv.) was added dropwise. The mixture was stirred 20 min at $-78\text{ }^{\circ}\text{C}$, then warmed to rt over 1 h. The mixture was quenched with H_2O , and extracted into CH_2Cl_2 (3 $\times$). The combined organic layers were washed with H_2O , brine, dried over MgSO_4 , filtered, and were concentrated. The crude residue was washed with a cold solution of 10% EtOAc/hexanes ($0\text{ }^{\circ}\text{C}$) and filtered through cotton. Further purification by flash chromatography (silica, 5% EtOAc/hexanes) provided desired aldehyde **19** as a yellow oil (1.82 g, 91.9%). ^1H NMR (400 MHz, CDCl_3) δ 0.12 (9H, s), 0.91 (3H, d, $J=7.5$), 1.00 (3H, d, $J=6.6$), 2.02 (2H, q, $J=7.5$), 2.32 (2H, dd, $J=2.5$, 6.8), 2.92 (1H, dtq, $J=6.6$, 6.8, 10.6), 5.69 (1H, d, $J=10.6$), 9.68 (1H, t, $J=2.5$); ^{13}C NMR (75 MHz, CDCl_3) δ 0.2, 15.2, 21.2, 30.7, 31.4, 50.9, 141.1, 144.9, 202.6; IR (neat) ν_{max} 2962, 2325, 1630, 1415, 1248, 1090, 890; HRMS(CI, NH_3) m/z calc'd for $\text{C}_{11}\text{H}_{23}\text{O}_1\text{Si}_1$ $[\text{M}+\text{H}]^+$ 199.1518, found: 199.1479; $[\alpha]_{\text{D}} = -18.6$ ($c=0.20$, CHCl_3).

Trimethyl((*R,3Z,7E*)-5-methyl-8-(tributylstannyl)octa-3,7-dien-3-yl)silane (**3**). A sample of CrCl_2 (6.20 g, 50.4 mmol, 10.0 equiv.) was flame-dried under reduced

pressure, and argon was introduced into the flask. The CrCl_2 sample was cooled to $0\text{ }^{\circ}\text{C}$, then degassed DMF (79.1 mL) was added dropwise over 10 min. At the completion of addition, the chromium/DMF flask was warmed to rt over 20 min. Separately, aldehyde **19** (1.00 g, 5.04 mmol), freshly prepared Bu_3SnCH_2 ¹³ (5.61 g, 10.1 mmol, 2.00 equiv.), and degassed DMF (10.9 mL) were combined. The aldehyde/Sn solution was added to the chromium mixture by cannula, followed by a DMF rinse (10.9 mL). The reaction mixture was stirred at rt 24 h (protected from light), until complete aldehyde consumption was achieved, as measured by TLC. The reaction was quenched with H_2O , and Et_2O was added. The aqueous layer was extracted into Et_2O (4 $\times$). The combined organic layers were washed with H_2O , brine, dried with MgSO_4 , filtered and concentrated. Analysis of the crude material revealed one olefin isomer (^1H NMR, 400 MHz, vinyl proton region). Purification by reverse phase chromatography³ ($\text{C}_{18}\text{-SiO}_2$, 5 $\rightarrow$ 40% $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{CN}$) provides desired vinyl stannane **3** as a yellow oil (1.66 g, 67.8%). ^1H NMR[†] (400 MHz, CDCl_3) δ 0.11 (9H, s), 0.74-0.94 (21H, m), 1.29 (6H, m), 1.46 (6H, m), 2.02 (2H, m), 2.07 (2H, m), 2.41 (1H, m), 5.68 (1H, d, $J=10.8$), 5.75-5.95 (2H, m); ^{13}C NMR (75 MHz, CDCl_3) δ 0.5, 9.2, 13.6, 15.6, 20.7, 27.2, 29.1, 30.8, 36.0, 46.1, 128.9, 139.1, 147.7, 147.8; IR (neat) ν_{max} 2956, 2926, 2871, 2843, 1464, 1376, 1368, 1248, 1071, 836;

(13) In accordance with literature reports, Bu_3SnCH_2 reactivity decreases within two weeks of preparation. However, precursor $\text{Bu}_3\text{SnCHBr}_2$ is stable at $-20\text{ }^{\circ}\text{C}$ for months. For details, see: Hodgson, D. M.; Foley, A. M.; Boulton, L. T.; Lovell, P. J.; Maw, G. N. *J. Chem. Soc., Perkin Trans. 1*, **1999**, 2911-2922.

HRMS (CI, NH₃) m/z calc'd for C₂₆H₅₉Si₁Sn₁ [M+H]⁺ 519.3408, found: 519.3420; [α]_D = -18.2 (c=0.15, CHCl₃).

(5*R*,6*R*,7*S*,*E*)-Methyl 6-hydroxy-5,7-dimethylnon-3-enoate (**20**). In a flask were combined (*S*)-2-methylbutanal (**8**)¹⁴ (6.84 g, 79.4 mmol), silane (**R**)-**7** (25.0 g, 95.4 mmol, 1.20 equiv.), and (265 mL, 0.300 M) CH₂Cl₂. The mixture was cooled to -78 °C, and then TiCl₄ (17.4 mL, 159 mmol, 2.00 equiv.) was added dropwise. Following addition, the reaction was warmed slowly to -30 °C. After 24 h at -30 °C, 20 mL EtOAc was added, followed by 20 mL H₂O. The mixture was allowed to warm to rt, then 10% aq. NaHCO₃ and additional EtOAc were added. Following separation of the layers, the aqueous layer was extracted into EtOAc (3×). The combined organic layers were washed with H₂O, brine, dried over MgSO₄, filtered and concentrated. ¹H NMR analysis of the crude residue revealed a 10:1 mixture of diastereomers.¹⁵ Purification by flash chromatography (silica, 10→50% EtOAc/hexanes) yields 14.3 g (84.0%) desired alcohol **20** as a single diastereomer. ¹H NMR (400 MHz, CDCl₃) δ 0.85 (3H, dd, J=6.6, 7.6), 0.88 (3H, d, J=6.8), 1.03 (3H, d, J=6.9), 1.20 (1H, ddq, J=1.6, 7.6, 7.2), 1.38 (1H, ddq, J=1.6, 6.6, 7.1), 1.47 (1H, dddq, J=6.8, 7.1, 7.2, 4.6), 1.62 (1H, bs), 2.33 (1H, ddq, J=6.9, 7.3, 7.9), 3.03 (2H, d, J=6.6), 3.24 (1H, dd, J=4.6, 7.3), 3.66 (3H, s), 5.43 (1H, dd, J=7.9, 15.5), 5.56 (1H, dt, J=6.6, 15.5); ¹³C NMR (75 MHz, CDCl₃) δ 11.6, 12.8, 16.6, 26.6, 36.9, 37.9, 40.1, 51.8, 77.8, 121.8, 137.4, 172.4; IR (neat) ν_{max} 3605, 2965, 2879, 1750, 1457, 1376, 1244, 1168, 1098, 974; HRMS(EI) m/z calc'd for C₁₂H₂₂O₃ [M]⁺ 214.1569, found: 214.1576; [α]_D = +12.4 (c=0.30, CHCl₃).

(5*R*,6*R*,7*S*,*E*)-Methyl 6-acetoxy-5,7-dimethylnon-3-enoate. To a flask charged with argon were added alcohol **20** (14.3 g, 66.7 mmol), CH₂Cl₂ (333 mL, 0.200 M), and pyridine (27.0 mL, 334 mmol, 5.00 equiv.), and the mixture was cooled to 0 °C. Ac₂O (25.2 mL, 227 mmol, 4.00 equiv.) was added by syringe, followed by DMAP (407 mg, 5.00 mol %). The reaction mixture was allowed to warm to rt, and stirred until consumption of alcohol starting material (~2 h). H₂O was introduced to the flask, and the resulting aqueous layer was extracted into CH₂Cl₂ (3×). The combined organic layers were washed with H₂O, brine, dried over MgSO₄, filtered and concentrated. Purification by flash chromatography (silica, 8% EtOAc/hexanes) provides desired acetate in 98.8% yield (16.9 g). ¹H NMR (400 MHz, CDCl₃) δ 0.83 (3H, d, J=6.8), 0.86 (3H, dd, J=7.2, 7.2), 0.93 (3H, d, J=6.9), 1.10 (1H, m), 1.27 (1H, m), 1.57 (1H, m), 2.04 (3H, s), 2.46 (1H, ddq, J=6.9, 7.8, 8.8), 3.02 (2H, d, J=6.9), 3.66 (3H, s), 4.77 (1H, dd, J=3.6, 8.8), 5.38 (1H, dd, J=15.6, 7.8), 5.57 (1H, dt, J=6.9, 15.6); ¹³C NMR (75 MHz, CDCl₃) δ 11.1, 12.5, 16.0, 20.2, 26.2, 35.7, 37.2, 38.4, 51.1, 78.1, 122.0, 135.5, 170.3, 171.5; IR (neat) ν_{max} 2950, 2880, 1738, 1668, 1437,

(14) White, J. D.; Bolton, G. L.; Dantanarayan, A. P.; Fox, C. M. J.; Hiner, R. N.; Jackson, R. W.; Sakuma, K.; Warrier, U. S. *J. Am. Chem. Soc.* **1995**, *117*, 1908-1939.

(15) Although not characterized, precedent indicates that the minor product is (3*E*, 5*R*, 6*S*, 7*S*)-**20**.

1376, 1244, 1168, 1098, 1020, 975; HRMS(EI) m/z calc'd for $C_{14}H_{24}O_4$ $[M]^+$ 256.1675, found: 256.1665; $[\alpha]_D = +10.8$ ($c=0.20$, $CHCl_3$).

(2*S*,3*R*,4*S*)-3-Acetoxy-2,4-dimethylhexanal (**21b**). A flask was charged with olefin (11.0 g, 42.8 mmol), pyridine (3.46 mL, 42.8 mmol, 1.00 equiv.), and 1:1 MeOH: CH_2Cl_2 (534 mL, 0.0800 M),

purged with argon, and cooled to $-78\text{ }^\circ\text{C}$. O_3 gas was bubbled into the solution until starting material was consumed and a blue color persisted. Me_2S (94.0 mL, 1280 mmol, 30.0 equiv.) was added to the flask, and the reaction was allowed to warm slowly to rt over a period of 18 h. The crude mixture was concentrated *in vacuo*, and EtOAc (750 mL) was added. The organic layer was washed with H_2O (2 $\times$), brine (1 $\times$), dried over $MgSO_4$, filtered and concentrated. The crude material (8.33 g, 105% theoretical yield) was tentatively assigned by 1H and ^{13}C NMR analysis and used without further purification. 1H NMR (400 MHz, $CDCl_3$) δ 0.90 (6H, m), 1.06 (3H, d, $J=6.9$), 1.17 (H, m), 1.40 (1H, m), 1.66 (1H, m), 2.03 (3H, s), 2.67 (1H, ddq, $J=5.2, 1.0, 6.9$), 5.22 (1H, dd, $J=6.8, 5.2$), 9.65 (1H, d, $J=1.0$); ^{13}C NMR (75 MHz, $CDCl_3$) δ 8.2, 11.1, 14.1, 20.6, 25.8, 36.3, 48.5, 74.9, 170.1, 202.6.

(5*R*,6*S*,7*R*,8*R*,9*S*,*E*)-Methyl 8-acetoxy-6-hydroxy-5,7,9-trimethylundec-3-enoate (**22b**). A flask was charged with crude aldehyde **21b** (8.33 g, ~42.8 mmol), silane (**R**)-**7** (14.6 g,

55.6 mmol, 1.30 equiv.), and CH_2Cl_2 (214 mL, 0.200 M). The flask was purged with argon and cooled to $-78\text{ }^\circ\text{C}$. To the flask was added dropwise, $TiCl_4$ (9.38 mL, 85.5 mmol, 2.00 equiv.), and the reaction mixture was warmed to $-30\text{ }^\circ\text{C}$. After 60 h, EtOAc was introduced to the flask, followed by H_2O . After warming to rt, an additional portion of EtOAc and 10% aq. $NaHCO_3$ was added. The layers were separated, and the aqueous layer was extracted into EtOAc (3 $\times$). The combined organic layers were washed with H_2O , brine, dried over $MgSO_4$, filtered, and concentrated. 1H NMR analysis of the crude reaction mixture revealed one product diastereomer. Purification by flash chromatography (silica, 15 $\rightarrow$ 50% EtOAc/hexanes) provides 9.14 g (67.9% yield, two steps) desired homoallylic alcohol **22b**. 1H NMR (400 MHz, $CDCl_3$) δ 0.90 (3H, d, $J=7.2$), 0.93 (3H, dd, $J=7.0, 7.0$), 0.98 (3H, d, $J=6.9$), 1.05 (3H, d, $J=7.0$), 1.12 (1H, m), 1.34 (1H, m), 1.62 (1H, m), 2.05 (3H, s), 2.34 (1H, ddq, $J=7.0, 7.6, 8.8$), 2.44 (1H, m), 3.01 (2H, d, $J=6.9$), 3.32 (1H, m), 3.63 (3H, s), 4.84 (1H, dd, $J=3.6, 4.0$), 5.39 (1H, dd, $J=7.6, 15.2$), 5.58 (1H, dt, $J=6.9, 15.2$); IR (neat) ν_{max} 3503, 3030, 2967, 2879, 1738, 1437, 1376, 1244, 1168, 1020, 974; HRMS(EI) m/z calc'd for $C_{17}H_{30}O_5$ $[M]^+$ 314.2093, found: 314.2107; $[\alpha]_D = +11.5$ ($c=0.10$, $CHCl_3$).

Stereochemical assignment of second (E)-crotylsilane addition product (22b). To a sample of homoallylic alcohol **22b** (150 mg, 0.477 mmol) in Et₂O (4.80 mL, 0.100 M) at 0 °C under an argon atmosphere was added LiAlH₄ (40.0 mg, 1.05 mmol, 2.20 equiv.) slowly over 1 min. After 10 min at 0 °C, the reaction was quenched with 10% aq. HCl to a pH of 5, and EtOAc (75 mL) and water (100 mL) were added. The layers were mixed and separated. The aqueous layer was extracted into EtOAc (3×). The combined organic layers were washed with H₂O, brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification by flash chromatography (silica, 50→100% EtOAc/hexanes) provides 93.1 mg desired triol (79.9% yield). The triol was dissolved in CH₂(OMe)₂ (3.80 mL, 0.100 M) under argon, and TsOH•H₂O (3.60 mg, 5.00 mol %) was added. The reaction was stirred at rt for 36 h, then concentrated *in vacuo*. Purification by flash chromatography (silica, 10→15% EtOAc/hexanes) provides desired acetonide (87.9 mg, 81.1%) as a colorless oil. Analysis of the ¹³C NMR spectrum of acetonide confirms 1,3-*syn* relationship of initial homoallylic alcohol starting material¹⁶ (key resonances are illustrated in figure above, underlined below). ¹H NMR (400 MHz, CDCl₃) δ 0.76 (3H, d, J=6.6), 0.85 (3H, d, J=5.93), 0.88 (3H, d, J=6.6), 0.99 (3H, d, J=6.8), 1.22-1.32 (2H, m), 1.34 (3H, s), 1.37 (3H, s), 1.40-1.54 (2H, m), 1.64 (1H, br), 2.23-2.28 (3H, m), 3.29-3.37 (2H, m), 3.60 (2H, t, J=6.4), 5.31 (1H, dd, J=9.2, 15.6), 5.45 (1H, dt, J=6.8, 15.6); ¹³C NMR† (75 MHz, CDCl₃) δ 4.6, 11.0, 15.5, 18.0, 19.5, 23.5, 30.0, 31.3, 35.4, 36.1, 39.1, 62.0, 77.5, 78.3, 98.8, 126.7, 134.9.

(2*S*,3*R*,4*R*,5*R*,6*S*)-5-Acetoxy-3-hydroxy-2,4,6-trimethyloctanal. To a flask charged with argon were added homoallylic alcohol **22b** (3.94 g, 12.5 mmol), pyridine (1.01 mL,

12.5 mmol, 1.00 equiv.), and 1:1 MeOH:CH₂Cl₂ (156 mL, 0.0800 M). The flask was cooled to -78 °C. O₃ was bubbled into the reaction mixture until starting material was consumed and a blue color persisted. Me₂S (18.4 mL, 251 mmol, 20.0 equiv.) was added cautiously. The reaction was allowed to warm to rt slowly overnight. After 12 h, the crude reaction mixture was concentrated *in vacuo*, then EtOAc was added. The organic layer was washed with H₂O (2×), brine (1×), dried over MgSO₄, filtered, and concentrated. The resulting material (3.30 g, 108% theoretical yield) was tentatively assigned by ¹H and ¹³C NMR analysis and used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 0.83 (3H, d, J=6.4), 0.90 (3H, t, J=7.4), 0.97 (3H, d, J=6.8), 1.07 (1H, m), 1.14 (3H, d, J=7.6), 1.41 (1H, m), 1.69 (1H, m), 1.92 (1H, m), 2.05 (3H, s), 2.19 (1H, d, J=4.8, exchangeable), 2.80 (1H, m), 3.86 (1H, m), 4.74 (1H, dd, J=3.4, 7.8), 9.65 (1H, s); ¹³C NMR (75 MHz, CDCl₃) δ 7.8, 8.8, 11.0, 14.4, 20.8, 25.4, 36.6, 37.0, 48.9, 71.7, 77.6, 171.7, 204.8.

(16) Evans, D. A.; Rieger, D. L.; Gage, J. R. *Tetrahedron Lett.* **1990**, 31, 7099-7100.

(*3S,4R,5R,6S,7R*)-9,9-Dibromo-6-hydroxy-3,5,7-trimethylnon-8-en-4-yl acetate (**23**). In a flask charged with argon were combined PPh_3 (13.1 g, 50.1 mmol, 4.00 equiv.) and CH_2Cl_2 (62.4 mL). The flask was cooled to 0 °C, and then CBr_4 (8.31 g, 25.1 mmol, 2.00 equiv.) was added. After 15 min, crude aldehyde (3.30 g, ~12.5 mmol) in CH_2Cl_2 (62.1 mL, total vol. = 125 mL, 0.100 M) was added by cannula, with a rinse. 2,6-Lutidine (2.92 mL, 25.1 mmol, 2.00 equiv.) was added by syringe, and the reaction was allowed to warm to rt slowly. After 12 h, the reaction was quenched with aq. NH_4Cl , and the layers were separated. The aqueous layers were extracted into CH_2Cl_2 (3 $\times$). The resulting organic layers were combined, washed with H_2O , brine, dried over MgSO_4 , filtered, and concentrated. Purification by flash chromatography (silica, 5 $\rightarrow$ 20% EtOAc/hexanes) provides 4.11 g **23** (82.2% over two steps) as a yellow oil. ^1H NMR (400 MHz, CDCl_3) δ 0.82-0.98 (9H, m), 1.09 (3H, d, $J=6.8$), 1.14 (1H, m), 1.22 (1H, m), 1.39 (1H, m), 1.64 (1H, m), 1.99 (1H, d, $J=4.0$), 2.07 (3H, s), 2.62 (1H, ddq, $J=6.4, 6.8, 9.6$), 3.63 (1H, m), 4.82 (1H, dd, $J=6.4, 6.6$), 6.18 (1H, d, $J=9.6$); ^{13}C NMR (75 MHz, CDCl_3) δ 5.2, 10.2, 13.8, 15.1, 21.0, 25.3, 35.5, 37.2, 40.8, 80.2, 82.9, 88.9, 138.8, 172.2; IR (neat) ν_{max} 3633, 2970, 2876, 1470, 1390, 1247, 1110; HRMS(EI) m/z calc'd for $\text{C}_{14}\text{H}_{24}\text{Br}_2\text{O}_3$ $[\text{M}]^+$ 398.0092, found: 397.9998; $[\alpha]_{\text{D}} = -23.6$ ($c=0.10$, CH_2Cl_2).

(*3R,4S,5S,6R,7S*)-1,1-Dibromo-3,5,7-trimethyl-1-nonene-4,6-diol (**24**). In a flask charged with argon were combined acetate **23** (1.70 g, 4.25 mmol), MeOH (42.5 mL, 0.100 M), K_2CO_3 (45.0 mg, 4.25 mmol, 1.00 equiv.), and H_2O (one drop). The reaction was stirred at rt 12 h, then quenched with sat. aq. NH_4Cl . The layers were separated, and the aqueous layer was extracted into CH_2Cl_2 (3 $\times$). The combined organic layers were washed with H_2O , brine, dried over MgSO_4 , filtered, and concentrated. Purification by flash chromatography (silica, 15 $\rightarrow$ 40% EtOAc/hexanes) yields 1.50 g desired diol **24** (98.6%) as a white solid: ^1H NMR (400 MHz, CDCl_3) δ 0.86 (3H, t, $J=7.3$), 0.88 (3H, d, $J=7.3$), 0.93 (3H, d, $J=6.6$), 1.03 (1H, m), 1.07 (3H, d, $J=6.6$), 1.37 (1H, m), 1.49 (1H, m), 1.72 (1H, m), 2.57 (1H, brs, exchangeable), 2.61 (1H, m), 3.39 (1H, m), 3.51 (1H, m), 3.58 (1H, brs, exchangeable), 6.11 (1H, d, $J=9.6$); ^{13}C NMR (75 MHz, CDCl_3) δ 5.0, 10.7, 14.8, 15.8, 25.1, 36.1, 37.4, 42.3, 79.5, 81.6, 88.8, 140.7; IR (neat) ν_{max} 3750, 2970, 2867, 1462, 1390, 1247, 1141, 1110, 991, 883; HRMS(Cl, NH_3) m/z calc'd for $\text{C}_{12}\text{H}_{23}\text{Br}_2\text{O}_2$ $[\text{M}+\text{H}]^+$ 357.0065, found: 357.0033; $[\alpha]_{\text{D}} = -57.2$ ($c=0.30$, CH_2Cl_2); mp = 94-95 °C.

(*3R,4S,5R,6R,7S*)-1,1-Dibromo-3,5,7-trimethyl-6-(triisopropylsilyloxy)non-1-en-4-ol (**25**). A flask was charged with diol **24** (1.26 g, 3.52 mmol), CH_2Cl_2 (35.2 mL, 0.100 M), and 2,6-lutidine (980 μL , 8.44 mmol, 2.40 equiv.), and the resulting solution was cooled to 0 °C. TIPSOTf (1.13 mL, 4.22 mmol, 1.20 equiv.) was added dropwise.

After 19 h at 0 °C, the reaction was quenched with 20% aq. NH₄Cl, and the layers were separated. The aqueous layer was extracted into CH₂Cl₂ (3×). The combined organic layers were washed with H₂O, brine, dried over MgSO₄, filtered, and concentrated. Analysis of the crude material revealed one product regioisomer.¹⁷ Purification by flash chromatography (silica, 5→20% EtOAc/hexanes) results in 1.62 g (89.5%) desired product as a colorless oil. ¹H NMR (400 MHz, CDCl₃) δ 0.86 (3H, d, J=6.9), 0.87 (3H, t, J=7.6), 0.93 (3H, d, J=6.9), 0.98-1.07 (4H, m), 1.08 (21H, m), 1.60 (1H, m), 1.66-1.74 (2H, m), 2.14 (1H, d, J=3.6, exchangeable), 2.59 (1H, m), 3.47 (1H, m), 3.82 (1H, m), 6.15 (1H, d, J=9.9); ¹³C NMR (75 MHz, CDCl₃) δ 8.4, 12.5, 13.4, 15.4, 18.1, 18.2, 24.8, 37.3, 41.1, 42.4, 78.4, 81.1, 88.6, 141.1; IR (neat) ν_{max} 3450, 2967, 2940, 2874, 1460, 1385, 1247, 1110, 1059, 991; HRMS(CI, NH₃) *m/z* calc'd for C₂₁H₄₃Br₂O₂Si₁ [M+H]⁺ 513.1399, found: 513.1411; [α]_D = -45.0 (c=0.20, CHCl₃).

(3*R*,5*S*,6*R*,7*S*)-1,1-Dibromo-3,5,7-trimethyl-6-(triisopropylsilyloxy)non-1-en-4-one (**4**). To a sample of alcohol **25** (500 mg, 0.972 mmol) in CH₂Cl₂ (4.86 mL, 0.200 M) was added Celite

filter agent (629 mg), followed by PCC (629 mg, 2.92 mmol, 3.00 equiv.). The reaction was stirred vigorously at rt for 20 h, then filtered, rinsed with CH₂Cl₂, and concentrated. Purification by flash chromatography (silica, 20% EtOAc/hexanes) provides desired ketone **4** (423 mg, 84.9%) as a yellow oil. ¹H NMR[†] (400 MHz, CDCl₃) δ 0.80 (3H, d, J=6.8), 0.87 (3H, t, J=7.6), 1.06-1.10 (22H, m), 1.12 (3H, d, J=7.2), 1.17 (3H, d, J=6.8), 1.42 (1H, m), 1.61 (1H, m), 2.85 (1H, dt, J=5.6, 7.2), 3.66 (1H, dt, J=6.8, 9.6), 4.04 (1H, dd, J=3.0, 5.6), 6.46 (1H, d, J=9.6); ¹³C NMR (75 MHz, CDCl₃) δ 12.3, 13.1, 13.8, 14.5, 16.0, 18.2, 25.0, 41.2, 48.2, 50.2, 76.1, 90.3, 137.7, 211.6; IR (neat) ν_{max} 2963, 2943, 2867, 1716, 1462, 1383, 1247, 1141, 1110, 1059, 991, 883, 679; HRMS(CI, NH₃) *m/z* calc'd for C₂₁H₄₁O₂Br₂Si₁ [M+H]⁺ 511.1243, found: 511.1253; [α]_D = -54.0 (c=0.10, CHCl₃).

(3*S*,4*R*,5*S*,7*R*,8*Z*,10*E*,13*R*,14*Z*)-9-Bromo-3,5,7,13-tetramethyl-4-(triisopropylsilyloxy)-15-(trimethylsilyl)heptadeca-8,10,14-trien-6-one (**26**). In a round-bottomed flask purged with argon were combined dibromide **4** (140 mg, 0.28 mmol), stannane **3** (150 mg, 0.31 mmol, 1.1 equiv.), and degassed PhMe (2.8 mL, 0.10 M). To the resulting mixture were added Pd₂dba₃ (8.0 mg, 0.0084 mmol, 3.0 mol %) and P(2-fur)₃ (10 mg, 0.042 mmol, 15 mol %), and the reaction was heated to 100 °C. After consumption of **4** (~40 min), the flask was cooled to rt, and the crude material was passed through a short silica plug, rinsed with 100 mL 20% EtOAc/hexanes, and concentrated *in vacuo*. Purification by flash chromatography (silica, 1% EtOAc/hexanes) provides the desired triene **26** (160 mg, 92%) as a yellow oil. ¹H NMR[†] (400 MHz, CDCl₃) δ 0.11 (9H, s), 0.74 (3H, d, J=6.8), 0.84 (3H, dd, J=7.2, 6.8), 0.93 (6H, m), 1.05 (22H, m), 1.13 (3H, d, J=6.8), 1.16

(17) At increased TIPSOTf stoichiometry (>2.00 eq., rt), partial silylation of both alcohols begins to occur.

(3H, d, J=7.2), 1.39 (1H, m), 1.60 (1H, m), 2.02 (2H, m), 2.09 (2H, dd, J=6.0, 6.8), 2.43 (1H, m), 2.87 (1H, dq, J=6.0, 6.8), 3.96 (1H, dq, J=7.2, 9.6), 4.07 (1H, dd, J=6.0, 6.0), 5.65 (1H, d, J=10.5), 5.79 (1H, d, J=9.6), 5.98 (2H, m); ^{13}C NMR (75 MHz, CDCl_3) δ 0.39, 12.3, 13.1, 14.0, 14.3, 15.6, 16.4, 18.2, 20.8, 24.8, 30.9, 36.3, 40.1, 41.4, 48.1, 48.6, 75.9, 126.4, 129.8, 130.6, 135.1, 140.0, 146.9, 213.3; IR (neat) ν_{max} 2961, 2929, 2868, 1714, 1671, 1457, 1372, 1248, 1059, 1013, 836; HRMS(CI, NH_3) m/z calc'd for $\text{C}_{37}\text{H}_{80}\text{Br}_1\text{O}_2\text{Si}_2$ $[\text{M}+\text{H}]^+$ 691.4880, found: 691.4888; $[\alpha]_{\text{D}} = -40.9$ ($c=0.08$, CH_2Cl_2).

(3*S*,4*R*,5*S*,7*R*,8*E*,10*E*,13*R*,14*Z*)-3,5,7,9,13-Pentamethyl-4-(triisopropylsilyloxy)-15-(trimethylsilyl)heptadeca-8,10,14-trien-6-one (**27**). To a solution of bromide **26** (40 mg, 0.064 mmol) in THF (0.64 mL, 0.10 M) at 0 °C was added $\text{Pd}(t\text{-Bu}_3)_2$ (1.6 mg, 3.1 μmol , 5.0 mol %), followed by Me_2Zn (2.0 M solution in toluene, 38 μL , 0.076 mmol, 1.2 equiv.). The reaction was stirred at 0 °C for 4 h, then quenched with 20% aq. NH_4Cl . Et_2O was added to the mixture, and the layers were separated. The aqueous layer was extracted into Et_2O (3 $\times$). The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and concentrated. Purification by flash chromatography (silica, 1% EtOAc /hexanes) provides 34 mg desired product (96%) as a brown oil. ^1H NMR † (400 MHz, CDCl_3) δ 0.10 (9H, s), 0.69 (3H, d, J=6.8), 0.82 (3H, dd, J=6.4, 7.2), 0.92 (6H, m), 1.06 (22H, m), 1.10 (6H, m), 1.37 (1H, m), 1.56 (1H, m), 1.79 (3H, s), 2.03 (4H, m), 2.39 (1H, m), 2.78 (1H, dt, J=6.8, 7.2), 3.66, (1H, dt, J=6.8, 10.0), 4.10 (1H, dd, J=6.8), 5.19 (1H, d, J=10.0), 5.56 (1H, dt, J=7.2, 15.6), 5.67 (1H, d, J=10.4), 5.98 (1H, d, J=15.6); ^{13}C NMR (75 MHz, CDCl_3) δ 0.4, 12.3, 13.2, 14.0, 14.3, 15.6, 16.8, 18.3, 20.7, 25.0, 29.6, 30.9, 36.5, 40.8, 41.2, 45.3, 47.9, 76.0, 127.7, 128.9, 129.2, 135.3, 139.5, 147.6, 214.4; IR (neat) ν_{max} 2962, 2868, 1712, 1462, 1385, 1372, 1248, 1060, 989, 836; HRMS(CI, NH_3) m/z calc'd for $\text{C}_{34}\text{H}_{67}\text{O}_2\text{Si}_2$ $[\text{M}+\text{H}]^+$ 563.4680, found: 563.4702; $[\alpha]_{\text{D}} = +69.2$ ($c=0.20$, CH_2Cl_2).

(3*S*,4*R*,5*S*,7*R*,8*Z*,10*E*,13*R*,14*Z*)-9-Bromo-15-iodo-3,5,7,13-tetramethyl-4-(triisopropylsilyloxy)heptadeca-8,10,14-trien-6-one (**28**). A sample of **26** (61.0 mg, 0.0971 mmol) was dissolved in (0.970 mL, 0.100 M) EtCN , and the flask was protected from light. To the solution was added NIS (31.0 mg, 0.136 mmol, 1.40 equiv.), and the reaction was stirred 3 h at rt. The reaction was quenched with aq. $\text{Na}_2\text{S}_2\text{O}_3$ and CH_2Cl_2 . After separation of the layers, the aqueous layer was extracted into CH_2Cl_2 (4 $\times$). The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and concentrated. Purification by flash chromatography (silica, 1% EtOAc /hexanes) provides desired iodobromide **28** (55.8 mg, 84.2%) as a brown oil. ^1H NMR † (400 MHz, CDCl_3) δ 0.75 (3H, d, J=7.2), 0.83 (3H, t, J=7.4), 0.90 (3H, t, J=7.3), 0.97 (3H, d, J=6.8), 1.5 (21H, m), 1.13 (3H, d, J=7.2), 1.16 (3H, d, J=7.2), 1.25 (1H, m), 1.61 (2H, m), 2.17 (2H, m), 2.49

(2H, q, J=7.3), 2.51 (1H, m), 2.88 (1H, dq, J=7.2, 5.6), 3.97 (1H, dq, J=6.9, 9.2), 4.07 (1H, dd, J=2.6, 5.6), 5.23 (1H, d, J=8.6), 5.81 (1H, d, J=9.2), 6.02 (2H, m); ^{13}C NMR (75 MHz, CDCl_3) δ 12.2, 13.1, 13.9, 14.3, 14.9, 16.3, 18.1, 18.9, 24.7, 26.7, 27.7, 38.8, 40.9, 48.0, 48.6, 75.8, 110.3, 126.2, 130.3, 130.8, 134.0, 138.1, 213.2; IR (neat) ν_{max} 2960, 2862, 1710, 1385, 1248, 1030, 980; HRMS(API, NH_3) m/z calc'd for $\text{C}_{30}\text{H}_{55}\text{BrI}_1\text{O}_2\text{Si}_1$ $[\text{M}+\text{H}]^+$ 681.2199, found: 681.2207; $[\alpha]_{\text{D}} = -35.4$ ($c=0.07$, CH_2Cl_2).

(3*S*,4*R*,5*S*,7*R*,8*E*,10*E*,13*R*,14*Z*)-15-Iodo-3,5,7,9,13-pentamethyl-4-(triisopropylsilyloxy)-heptadeca-8,10,14-trien-6-one (**29**). A sample of **28** (29 mg, 0.046 mmol) in THF (0.46 mL, 0.10 M) was cooled to 0 °C. To the solution was added $\text{Pd}(t\text{-Bu}_3\text{P})_2$ (1.2 mg, 0.0023 mmol, 5.0 % mol), followed by Me_2Zn (2.0 M in toluene, 28 μL , 0.055 mmol, 1.2 equiv.). The reaction was stirred at 0 °C for 8 h. To the brown mixture was added H_2O and Et_2O , and the resulting layers were separated. The aqueous layer was extracted into Et_2O (3 $\times$). The combined organic layers were dried with MgSO_4 , filtered, and concentrated. Flash chromatography (silica, 1% EtOAc /hexanes) provides 26 mg (93%) desired **29** as a dark yellow oil. Regiochemistry of addition was determined by NOE analysis upon irradiation of C-14 Me (key signals shown in figure above, underlined below). ^1H NMR † (400 MHz, CDCl_3) δ 0.69 (3H, d, J=7.2), 0.82 (3H, t, J=7.2), 0.90 (3H, d, J=6.4), 0.95 (3H, t, J=7.2), 1.06 (22H, m), 1.09 (3H, d, J=7.6), 1.11 (3H, d, J=6.0), 1.34 (1H, m), 1.51 (1H, m), 1.79 (3H, s), 1.95 (2H, dd, J=7.2, 7.6), 2.01 (2H, J=7.2), 2.38 (1H, ddq, J=6.4, 7.2, 9.2), 2.78 (1H, dq, J=6.4, 7.6), 3.66 (1H, dq, J=6.0, 10.0), 4.09 (1H, dd, J=6.4, 6.8), 4.89 (1H, d, J=9.2), 5.17 (1H, d, J=10.0), 5.58 (1H, dt, J=7.6, 16.0), 5.98 (1H, d, J=16.0); ^{13}C NMR (75 MHz, CDCl_3) δ 12.3, 13.1, 14.0, 14.5, 15.0, 16.8, 18.2, 18.5, 24.8, 26.5, 27.8, 29.3, 38.6, 40.3, 46.2, 46.6, 96.1, 112.4, 125.5, 129.2, 129.6, 130.0, 139.2, 214.0; IR (neat) ν_{max} 2960, 2935, 1730, 1393, 1372, 1248, 1060, 1030, 981, 825; HRMS(API, NH_3) m/z calc'd for $\text{C}_{31}\text{H}_{58}\text{I}_1\text{O}_2\text{Si}_1$ $[\text{M}+\text{H}]^+$ 617.3251, found: 617.3266; $[\alpha]_{\text{D}} = -22.4$ ($c=0.08$, CH_2Cl_2).

(3*S*,4*R*,5*S*,7*R*,8*E*,10*E*,13*R*,14*Z*,16*E*)-15-Ethyl-17-((2*R*,6*R*)-6-isopropoxy-3,6-dihydro-2H-pyran-2-yl)-3,5,7,9,13-pentamethyl-4-(triisopropylsilyloxy)heptadeca-8,10,14,16-tetraen-6-one (**30**). To a sample of alkyne **2** (28 mg, 0.13 mmol, 4.0 equiv.) in THF (1.3 mL, 0.10 M vs. **2**) at 0 °C was added Cp_2ZrHCl (40 mg, 0.16 mmol, 4.8 equiv.). The reaction was stirred at 0 °C 30 min, then warmed to rt until consumption of **2** was complete as monitored by TLC (~60 min). Separately, a sample of ZnCl_2 (24 mg, 0.17 mmol, 5.3 equiv.) was fused and dissolved in THF (1.7 mL, 0.10 M vs. ZnCl_2). The ZnCl_2

solution was added to the organozirconium mixture by cannula, and the reaction was stirred at rt. After 15 min, a solution containing iodide **29** (20 mg, 0.033 mmol), Pd(PPh₃)₄ (1.8 mg, 15 μmol, 5.0 mol %), and THF (0.50 mL, 0.070 M vs. **29**) was added to the organozinc solution by cannula. The reaction was stirred at rt, monitored by TLC. After 120 min, the reaction was quenched with H₂O and Et₂O, and the layers were separated. The aqueous layers were extracted into Et₂O (3×). The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification by flash chromatography (silica, 1→10% EtOAc/hexanes) provides 11 mg (51%) desired product as a pale yellow oil.¹⁸ ¹H NMR (400 MHz, CDCl₃) δ 0.76 (3H, d, J=7.2), 0.86 (3H, t, J=7.2), 0.91 (3H, d, J=6.4), 0.95 (3H, t, J=7.2), 1.06 (22H, m), 1.09 (3H, d, J=7.6), 1.11 (3H, d, J=6.0), 1.15 (3H, d, J=6.2), 1.24 (3H, d, J=6.2), 1.34 (1H, m), 1.51 (1H, m), 1.79 (3H, s), 1.95 (2H, m), 2.01 (2H, m), 2.19-2.36 (2H, m), 2.42 (1H, ddq, J=6.4, 7.2, 9.2), 2.78 (1H, dq, J=6.4, 7.4), 3.66 (1H, dq, J=6.4, 10.0), 3.99 (1H, qq, J=6.2, 6.2), 4.09 (1H, dd, J=6.4, 6.8), 4.66 (1H, m), 4.89 (1H, d, J=9.2), 5.09 (1H, m), 5.17 (1H, d, J=10.0), 5.58 (1H, dt, J=7.6, 16.0), 5.67 (1H, ddd, J=1.2, 2.8, 10.0), 5.75 (1H, m), 5.92 (1H, dd, J=5.6, 10.0), 6.02 (1H, d, J=16.0), 6.05 (1H, d, J=16.1); ¹³C NMR (75 MHz, CDCl₃) δ 12.7, 13.4, 14.4, 14.5, 15.3, 16.1, 16.4, 17.4, 18.2, 18.5, 22.0, 23.4, 25.8, 26.5, 27.8, 30.6, 32.5, 38.6, 40.3, 46.2, 46.6, 58.2, 96.1, 112.4, 122.0, 125.5, 128.2, 128.8, 129.2, 129.6, 130.0, 135.2, 135.7, 143.7, 221.2; IR (neat) ν_{max} 3030, 2968, 2875, 2865, 1460, 1415, 1380, 1248, 1170, 1025, 989; HRMS(CI, NH₃) *m/z* calc'd for C₄₁H₇₃O₄Si₁ [M+H]⁺ 657.5278, found: 657.5281; [α]_D = +68.2 (c=0.10, CHCl₃).

(–)Callystatin A (**1**). To a round bottom flask charged with lactol **30** (6.0 mg, 0.0091 mmol) were added sequentially THF (0.46 mL, 0.020 M), H₂O (0.18 mL, 0.050 M), and glacial AcOH (0.18 mL, 0.050 M). The reaction was stirred at rt 24 h, until starting material was consumed by TLC. EtOAc and H₂O were added to the reaction mixture. The layers were separated, and the aqueous layer was extracted (3×) into EtOAc. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. The crude material was dissolved in CH₂Cl₂ (0.54 mL, 0.017 M), and Celite (14 mg) and PDC (14 mg, 0.036 mmol, 4.0 equiv.) were added. The mixture was stirred vigorously at rt 24 h, until starting material was consumed. The material was filtered through a 1/4 inch silica plug, rinsed with 30% EtOAc/hexanes, and concentrated. Purification by flash chromatography (silica, 0→10% EtOAc/hexanes) provided 4.1 mg (74%, two steps) desired lactone.¹⁹ The lactone was placed in a plastic

(18) Elution with 1% EtOAc/hexanes provides 5.5 mg unreacted starting material **29**.

(19) Further elution (silica, 10→30% EtOAc/hexanes) provided an (*E*)-enal (1.2 mg) which presumably resulted from isomerization of the C2-C3 alkene during lactol deprotection. For literature precedent and a

vial, and THF (0.20 mL) and HF•pyr (0.20 mL) were added. The reaction mixture was stirred at rt 48 h. To the reaction mixture were added EtOAc and H₂O, and the layers were separated. The aqueous layer was extracted 3× into EtOAc. The combined organic layers were washed with brine, dried over MgSO₄, filtered, and concentrated. Purification by flash chromatography (silica, 10→30% EtOAc/hexanes) provides 2.6 mg (88%) (-)-callystatin A as a colorless oil. ¹H NMR† (400 MHz, CDCl₃) δ 0.85 (3H, t, J=7.2), 0.89 (3H, d, J=6.4), 0.97 (3H, d, J=7.4), 1.05 (3H, t, J=6.9), 1.12 (3H, d, J=7.2), 1.14 (3H, d, J=6.9), 1.35 (2H, m), 1.80 (3H, s), 1.93-2.08 (3H, m), 2.19 (2H, q, J=6.9), 2.46 (2H, m), 2.64-2.70 (2H, m), 2.86 (1H, m), 3.56 (1H, m), 3.64 (1H, dq, J=6.9, 9.6), 4.98 (1H, ddd, J=5.6, 6.7, 7.2), 5.12 (1H, d, J=9.6), 5.25 (1H, d, J=9.6), 5.58 (1H, m), 5.75 (1H, dd, J=6.7, 16.1), 6.00 (1H, d, J=16.1), 6.02 (1H, dd, J=1.4, 9.6), 6.64 (1H, d, J=16.1), 6.89 (1H, m); ¹³C NMR (75 MHz, CDCl₃) δ 10.8, 11.2, 13.0, 13.3, 14.1, 16.0, 20.7, 25.8, 26.3, 30.0, 32.1, 36.7, 40.7, 45.5, 45.7, 74.5, 78.9, 121.7, 124.5, 127.6, 128.3, 129.8, 135.2, 135.4, 136.0, 137.1, 144.6, 163.9, 216.3; IR (neat) ν_{max} 3510, 2965, 2930, 1715, 1432, 1380, 1248, 1030, 989, 825; HRMS(CI, NH₃) *m/z* calc'd for C₂₉H₄₅O₄ [M+H]⁺ 457.3318, found: 457.3301; [α]_D = -98.4 (c=0.05, MeOH).

procedure for converting this material to desired lactone, see: Lautens, M.; Stammers, T. A. *Synthesis* **2002**, 1993-2012.

*Callystatin A Spectral Data Compared to Reported Synthetic Data*²⁰

<i>¹H NMR Data</i>				
Panek		Crimmins		
Chem. Shift	Integration	Chem. Shift	Integration	(Δ)δ
0.85	3 H	0.83	3 H	0.02
0.89	3 H	0.88	3 H	0.01
0.97	3 H	0.95	3 H	0.02
1.05	3 H	1.02	3 H	0.03
1.12	3 H	1.10	3 H	0.02
1.14	3 H	1.12	3 H	0.02
1.35	2 H	1.23-1.42	2 H	---
1.80	3 H	1.79	3 H	0.01
1.93-2.08	3 H	1.93-2.10	3 H	---
2.19	2 H	2.17	2 H	0.02
2.46	2 H	2.46	2 H	0.00
2.64-2.70	2 H	2.58-2.70	2 H	---
2.86	1 H	2.83	1 H	0.03
3.56	1 H	3.54	1 H	0.02
3.64	1 H	3.63	1 H	0.01
4.98	1 H	4.98	1 H	0.00
5.12	1 H	5.11	1 H	0.01
5.25	1 H	5.23	1 H	0.02
5.58	1 H	5.57	1 H	0.01
5.75	1 H	5.73	1 H	0.02
6.00	1 H	5.98	1 H	0.02
6.02	1 H	6.04	1 H	-0.02
6.64	1 H	6.63	1 H	0.01
6.89	1 H	6.88	1 H	0.01

(20) Comparison data from total synthesis of callystatin A, as disclosed by Crimmins and King: Crimmins, M. T.; King, B. W. *J. Am. Chem. Soc.* **1998**, *120*, 9084-9085.

<i>¹³C NMR Data</i>		
Panek Chem. Shift	Crimmins Chem. Shift	(Δ)δ
10.8	10.9	-0.1
11.2	11.2	0.0
13.0	13.0	0.0
13.3	13.4	-0.1
14.1	14.2	-0.1
16.0	16.1	-0.1
20.7	20.7	0.0
25.8	25.8	0.0
26.3	26.4	-0.1
30.0	30.1	-0.1
32.1	32.1	0.0
36.7	36.7	0.0
40.7	40.8	-0.1
45.5	45.6	-0.1
45.7	45.7	0.0
74.5	74.4	0.1
78.9	78.9	0.0
121.7	121.7	0.0
124.5	124.7	-0.2
127.6	127.7	-0.1
128.3	128.3	0.0
129.8	129.9	-0.1
135.2	135.3	-0.1
135.4	135.4	0.0
136.0	136.2	-0.2
137.1	137.2	-0.1
144.6	144.7	-0.1
163.9	164.1	-0.2
216.3	216.4	-0.1

(R)-OAc-mandelate of (S)-10