Total Synthesis of Carbazaquinocin-C:
Application of the *ortho*-benzannulation of Fischer Carbene
Complexes to carbazole 3, 4-quinone alkaloids

Manish Rawat and William D. Wulff*

Department of Chemistry

Michigan State University

East Lansing, MI 48824

General Information

Unless and otherwise indicated, all common reagents and solvents were used as obtained from commercial suppliers without further purification. Prior to use, tetrahydrofuran and diethyl ether were distilled from Na/benzophenone ketyl, toluene was distilled from Na and methylene chloride was prepared by distillation from calcium hydride. Routine NMR spectra were recorded on 300MHz, 500MHz INOVA and VXR instruments. Both low and high resolution mass spectra were obtained from the Michigan State University Mass Spectrometry Facility. Elemental analyses were done by the Michigan State University Elemental Analysis Facility.

***N*-Methoxy-*N*-methyl-1*H*-indole-2-carboxamide (11).¹**

To a flame dried round-bottomed flask under a N₂ atmosphere was added indole-2-carboxylic acid (500 mg, 3.1 mmol), N,O-dimethylhydroxylamine hydrochloride (300 mg, 3.1 mmol), triethylamine (0.44 mL, 3.1 mmol) and anhydrous methylene chloride (20 mL). After stirring at room temperature for 5 min, the reaction mixture was cooled to

0 °C and solid dicyclohexylcarbodiimide (640 mg, 3.1 mmol) was added. The reaction mixture was allowed to warm to room temperature over 15 min and then stirring was continued at room temperature for 3 h. The solvent was removed under vacuum and the crude product was purified by silica gel chromatography (30 % acetone/hexanes) to give 475 mg (75 %) of the desired amide **11** as a bright yellow solid; mp 147–149 °C (acetone/hexanes) (lit: mp 147-149 °C).¹ Spectral data for **11**: ¹H-NMR (500 MHz, CDCl₃) δ = 3.44 (s, 3H), 3.84 (s, 3H), 7.12 (ddd, 1H, J = 8.1, 7.1, 1.1 Hz), 7.23 (dd, 1H, J = 2.0, 0.9 Hz), 7.29 (ddd, 1H, J = 8.1, 7.1, 1.1 Hz), 7.42 (dd, 1H, J = 8.2, 0.9 Hz), 7.69 (dd, 1H, J = 8.2, 0.9 Hz), 9.56 (br, s, 1H); ¹³C-NMR (125 MHz, CDCl₃) δ = 33.2, 61.3, 108.0, 111.7, 120.4, 122.5, 124.8, 128.0, 128.2, 135.8, 161.7; mass spectrum (EI) m/z (% rel. intensity) 204 M⁺ (57), 144 (100), 116 (91), 889 (100); IR (CH₂Cl₂, cm⁻¹) 3276 (s, NH), 1605 (s, CO). Anal calcd for C₁₁H₁₂N₂O₂: C, 64.69; H, 5.92; N, 13.72. Found C, 64.98; H, 5.79; N, 13.69.

1-(1*H*-Indol-2-yl)octan-1-one (**12**)

12

To a stirred suspension of Mg (4.8 g, 196 mmol) in 20 mL THF under N₂ atmosphere was added 1-chlorooctane (7.5 mL, 49 mmol) and a pinch of I₂ to give a brown colored solution. The mixture was refluxed until the solution became colorless. At this point, the reaction mixture was warmed to room temperature and the remainder of the 1-chlorooctane (7.5 mL, 49 mmol) was added dropwise over 15 min. The solution was stirred at room temperature for 1 h and then the reaction mixture was transferred to a solution of amide **11a** (4.00 g, 19.6 mmol) in anhydrous THF (20 mL) at 0 °C. The resulting solution was stirred for 12 h, quenched with 5 % HCl/ice-H₂O at 0 °C and extracted with 3x50 mL of ether. The combined organic layer was dried over Na₂SO₄, evaporated to a yellow oil and loaded onto a silica gel chromatography and eluted (10% ethyl acetate/hexanes) to give the desired ketone **12** (4.05 g, 90 %) as white solid; mp

115–117 °C (ethyl acetate/hexanes). Spectral data for **12**: ^1H NMR (500 MHz, CDCl_3) δ = 0.87 (t, 3 H, J = 6.9 Hz), 1.22–1.43 (m, 8 H), 1.73–1.81 (m, 2 H), 2.92 (t, 2 H, J = 7.4 Hz), 7.13 (ddd, 1 H, J = 8.0, 7.1, 1.1 Hz), 7.17–7.19 (m, 1 H), 7.32 (ddd, 1 H, J = 8.3, 7.3, 1.1 Hz), 7.41 (dt, 1 H, J = 8.4, 0.9 Hz), 7.69 (dd, 1 H, J = 7.9, 1.1 Hz), 9.10 (br, s, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ = 14.03, 22.60, 25.19, 29.10, 29.37, 31.68, 38.39, 109.09, 112.24, 120.84, 122.96, 126.1, 127.58, 135.27, 137.27, 193.77; IR (CH_2Cl_2 , cm^{-1}) 3316s, 2928s, 2857s, 1653s; mass spectrum (EI) m/z (% rel. intensity) 243 M^+ (31), 159 (100), 144 (70), 118 (20), 89 (47), 84 (58). Anal calcd for $\text{C}_{16}\text{H}_{21}\text{NO}$: C, 78.97; H, 8.70; N, 5.76. Found C, 78.88; H, 8.37; N, 5.80.

1-(3-Bromo-1*H*-indol-2-yl)-octan-1-one (**13**)

An ice-cooled solution of Br_2 (0.221 μL , 4.3 mmol) in DMF (10 mL) was added to the solution of **12** (400 mg, 1.7 mmol) and powdered KOH (58 mg, 12.9 mmol) in DMF (10 mL). After stirring at room temperature for 6 h, the mixture was poured into a solution of 28 % NH_4OH (25 mL) and NaHSO_3 (230 mg) in water (375 mL) to give a yellow precipitate. The precipitate separated by filtration and purified using silica gel chromatography (10 % ethylacetate/hexane) to give **13** (370 mg, 71 %) as light yellow solid. R_f (hexanes/ethylacetate 10:1) = 0.50; mp 125 – 127 °C (hexane/EtOAc). Spectral data for **13**: ^1H NMR (500MHz, CDCl_3): δ 0.88 (t, 3H, J = 6.9 Hz), 1.23–1.46 (m, 8H), 1.75–1.84 (m, 2H), 3.17 (t, 2H, J = 7.4 Hz), 7.20 (ddd, 1H, J = 1.3, 6.7, 8.1, 6.7, 1.3 Hz), 7.34 - 7.42 (m, 2H), 7.66 (qd, 1H, J = 8.2, 0.9 Hz), 9.50 (br, s, 1H); ^{13}C -NMR (125 MHz, CDCl_3): δ = 14.06, 22.62, 24.27, 29.14, 29.32, 31.71, 40.91, 97.4, 112.24, 121.51, 121.76, 127.15, 128.25, 132.11, 135.42, 192.97; mass spectrum (EI) m/z (% rel. intensity) 323 (17, ^{81}Br), 321 (19, ^{79}Br), 242 (23), 239 (94, ^{81}Br), 237 (100, ^{79}Br), 224 (64, ^{81}Br), 222 (61, ^{79}Br), 197 (17, ^{81}Br), 195 (19, ^{79}Br), 171 (28), 144 (21), 88 (31); IR (CH_2Cl_2 , cm^{-1}) 3304s, 2922s, 2855s, 1642s. Anal calcd for $\text{C}_{16}\text{H}_{20}\text{BrNO}$: C, 59.64; H, 6.26; N, 4.35. Found C, 59.60; H, 6.18; N, 4.36.

2-(dec-2-en-3-yl)-1H-indole (**16**)

16

To a stirred suspension of ethyltriphenylphosphonium bromide (78 g, 210 mmol) in toluene (210 mL) under N₂ at room temperature was added solid KHMDS (44g, 210mmol) over 30 min. The solution was stirred at room temperature for 1 h and then the solid ketone **15** (13 g, 42 mmol) was added. After stirring vigorously for 3 days at room temperature, the reaction mixture was quenched with 10 mL of H₂O at 0 °C. At this point the reaction mixture was filtered and washed with 3 x 150 mL of ether. The combined organic layers were evaporated and purified by silica gel chromatography (10 % EtOAc/hexanes) to give **16** (12.1 g, 93 %) as light pink oil and as a 1.0 : 0.8 ratio of E : Z isomers. These isomers were not separable and the following spectral data were collected on the mixture. The ¹H and ¹³C spectral data of each isomer were determined with the aid of samples of **16** with different ratios of E : Z isomers. The stereochemical assignment was made based on the NOE data shown below. Spectral data of **16**: ¹H NMR (500 MHz, CDCl₃) (Z isomer) δ 0.87 (t, 3H, J = 7.1 Hz), 1.18-1.60 (m, 10H), 1.90 (dd, 3H, J = 1.3, 7.1 Hz), 2.40 (td, 2H, J = 7.6, 1.1 Hz), 5.68 (qd, 1H, J = 7.1, 1.3 Hz), 6.42-6.48 (m, 1H), 7.01-7.20 (m, 2H), 7.32-7.36 (m, 1H), 7.59 (dd, 1H, J = 0.9, 7.7 Hz), 7.98 (br, s, 1H); (E isomer) δ 0.87 (t, 3H, J = 7.1 Hz), 1.18-1.40 (m, 8H), 1.46-1.56 (m, 2H), 1.83 (d, 3H, J = 7.1 Hz), 2.48 (t, 2H, J = 7.8 Hz), 5.88 (q, 1H, J = 7.1 Hz), 6.38 (d, 1H, J = 1.5 Hz), 7.04 (ddd, 1 H, J = 7.5, 7.1, 1.1 Hz), 7.11 (ddd, 1H, J = 8.1, 7.1, 1.1 Hz), 7.27 (dd, 1H, J = 8.8, 0.8 Hz), 7.53 (d, 1H, J = 7.7 Hz), 8.02 (br, s, 1H); ¹³C NMR (125 MHz, CDCl₃) (Z isomer) δ 14.05, 15.24, 22.62, 29.04, 29.09, 29.28, 31.82, 37.72, 102.29, 110.513, 119.758, 120.25, 121.66, 123.73, 128.50, 132.72, 135.62, 137.13; (E isomer) δ 13.71, 14.07, 22.65, 28.87, 29.20, 29.29, 29.76, 31.86, 99.49, 110.32, 119.75, 120.06, 120.28, 121.88, 129.08, 133.08, 136.26, 139.73; IR (CH₂Cl₂) 3422s, 3057s, 3032s, 2930brs, 2855s, 1651s, cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 255 M⁺

(90), 240 (90), 171 (100), 130 (100), 117 (100). Anal calcd for C₁₈H₂₅N: C, 84.65; H 9.87; N 5.48. Found C, 84.24; H, 9.90; N, 5.43.

1-(*tert*-Butyldimethylsilyl)-3-bromo-2-(dec-2-en-3-yl)-1*H*-indole (15)

To a solution of vinyl indole **16** (230 mg, 0.96 mmol) in 10 mL of DMF at room temperature was added crushed KOH (59 mg, 1.05 mmol). After stirring for 30 min at room temperature, NBS (187 mg, 1.05 mmol) was added, and the reaction mixture was again stirred for 1 h at room temperature. At this point the reaction mixture was diluted with CCl₄ and washed with 3x10 mL of brine and 2x10 mL of water. The organic extract was dried over MgSO₄ and evaporated to a brown residue. This residue containing **14** was then dissolved in 10 mL THF and transferred to a flame-dried round-bottomed flask containing a suspension of NaH (50 mg, 1.25 mmol) in THF (5 mL). Then, a *tert*-butyldimethylsilyl chloride (173 mg, 1.15 mmol) solution in THF (5 mL) was added and the resulting solution was stirred for 6 h at room temperature. The reaction mixture was diluted with ether and washed with three portions of water and brine. The organic fraction was dried over MgSO₄, filtered and evaporated to give a brown residue. The residue was purified by column chromatography using ethyl acetate/hexanes (10 %) to give **15** (183 mg, 45 %) as light yellow oil. *R*_f (Hexanes/benzene 5:1) = 0.62.

1-Benzyl-2-(dec-2-en-3-yl)-1*H*-indole (**17**)

To a suspension of sodium hydride (132 mg, 1.5 mmol) in 11 mL of DMF at room temperature was added a solution of 2-vinyl indole **16** (562 mg, 2.2 mmol, E/Z ratio of 1:0.7) in 5 mL DMF. The reaction mixture was stirred for 30 min and then benzyl bromide (376 mg, 2.2 mmol) was added in small portions. After reaction mixture had stirred for 3 h at room temperature, it was carefully quenched with water and extracted with 3x20 mL of ethyl acetate. The combined organic extract was washed with 20 mL of brine and 20 mL of H₂O. The organic fraction was dried over MgSO₄, filtered, evaporated to a yellow oil, and purified by silica gel chromatography (10 % EtOAc/hexane) to give 531 mg of **17** (70 %) as colorless oil and as a 1.0 : 0.5 ratio of E : Z isomers. These isomers were not separable and the following spectral data were collected on the mixture. The ¹H and ¹³C spectral data of each isomer were determined with the aid of samples of **17** with different ratios of E : Z isomers. The stereochemical assignment was made based on the NOE data shown below. Spectral data of **17**: ¹H NMR (500 MHz, CDCl₃) (Z isomer) δ 0.84 (t, 3H, J = 7.1 Hz), 1.10-1.35 (m, 10H), 1.48 (dt, 3H, J = 6.8, 1.1 Hz), 2.13 (tt, 2H, J = 1.1, 7.7 Hz), 5.2 (s, 2H), 5.77 (qt, 1H, J = 7.6, 1.2 Hz), 6.30 (d, 1H, J = 0.8 Hz), 6.98 (d, 2 H, J = Hz), 7.06-7.11(m, 2H), 7.11-7.16(m, 1H), 7.16-7.25(m, 3H), 7.57-7.63(m, 1H); (E-isomer): δ 0.86 (t, 3H, J = 7.1 Hz), 1.14-1.36 (m, 10H), 1.76 (d, 3H, J = 6.8 Hz), 2.35 (t, 2H, J = 7.7 Hz), 5.31 (s, 2H), 5.62 (q, 1H, J = 6.8 Hz), 6.64 (s, 1H), 7.00-7.04 (m, 2H), 7.07-7.11 (m, 3H), 7.18-7.29 (m, 3H), 7.57-7.64 (m, 1H); ¹³C NMR (125 MHz, CDCl₃) (Z isomer): δ 14.04, 15.20, 22.61, 28.28, 29.07, 29.14, 31.81, 38.56, 47.26, 101.42, 110.25, 119.51, 120.14, 120.98, 126.23, 127.00, 127.58, 128.44, 128.47, 133.543, 136.96, 138.15, 139.68; (E isomer) δ 13.09, 14.04, 22.61, 28.419, 29.11, 29.47, 31.31, 31.81, 47.68, 100.83, 110.30, 119.73, 120.147,

121.24, 125.97, 126.96, 127.30, 128.24, 128.57, 133.30, 137.59, 138.48, 143.95; IR (CH₂Cl₂) 3058s, 3031s, 2955m, 2926m, 2857s cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 345 M⁺ (65), 330 (23), 260 (47), 246 (31), 91 (100). Anal calcd for C₂₅H₃₁N: C, 86.90; H, 9.04; N, 4.05. Found C, 86.95; H, 8.84; N, 3.85.

1-(4-Methoxybenzyl)-2-(dec-2-en-3-yl)-1H-indole (**18**)

To a suspension of sodium hydride (95.6 mg, 2.39 mmol) in 8 mL of DMF at room temperature was added a solution of vinyl indole **16** (554 mg, 2.17 mmol, E/ Z ratio of 1:0.6) in 5 mL DMF. The reaction mixture was stirred for 30 min and then PMBCl (357 mg, 2.46 mmol) was added in small portions. After the reaction mixture was stirred for 3 h at room temperature, it was carefully quenched with water and extracted with 3x10 mL of ethyl acetate. The combined organic extract was washed with 10 mL of brine and 10 mL of H₂O. The organic fraction was dried over MgSO₄, filtered, evaporated to a yellow oil, and purified by silica gel chromatography (10 % EtOAc/hexane) to give 586 mg of **18** (72 % yield) as a colorless oil and as a 1.0 : 0.5 ratio of E : Z isomers. These isomers were not separable and the following spectral data were collected on the mixture. Spectral data of **18**: ¹H NMR (300 MHz, CDCl₃): (E-isomer) δ 0.87 (t, 3H, J = 6.6 Hz),

1.00-1.40 (m, 10H), 1.78 (d, 3H, $J = 6.9$ Hz), 2.36 (t, 2H, $J = 6.0$ Hz), 3.76 (s, 3H), 5.26 (s, 2H), 5.64 (q, 1H, $J = 6.9$ Hz), 6.41 (s, 1 H), 6.78-7.20 (m, 7H), 7.58 - 7.64 (m, 1H); (Z-isomer) δ 0.87 (t, 3H, $J = 6.6$ Hz), 1.00-1.40 (m, 10H), 1.50 (d, 3H, $J = 6.9$ Hz), 2.17 (t, 2H, $J = 6.0$ Hz), 3.76 (s, 3H), 5.18 (s, 2H), 5.80 (q, 1H, $J = 6.4$ Hz), 6.31 (s, 1 H), 6.78-7.20 (m, 7H), 7.58 - 7.64 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 22.62, 15.24, 14.08, 13.94, 22.62, 28.26, 28.40, 29.10, 29.14, 29.47, 31.28, 31.81, 38.57, 55.16, 47.08, 46.67, 100.65, 101.25, 110.27, 110.32, 113.79, 113.90, 119.42, 119.65, 120.07, 120.87, 121.15, 127.08, 127.19, 127.40, 127.48, 128.15, 128.36, 130.14, 130.42, 133.13, 133.48, 136.79, 137.46, 139.60, 143.89, 158.48, 158.53, 5 aliphatic and 1 aryl carbons not located; IR (CH_2Cl_2) 3031s, 2997s, 2928brs, 2855s, 1613s, 1512s, 1482s, 1443s cm^{-1} ; mass spectrum (EI) m/z (% rel intensity) 375 M^+ (80), 290 (27), 254 (29), 227 (86), 121 (100). Anal calcd for $\text{C}_{26}\text{H}_{33}\text{NO}$: C, 83.15; H, 8.86; N, 3.73. Found C, 82.78; H, 9.08; N, 3.66.

1-Benzyl-3-bromo-2-(dec-2-en-3-yl)-1H-indole (19)

To a stirred solution of **17** (1.00 g, 2.89 mmol, E/Z ratio of 1:0.5) in 20 mL of DMF under a N_2 atmosphere was added NBS (540 mg, 3.03 mmol). The reaction

mixture was stirred for 15 min and quenched with water and diluted with ether. The water layer was extracted with 2x20 mL of ether. The combined organic extract was washed with 2x10 mL of brine, 2x10 mL of water and dried over MgSO₄. The solvent was evaporated to give a light green residue which was purified using silica gel chromatography (5 % EtOAc/hexanes) to give the desired compound **19** (1.09 g, 90 %) as a light green oil and as a 1.0 : 0.4 ratio of E : Z isomers. These isomers were not separable and the following spectral data were collected on the mixture. Spectral data for **19**: ¹H NMR (500 MHz, CDCl₃) (E isomer) δ 0.86 (t, 3H, J = 7.2 Hz), 1.10-1.41 (m, 10H), 1.8 (d, 3H, J = 7.1 Hz), 2.37 (t, 2H, J = 7.5 Hz), 5.28 (s, 2H), 5.69 (q, 1H, J = 6.9 Hz), 6.92-7.32 (m, 8H), 7.54-7.64 (m, 1H); (Z isomer) δ 0.86 (t, 3H, J = 7.1 Hz), 1.10-1.41 (m, 10H), 1.48 (dt, 3H, J = 6.9, 1.1 Hz), 2.15-2.25 (m, 2H), 5.20 (d, 1 H, J = 6.8 Hz), 5.30(d, 1H, J = 6.8 Hz), 5.92 (qt, 1H, J = 6.8, 1.3 Hz), 6.92-7.32 (m, 8H), 7.54-7.64 (m, 1H); ¹³C-NMR (125 MHz, CDCl₃) δ 13.91, 13.99, 15.13, 22.59, 28.10, 29.05, 29.07, 29.33, 29.63, 30.93, 31.81, 37.63, 48.08, 48.19, 90.32, 91.02, 110.38, 110.45, 119.05, 119.15, 120.28, 120.38, 122.36, 122.44, 125.99, 126.23, 127.23, 127.28, 127.54, 128.58, 128.63, 129.93, 131.18, 131.22, 131.58, 136.03, 136.33, 137.55, 137.61, 137.87, 140.91, 2 aliphatic and 3 aryl carbons not located; IR (CH₂Cl₂,) 3061s, 3031s, 2953s, 2924brs, 2835brs, 1497s, 1455s cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 425 M⁺ (52, ⁸¹Br), 423 M⁺ (45, ⁷⁹Br), 344 (24), 91 (100); HRMS (FAB⁺) calcd for C₂₅H₃₀BrN 423.1562 (⁷⁹Br) found 423.1560 (⁷⁹Br).

1-(4-Methoxybenzyl)-3-bromo-2-(dec-2-en-3-yl)-1H-indole (**20**)

To a stirred solution of **18** (1 g, 2.66 mmol, E/Z ratio of 1:0.6) in 20 mL of DMF under N₂ atmosphere added NBS (480 mg, 2.70 mmol). The reaction mixture was stirred for 30 min, quenched with water (20 mL) and diluted with ether. The water layer was

extracted with 2x30 mL of ether. The combined organic extract was washed with 2x10 mL of brine, 2x10 mL of water and dried over MgSO_4 . The solvent was evaporated to give a light green residue, which was purified using silica gel chromatography (5% EtOAc/hexanes) to give the desired compound **20** (1.07 g, 89 %) as a light green oil and as a 1.0 : 0.5 ratio of E : Z isomers. These isomers were not separable and the following spectral data were collected on the mixture. Spectral data of **20**: ^1H NMR (300 MHz, CDCl_3) (E-isomer) δ 0.88 (t, 3H, J = 7.1 Hz), 1.10-1.44 (m, 10H), 1.85 (d, 3H, J = 7.0 Hz), 2.38 (t, 2H, J = 7.5 Hz), 3.76 (s, 3H), 5.23 (s, 2H), 5.73 (q, 1H, J = 6.9 Hz), 7.08-7.25 (m, 7H), 7.58-7.65 (m, 1H); (Z-isomer) δ 0.89 (t, 3H, J = 7.1 Hz), 1.10-1.44 (m, 10H), 1.51 (td, 3H, J = 6.6, 1.1 Hz), 2.20-2.28 (m, 2H), 3.77 (s, 3H), 5.16 (d, 1H, J = 6.4 Hz), 5.25 (d, 1H, J = 6.4 Hz), 5.95 (qt, 1H, J = 6.6, 1.3 Hz), 7.08-7.25 (m, 7H), 7.58-7.65 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ 13.95, 14.04, 15.18, 22.60, 28.07, 29.07, 29.09, 29.32, 29.61, 30.91, 31.81, 37.64, 47.52, 47.62, 55.16, 90.12, 90.85, 110.41, 110.49, 113.95, 113.99, 118.95, 119.04, 120.17, 120.29, 122.25, 122.33, 127.14, 127.42, 129.57, 129.80, 129.83, 131.12, 131.15, 131.48, 135.86, 136.15, 137.42, 140.81, 158.76, 158.81, 3 aliphatic and 4 aryl carbons not located; IR (CH_2Cl_2) 2998s, 2955s, 2926brs, 2855s, 1613s, 1512s cm^{-1} ; mass spectrum (EI) m/z (% rel intensity) 455 M^+ (36, ^{81}Br), 453 M^+ (33, ^{79}Br), 289 (5), 121 (100); HRMS (FAB^+) calcd for $\text{C}_{26}\text{H}_{32}\text{BrNO}$ m/z 453.1667 (^{79}Br), measd 453.1668 (^{79}Br).

(1-benzyl-2-(dec-2-en-3-yl)-1*H*-indol-3-yl)(methoxy)carbene pentacarbonyl chromium (0) (21)

To a flame-dried round-bottomed flask filled with Argon was added bromoindole **19** (1.23 g, 2.90 mmol, E:Z 1/0.58) in 20 mL of ether. The solution was cooled to -78°C and *t*-butyl lithium (3.4 mL, 1.7 M, 5.80 mmol) was added drop wise. The resulting

solution was stirred at $-78\text{ }^{\circ}\text{C}$ for 30 min, and then at $0\text{ }^{\circ}\text{C}$ for 1h. This solution was then cooled to $-78\text{ }^{\circ}\text{C}$ and transferred by cannula into a side arm flask containing $\text{Cr}(\text{CO})_6$ (702 mg, 3.19 mmol) and 15 mL of ether. The system was deoxygenated by freeze-thaw method (3 cycles) and then stirred at room temperature for 18 h. The reaction mixture was then cooled to $0\text{ }^{\circ}\text{C}$ and Me_3OBF_4 (858 mg, 5.8 mmol) was added. After stirring for 30 min at $0\text{ }^{\circ}\text{C}$, the solvent was evaporated to give a red solid residue, which contained carbene complex **21** (E/Z 1:0.18) along with the reduced product **17** (E/Z 1:1.8). The ratio of **21**:**17** in the crude mixture was 1:0.18. The residue was purified by column chromatography using ethyl acetate/hexanes (3 : 97) to give 860 mg of a red oil that was a mixture of **21** and **17**. The ratio of **21** to **17** was determined to be 1.0 : 0.3 by integration of the 3-indole hydrogens of the E and Z-isomers of **17** versus the methoxy hydrogens of the E- and Z-isomers of **21**. This reveals that this reaction gave 730 mg (43 %) of carbene complex **21** which was used in subsequent reactions without further purification. R_f of **21** (hexanes/EtOAc 10:1) = 0.60. Spectral data for **21**: (E-isomer), 300 MHz ^1H NMR(CDCl_3): δ 0.74-0.92 (m, 3H), 1.00-1.80 (m, 13 H), 1.80-2.00 (m, 1H), 2.06-2.25 (m, 1H), 4.46 (s, 3H), 5.24 (s, 2H), 5.81 (q, 1H, $J = 7.4\text{ Hz}$), 6.80-7.40 (m, 8H), 8.12 (d, 1H, $J = 8.0\text{ Hz}$). (Z isomer), 300 MHz ^1H NMR(CDCl_3): δ 0.74 – 0.92 (m, 3H), 1.00 – 1.80 (m, 13H), 2.40-3.04 (m, 2H), 4.55 (s, 3H), 4.96 (q, 1H, $J = 6.2\text{ Hz}$), 5.32 (d, 1H, $J = 17.0\text{ Hz}$), 5.42 (d, 1H, $J = 17\text{ Hz}$), 6.80-7.40 (m, 8H), 7.86 (d, 1H, $J = 8.7\text{ Hz}$).

(1-(4-Methoxybenzyl)-2-(dec-2-en-3-yl)-1H-indol-3-yl)(methoxy)carbene pentacarbonyl chromium (0) (22**)**

To a flame-dried round-bottomed flask filled with Argon was added bromoindole **20** (1.07 g, 2.36 mmol, E/Z 1:0.5) in 20 mL of ether (degenated by the freeze-thaw method). The solution was cooled to $-78\text{ }^{\circ}\text{C}$ and *tert*-butyllithium (2.8 mL, 1.7 M, 4.76

mmol) was added drop wise. The resulting solution was stirred at $-78\text{ }^{\circ}\text{C}$ for 30 min and then at $0\text{ }^{\circ}\text{C}$ for 1 h. This solution was then cooled to $-78\text{ }^{\circ}\text{C}$ and transferred by cannula into a side arm flask containing $\text{Cr}(\text{CO})_6$ (529 mg, 2.8 mmol) and 15 mL of ether. The system was deoxygenated by the freeze-thaw method (3 cycles) and then stirred at room temperature for 18 h. The reaction mixture was then cooled to $0\text{ }^{\circ}\text{C}$ and Me_3OBF_4 (698 mg, 4.72 mmol) was added. After stirring for 30 min at $0\text{ }^{\circ}\text{C}$, the solvent was evaporated to give a red solid residue which contained carbene complex **22** (E/Z 1:0.24) along with the reduced product **18** (E/Z 1:2.00). The ratio of **22:18** in the crude mixture was 1:0.29. The reduced product could be separated by column chromatography using (3 % ethyl acetate/hexanes) to give compound **22** (618 mg, 43%) as dark red oil. Spectral data for **22**: (E isomer), 300 MHz ^1H NMR(CDCl_3): δ 0.83 (t, 3H, $J = 6.9\text{ Hz}$), 1.05 – 1.80 (m, 13H), 1.80-2.30 (m, 1H), 2.06-2.26 (m, 1H), 3.77 (s, 3H), 4.57 (s, 3H), 5.18 (s, 2H), 5.82 (q, 1H, $J = 6.9\text{ Hz}$), 6.76 (d, 2H, $J = 8.8\text{ Hz}$), 6.86 (d, 2H, $J = 8.5\text{ Hz}$), 7.10 – 7.30 (m, 3H), 8.12 (d, 1H, $J = 8.0\text{ Hz}$); (Z isomer), 300 MHz ^1H NMR(CDCl_3): δ 0.85 (t, 3H, $J = 7.2\text{ Hz}$), 1.05 – 1.80 (m, 13H), 2.48 – 2.64 (m, 1H), 2.90-3.06 (m, 1H), 3.76 (s, 3H), 4.61 (s, 3H), 4.96 (q, 1H, $J = 6.3\text{ Hz}$), 5.25 (d, 1H, $J = 17.0\text{ Hz}$), 5.36 (d, 1H, $J = 16.7\text{ Hz}$), 6.82 (d, 2H, $J = 8.8\text{ Hz}$), 6.97 (d, 2H, $J = 8.8\text{ Hz}$), 7.80-7.32(m, 3H), 7.85 (d, 1H, $J = 5.8\text{ Hz}$).

(1-Benzyl-2-(dec-2-en-3-yl)-1H-indol-3-yl)(methoxymethoxy)carbene pentacarbonyl chromium (0) (23)

To a flame-dried round-bottomed flask filled with argon was added bromoindole **19** (1.01g, 2.38 mmol) in 20 mL of ether which had been deoxygenated by the freeze-thaw method. The solution was cooled to $-78\text{ }^{\circ}\text{C}$ and *tert*-butyl lithium (2.8 mL, 1.7 M, 4.76 mmol) was added drop wise. The resulting solution was stirred at $-78\text{ }^{\circ}\text{C}$ for 30 min

and then at 0 °C for 1 h. This solution was then cooled to –78 °C and transferred by cannula into a side-armed flask containing Cr(CO)₆ (529 mg, 2.85 mmol) and 15 mL of ether. The system was deoxygenated by the freeze-thaw method (3 cycles) and then stirred at room temperature for 20 h. The reaction mixture was then cooled to 0 °C and MOMCl (271 μL, 3.57 mmol) was added. After stirring for 30 min at 0 °C, the solvent was evaporated to give a red solid residue. The residue was purified by column chromatography using (10 % ethyl acetate/hexanes) to give a dark red oil which mainly consisted of desired carbene complex **23** (747 mg, 48 % yield) with trace amount of impurities. This compound was used in subsequent steps without further purification. Spectral data for **23**: (E-isomer); 300 MHz ¹H NMR(CDCl₃): δ 0.68-0.90 (m, 3H), 1.00-1.80 (m, 13H), 1.80-2.24 (m, 2H), 3.73 (s, 3H), 4.45 (s, 2H), 5.17 (s, 2H), 5.81 (q, 1H, J = 7.1 Hz), 6.64-7.32 (m, 8H), 8.11 (d, 1H, J = 7.8 Hz), (Z-isomer), 300 MHz ¹H NMR(CDCl₃): δ 0.68-0.90 (m, 3H), 1.00-1.80 (m, 13H), 2.48-2.64 (m, 1H), 2.88-3.04 (m, 1H), 3.76 (s, 3H), 4.56 (s, 2H), 4.96 (q, 1H, J = 6.3 Hz), 5.26 (d, 1H, J = 6.7 Hz), 5.34 (d, 1H, J = 6.7 Hz), 6.82 (d, 2 H, J = 6.6 Hz), 6.97 (d, 2 H, J = 8.5 Hz), 7.10-7.28 (m, 4H), 7.84 (dd, 1H, J = 6.9, 1.1 Hz).

N-Benzyl-1-heptyl-4-methoxy-2-methyl-9*H*-carbazol-3-ol (**25**)

A solution of carbene **21** (472 mg, 0.815 mmol) in 150 mL THF in a quartz photoreactor was purged with nitrogen for 15 min and then purged with carbon monoxide for another 15 min. The solution was irradiated with a 450 W medium pressure mercury lamp for 30 min at room temperature. The resulting solution was kept under a CO atmosphere for 12 h. The solvent was then evaporated to give red oily residue. The residue was purified by column chromatography (3 % ethyl acetate/hexanes) to give **25** (223 mg, 65 %) as a light brown solid; mp 102 - 104 °C (ethyl acetate/hexanes). Spectral data for **25**: ¹H NMR (500 MHz, CDCl₃) δ 0.91 (t, 3H, J = 6.9 Hz), 1.20-1.40 (m, 8H),

1.42 -1.64 (m, 2H), 2.38 (s, 3H), 2.74-2.84 (m, 2H), 4.06 (s, 3H), 5.65 (s, 1H), 5.66 (s, 2 H), 7.04 (d, 2H, J = 6.8 Hz), 7.19-7.30 (m, 5H), 7.34-7.40 (1H, m), 8.20 (dd, 1H, J = 7.8, 1.3 Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 11.98, 14.06, 22.62, 28.18, 29.10, 29.78, 31.79, 31.85, 48.68, 60.53, 108.75, 114.71, 119.23, 121.02, 122.13, 122.98, 124.34, 125.41, 127.11, 128.82, 134.22, 138.66, 138.79, 140.62, 142.10, 1 aryl C not located; IR (CH_2Cl_2) 3544s, 2959s, 2928s, 2857s cm^{-1} ; mass spectrum (EI) m/z (% rel intensity) 415 (65), 330 (20), 240 (80), 91 (100). Anal calcd for $\text{C}_{28}\text{H}_{33}\text{NO}_2$: C, 80.93; H, 8.00; N, 3.37. Found C, 81.23; H, 7.84; N, 3.48.

9-(4-Methoxybenzyl)-1-heptyl-4-methoxy-2-methyl-9H-carbazol-3-ol (26)

The *ortho*-benzannulation of dienyl-chromium carbene complex **22** was performed following the procedure described above for complex **21**. The desired carbazole **26** was obtained after purification using column chromatography on silica gel (20 % ether/hexane) in 62 % yield as a white solid; mp 108 - 110 $^{\circ}\text{C}$. Spectral data for **26**: ^1H NMR (300 MHz, CDCl_3) δ 0.87 (t, 3H, J = 6.9 Hz), 1.18-1.42 (m, 8H), 1.44-1.62 (m, 2H), 2.34 (s, 3H), 2.68-2.82 (m, 2H), 3.73 (s, 3H), 4.03 (s, 3H), 5.64 (s, 2H), 5.60 (s, 1H), 6.78 (d, 2H, J = 8.7 Hz), 6.92 (d, 2H, J = 9 Hz), 7.14-7.28 (m, 2H), 7.30-7.42 (m, 1 H), 8.17 (d, 1H, J = 7.8 Hz); ^{13}C NMR (125 MHz, CDCl_3) δ 11.97, 14.08, 22.64, 28.18, 29.14, 29.81, 31.79, 31.86, 48.11, 55.21, 60.54, 108.78, 114.25, 114.67, 119.17, 120.98, 121.04, 122.11, 122.93, 125.32, 126.53, 130.65, 134.21, 138.76, 140.56, 142.07, 158.73; IR (CH_2Cl_2) 3544s, 2959s, 2928s, 2857s., 2255s, 1512s, 1458s cm^{-1} ; mass spectrum (EI) m/z (% rel intensity) 445 M^+ (30), 240 (14), 121 (100). Anal calcd for $\text{C}_{29}\text{H}_{35}\text{NO}_3$: C, 78.17; H, 7.92; N, 3.14. Found C, 77.70; H, 8.07; N, 3.16.

9-Benzyl-1-heptyl-3,4-dimethoxy-2-methyl-9H-carbazole (26a)

To a stirred suspension of NaH (57 mg, 1.420 mmol) in THF (3.5 mL) was added a solution of **25** (295 mg, 0.710 mmol) in 1.5 mL of THF. After 15 min, Me₂SO₄ (336 μL, 3.550 mmol) was added. The reaction mixture was further stirred for 6 h, quenched with water and extracted with ether (2x20 mL). The organic extract was washed with 1x10 mL of brine and 1x10 mL of water, dried over MgSO₄ and evaporated to give an oily residue. The residue was purified using silica gel column chromatography (5 % ethyl acetate/hexanes) to give 291 mg (96 % yield) of compound **26a** as colorless wax. Spectral data for **26a**: ¹H NMR (500 MHz, CDCl₃) δ 0.88 (t, 3H, J = 7.0 Hz), 1.18-1.38 (m, 8H), 1.52-1.62 (m, 2H), 2.35 (s, 3H), 2.70-2.78 (m, 2H), 3.89 (s, 3H), 4.12 (s, 3H), 5.63 (s, 2H), 7.02 (d, 2H, J = 7.0 Hz), 7.16-7.30 (m, 5H), 7.30-7.38 (m, 1H), 8.32 (dd, 1H, J = 8.1, 1.1 Hz); ¹³C NMR (125 MHz, CDCl₃) δ 12.16, 14.07, 22.63, 28.43, 29.11, 29.82, 31.78, 31.86, 48.62, 60.24, 60.77, 108.56, 115.94, 119.42, 120.07, 122.15, 122.47, 125.26, 125.45, 127.11, 128.82, 129.47, 136.59, 138.65, 142.02, 144.38, 146.04; IR (CH₂Cl₂) 2960m, 2928m, 2855m cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 429 M⁺ (61), 414 (28), 344 (23), 254 (47), 91 (100). Anal calcd for C₂₉H₃₅NO₂: C, 81.08; H, 8.21; N, 3.26. Found C, 80.90; H, 8.27; N, 3.29.

1-Heptyl-3,4-dimethoxy-2-methyl-9H-carbazole (27)

To a stirred of **26a** (0.521 mmol, 224 mg) in 15 mL DMSO was added potassium *tert*-butoxide (6.63 mmol, 6.63 mL of 1 M of potassium *tert*-butoxide in THF) at room temperature. The reaction mixture was stirred for 5 min and O₂ was bubbled through the reaction mixture for 30 min. The reaction mixture was quenched with a saturated solution of NH₄Cl (10 mL) and extracted with 3x10 mL of ether. The organic extract was washed with 10 ml of water, dried over MgSO₄ and evaporated to give the an oily residue. This residue was purified using silica gel column chromatography (5 % ethyl acetate/hexanes) to give 173 mg of compound **27** (98 % yield) as white solid; mp 80 - 82 °C. Spectral data for **27**: ¹H NMR (500 MHz, CDCl₃) δ 0.89 (t, 3H, J = 6.8 Hz), 1.20-1.40 (m, 6 H), 1.40-1.50 (m, 2H), 1.55-1.70 (m, 2H), 2.39 (s, 3H), 2.82 (m, 2H, J = 7.9 Hz), 3.90 (s, 3H), 4.11 (s, 3H), 7.21 (t, 1H, J = 7.3 Hz), 7.30-7.45 (m, 2H), 7.80 (s, 1H), 8.23 (d, 1H, J = 7.7 Hz); ¹³C NMR(125 MHz, CDCl₃) δ 12.20, 14.07, 22.66, 28.64, 29.29, 29.52, 30.00, 31.85, 60.40, 60.92, 110.17, 114.64, 118.79, 119.39, 122.50, 122.87, 125.01, 128.35, 136.11, 139.33, 144.56, 146.03; IR (CH₂Cl₂) 3467s, 3054s, 2957s, 2930s, 2857s cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 339 M⁺ (84), 324 (100), 296 (12), 254 (72). Anal Calcd for C₂₂H₂₉NO₂: C, 77.84; H, 8.61; N, 4.13. Found C, 77.81; H, 8.76; N, 4.04.

9-benzyl-1-heptyl-2-methyl-9H-carbazole-3,4-dione 28:

Carbazole derivative **25** (100 mg, 0.24 mmol) was dissolved in 10 mL of THF/H₂O (3:1). This solution was stirred for 5 min and CAN (394 mg, 0.72 mmol) was added. The mixture was stirred for 2 h, diluted with H₂O and extracted with three portions of ethyl acetate. The solvent was removed under vacuuo and residue was purified using flash chromatography on silica gel (hexanes:EtOAc = 5 : 1) to give **28** (72

mg, 75 %). $R_f = 0.16$ (hexanes/EtOAc = 9 : 2). Spectral data for **28**: ^1H NMR(300 MHz, CDCl_3): δ 0.87 (t, 3H, $J = 6.6$ Hz), 1.15-1.39 (m, 8H), 1.48-1.62 (m, 2H), 1.98 (s, 3H), 2.47-2.54 (m, 2H, $J = 8.7$ Hz), 5.57(s, 2H), 6.94 (dd, 2H, $J = 7.5, 1.8$ Hz), 7.19-7.39 (m, 6H), 8.25 (dd, 1H, $J = 7.2, 1.2$ Hz); ^{13}C NMR (75 MHz, CDCl_3) δ 11.9, 14.0, 22.5, 28.8, 28.9, 29.4, 29.7, 31.7, 48.9, 110.8, 122.1, 125.0, 125.1, 125.7, 128.1, 129.33, 134.67, 135.7, 139.5, 142.4, 144.6, 174.0, 183.2, 2 aryl carbons not located.

Carbazaquinocin-C (1C) from **27**.

To a stirred solution of **27** (0.074 mmol, 25 mg) in CH_2Cl_2 added BBr_3 (221 μL , 0.221 mmol, 1 M in THF) at -78°C . The reaction mixture was brought to room temperature over 30 min. After 30 min, the reaction mixture was diluted with CH_2Cl_2 (20 mL) and washed with 2 x 5 mL of water. To the organic extract was added excess NaIO_4 and the resulting solution was stirred for 20 min. At this point, the organic extract was filtered through a small pad of Celite. The resultant filtrate was evaporated to give the desired compound carbazaquinocin-C (17.6 mg) in 77 % yield which was clean by ^1H NMR and whose mp ($226\text{-}228^\circ\text{C}$) matches that in the literature ($227\text{-}228^\circ\text{C}$).² The ^1H and ^{13}C NMR spectra matched the published spectra.² This compound is dark green as a solid and is red in solution and is stable in air and at room temperature. The compound binds strongly to silica gel and is difficult to purify by chromatograph. Spectral data for carbazaquinocin C: ^1H NMR (300 MHz, DMSO) δ 0.86 (t, 3H, $J = 6.4$ Hz), 1.20-1.40 (m, 6 H), 1.40-1.62 (m, 4H), 1.90 (s, 3H), 2.66 (t, 2H, $J = 7.6$ Hz), 7.20-7.30 (m, 2H), 7.48-7.56 (m, 1H), 7.82-7.92 (m, 1H), 12.35 (brs, 1H); ^{13}C NMR (125 MHz, DMSO), δ 11.41, 13.91, 22.02, 28.02, 28.48, 28.59, 28.96, 31.28, 110.98, 113.33, 120.21, 123.92, 124.15, 125.60, 133.06, 137.00, 142.09, 145.57, 172.68, 183.41; mass spectrum (FAB)

m/z (% rel intensity) 311 (8), 310 (M+1)⁺ (11), 231 (17), 215 (11), 117 (100), 95 (60), 81 (60), 69 (75), 55 (85), 57 (70), 55 (85); HRMS (FAB) calcd for C₂₀H₂₄NO₂ m/z 310.1807, meas 310.1808.

N-tert-Butyl-9-benzyl-1-heptyl-4-methoxy-2-methyl-9H-carbazol-3-amine (30)

To a solution of **21** (710 mg, 1.23 mmol) in 1 mL THF was added *t*-BuNC (974 μ L, 8.61 mmol). The reaction mixture was stirred at room temperature for 24 h. After 24 h, the reaction mixture was diluted with 2 ml of THF and heated to 75 °C for another 18 h. The reaction mixture was concentrated and the desired compound **30** was isolated in 86 % yield upon silica gel chromatography (5 %, ethyl acetate/hexanes) as a light yellow waxy liquid. Spectral data for **30**: ¹H NMR (300 MHz, CDCl₃): δ 0.90 (t, 3H, J = 7.0 Hz), 1.21 (s, 9H), 1.24-1.42 (m, 8H), 1.50-1.65 (m, 2H), 2.47 (s, 3 H), 2.73-2.85 (m, 2H), 2.95-3.20 (brs, 1H), 3.98 (s, 3H), 5.65 (s, 2H), 7.09 (d, 2 H, J = 7.1 Hz), 7.23 (t, 3H, J = 7.7 Hz), 7.26-7.32 (m, 2H), 7.32-7.39 (m, 1H), 8.26 (dd, 1H, J = 8.6, 1.3 Hz); ¹³C NMR (125 MHz, CDCl₃) δ 14.07, 16.41, 22.62, 28.82, 29.12, 29.84, 30.38, 31.62, 31.87, 48.78, 55.40, 59.69, 108.59, 115.27, 119.40, 120.44, 121.72, 122.44, 125.08, 125.52, 127.07, 128.79, 130.38, 134.43, 137.43, 138.75, 142.01, 149.52; IR (CH₂Cl₂) 2959s, 2928s, 2859s, 1453s cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 472 (23), 470 (82), 455 (13), 414 (27), 413 (79), 399 (12), 239 (22), 149 (37), 130 (14), 91 (100), 57 (50). Anal calcd for C₃₂H₄₂N₂O: C, 81.66; H, 8.99; N, 5.95. Found C, 81.80; H, 9.12; N, 5.85.

***N*-tert-Butyl-1-heptyl-4-methoxy-2-methyl-9*H*-carbazol-3-amine (32)**

To a stirred of **30** (0.038mmol, 17.8mg) in 15ml DMSO added potassium *tert*-butoxide (0.38mmol, 380 μ l, 1M sol of potassium *tert*-butoxide in THF) at room temperature. The reaction mixture was stirred for 5 min and O₂ was bubbled through the reaction mixture for 30min. The reaction mixture was quenched with saturated solution of NH₄Cl (1*5ml), extracted with 3*5ml of ether. The organic extract was washed with 1* 10ml of water, dried over MgSO₄ and evaporated to give the oily residue. This residue was purified using silica gel column chromatography (20%, Ethylacetate/hexanes) to give 173mg of compound **32** (76% yield) as light yellow viscous liquid. Spectral data for **32**: ¹H NMR (300MHz, CDCl₃): δ 0.91 (t, 3H, J = 6.6 Hz), 1.23 (s, 9H), 1.26-1.56 (m, 8H), 1.58-1.74 (m, 2H), 2.50 (s, 3H), 2.85 (t, 2H, J = 7.8 Hz), 3.10 (brs, 1H), 3.98 (s, 3H), 7.16-7.28 (m, 1H), 7.32-7.50 (m, 2H), 7.84 (s, 1H), 8.20 (d, 1H, J = 7.8 Hz); ¹³C NMR (125 Mhz, CDCl₃) δ 14.08, 16.34, 22.63, 29.04, 29.29, 30.00, 30.36, 31.85, 55.34, 59.78, 110.16, 113.90, 119.07, 119.34, 122.37, 122.50, 124.82, 130.19, 132.74, 136.77, 139.23, 149.24, one aliphatic C missing; IR (CH₂Cl₂) 3438s, 2926brm, 2857s, 1709), 1611s, 1502s, 1455s cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 381 (21), 380 M⁺ (90), 365 (16), 323 (47), 323 (100), 311 (12), 309 (62), 295 (10), 239 (24), 233 (14), 210 (15), 197 (18), 196 (26), 195 (17), 180 (8). Anal calcd for C₂₅H₃₆N₂O: C, 78.90; H, 9.53; N, 7.36. Found C, 78.88; H, 9.54; N, 7.08.

***N*-tert-Butyl-9-benzyl-1-heptyl-4-(methoxymethoxy)-2-methyl-9*H*-carbazol-3-amine
(**31**)**

To a solution of **23** (0.580 mg, 0.93 mmol) in 750 μ L THF was added *t*-BuNC (974 μ L, 8.61 mmol). The reaction mixture was stirred at room temperature for 24 h. After 24 h, the reaction mixture was diluted with 2 mL of THF and heated to 75 $^{\circ}$ C for another 18 h. The reaction mixture was concentrated and the desired compound **31** was isolated in 72 % yield as a yellow waxy solid after silica gel chromatography (5 % ethyl acetate/hexanes). Spectral data for **31**: ^1H NMR (300 MHz, CDCl_3) δ 0.78-0.92 (m, 3H), 1.17 (s, 9H), 1.20-1.42 (m, 8H), 1.46-1.64 (m, 2H), 2.47 (s, 3H), 2.70-2.89 (m, 2 H), 3.59-3.63 (m, 3H), 5.27 (s, 2H), 5.63 (s, 2H), 7.05 (d, 2H, $J = 7.7$ Hz), 7.12-7.40 (m, 6H), 8.15 (d, 2H, $J = 8.0$ Hz); ^{13}C NMR (125 MHz CDCl_3) δ 14.06, 16.78, 22.61, 28.83, 29.11, 29.85, 30.19, 31.51, 31.87, 48.74, 55.37, 57.80, 100.00, 108.71, 115.05, 119.33, 120.93, 121.66, 121.71, 125.14, 125.51, 127.19, 128.80, 130.90, 134.75, 137.56, 138.68, 142.03, 147.46; IR (CH_2Cl_2) 3350s, 3054s, 2959s, 2928s, 1701s, 1593s, 1497s, 1453s cm^{-1} ; mass spectrum (EI) m/z (% rel intensity) 500 M^+ (19), 455 (8), 433 (7), 411 (8), 400 (19), 399 (72), 388 (15), 371 (8), 361 (7), 287 (12), 91 (100). Anal calcd for $\text{C}_{33}\text{H}_{44}\text{N}_2\text{O}_2$: C, 79.16; H, 8.86; N, 5.59. Found C, 79.06; H, 8.99; N, 5.23.

***N*-tert-Butyl-1-heptyl-4-(methoxymethoxy)-2-methyl-9*H*-carbazol-3-amine (33)**

To a stirred of **31** (0.16 mmol, 82 mg) in 5 mL DMSO was added potassium *tert*-butoxide (1.6 mmol, 1.6 mL, 1 M potassium *tert*-butoxide in THF) at room temperature. The reaction mixture was stirred for 5 min and O₂ was bubbled through the reaction mixture for 30 min. The reaction mixture was quenched with saturated aqueous NH₄Cl (5 mL) and extracted with 3x5 mL of ether. The organic extract was washed with 10 mL of water, dried over MgSO₄ and evaporated to give the an oily residue. This residue was purified by silica gel column chromatography (20 % ethyl acetate/hexanes) to give 173 mg of compound **33** (72 % yield) as a light yellow viscous oil. Spectral data for **33**: ¹H NMR (300 MHz, CDCl₃) δ 0.88 (t, 3H, J = 7.0 Hz), 1.19 (s, 9H), 1.22-1.50 (m, 8H), 1.56-1.67 (m, 2H), 2.49 (s, 3H), 2.82 (t, 2H, J = 8.1 Hz), 3.59 (s, 3H), 5.82 (s, 2H), 7.17-7.21 (m, 1H), 7.32-7.39 (m, 1H), 7.41 (d, 1H, J = 8.2 Hz), 7.86 (s, 1H), 8.08 (d, 1H, J = 8.0 Hz), 1 NH proton not located; ¹³C NMR (125 MHz, CDCl₃) δ 14.11, 16.67, 22.64, 29.06, 29.21, 29.29, 30.04, 30.16, 31.86, 55.39, 57.82, 99.84, 110.28, 113.66, 119.28, 119.55, 121.74, 122.27, 124.90, 130.59, 133.42, 136.88, 139.24, 147.27; IR (CH₂Cl₂) 3357brs, 3058s, 2926brm, 2857brs, 1709s, 1611s, 1455s, 1406s cm⁻¹; mass spectrum (EI) m/z (% rel intensity) 410 M⁺ (18), 365 (6), 321 (11), 309 (100). Anal calcd for C₂₆H₃₈N₂O₂: C, 76.06; H, 9.33; N, 6.82. Found C, 76.54; H, 9.64; N, 6.64.

Carbazaquinocin-C from **33**.

To a stirred solution of **33** (12 mg, 0.03 mmol) in 1 mL of MeOH and 300 μ L of H₂O was added 200 μ L of conc. HCl. The reaction mixture was refluxed for 45 min. The resulting mixture was concentrated in vacuo and purified by silica gel column chromatography (30 % ethyl acetate/hexanes). The resulting compound was dissolved in 1 mL CH₂Cl₂ and 300 μ L of H₂O. To this mixture was added excess NaIO₄ (200 mg) and the reaction mixture stirred for 24 h at rt. After 24h, the reaction mixture was diluted with 2 mL of CH₂Cl₂ and washed with 3x2 mL of water. The resulting organic layer was passed through Celite and washed with CH₂Cl₂. The organic solvent was then evaporated to give carbazaquinocin C as dark green solid in 82 % (7.5 mg) yield. This compound had spectral data identical to the compound obtained from **28** as described above.

References:

- 1) Mouloud, D.; Dodd, R. H. *Tetrahedron* **1994**, *50*, 6299.
- 2) Knolker, H.-J.; Frohner, W.; Reddy, K. R., *Synthesis*, **2002**, 557.