Supporting Information

Total Synthesis of Cystothiazole A & B

Jian Shao and James S. Panek*

*Department of Chemistry and Center for Chemical Methodology and Library Development,
Boston University, 590 Commonwealth Avenue, Boston, Massachusetts 02215*

General Information

¹H- and ¹³C-NMR were recorded in CDCl₃ at 400 MHz and 100 MHz respectively unless specified otherwise. Chemical shifts are reported in parts per million using the solvent resonance internal standard chloroform (7.24 and 77.0 ppm, unless specified otherwise). Data are reported as follows: chemical shift, multiplicity (app = apparent, ovrlp = overlapping, s = singlet, d = doublet, t = triplet, q = quartet, sept = septuplet, m = multiplet, br = broad, abq = ab quartet), integration, and coupling constant. Ratios of diastereomers (dr) were determined by ¹H-NMR (400 MHz) operating at a signal/noise ratio of >200:1. Infrared Resonance (IR) spectra were recorded on a Nicolet Impact 400 FT-IR spectrometer. Optical rotations were recorded on an AUTOPOL III digital polarimeter at 589 nm, and are reported as [α]_D (concentration in grams/100 mL solvent). Tetrahydrofuran (THF), ethyl ether (Et₂O), methylene chloride (CH₂Cl₂), benzene and toluene were dried using a solvent drying system (Al₂O₃). Dioxane was distilled under Argon from sodium. All other reagents were used as supplied from commercial sources. All reactions were carried out in oven-dried glassware under argon or N₂ atmosphere unless otherwise noted. Analytical thin layer chromatography was performed using Whatman Reagent silica gel 60-A plates. Flash chromatography¹ was performed using E. Merck silica gel (230-400 mesh).

Spectral data

β -Hydroxy ester 15: Homoallylic ether **12** (226 mg, 0.843 mmol) was dissolved in CH₂Cl₂ (12 mL) and cooled to -78 °C. Ozone was bubbled through the solution for about 5 min, and the reaction was monitored by TLC. Excess dimethyl sulfide (DMS) was added to quench the ozonides, and the reaction was allowed to stir at room temperature for 16 hours. Aqueous workup was followed by extraction with ethyl acetate, washing with brine and finally drying with MgSO₄. Removal of solvent under reduced pressure and purification by flash chromatography (silica gel, 2% Et₂O in pentane) afforded aldehyde **14** (114 mg, 68%) as a pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 9.75 (s, 1H), 4.24 (d, 1H, *J* = 4.8 Hz), 3.40 (s, 3H),

¹ Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923-2925

2.59 (dq, 1H, $J = 7.2, 4.8$ Hz), 1.19 (d, 3H, $J = 7.2$ Hz), 0.17 (s, 9H). Aldehyde **14** is volatile and unstable at room temperature, especially in high concentration, and was submitted to the next step immediately.

To a solution of diisopropylamine (0.130 mL, 0.93 mmol) in THF (2.8 mL) at -78°C was added *n*-butyl lithium (0.465 mL of a 2.5 M solution in hexane, 1.16 mmol). The solution was warmed up to 0°C and stirred for 15 min. After the solution was cooled back to -78°C , methyl acetate (0.5 M in THF, 1.8 mL, 0.93 mmol) was added dropwise. The mixture was stirred for 20 min at -78°C followed by the addition of aldehyde **14** (46 mg, 0.23 mmol) in THF (1.0 mL). After stirring for 45 min at -78°C , the reaction mixture was diluted with saturated NH_4Cl (10 mL). The mixture was extracted with Et_2O (3 x 10 mL) and the combined organic layers were dried over MgSO_4 . Removal of solvent under reduced pressure and purification by flash chromatography (silica gel, 15% EtOAc in hexane) afforded the ester **15** as a (~2:1 ratio) mixture of diastereomers (54 mg, 90%). IR (neat) ν_{max} : 3511, 2958, 2169, 1739, 1251, 846 cm^{-1} ; CIHRMS $[\text{M}]^+$ calculated for $\text{C}_{13}\text{H}_{24}\text{O}_4\text{Si}$ 272.1444, found 272.1427.

β -Ketoester 6: To a solution of β -hydroxy ester **15** (62 mg, 0.23 mmol) in CH_2Cl_2 (2.3 mL) was added PCC (147 mg, 0.69 mmol). The solution was stirred for 16 h at room temperature. The reaction mixture was filtered through a plug of silica gel, and removal of solvent under reduced pressure afforded 52 mg (85%) of desired β -ketoester **6**. This material was used without further purification. ^1H NMR (400 MHz, CDCl_3) δ 4.12 (d, 1H, $J = 5.6$ Hz), 3.71 (s, 3H), 3.62 (abq, 1H, $J = 16$ Hz), 3.56 (abq, 1H, $J = 16$ Hz), 3.37 (s, 3H), 2.93 (dq, 1H, $J = 6.8, 5.6$ Hz), 1.21 (d, 3H, $J = 6.8$ Hz), 0.16 (s, 9H); Enol form distinguished by ^1H NMR (400 MHz, CDCl_3) 5.09 (s, 1H), 4.16 (d, 1H, $J = 5.2$ Hz), 1.16 (d, 3H, $J = 7.2$ Hz); ^{13}C NMR (100 MHz, CDCl_3) 203.1, 167.8, 101.5, 93.1, 72.3, 56.8, 52.2, 51.1, 48.7, 11.9, -0.3; IR (neat) ν_{max} : 2957, 2171, 1751, 1719, 1252, 846 cm^{-1} ; $[\alpha]_D^{23} = +14.7^{\circ}$ ($c = 0.75$, CHCl_3); CIHRMS $[\text{M}]^+$ calculated for $\text{C}_{13}\text{H}_{22}\text{O}_4\text{Si}$ 270.1287, found 270.1249.

(E)- β -Methoxyacrylate 16: A solution of trimethylorthoformate (0.215 mL, 1.97 mmol), β -ketoester **6** (125 mg, 0.46 mmol), sulfuric acid (10 mg, 0.01 mmol) in MeOH (3.2 mL, 0.15 M) was stirred for 24h at room temperature. The reaction was treated with saturated NaHCO_3 (3 mL). The mixture was extracted with EtOAc (3 x 5 mL) and the combined organic layers were dried over MgSO_4 . Removal of solvent under reduced pressure and purification by flash

chromatography (silica gel, 5% EtOAc in hexane) afforded methoxyacrylate **16** (113 mg, 86%) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 5.00 (s, 1H), 4.23 (dq, 1H, $J = 8.8, 6.8$ Hz), 4.02 (d, 1H, $J = 8.8$ Hz), 3.66 (s, 3H), 3.60 (s, 3H), 3.37 (s, 3H), 1.17 (d, 3H, $J = 6.8$ Hz), 0.11 (s, 9H); ^{13}C NMR (67.5 MHz, CDCl_3) 175.9, 167.6, 103.0, 91.3, 91.0, 73.6, 56.0, 55.5, 50.8, 39.6, 14.0, -0.2; IR (neat) ν_{max} : 2958, 2172, 1716, 1629, 1145 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = +158.4^\circ$ ($c = 0.80$, CHCl_3); CIHRMS $[\text{M}]^+$ calculated for $\text{C}_{14}\text{H}_{24}\text{O}_4\text{Si}$ 284.1444, found 284.1485.

Alkyne 17: To a solution of (E)- β -methoxyacrylate **16** (47 mg, 0.17 mmol) in MeOH (3.3 mL) was added K_2CO_3 (46 mg, 0.33 mmol). The mixture was stirred for 12 h at room temperature before water (5 mL) was added. The mixture was extracted with CH_2Cl_2 (3 x 10 mL). The combined organic layers were dried over MgSO_4 . Removal of solvent under reduced pressure afforded alkyne **17** as a colorless oil (35 mg, 100%). This material was used without further purification. ^1H NMR (400 MHz, CDCl_3) δ 5.02 (s, 1H), 4.23 (dq, 1H, $J = 8.8, 6.8$ Hz), 4.05 (d, 1H, $J = 8.8$ Hz), 3.66 (s, 3H), 3.61 (s, 3H), 3.39 (s, 3H), 2.36 (s, 1H), 1.19 (d, 2H, $J = 6.8$ Hz); ^{13}C NMR (67.5 MHz, CDCl_3) 175.8, 167.6, 91.2, 81.1, 74.1, 73.0, 56.2, 55.7, 50.9, 39.6, 14.1; IR (neat) ν_{max} : 3291, 2941, 1712, 1627, 1148 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = +126.2^\circ$ ($c = 0.60$, CHCl_3); CIHRMS $[\text{M}]^+$ calculated for $\text{C}_{11}\text{H}_{16}\text{O}_4$ 212.1049, found 212.1046.

Vinyl stannane 4: To a solution of alkyne **17** (18 mg, 0.085 mmol) in CH_2Cl_2 (1.7 mL) at 0 $^\circ\text{C}$ was added bis(triphenylphosphine) palladium(II) chloride (6 mg, 0.0085 mmol). Tributyltinhydride (0.034 mL, 0.13 mmol) was added dropwise over 3 min; the reaction mixture was then stirred at 0 $^\circ\text{C}$ for 10 min and warmed to room temperature (20 min). Concentration and flash chromatography (1% EtOAc in hexane, 1% Et_3N was added to the eluent) afforded vinyl stannane **4** as a colorless oil (36 mg, 84%). ^1H NMR (400 MHz, CDCl_3) δ 6.01 (d, 1H, $J = 19.2$), 5.74 (dd, 1H, $J = 19.2, 7.6$ Hz), 4.91 (s, 1H), 4.08 (dq, 1H, $J = 8.4, 6.8$ Hz), 3.62 (s, 3H), 3.52 (s, 3H), 3.50 (dd, 1H, $J = 8.4, 8.0$ Hz), 3.24 (s, 3H), 1.52-1.36 (m, 6H), 1.31-1.22 (m, 6H), 1.13 (d, 3H, $J = 6.8$ Hz), 0.92-0.80 (m, 15H); ^{13}C NMR (67.5 MHz, CDCl_3) δ 176.8, 167.6, 147.1, 131.9, 91.0, 88.3, 56.5, 55.2, 50.7, 39.2, 29.1, 27.2, 14.3, 13.7, 9.4; IR (neat) ν_{max} : 2956, 2926, 1717, 1625, 1143 cm^{-1} ; $[\alpha]_{\text{D}}^{23} = +94.5^\circ$ ($c = 1.0$, CHCl_3); CIHRMS $[\text{M}]^+$ calculated for $\text{C}_{23}\text{H}_{44}\text{O}_4\text{Sn}$ 504.2262, found 504.2308.

Bisthiazole 5a: To a solution of 4-bromo-2-isopropyl thiazole **8a** (385 mg, 1.88 mmol) in diethyl ether (12.5 mL) at -78 °C, was added ^tBuLi (2.32 mL of a 1.7 M solution in hexane, 3.94 mmol). After the solution was stirred at -78 °C for 15 min, tributyltin chloride (0.611 mL, 1.88 mmol) was added slowly. The reaction mixture was stirred at room temperature for 1.5 h, and water (10 mL) was added. The mixture was extracted with diethyl ether (3 x 15 mL). The combined organic layers were dried over MgSO₄. Removal of solvent under reduced pressure afforded the crude thiazolyl stannane **19a** as an orange oil (789 mg, 100%). To a solution of crude thiazolyl stannane **19a** (48 mg, 0.115 mmol) and ditriflate **7** (40 mg, 0.115 mmol) in a sealed tube was added LiCl (15 mg, 0.35 mmol), Pd(PPh₃)₄ (13 mg, 0.012 mmol) and dioxane (1.2 mL, freeze pump thawed for two cycles). The reaction mixture was heated to 100 °C and stirred for 16 h. Concentration and flash chromatography (1→5% EtOAc in hexane) afforded the bisthiazole **5a** as a white solid (30 mg, 72 %). ¹H NMR (400 MHz, CDCl₃) δ 7.84 (s, 1H), 7.04 (s, 1H), 3.33 (sept, 1H, *J* = 6.8 Hz), 1.42 (d, 6H, *J* = 6.8 Hz); ¹³C NMR (67.5 MHz, CDCl₃) 179.2, 161.3, 150.7, 147.2, 118.7 (q, *J* = 319 Hz), 116.4, 106.3, 33.3, 23.0; IR (neat) ν_{max}: 3115, 2922, 1429, 1220, 1136 cm⁻¹; CIHRMS [*M*]⁺ calculated for C₁₀H₉F₃N₂O₃S₃ 357.9727, found 357.9748.

Cystothiazole A 1: To a solution of vinyl stannane **4** (15 mg, 0.030 mmol) and bisthiazolyl triflate **5a** (11 mg, 0.031 mmol) in a sealed tube was added LiCl (5 mg, 0.12 mmol), Pd(PPh₃)₄ (3.5 mg, 0.0030 mmol) and dioxane (1.5 mL, freeze pump thawed for two cycles). The reaction mixture was heated to 100 °C and stirred for 16 h. Concentration and flash chromatography (5→10% EtOAc in hexane) afforded cystothiazole A **1** as a white solid (11 mg, 85 %). ¹H NMR (400 MHz, CDCl₃) δ 7.85 (s, 1H), 7.07 (s, 1H), 6.55 (d, 1H, *J* = 15.6 Hz), 6.39 (dd, 1H, *J* = 15.6, 7.6 Hz), 4.94 (s, 1H), 4.15 (dq, 1H, *J* = 8.0, 6.8 Hz), 3.79 (dd, 1H, *J* = 8.0, 7.2 Hz), 3.64 (s, 3H), 3.58 (s, 3H), 3.35 (sept, 1H, *J* = 6.8 Hz), 3.31 (s, 3H), 1.42 (d, 6H, *J* = 6.8 Hz), 1.19 (d, 3H, *J* = 6.8 Hz); ¹³C NMR (100 MHz, CDCl₃) 178.6, 176.7, 167.7, 162.5, 154.4, 148.6, 131.5, 125.5, 115.0, 114.8, 91.0, 84.3, 57.0, 55.5, 50.8, 39.8, 33.3, 23.1, 14.0; IR (neat) ν_{max}: cm⁻¹ 3110, 2926, 1711, 1624, 1146, 1094; [α]_D²³ = +104.0° (c = 0.55, CHCl₃); CIHRMS [*M*]⁺ calculated for C₂₀H₂₆N₂O₄S₂ 422.1334, found 422.1328.

Thiazole Alcohol 8c: A solution of the dibromothiazole **18** (320 mg, 1.32 mmol) in diethyl ether (6.6 mL) was cooled to -78 °C. ^tBuLi (1.62 mL, 1.7 M in hexane, 2.75 mmol) was then added dropwise, and the solution was allowed to stir for 30 min at -78 °C. Anhydrous acetone (0.194 mL, 2.64 mmol) was then added, and the solution was stirred for an additional 1 hour at -78 °C before saturated NaHCO₃ (4 mL) was added. The mixture was extracted with EtOAc (3 x 10 mL) and the combined organic layers were dried over MgSO₄. Removal of solvent under reduced pressure and purification by flash chromatography (silica gel, 10% EtOAc in hexane) afforded thiazole alcohol **8c** (189 mg, 64%) as a pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 7.13 (s, 1H), 2.54 (s, 1H), 1.66 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 180.6, 124.2, 116.8, 73.5, 30.5; IR (neat) ν_{max}: 3397, 2979, 1482, 1366, 1253, 1080 cm⁻¹; CIHRMS [M]⁺ calculated for C₆H₈BrNOS 220.9510, found 220.9500.

TBS Protected Thiazole Alcohol 8b: To a solution of Alcohol **8c** (287 mg, 1.29 mmol) in CH₂Cl₂ (12.9 mL) at 0 °C was added 2,6-lutidine (0.454 mL, 3.88 mmol), followed by dropwise addition of TBSOTf (0.362 mL, 2.07 mmol). The reaction mixture was warmed to room temperature and stirred 12 h before saturated NH₄Cl (10 mL) was added. The mixture was extracted with Et₂O (3 x 20 mL) and the combined organic layers were dried over MgSO₄. Removal of solvent under reduced pressure and purification by flash chromatography (silica gel, 2 % EtOAc in hexane) afforded the TBS protected alcohol **8b** as pale yellow oil (424 mg, 98%). ¹H NMR (400 MHz, CDCl₃) δ 7.07 (s, 1H), 1.64 (s, 6H), 0.93 (s, 9H), 0.15 (s, 6H). ¹³C NMR (100 MHz, CDCl₃) δ 182.7, 123.7, 116.6, 76.6, 31.1, 25.7, 18.1, -2.3; IR (neat) ν_{max}: 2956, 2931, 2858, 1254, 1187, 1038, 834 cm⁻¹; CIHRMS [M]⁺ calculated for C₁₂H₂₂BrNOSSi 335.0375, found 335.0402.

Bisthiazole 5b: To a solution of 4-bromo bisthiazole **8b** (150 mg, 0.446 mmol) in diethyl ether (4.5 mL) at -78 °C, was added ^tBuLi (0.551 mL of a 1.7 M solution in hexane, 0.937 mmol). After the solution was stirred at -78 °C for 15 min, tributyltin chloride (0.121 mL, 0.446 mmol)

was added. The reaction mixture was stirred at room temperature for 1.5 h, and water (5 mL) was added. The mixture was extracted with diethyl ether (3 x 10 mL). The combined organic layers were dried over MgSO₄. Removal of solvent under reduced pressure afforded the crude thiazolyl stannane **19b** as an orange oil (253 mg, 100%). To a solution of crude thiazolyl stannane **19b** (52 mg, 0.095 mmol) and ditriflate **7** (33 mg, 0.095 mmol) in a sealed tube was added LiCl (12 mg, 0.28 mmol), Pd(PPh₃)₄ (11 mg, 0.0095 mmol) and dioxane (1.0 mL, freeze pump thawed for two cycles). The reaction mixture was heated to 100 °C and stirred for 16 h. Concentration and flash chromatography (1%→5% EtOAc in hexane) afforded the bisthiazole **5b** as a colorless oil (32 mg, 68 %). ¹H NMR (400 MHz, CDCl₃) δ 7.84 (s, 1H), 7.03 (s, 1H), 1.68 (s, 6H), 0.95 (s, 9H), 0.16 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 183.0, 161.6, 150.6, 147.4, 118.6 (q, *J* = 320 Hz), 117.3, 106.1, 76.5, 31.2, 25.7, 18.1, -2.3; IR (neat) ν_{max}: 2957, 2931, 2859, 1431, 1243, 1214, 1139, 1041 cm⁻¹; CIHRMS [M]⁺ calculated for C₁₆H₂₃F₃N₂O₄S₃Si 488.0541, found 488.0538.

Protected Cystothiazole B 20: To a solution of vinyl stannane **4** (9 mg, 0.018 mmol) and bisthiazole **5b** (9 mg, 0.018 mmol) in a sealed tube was added LiCl (2.5 mg, 0.051 mmol), Pd(PPh₃)₄ (2 mg, 0.0017 mmol) and dioxane (1 mL, freeze pump thawed for two cycles). The reaction mixture was heated to 100 °C and stirred for 16 h. Concentration and flash chromatography (5%→10% EtOAc in hexane) afforded a pale yellow oil **20** (7 mg, 72 %). ¹H NMR (400 MHz, CDCl₃) δ 7.82 (s, 1H), 7.06 (s, 1H), 6.55 (d, 1H, *J* = 15.6 Hz), 6.38 (1H, dd, *J* = 16.0, 8.0 Hz), 4.94 (s, 1H), 4.15 (dq, 1H, *J* = 7.6, 6.8 Hz), 3.79 (dd, 1H, *J* = 8.0, 7.6 Hz), 3.64 (s, 3H), 3.58 (s, 3H), 3.31 (s, 3H), 1.69 (s, 6H), 1.19 (d, 3H, *J* = 6.8 Hz), 0.94 (s, 9H), 0.15 (s, 6H); ¹³C NMR (67.5 MHz, CDCl₃) δ 182.1, 176.7, 167.7, 162.8, 154.3, 148.9, 131.5, 125.6, 115.9, 114.9, 91.0, 84.4, 76.5, 57.0, 55.5, 50.8, 39.8, 31.3, 25.8, 18.2, 14.1, -2.2; IR (neat) ν_{max}: 2930, 2856, 1713, 1624, 1145 cm⁻¹; [α]_D²³ = +65.4° (c = 0.50, CHCl₃); CIHRMS [M+1]⁺ calculated for C₂₆H₄₁N₂O₅S₂Si 553.2227, found 553.2294.

Cystothiazole B 2: TBAF (0.049 mL, 0.049 mmol) was added to a solution of TBS ether **20** (9 mg, 0.016 mmol) in THF (0.8 mL). The reaction mixture was stirred for 20 min and filtered through a plug of silica gel. Concentration and flash chromatography (10→25% EtOAc in

hexane) afforded cystothiazole B (7.0 mg, 98%) as a white power **2**. ^1H NMR (400 MHz, CDCl_3) δ 7.91 (s, 1H), 7.08 (s, 1H), 6.56 (d, 1H, $J = 15.6$ Hz), 6.40 (dd, 1H, $J = 15.6, 8.0$ Hz), 4.94 (s, 1H), 4.15 (dq, 1H, $J = 7.6, 6.8$ Hz), 3.80 (dd, 1H, $J = 8.0, 8.0$ Hz), 3.64 (s, 3H), 3.58 (s, 3H), 3.31 (s, 3H), 2.84 (s, 1H), 1.70 (s, 6H), 1.19 (d, 3H, $J = 7.2$ Hz); ^{13}C NMR (100 MHz, CDCl_3) δ 179.5, 176.7, 167.7, 162.3, 154.3, 148.7, 131.8, 125.3, 116.1, 115.2, 91.0, 84.3, 73.3, 57.0, 55.5, 50.8, 39.8, 30.9, 14.0; IR (neat) ν_{max} : 3404, 3106, 2927, 1710, 1622, 1148; $[\alpha]_{\text{D}}^{23} = +110.4^\circ$ ($c = 0.10$, CHCl_3); CIHRMS $[\text{M}]^+$ calculated for $\text{C}_{20}\text{H}_{26}\text{N}_2\text{O}_5\text{S}_2$ 438.1283, found 438.1251.

**¹H NMR (400 MHz, CDCl₃) of
Natural Cystothiazole B**

**¹H NMR (400 MHz, CDCl₃) of
Synthetic Cystothiazole B**

**¹³C NMR (100 MHz, CDCl₃) of
Synthetic Cystothiazole A**

**¹³C NMR (100 MHz, CDCl₃) of
Synthetic Cystothiazole B**