Total Synthesis of Jerangolid D

Jiří Pospíšil and István E. Markó*

*Université catholique de Louvain, Département de Chimie, Bâtiment Lavoisier,
Place Louis Pasteur 1, B-1348 Louvain-la-Neuve, Belgium*

marko@chim.ucl.ac.be

Table of contents

Copy of ^1H and ^{13}C NMR spectra of 10	2
Copy of ^1H and ^{13}C NMR spectra of 11	4
Copy of ^1H and ^{13}C NMR spectra of 12	6
Copy of ^1H and ^{13}C NMR spectra of 6	8
Copy of ^1H and ^{13}C NMR spectra of 18	10
Copy of ^1H and ^{13}C NMR spectra of 19	12
Copy of ^1H and ^{13}C NMR spectra of 19A	14
Copy of ^1H and ^{13}C NMR spectra of 8	16
Copy of ^1H and ^{13}C NMR spectra of 23	18
Copy of ^1H and ^{13}C NMR spectra of 25	20
Copy of ^1H and ^{13}C NMR spectra of synthetic jerangolid D (2).....	22
Copy of ^1H and ^{13}C NMR spectra of natural jerangolid D (2).....	24
Copy of ^1H NMR spectra of commixed natural and synthetic jerangolid D (2).....	27

Copy of ^1H and ^{13}C NMR spectra of 10

Chemical Formula: C₁₁H₁₃NO₂
Molecular Weight: 191.23

Copy of ^1H and ^{13}C NMR spectra of 11

Copy of ^1H and ^{13}C NMR spectra of 12

Copy of ^1H and ^{13}C NMR spectra of 6

JUP-AA-01, CDCl₃, 300 MHz, after the CC

Copy of ^1H and ^{13}C NMR spectra of 19

JUP-AB-07, CDCl₃, 300 MHz, after the CC

Copy of ^1H and ^{13}C NMR spectra of 8

JP-M-7, CDCl₃, 300 MHz, after the CC

Chemical Formula: $C_{20}H_{38}O_2Si$
Molecular Weight: 338.60

Chemical Formula: $C_{20}H_{38}O_2Si$
 Molecular Weight: 338.60

JP-M-5, CDCl₃, 500 MHz, after the CC

Copy of ^1H and ^{13}C NMR spectra of synthetic jerangolid D (2)

Copy of ^1H and ^{13}C NMR spectra of natural jerangolid D (2)

Copy of ^1H NMR spectra of commixed natural and synthetic jerangolid D (2)