Supporting Information

Experimental procedures, spectroscopic data, and ^1H and ^{13}C NMR spectra

Total Synthesis of Lycopallhine A

Yuji Ochi, Satoshi Yokoshima, Tohru Fukuyama

Graduate School of Pharmaceutical Sciences, Nagoya University

Furo-cho, Chikusa-ku, Nagoya, 464-8601, Japan

Table of Contents

General Remarks	page S2
Experimental Procedure for Compounds	page S3
Spectra for Compounds.....	page S38

General Remarks: Nuclear magnetic resonance (^1H NMR (400 MHz), ^{13}C NMR (100 MHz)) spectra were determined on a JEOL-ECS400 instrument unless otherwise noted. Chemical shifts for ^1H NMR are reported in parts per million (ppm) downfield from tetramethylsilane (δ) or relative to the singlet for residual chloroform at 7.26 ppm or pyridine at 7.19 ppm as the internal standard. Coupling constants are reported in hertz (Hz). The following abbreviations are used for spin multiplicity: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad. Chemical shifts for ^{13}C NMR were reported in ppm relative to the center line of a triplet at 77.0 ppm for deuteriochloroform or a triplet at 123.4 ppm for pyridine- d_5 . Infrared (IR) spectra were recorded on a JASCO FT/IR-4100 Fourier Transform Infrared Spectrophotometer and were reported in wavenumbers (cm^{-1}). High resolution mass spectra (HRMS) were obtained on a JEOL JMS-T100LP AccuTOF LC-plus either in positive electrospray ionization (ESI) method or in positive direct analysis in real time (DART) ionization method, using PEG as the internal standard. Melting points (mp) were determined on a Yanaco Micro Melting Point Apparatus. Analytical thin layer chromatography (TLC) was performed on Merck precoated analytical plates, 0.25 mm thick, silica gel 60 F₂₅₄. Preparative TLC separations were performed on Merck analytical plates (0.25 or 0.50 mm thick) precoated with silica gel 60 F₂₅₄. Flash chromatography separations were performed on KANTO CHEMICAL Silica Gel 60 (spherical, 40-100 mesh) unless otherwise noted. Reagents were commercial grades and were used without any purification. Dehydrated tetrahydrofuran, diethyl ether, toluene, and dichloromethane were purchased from Kanto Chemicals Co., Inc., and were purified using a Glass Contour Solvent System. Dehydrated benzene and *N,N*-dimethylformamide were purchased from Kanto Chemicals Co., Inc. and stored over activated MS4A*. Dehydrated methanol, ethanol and acetonitrile were also purchased from Kanto Chemicals Co., Inc. and stored over activated MS3A*. All reactions sensitive to oxygen or moisture were conducted under an argon atmosphere.

* Molecular sieves were "activated" in the following manner: A round-bottom flask containing molecular sieves was heated in a regular microwave for 1.5-2.0 minute and the flask was immediately evacuated. When cooled to room temperature, the flask was backfilled with argon. The above procedure was repeated three times.

(1*R*,3*S*,5*S*)-1-Acetoxy-3-(*p*-toluenesulfonyloxy)-5-(*tert*-butyldimethylsilyloxy)-cyclohexane (39)

To a solution of **9** (29.7 g, 102 mmol), triethylamine (35.4 ml, 254 mmol) and trimethylammonium hydrochloride (9.7 g, 102 mmol) in CH₂Cl₂ (300 mL) was added TsCl (29.0 g, 152 mmol) at 0 °C. The mixture was stirred for 1 h. To decompose the remaining TsCl, *N,N*-dimethylethylenediamine (ca. 18.0 ml) was added to the reaction mixture. After stirring for another 10 min, water was added to the mixture. The resulting solution was extracted three times with CH₂Cl₂. The combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 10:1) to give **39** (44.9 g, 99%) as a colorless oil. $[\alpha]_D^{24} -1.88^\circ$ (*c* = 1.00, CHCl₃); IR (film) 2954, 2930, 2857, 1738, 1598, 1471, 1365, 1241, 1178, 1099, 1036, 935, 861, 813, 778 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.78 (d, *J*=8.2 Hz, 2H), 7.34 (d, *J*=8.2 Hz, 2H), 4.61 (dddd, *J*=11.5, 11.5, 4.6, 4.6 Hz, 1H), 4.36 (dddd, *J*=11.5, 11.5, 4.6, 4.6 Hz, 1H), 3.53 (dddd, *J*=11.5, 11.5, 4.6, 4.6 Hz, 1H), 2.43 (s, 3H), 2.18 (ddd, *J*=11.5, 4.6, 4.6 Hz, 1H), 2.15-2.03 (m, 2H), 2.00 (s, 3H), 1.48 (ddd, *J*=11.5, 11.5, 11.5 Hz, 1H), 1.43 (ddd, *J*=11.5, 11.5, 11.5 Hz, 1H), 1.29 (ddd, *J*=11.5, 11.5, 11.5 Hz, 1H), 0.82 (s, 9H), -0.01 (s, 3H), -0.02 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 170.0 (C), 144.8 (C), 134.1 (C), 129.9 (CH), 127.6 (CH), 75.2 (CH), 66.6 (CH), 65.3 (CH), 41.4 (CH₂), 40.2 (CH₂), 37.1 (CH₂), 25.6 (CH₃), 21.6 (CH₃), 21.0 (CH₃), 17.9 (C), -4.9 (CH₃); HRMS (ESI+) 465.1744 (calcd for C₂₁H₃₄NaO₆SSi 465.1743).

(1*R*,3*S*,5*S*)-1-Hydroxy-3-(*p*-toluenesulfonyloxy)-5-(*tert*-butyldimethylsilyloxy)-cyclohexane (10)

To a solution of **39** (44.9 g, 101 mmol) in MeOH (1.00 L) was added potassium carbonate (21.4 g, 152 mmol) at room temperature and stirring was continued at room temperature for 2 h. The solution was evaporated and the residue was partitioned between EtOAc and water. The aqueous phase was further extracted with EtOAc and the combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 5:1~2:1) to afford **10** (39.6 g, 98%, >99.5% ee) as a colorless oil. The enantiomeric excess was determined by HPLC analysis with a chiral HPLC column (DAICEL CHIRALCEL AD-H, 4% 2-propanol in *n*-hexane, 1.0 mL/min at 25 °C, 210 nm). The retention times corresponding to **10** and its enantiomer are 30.9 and 27.1 min, respectively. $[\alpha]_D^{25} -1.32^\circ$ ($c = 1.00$, CHCl₃); IR (film) 2953, 2930, 2857, 1598, 1470, 1360, 1256, 1176, 1099, 1049, 956, 931, 873, 836, 777 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.82 (d, $J=8.2$ Hz, 2H), 7.33 (d, $J=8.2$ Hz, 2H), 4.42 (dddd, $J=10.5, 10.5, 4.6, 4.6$ Hz, 1H), 3.57 (dddd, $J=10.5, 10.5, 4.6, 4.6$ Hz, 1H), 3.57 (dddd, $J=10.5, 10.5, 4.6, 4.6$ Hz, 1H), 2.44 (s, 3H), 2.18-2.08 (m, 1H), 2.08-1.94 (m, 2H), 1.91 (d, $J=5.0$ Hz, 1H), 1.50 (ddd, $J=12.0, 10.5, 10.5$ Hz, 1H), 1.48 (ddd, $J=12.2, 10.5, 10.5$ Hz, 1H), 1.33 (ddd, $J=12.0, 10.5, 10.5$ Hz, 1H), 0.83 (s, 9H), 0.00 (s, 3H), -0.01 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 144.7 (C), 134.3 (C), 129.8 (CH), 127.6 (CH), 75.8 (CH), 65.7 (CH), 65.1 (CH), 43.4 (CH₂), 40.8 (CH₂), 40.5 (CH₂), 25.7 (CH₃), 21.6 (CH₃), 18.0(C), -4.9(CH₃); HRMS (ESI+) 423.1622 (calcd for C₁₉H₃₂NaO₅SSi 423.1637).

(5*S*)-5-*tert*-Butyldimethylsilyloxycyclohex-2-enone (11)

To a solution of **10** (56.0 g, 140 mmol) and PhI(OAc)₂ (98.0 g, 304 mmol) in acetonitrile (580 mL) and phosphate buffer (pH7.4, 580 mL) was added AZADO (1.58 g, 10.4 mmol) at 0 °C. The reaction mixture was allowed to warm to room temperature. After stirring for 2 h at room temperature, the reaction was quenched with aqueous Na₂S₂O₃. The resulting solution was extracted three times with EtOAc. The combined organic

phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residue was dissolved in CH₂Cl₂ (1.00 L) and DBU (25.5 mL, 171 mmol) was added at 0 °C. After stirring for 1.5 h at 0 °C, aqueous NH₄Cl was added. The resulting mixture was extracted three times with CH₂Cl₂. The combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:0~10:1) to give **11** (28.5 g, 91%) as a pale yellow oil. $[\alpha]_D^{25}$ 10.4° (*c* = 1.00, CHCl₃); IR (film) 2955, 2930, 2894, 2857, 1682, 1472, 1389, 1253, 1103, 939, 836, 777 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.85 (ddd, *J*=10.1, 5.0, 3.2 Hz, 1H), 6.03 (d, *J*=10.1 Hz, 1H), 4.22 (dddd, *J*=9.6, 7.5, 4.7, 4.1 Hz, 1H), 2.64 (dd, *J*=16.0, 4.1 Hz, 1H), 2.58 (ddd, *J*=18.3, 5.0, 4.7 Hz, 1H), 2.46 (dd, *J*=16.0, 9.6 Hz, 1H), 2.36 (dddd, *J*=18.3, 7.5, 3.2, 2.7 Hz 1H), 0.86 (s, 9H), 0.05 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 198.6 (C), 146.8 (CH), 130.0 (CH), 67.5 (CH), 48.0 (CH₂), 35.5 (CH₂), 25.7 (CH₃), 17.9 (C), -4.8 (CH₃), -4.9 (CH₃); HRMS (ESI+) 249.1284 (calcd for C₁₂H₂₂NaO₂Si 249.1287).

(5*S*)-2-Iodo-5-*tert*-butyldimethylsilyloxycyclohex-2-enone (12**)**

To a solution of **11** (16.4 g, 72.4 mmol) and DMAP (17.7 g, 145 mmol) in CH₂Cl₂ (40.0 mL) was added iodine (20.5 g, 80.8 mmol) in CH₂Cl₂ (500 mL) slowly at 0 °C. The reaction mixture was allowed to warm to room temperature. After stirring for 2 h at room temperature, the reaction was quenched with aqueous Na₂S₂O₃. The resulting solution was extracted with CH₂Cl₂. The combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 50:1) to afford **12** (22.5 g, 88%) as a pale yellow oil. $[\alpha]_D^{26}$ 11.0° (*c* = 1.00, CHCl₃); IR (film) 2953, 2929, 2893, 2856, 1690, 1591, 1471, 1322, 1254, 1106, 962, 837, 776 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.61 (dd, *J*=5.3, 3.8 Hz, 1H), 4.27 (dddd, *J*=9.2, 6.9, 5.0, 3.7 Hz, 1H), 2.85 (dd, *J*=15.6, 5.0 Hz, 1H), 2.67 (dd, *J*=15.6, 9.2 Hz, 1H), 2.63 (ddd, *J*=18.3, 5.3, 3.7 Hz, 1H), 2.45 (ddd, *J*=18.3, 6.9, 3.8 Hz, 1H), 0.85 (s, 9H), 0.05 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 191.2 (C), 155.1 (CH), 103.5 (C), 67.2 (CH), 46.7 (CH₂),

39.3 (CH₂), 25.6 (CH₃), 17.9 (C), -4.8 (CH₃), -4.9 (CH₃); HRMS (ESI+) 375.0249 (calcd for C₁₂H₂₁INaO₂Si 375.0253).

(S)-5-((*tert*-Butyldimethylsilyl)oxy)-2-(3-(4-methoxyphenoxy)propyl)cyclohex-2-ene (14)

To a solution of 1-allyloxy-4-methoxybenzene (14.2 g, 86.5 mmol) in THF (120 mL) was added 9-BBN (0.5 M solution in THF, 152 mL, 76.0 mmol) at room temperature and the solution was heated at reflux for 2 h. In another flask **12** (20.6 g, 58.5 mmol) were dissolved in THF (240 mL). To this solution were added the above solution containing the boron reagent, PdCl₂(dppf)·CH₂Cl₂ (4.74 g, 5.80 mmol), potassium phosphate (24.0 g 113 mmol), triphenylarsine (1.81 g, 5.91 mmol) and water (120 mL) at 0 °C. After stirring for 2 h at room temperature, the reaction was quenched with aqueous NH₄Cl. The resulting solution was extracted with EtOAc. The combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified roughly by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 50:1~10:1) to afford 5:1 mixture of **14** and **11** (20.86 g) as a yellow oil, which was used for the next step without further purification. The spectroscopic data for **14** were collected after purification by PTLC (SiO₂; toluene:acetone = 100:1). [α]_D²⁶ 1.98° (*c* = 1.00, CHCl₃); IR (film) 2953, 2929, 2896, 2856, 1676, 1508, 1471, 1380, 1231, 1106, 1041, 866, 835, 777 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.85 (s, 4H), 6.61 (dd, *J*=4.6, 3.2 Hz, 1H), 4.17 (dddd, *J*=9.6, 7.3, 4.6, 4.1 Hz, 1H), 3.88 (t, *J*=6.4 Hz, 2H), 3.75 (s, 3H), 2.66 (dd, *J*=16.0, 4.1 Hz, 1H), 2.57 (ddd, *J*=17.8, 4.6, 4.6 Hz, 1H), 2.47 (dd, *J*=16.0, 9.6 Hz, 1H), 2.42-2.30 (m, 3H), 1.86 (tt, *J*=7.6, 6.4 Hz, 2H), 0.86 (s, 9H), 0.05 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 198.0 (C), 153.7 (C), 153.0 (C), 141.9 (CH), 139.3 (C), 115.4 (CH), 114.6 (CH), 67.8 (CH₂), 67.7 (CH), 55.7 (CH₃), 48.2 (CH₂), 35.9 (CH₂), 28.1 (CH₂), 26.0 (CH₂), 25.8 (CH₃), 17.9 (C), -4.8 (CH₃), -4.9 (CH₃); HRMS (ESI+) 413.2112 (calcd for C₂₂H₃₄NaO₄Si 413.2124).

(1*S*,5*S*)-5-((*tert*-Butyldimethylsilyl)oxy)-2-(3-(4-methoxyphenoxy)propyl)cyclohex-2-enol (15)

To a solution of **14** containing a small amount of **10** in MeOH (500 mL) was added $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (28.3 g, 76.0 mmol) and NaBH_4 (2.43 g, 64.3 mmol) at 0 °C. After stirring for 2 h at 0 °C, the reaction was quenched with aqueous NH_4Cl . The solution was evaporated and the residue was partitioned between EtOAc and water. The aqueous phase was further extracted with EtOAc and the combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified roughly by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 20:1~10:1) to afford 5:1 mixture of **15** and **40** (20.4 g) as a yellow oil, which was used for the next step without further purification. The spectroscopic data for **15** were collected after purification by PTLC (SiO_2 ; toluene:acetone = 100:1). $[\alpha]_D^{26} -3.04^\circ$ ($c = 1.00$, CHCl_3); IR (film) 3445, 2950, 2928, 2857, 1508, 1470, 1322, 1231, 1044, 962, 835, 777 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 6.82 (s, 4H), 5.38 (m, 1H), 4.23 (d, $J=3.2$ Hz, 1H), 3.98 (m, 1H), 3.91 (t, $J=6.4$ Hz, 2H), 3.75 (s, 3H), 3.38 (d, $J=10.1$ Hz, 1H), 2.35-2.26 (m, 2H), 2.23-2.16 (m, 2H), 2.13 (ddd, $J=13.8, 4.1, 3.7$ Hz, 1H), 1.94 (m, 1H), 1.82 (ddd, $J=13.8, 5.0, 1.8$ Hz, 1H), 0.88 (s, 9H), 0.09 (s, 3H), 0.07 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 153.7 (C), 153.2 (C), 139.2 (C), 119.2 (CH), 115.5 (CH), 115.5 (CH), 114.6 (CH), 114.6 (CH), 68.3 (CH_2), 66.7 (CH), 66.7 (CH), 55.7 (CH_2), 37.5 (CH_2), 34.3 (CH_2), 30.7 (CH_2), 27.6 (CH_3), 25.8 (CH_3), 18.0 (C), -4.9 (CH_3), -5.1 (CH_3); HRMS (ESI+) 415.2264 (calcd for $\text{C}_{22}\text{H}_{36}\text{NaO}_4\text{Si}$ 415.2281).

Ethyl 2-((1*S*,5*R*)-5-((*tert*-butyldimethylsilyl)oxy)-2-(3-(4-methoxyphenoxy)propyl)cyclohex-2-en-1-yl)acetate (16)

To a solution of **15** containing a small amount of **40** in *m*-xylene (200 mL) was added triethyl orthoacetate (200 mL) and *o*-nitrophenol (400 mg, 2.88 mmol) at room

temperature and the solution was heated at reflux for 13.5 h. After cooling to room temperature, the solvent was evaporated. The residual oil was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:0~20:1) to afford **16** (16.4 g, 58% for 3 steps) as a pale yellow oil. $[\alpha]_D^{26} -34.8^\circ$ ($c = 1.00$, CHCl₃); IR (film) 2951, 2929, 2856, 1733, 1508, 1471, 1232, 1105, 1039, 835, 775 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.82 (s, 4H), 5.38-5.37 (m, 1H), 4.14 (q, $J=7.4$ Hz, 2H), 3.96-3.80 (m, 2H), 3.76 (s, 3H), 2.80-2.65 (m, 1H), 2.61 (dd, $J=15.1, 4.1$ Hz, 1H), 2.28 (dd, $J=15.1, 9.6$ Hz, 1H), 2.25-2.15 (m, 1H), 2.13 (t, $J=7.6$ Hz, 1H), 2.05-19.2 (m, 2H), 1.90-1.85 (m, 1H), 1.85-1.75 (m, 1H), 1.46 (ddd, $J=12.4, 10.1, 10.1$ Hz, 1H), 1.25 (t, $J=7.4$ Hz, 3H), 0.88 (s, 9H), 0.06 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 172.9 (C), 153.7 (C), 153.1 (C), 137.8 (C), 120.9 (CH), 115.4 (CH), 114.6 (CH), 68.0 (CH₂), 67.9 (CH), 60.2 (CH₂), 55.6 (CH₃), 38.4 (CH₂), 38.2 (CH₂), 35.3 (CH₂), 34.6 (CH), 30.5 (CH₂), 27.5 (CH₂), 25.9 (CH₃), 18.1 (C), 14.2 (CH₃), -4.7 (CH₃), -4.7 (CH₃); HRMS (ESI+) 485.2714 (calcd for C₂₆H₄₂NaO₅Si 485.2699).

2-((1*R*,5*R*)-5-((*tert*-Butyldimethylsilyl)oxy)-2-(3-(4-methoxyphenoxy)propyl)cyclohex-2-en-1-yl)ethanol (41**)**

To a solution of **16** (28.0 g, 60.5 mmol) in THF (900 mL) was added LiAlH₄ (2.53 g, 66.6 mmol) at 0 °C. After stirring for 2 h at 0 °C, H₂O (2.60 mL), 3M aqueous NaOH (2.60 mL) and H₂O (7.80 mL) were successively added at the same temperature, and then the resulting mixture was filtered through a pad of Celite[®]. The filtrate was concentrated *in vacuo*. The residual oil was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 5:1~2:1) to afford **41** (23.1 g, 92%) as a pale yellow oil. $[\alpha]_D^{27} -38.1^\circ$ ($c = 1.00$, CHCl₃); IR (film) 3409, 2950, 2929, 2856, 1508, 1231, 1105, 1041, 835, 775 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.82 (s, 4H), 5.36 (d, $J=5.5$ Hz, 1H), 3.89 (dd, $J=9.2, 6.4$ Hz, 2H), 3.85-3.78 (m, 1H), 3.76 (s, 3H), 3.74-3.60 (m, 2H), 2.50-2.35 (m, 1H), 2.25-2.05 (m, 3H), 2.05-1.85 (m, 4H), 1.85-1.70 (m, 1H), 1.56-1.44 (m, 2H), 1.37 (ddd, $J=12.3, 11.0, 10.6$ Hz, 1H), 0.89 (s, 9H), 0.07 (s, 3H), 0.06 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 153.7 (C), 153.1 (C), 137.8 (C), 120.3 (CH), 115.4 (CH),

Ethyl 4-((1*S*,5*R*)-5-((*tert*-butyldimethylsilyl)oxy)-2-(3-(4-methoxyphenoxy)propyl)cyclohex-2-en-1-yl)-3-oxobutanoate (18**)**

To a solution of **17** (22.2 g, 53.0 mmol) in CH₂Cl₂ (750 mL) was added SnCl₂ (1.12 g, 5.91 mmol) and ethyl diazoacetate (15% solution in toluene, 61.0 mL, 80.2 mmol) at 0 °C. After stirring for 1 h at room temperature, aqueous NH₄Cl was added at 0 °C. The resulting solution was extracted three times with CH₂Cl₂. The combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 10:1~5:1) to give **18** (26.1 g, 98%) as a pale yellow oil. $[\alpha]_D^{26}$ –39.8° (*c* = 1.00, CHCl₃); IR (film) 2951, 2928, 2856, 1724, 1508, 1231, 1106, 1040, 835, 775 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 12.13 (s, 0.13H), 6.81 (s, 4H), 5.42–5.30 (m, 1H), 5.00 (s, 0.14H), 4.19 (q, *J*=7.1 Hz, 0.28H), 4.18 (q, *J*=7.1 Hz, 2.72H), 3.95–3.80 (m, 3H), 3.76 (s, 3H), 3.42 (s, 1.66H), 2.85–2.55 (m, 1.67H), 2.54–2.51 (m, 1.12H), 2.30–1.70 (m, 7H), 1.45–1.30 (m, 1H), 1.26 (t, *J*=7.1 Hz, 3H), 0.88 (s, 9H), 0.05 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 202.1 (C), 177.2 (C), 172.5 (C), 167.0 (C), 153.7 (CH), 153.1 (CH), 138.2 (C), 137.6 (C), 120.9 (CH), 115.4 (CH), 114.6 (CH), 90.7 (CH), 68.0 (CH), 68.0 (CH₂), 67.4 (CH), 61.3 (CH₂), 59.9 (CH₂), 55.7 (CH₃), 50.0 (CH₂), 46.9 (CH₂), 39.1 (CH₂), 38.2 (CH₂), 37.9 (CH₂), 35.4 (CH₂), 35.2 (CH₂), 34.9 (CH), 33.0 (CH), 30.6 (CH₂), 27.5 (CH₂), 25.9 (CH₃), 18.2 (C), 14.2 (CH₂), 14.1 (CH₃), –4.7 (CH₃); HRMS (ESI+) 527.2789 (calcd for C₂₈H₄₄NaO₆Si 527.2804).

Ethyl 4-((1*S*,5*R*)-5-((*tert*-butyldimethylsilyl)oxy)-2-(3-(4-methoxyphenoxy)propyl)cyclohex-2-en-1-yl)-2-diazo-3-oxobutanoate (20**)**

To a solution of **18** (26.0 g, 51.5 mmol) and triethylamine (14.9 mL, 107 mmol) in acetonitrile (93.0 mL) and THF (370 mL) was added 2-azido-1,3-dimethylimidazolium hexafluorophosphate (17.6 g, 61.8 mmol) at 0 °C. After stirring for 1 h at the same temperature, EtOAc and aqueous NH₄Cl were added. The resulting solution was extracted with EtOAc. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 10:1) to give **20** (26.68 g, 98%) as a pale yellow oil. $[\alpha]_D^{26} -27.3^\circ$ ($c = 1.00$, CHCl₃); IR (film) 2952, 2929, 2856, 2132, 1718, 1654, 1508, 1470, 1306, 1231, 1106, 1070, 835, 775 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.81 (s, 4H), 5.46-5.32 (m, 1H), 4.28 (q, $J=7.3$ Hz, 2H), 3.94-3.78 (m, 3H), 3.75 (s, 3H), 3.10 (dd, $J=16.0, 3.2$ Hz, 1H), 2.93 (dd, $J=16.0, 9.6$ Hz, 1H), 2.92-2.80 (m, 1H), 2.20 (ddd, $J=17.0, 5.0, 5.0$ Hz, 1H), 2.12 (t, $J=7.6$ Hz, 2H), 2.22-1.94 (m, 2H), 1.93-1.84 (m, 1H), 1.84-1.72 (m, 1H), 1.42 (ddd, $J=12.4, 10.1, 10.1$ Hz, 1H), 1.32 (t, $J=7.3$ Hz, 3H), 0.87 (s, 9H), 0.05 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 192.0 (C), 161.2 (C), 153.6 (C), 153.1 (C), 138.1 (C), 120.8 (CH), 115.4 (CH), 114.5 (CH), 76.3 (C), 68.1 (CH₂), 67.9 (CH), 61.3 (CH₂), 55.7 (CH₃), 43.6 (CH₂), 38.6 (CH₂), 35.4 (CH₂), 33.8 (CH), 30.5 (CH₂), 27.6 (CH₂), 25.9 (CH₃), 18.1 (C), 14.3 (CH₃), -4.6 (CH₃), -4.7 (CH₃); HRMS (ESI+) 553.2693 (calcd for C₂₈H₄₂N₂NaO₆Si 553.2710).

(4*S*,5*aS*)-Ethyl 4-((*tert*-butyldimethylsilyl)oxy)-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-oxooctahydro-1*H*-cyclopropa[*cd*]indene-2*a*-carboxylate (22**)**

0.04 (s, 3H), 0.02 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 216.1 (C), 169.2 (C), 153.8 (C), 152.9 (C), 115.3 (CH), 114.6 (CH), 68.2 (CH_2), 64.4 (CH), 55.6 (CH_3), 52.4 (C), 48.4 (C), 46.1 (CH_2), 41.5 (CH), 33.6 (CH_2), 29.9 (CH), 29.2 (CH_2), 27.5 (CH_2), 27.3 (CH_2), 25.7 (CH_3), 17.9 (C), -4.9 (CH_3), -5.0 (CH_3); HRMS (ESI+) 519.2164 (calcd for $\text{C}_{26}\text{H}_{37}\text{Na}_2\text{O}_6\text{Si}$ 519.2154).

(4*S*,5*aS*)-4-((*tert*-Butyldimethylsilyl)oxy)-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-oxo-*N*-((4-(trifluoromethyl)phenyl)sulfonyl)octahydro-1*H*-cyclopropa[*cd*]indene-2*a*-carboxamide (43**)**

To a solution of **42** (9.00 g, 19.0 mmol) in THF (130 mL) was added 1,1'-carbonyldiimidazole (6.15 g, 38.0 mmol) at room temperature and the solution was stirred at room temperature for 2 h. In another flask 4-(trifluoromethyl)benzenesulfonamide (6.40 g, 28.5 mmol) and DBU (4.00 mL, 26.6 mmol) were dissolved in THF (260 mL). To this solution was added the above solution containing the acyl imidazole reagent. After stirring for 4 h at the same temperature, EtOAc and aqueous NH_4Cl were added. The resulting solution was extracted with EtOAc. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified roughly by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 4:1) to give a mixture of **43** and 4-(trifluoromethyl)benzenesulfonamide as a yellow oil, which was used for the next step without further purification. The spectroscopic data for **43** were collected after purification by PTLC (SiO_2 ; *n*-hexane:EtOAc = 3:1). $[\alpha]_{\text{D}}^{27} -13.1^\circ$ ($c = 1.00$, CHCl_3); IR (film) 3105, 2952, 2931, 2857, 1710, 1508, 1322, 1231, 1175, 1062, 835, 777 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 11.02 (s, 1H), 8.19 (d, $J=8.3$ Hz, 1H), 7.76 (d, $J=8.3$ Hz, 1H), 6.82 (dd, $J=6.8, 2.8$ Hz, 2H), 6.74 (dd, $J=6.8, 2.8$ Hz, 2H), 4.02 (dddd, $J=6.5, 6.5, 3.2$ Hz, 1H), 3.80-3.70 (m, 5H), 2.73 (dd, $J=18.1, 10.9$ Hz, 1H), 2.68-2.63 (m, 1H), 2.55-2.45 (m, 2H), 2.32 (ddd, $J=15.8, 8.0, 8.0$ Hz, 1H), 2.10 (ddd, $J=14.2, 10.1, 4.6$ Hz, 1H), 2.00-1.81 (m, 2H), 1.80-1.70 (m, 1H), 1.62-1.45 (m, 3H), 0.85 (s, 9H), 0.02 (s, 3H),

0.00 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 212.9 (C), 165.9 (C), 153.8 (C), 152.8 (C), 142.7 (C), 135.5 (C, q, $J=32.4$ Hz), 129.0 (CH), 125.9 (CH, q, $J=3.8$ Hz), 122.8 (CF_3 , q, $J=271.7$ Hz), 115.2 (CH), 114.7 (CH), 67.8 (CH_2), 64.3 (CH), 55.7 (CH_3), 53.5 (C), 49.3 (C), 46.6 (CH_2), 40.7 (CH), 33.6 (CH_2), 29.7 (CH), 29.2 (CH_2), 26.9 (CH_2), 26.8 (CH_2), 25.7 (CH_3), 17.9 (C), -4.9 (CH_3), -5.0 (CH_3); HRMS (ESI+) 726.2147 (calcd for $\text{C}_{33}\text{H}_{41}\text{F}_3\text{NNa}_2\text{O}_7\text{SSi}$ 726.2121).

(4*S*,5*aS*)-4-Hydroxy-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-oxo-*N*-((4-(trifluoromethyl)phenyl)sulfonyl)octahydro-1*H*-cyclopropa[*cd*]indene-2*a*-carboxamide (23**)**

To a solution of a mixture containing **43** and 4-(trifluoromethyl)benzenesulfonamide in THF (250 mL) was added TBAF (190 mL, 190 mmol, 1.0 M in THF) at 0 °C. After stirring for 1 h at room temperature, EtOAc and aqueous NH_4Cl were added. The resulting solution was extracted with EtOAc. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 0:1) to give **23** (9.25 g, 86% for 2 steps) as a pale yellow amorphous solid. $[\alpha]_D^{27} -32.1^\circ$ ($c = 1.00$, CHCl_3); IR (film) 3538, 2938, 1709, 1508, 1433, 1356, 1322, 1230, 1133, 1062, 826 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 10.98 (s, 1H), 8.18 (d, $J=8.2$ Hz, 2H), 7.76 (d, $J=8.2$ Hz, 2H), 6.84-6.80 (m, 2H), 6.78-6.70 (m, 2H), 4.15-4.05 (m, 1H), 3.80-3.70 (m, 4H), 2.81 (dd, $J=19.3, 11.7$ Hz, 1H), 2.74-2.62 (m, 1H), 2.54 (dd, $J=8.3, 4.6$ Hz, 1H), 2.44-2.34 (m, 2H), 2.10 (ddd, $J=14.2, 9.6, 4.6$ Hz, 1H), 2.04-1.90 (m, 2H), 1.88-1.69 (m, 2H), 1.65-1.50 (m, 2H), 1.44 (ddd, $J=15.6, 7.8, 4.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 212.9 (C), 165.7 (C), 153.8 (C), 152.7 (C), 142.3 (C), 135.0 (C, q, $J=32.4$ Hz), 125.9 (CH, q, $J=3.8$ Hz), 123.0 (CF_3 , q, $J=271.7$ Hz), 115.2 (CH), 114.7 (CH), 67.8 (CH_2), 64.1 (CH), 55.7 (CH_3), 53.1 (C), 49.3 (C), 47.0 (CH_2), 40.3 (CH), 33.0 (CH_2), 29.8 (CH), 28.5 (CH_2), 26.8 (CH_2); HRMS (ESI+) 612.1231 (calcd for $\text{C}_{27}\text{H}_{27}\text{F}_3\text{NNa}_2\text{O}_7\text{S}$ 612.1258).

(2a*S*,2a¹*S*,4a*S*,7a*S*)-2a¹-(3-(4-Methoxyphenoxy)propyl)-1-((4-(trifluoromethyl)phenyl)sulfonyl)tetrahydro-1*H*-cyclopenta[*cd*]indole-2,3,6(2a*H*,2a¹*H*,4*H*)-trione (25**)**

To a solution of **23** (2.20 g, 3.88 mmol) and pyridine (3.80 ml) in CH_2Cl_2 (130 mL) was added Dess-Martine periodinane (1.73 g, 4.07 mmol) at 0 °C and the solution was stirred for 1h at room temperature. After checking the reaction by TLC, the reaction mixture was stirred at 40 °C for 30 min. After cooling to 0 °C, EtOAc and aqueous NH_4Cl were added. The resulting solution was extracted with EtOAc. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 1:1) to give **25** (1.75 g, 80%) as a pale orange amorphous solid. $[\alpha]_D^{27}$ 32.9° ($c = 1.00$, CHCl_3); IR (film) 2934, 1768, 1720, 1508, 1405, 1370, 1322, 1231, 1174, 1133, 1062, 827 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.13 (d, $J=8.2$ Hz, 2H), 7.77 (d, $J=8.2$ Hz, 2H), 6.87-6.75 (m, 4H), 4.65 (dd, $J=6.0$, 3.6 Hz, 1H), 3.95 (t, $J=5.5$ Hz, 2H), 3.77 (s, 3H), 3.35 (s, 1H), 3.16 (dd, $J=18.3$, 6.0 Hz, 1H), 2.88 (dd, $J=18.3$, 3.6 Hz, 1H), 2.80 (dd, $J=19.0$, 9.2 Hz, 1H), 2.64-2.54 (m, 1H), 2.42 (dd, $J=18.3$, 4.8 Hz, 1H), 2.15 (dd, $J=19.0$, 3.0 Hz, 1H), 2.10-1.95 (m, 2H), 1.89 (dd, $J=18.3$, 11.0 Hz, 1H), 1.95-1.55 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 205.6 (C), 205.0 (C), 166.1 (C), 154.1 (C), 152.5 (C), 141.1 (C), 136.0 (C, q, $J=32.4$ Hz), 129.1 (CH), 126.3 (CH, q, $J=3.8$ Hz), 122.9 (CF_3 , q, $J=271.7$ Hz), 115.4 (CH), 114.7 (CH), 67.6 (CH_2), 60.1 (CH), 59.4 (CH), 55.7 (CH_3), 46.6 (C), 44.1 (CH_2), 41.6 (CH_2), 41.0 (CH_2), 36.4 (CH), 34.4 (CH_2), 24.6 (CH_2); HRMS (ESI⁺) 588.1253 (calcd for $\text{C}_{27}\text{H}_{26}\text{F}_3\text{NNaO}_7\text{S}$ 588.1280).

(2*R*,2*aS*,2*a*¹*S*,3*R*,4*aS*,6*S*,7*aS*)-2*a*¹-(3-(4-methoxyphenoxy)propyl)-1-((4-(trifluoromethyl)phenyl)sulfonyl)decahydro-1*H*-cyclopenta[*cd*]indole-2,3,6-triol (44)

To a solution of **25** (2.20 g, 4.01 mmol) in THF (100 mL) was added DIBAL (37 mL, 37 mmol, 1 M in *n*-hexane) at -78°C . After stirring for 10 min at the same temperature, the mixture was stirred for 1 h at -10°C . aqueous NaOH (1 M, 2.00 mL) was added at -40°C . The resulting solution was extracted with EtOAc. The organic phase was washed with brine, dried over sodium sulfate and filtered through a pad of Celite[®]. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:3~0:1) to afford an inseparable mixture of diastereomer (3:1) (2.20 g, 88%) as a white amorphous solid. $[\alpha]_D^{27} 20.5^\circ$ ($c = 1.00$, CHCl₃); IR (film) 3389, 2945, 1508, 1469, 1403, 1355, 1323, 1231, 1164, 1132, 1062, 826 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.13 (d, $J=8.3$ Hz, (3/4)2H), 8.03 (d, $J=8.3$ Hz, (1/4)2H), 7.83-7.73 (m, 2H), 6.87-6.71 (m, 4H), 6.10-5.96 (m, (3/4)1H), 5.85 (d, $J=5.5$ Hz, (1/4)1H), 5.76-5.65 (m, (1/4)1H), 4.90-4.76 (m, (3/4)1H), 4.32-4.15 (m, 1H), 4.15-4.03 (m, (3/4)1H), 4.02-3.89 (m, (1/4)1H+(1/4)1H), 3.89-3.80 (m, (3/4)2H), 3.77 (s, (1/4)3H), 3.74 (s, (3/4)3H), 3.72-3.61 (m, (1/4)2H+(1/4)1H), 3.51 (br s, (3/4)1H), 3.48-3.42 (m, (3/4)1H), 3.33-3.21 (br s, (1/4)1H), 2.61 (d, $J=17.0$ Hz, (3/4)10.69H), 2.42 (d, $J=8.2$ Hz, (3/4)1H), 2.40-2.28 (m, (3/4)3H), 2.27-2.18 (m, (1/4)2H), 2.14 (s, 1H), 2.11-1.73 (m, (3/4)6H+(1/4)4H), 1.73-1.54 (m, (3/4)2H+(1/4)1H), 1.54-1.38 (m, (1/4)2H), 1.38-1.20 (m, (1/4)2H), 1.15-0.99 (m, (1/4)1H); ¹³C NMR (100 MHz, CDCl₃) δ 153.9 (C), 153.8 (C), 152.8 (C), 152.7 (C), 142.8 (C), 141.9 (C), 134.4 (C, q, $J=32.4$ Hz), 134.5 (C, q, $J=32.4$ Hz), 129.0 (CH), 127.7 (CH), 126.3 (CH, q, $J=3.8$ Hz), 126.1 (CH, q, $J=3.8$ Hz), 123.2 (CF₃, q, $J=271.7$ Hz), 123.1 (CF₃, q, $J=271.7$ Hz), 115.4 (CH), 115.3 (CH), 114.6 (CH), 114.6 (CH), 86.6 (CH), 85.3 (CH), 73.2 (CH), 71.9 (CH), 68.6 (CH₂), 68.2 (CH₂), 66.0 (CH), 64.3 (CH), 63.0 (CH), 60.1 (CH), 58.9 (CH), 58.4 (CH), 55.7 (2CH₃), 51.9 (C), 51.4 (C), 44.8 (CH₂), 41.2 (CH₂), 41.0 (CH₂), 37.8 (CH₂), 37.2

(CH), 36.4 (CH), 34.8 (CH₂), 34.5 (CH₂), 30.4 (CH₂), 30.0 (CH₂), 25.5 (CH₂), 25.1 (CH₂); HRMS (ESI⁺) 594.1746 (calcd for C₂₇H₃₂F₃NNaO₇S 594.1745).

(3a*R*,3a¹*S*,4a*S*,4a¹*S*,6*S*,7a*S*,8a*R*)-4a¹-(3-(4-Methoxyphenoxy)propyl)-2,2-dimethyl-4-((4-(trifluoromethyl)phenyl)sulfonyl)decahydro-3a*H*-1,3-dioxo-4-azacyclopenta[*def*]fluoren-6-ol (26)

To a solution of a mixture of **44** and **45** (1.93 g, 3.38 mmol) in acetone (140 mL) was added anhydrous copper (II) sulfate (1.8 g, 11.3 mmol) and PPTS (850 mg, 3.38 mmol) at room temperature and the solution was stirred for 16 h at the same temperature. The solution was evaporated and the residue was partitioned between EtOAc and aqueous NaHCO₃. The aqueous phase was further extracted with EtOAc and the combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:1) to give **26** (1.65 g, 80%) as a white amorphous solid. $[\alpha]_D^{27}$ -43.4° (*c* = 1.00, CHCl₃); IR (film) 3504, 2939, 1508, 1470, 1403, 1355, 1323, 1232, 1166, 1107, 1062, 948, 826 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.13 (d, *J*=8.5 Hz, 2H), 7.72 (d, *J*=8.5 Hz, 2H), 6.85-6.77 (m, 4H), 5.89 (d, *J*=6.4 Hz, 1H), 4.42 (dd, *J*=6.4, 4.4 Hz, 1H), 4.03 (dd, *J*=10.5, 8.2 Hz, 1H), 3.87 (t, *J*=6.0 Hz, 2H), 3.77 (s, 3H), 3.42-3.30 (m, 1H), 2.33 (dd, *J*=6.4, 6.4 Hz, 1H), 2.20 (ddd, *J*=14.0, 7.8, 4.4 Hz, 1H), 2.13-2.07 (m, 1H), 1.94 (d, *J*=14.0 Hz, 1H), 1.91-1.50 (m, 9H), 1.45 (s, 3H), 1.31 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 153.8 (C), 153.0 (C), 145.5 (C), 134.1 (C, q, *J*=32.4 Hz), 128.3 (CH), 125.6 (CH, q, *J*=3.8 Hz), 123.3 (CF₃, q, *J*=271.7 Hz), 115.4 (CH), 114.6 (CH), 97.6 (C), 82.3 (CH), 72.4 (CH), 68.4 (CH₂), 66.9 (CH), 62.8 (CH), 55.7 (C), 55.7 (CH₃), 51.7 (CH), 42.0 (CH₂), 41.7 (CH), 40.8 (CH₂), 38.8 (CH₂), 38.7 (CH₂), 29.2 (CH₃), 25.6 (CH₂), 20.6 (CH₃); HRMS (ESI⁺) 634.2059 (calcd for C₃₀H₃₆F₃NNaO₇S 634.2062).

(3a*R*,3a¹*S*,4a*S*,4a¹*S*,6*S*,7a*R*,8a*R*)-4a¹-(3-(4-Methoxyphenoxy)propyl)-2,2-dimethyl-4-((4-(trifluoromethyl)phenyl)sulfonyl)decahydro-3a*H*-1,3-dioxo-4-azacyclopenta[*de*]fluoren-6-yl methanesulfonate (**46**)

To a solution of **26** (1.54 g, 2.52 mmol) and triethylamine (528 μL , 3.78 mmol) in CH_2Cl_2 (70 mL) was added methanesulfonyl chloride (216 μL , 2.77 mmol) at 0 °C. After stirring for 30 min at the same temperature, aqueous NaHCO_3 was added. The resulting solution was extracted with CH_2Cl_2 . The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 1:1) to afford **46** (1.62 g, 93%) as a white amorphous solid. $[\alpha]_D^{27} -37.2^\circ$ ($c = 1.00$, CHCl_3); IR (film) 3449, 2940, 1508, 1350, 1232, 1172, 1062, 943, 827 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.12 (d, $J=8.2$ Hz, 2H), 7.74 (d, $J=8.2$ Hz, 2H), 6.86-6.75 (m, 4H), 5.89 (d, $J=6.9$ Hz, 1H), 4.44 (dd, $J=6.0, 4.6$ Hz, 1H), 4.38-4.27 (m, 1H), 4.06 (dd, $J=10.1, 8.2$ Hz, 1H), 3.86 (t, $J=5.9$ Hz, 2H), 3.77 (s, 3H), 2.91 (s, 3H), 2.35 (dd, $J=6.9, 6.0$ Hz, 1H), 2.25-2.06 (m, 5H), 1.95 (d, $J=14.2$ Hz, 1H), 1.97-1.87 (m, 1H), 1.85-1.64 (m, 2H), 1.58-1.52 (m, 2H), 1.45 (s, 3H), 1.32 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 153.9 (C), 152.9 (C), 145.2 (C), 134.1 (C, q, $J=32.4$ Hz), 128.3 (CH), 125.6 (CH, q, $J=3.8$ Hz), 123.3 (CF_3 , q, $J=271.7$ Hz), 115.4 (CH), 114.6 (CH), 97.7 (C), 82.3 (CH), 72.4 (CH), 68.4 (CH_2), 66.9 (CH), 62.8 (CH), 55.7 (C), 55.7 (CH_3), 51.7 (CH), 42.0 (CH_2), 41.7 (CH), 40.8 (CH_2), 38.8 (CH_2), 38.7 (CH_2), 29.2 (CH_3), 25.5 (CH_2), 20.5 (CH_3); HRMS (ESI+) 712.1848 (calcd for $\text{C}_{31}\text{H}_{38}\text{F}_3\text{NNaO}_9\text{S}_2$ 712.1838).

(3a*R*,3a¹*S*,4a*S*,4a¹*S*,6*R*,7a*S*,8a*R*)-4a¹-(3-(4-Methoxyphenoxy)propyl)-2,2-dimethyl-4-((4-(trifluoromethyl)phenyl)sulfonyl)decahydro-3a*H*-1,3-dioxo-4-azacyclopenta[*de*]fluorene-6-carbonitrile (**27**)

To a solution of **46** (1.83 g, 2.52 mmol) in DMSO (30 mL) was added 18-crown-6 (4.36 g, 16.5 mmol) and potassium cyanide (4.36 g, 66.7 mmol) at room temperature and the solution was stirred for 5 h at 80 °C. After cooling to room temperature, EtOAc and aqueous NaHCO₃ were added. The resulting solution was extracted with EtOAc. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residual oil was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 3:1~2:1) to afford **27** (1.69 g, 97%) as a white amorphous solid. $[\alpha]_{\text{D}}^{27} -79.4^{\circ}$ ($c = 1.00$, CHCl₃); IR (film) 3483, 2990, 2939, 1508, 1468, 1403, 1383, 1350, 1323, 1232, 1172, 1132, 1062, 943, 828 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.13 (d, $J=8.3$ Hz, 2H), 7.73 (d, $J=8.3$ Hz, 2H), 6.85-6.76 (m, 4H), 5.89 (d, $J=6.4$ Hz, 1H), 4.45 (dd, $J=6.4$, 5.0 Hz, 1H), 4.10 (dd, $J=10.1$, 8.2 Hz, 1H), 3.90-3.83 (m, 2H), 3.77 (s, 3H), 2.82-2.75 (m, 1H), 2.43 (dd, $J=6.4$, 6.4 Hz, 1H), 2.31-2.20 (m, 2H), 2.09 (ddd, $J=13.3$, 11.5, 3.2 Hz, 1H), 2.00-1.65 (m, 8H), 1.46 (s, 3H), 1.29 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 153.8 (C), 153.0 (C), 145.3 (C), 134.3 (C, q, $J=32.4$ Hz), 128.3 (CH), 125.8 (CH, q, $J=3.8$ Hz), 123.2 (CF₃, q, $J=271.7$ Hz), 120.8 (C), 115.5 (CH), 114.6 (CH), 97.8 (C), 82.5 (CH), 72.6 (CH), 68.4 (CH₂), 60.0 (CH), 55.7 (CH₃), 55.6 (C), 55.4 (CH), 41.8 (CH₂), 41.3 (CH₂), 40.2 (CH), 31.5 (CH₂), 30.9 (CH₂), 29.3 (CH₃), 24.8 (CH₂), 24.0 (CH), 20.5 (CH₃); HRMS (ESI+) 643.2058 (calcd for C₃₁H₃₅F₃N₂NaO₆S 643.2066).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-2-Allyl-3-hydroxy-2*a*¹-(3-(4-methoxyphenoxy)propyl)-1-((4-(trifluoromethyl)phenyl)sulfonyl)decahydro-1*H*-cyclopenta[*cd*]indole-6-carbonitrile (28**)**

To a solution of **27** (1.21 g, 4.01 mmol) in CH_2Cl_2 (60 mL) was added allyltrimethylsilane (6.30 mL, 39.6 mmol) and $\text{BF}_3\cdot\text{OEt}_2$ (2.90 mL, 22.9 mmol) at -78°C . After stirring for 10 min at -78°C , the mixture was stirred for 30 min at 0°C . The reaction was quenched with aqueous NaHCO_3 . The resulting solution was extracted with CH_2Cl_2 . The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residual oil was purified by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 1:1) to afford **28** (1.13 g, 96%) as a white amorphous solid. $[\alpha]_{\text{D}}^{27} -0.812^\circ$ ($c = 1.00$, CHCl_3); IR (film) 3504, 2947, 2388, 2241, 1508, 1366, 1402, 1323, 1231, 1161, 1134, 1062, 923, 827 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.04 (d, $J=8.2$ Hz, 2H), 7.84 (d, $J=8.2$ Hz, 2H), 6.90-6.76 (m, 4H), 5.63 (dddd, $J=16.9, 11.5, 3.2, 1.8$ Hz, 1H), 5.10 (dd, $J=11.5, 0.9$ Hz, 1H), 5.06 (dd, $J=16.9, 0.9$ Hz, 1H), 4.68 (dd, $J=10.1, 3.2$ Hz, 1H), 4.24-4.10 (m, 1H), 3.96-3.85 (m, 2H), 3.77 (s, 3H), 3.65-3.60 (m, 1H), 2.78 (dd, $J=15.3, 2.1$ Hz, 1H), 2.70-2.63 (m, 1H), 2.52-2.10 (m, 2H), 2.28 (d, $J=7.8$ Hz, 1H), 2.04 (d, $J=4.1$ Hz, 1H), 2.02-2.00 (m, 1H), 1.97-1.86 (m, 1H), 1.82-1.57 (m, 8H); ^{13}C NMR (100 MHz, CDCl_3) δ 154.0 (C), 152.7 (C), 145.2 (C), 134.7 (C, q, $J=33.4$ Hz), 133.7 (CH), 127.6 (CH), 126.7 (CH, q, $J=2.9$ Hz), 123.2 (CF_3 , q, $J=271.7$ Hz), 121.6 (C), 118.7 (CH_2), 115.3 (CH), 114.7 (CH), 71.9 (CH), 68.1 (CH_2), 61.7 (CH), 60.2 (CH), 55.7 (CH_3), 52.1 (CH), 51.7 (C), 38.4 (CH_2), 37.9 (CH_2), 36.4 (CH), 34.2 (CH_2), 28.8 (CH_2), 28.1 (CH_2), 25.8 (CH_2), 19.0 (CH); HRMS (ESI+) 627.2091 (calcd for $\text{C}_{31}\text{H}_{35}\text{F}_3\text{N}_2\text{NaO}_5\text{S}$ 627.2117).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-2-Allyl-3-((*tert*-butyldimethylsilyl)oxy)-2*a*¹-(3-(4-methoxyphenoxy)propyl)-1-((4-(trifluoromethyl)phenyl)sulfonyl)decahydro-1*H*-cyclopenta[*cd*]indole-6-carbonitrile (47**)**

To a solution of **28** (2.00 g, 3.30 mmol) in CH₂Cl₂ (60 mL) were added 2,6-lutidine (6.90 mL, 59.6 mmol) and *tert*-butyldimethylsilyl trifluoromethanesulfonate (1.53 mL, 6.60 mmol) at 0 °C. After stirring for 1 h at the same temperature, aqueous NaHCO₃ was added. The resulting solution was extracted with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residual oil was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 20:1~1:1) to afford **47** (2.20 g, 93%) as a white amorphous solid. [α]_D²⁷ −5.20° (*c* = 1.00, CHCl₃); IR (film) 2952, 2857, 2239, 1508, 1471, 1402, 1323, 1232, 1163, 1135, 1062, 835, 778 cm^{−1}; ¹H NMR (400 MHz, CDCl₃) δ 8.03 (d, *J*=8.2 Hz, 2H), 7.83 (d, *J*=8.2 Hz, 2H), 6.85-6.76 (m, 4H), 5.60 (dddd, *J*=16.5, 10.1, 3.2, 1.8 Hz, 1H), 5.08 (d, *J*=10.1 Hz, 1H), 5.04 (d, *J*=16.5 Hz, 1H), 4.64 (dd, *J*=10.8, 3.0 Hz, 1H), 4.04 (ddd, *J*=17.8, 8.2, 2.8 Hz, 1H), 3.95-3.85 (m, 2H), 3.77 (s, 3H), 3.68-3.63 (m, 1H), 2.82 (dd, *J*=15.4, 2.1 Hz, 1H), 2.74-2.66 (m, 1H), 2.30-2.20 (m, 1H), 2.22 (d, *J*=8.2 Hz, 1H), 2.05-1.92 (m, 2H), 1.90-1.85 (m, 1H), 1.85-1.46 (m, 8H), 0.92 (s, 9H), 0.07 (s, 3H), 0.05 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 154.0 (C), 152.7 (C), 145.4 (C), 134.6 (C, q, *J*=32.4 Hz), 134.0 (CH), 127.6 (CH), 126.5 (CH, q, *J*=2.9 Hz), 124.4 (CH), 123.0 (CF₃, q, *J*=271.7 Hz), 121.6 (C), 118.4 (CH₂), 115.3 (CH), 114.7 (CH), 72.3 (CH), 68.1 (CH₂), 61.7 (CH), 55.7 (CH₃), 52.5 (CH), 50.9 (C), 39.0 (CH₂), 38.2 (CH₂), 36.3 (CH), 34.2 (CH₂), 28.7 (CH₂), 28.1 (CH₂), 25.8 (CH₂), 25.7 (CH₃), 19.0 (CH), 18.0 (C), −4.6 (CH₃), −5.0 (CH₃); HRMS (ESI⁺) 741.3003 (calcd for C₃₇H₄₉F₃N₂NaO₅SSi 741.2981).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-2-Allyl-3-((*tert*-butyldimethylsilyl)oxy)-2*a*¹-(3-(4-methoxyphenoxy)propyl)decahydro-1*H*-cyclopenta[*cd*]indole-6-carbonitrile (48**)**

To a solution of **47** (1.84 g, 2.56 mmol) in MeOH (200 mL) was added Mg (6.00 g, 247 mmol) at room temperature and the solution was heated at 40 °C for 3 h. After cooling to room temperature, EtOAc was added and then the solution was concentrated. The residue was partitioned between EtOAc and water. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 2:1) to afford **48** (1.26 g, 96%) as a pale yellow oil. $[\alpha]_D^{27}$ 5.47° (*c* = 1.00, CHCl₃); IR (film) 3464, 2929, 2856, 2237, 1639, 1508, 1470, 1413, 1359, 1231, 1143, 1044, 914, 835, 776 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.83 (s, 4H), 5.70 (dddd, *J*=17.0, 13.4, 8.2, 1.4 Hz, 1H), 5.07 (dd, *J*=8.2, 1.4 Hz, 1H), 5.02 (dd, *J*=17.0, 1.4 Hz, 1H), 4.05 (ddd, *J*=10.8, 8.0, 2.8 Hz, 1H), 3.91 (t, *J*=6.2 Hz, 2H), 3.77 (s, 3H), 3.66 (dd, *J*=6.9, 6.9 Hz, 1H), 3.36-3.32 (m, 1H), 3.22 (dddd, *J*=12.4, 12.4, 2.8, 2.8 Hz, 1H), 2.14-2.05 (m, 2H), 2.02-1.86 (m, 5H), 1.85-1.68 (m, 4H), 1.65-1.51 (m, 3H), 0.88 (s, 9H), 0.04 (s, 6H), 0.03 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 153.9 (C), 152.9 (C), 136.2 (CH), 123.8 (C), 117.2 (CH₂), 115.4 (CH), 114.7 (CH), 72.9 (CH), 68.8 (CH₂), 59.0 (CH), 58.5 (CH), 55.7 (CH₃), 54.3 (CH), 48.6 (C), 41.0 (CH₂), 39.6 (CH₂), 36.3 (CH), 35.3 (CH₂), 31.7 (CH₂), 29.4 (CH₂), 26.3 (CH₂), 25.8 (CH₂), 18.2 (CH), 18.0 (C), -4.7 (CH₃), -4.9 (CH₃); HRMS (ESI+) 511.3354 (calcd for C₃₀H₄₇N₂O₃Si 511.3356).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-*tert*-Butyl-2-allyl-3-((*tert*-butyldimethylsilyl)oxy)-6-cyano-2*a*¹-(3-(4-methoxyphenoxy)propyl)decahydro-1*H*-cyclopenta[*cd*]indole-1-carboxylate (29**)**

To a solution of **48** (1.74 g, 3.41 mmol) and triethylamine (1.43 mL, 10.2 mmol) in *N,N*-dimethylformamide (20.0 mL) was added *N*-(*tert*-butoxycarbonyloxy)succinimide (1.47 g, 6.82 mmol) at room temperature and the solution was heated at 40 °C for 3 h. After cooling to 0 °C, EtOAc and aqueous NaHCO₃ were added. The resulting solution was extracted with EtOAc. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 5:1) to give **29** (1.91 g, 92%) as a white amorphous solid. $[\alpha]_{\text{D}}^{27}$ 17.9° (*c* = 1.00, CHCl₃); IR (film) 2932, 2857, 2362, 2238, 1695, 1508, 1472, 1394, 1232, 1174, 1130, 1045, 884, 835, 776 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.82 (s, 4H), 5.67 (dddd, *J*=18.4, 13.4, 7.3, 2.3 Hz, 1H), 5.05 (d, *J*=15.6 Hz, 1H), 5.04 (d, *J*=12.4 Hz, 1H), 4.43 (ddd, *J*=8.2, 4.1, 2.3 Hz, 1H), 4.04 (ddd, *J*=13.2, 8.2, 7.3 Hz, 2H), 3.91 (t, *J*=6.0 Hz, 2H), 3.76 (s, 3H), 3.58 (dd, *J*=3.2, 2.7 Hz, 1H), 2.68 (dddd, *J*=12.4, 12.4, 3.0, 3.0 Hz, 1H), 2.36-2.20 (m, 2H), 2.07 (dd, *J*=7.3, 2.3 Hz, 1H), 2.02-1.90 (m, 2H), 1.85-1.70 (m, 3H), 1.69-1.54 (m, 6H), 1.47 (s, 9H), 0.89 (s, 9H), 0.04 (s, 3H), 0.03 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 153.9 (C), 153.5 (C), 152.8 (C), 134.7 (CH), 123.0 (C), 117.2 (CH₂), 115.4 (CH), 114.6 (CH), 79.7 (C), 72.6 (CH), 68.5 (CH₂), 60.5 (CH), 57.0 (CH), 55.7 (CH₃), 53.6 (CH), 50.2 (C), 39.2 (CH₂), 38.6 (CH₂), 36.7 (CH), 34.8 (CH₂), 29.3 (CH₂), 29.0 (CH₂), 28.4 (CH₃), 25.8 (CH₂), 25.8 (CH₃), 19.1 (CH), 18.0 (C), -4.6 (CH₃), -4.9 (CH₃); HRMS (ESI+) 633.3719 (calcd for C₃₅H₅₄N₂NaO₅Si 633.3700).

**(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-*tert*-Butyl-3-((*tert*-butyldimethylsilyl)oxy)-6-cyano-2*a*¹-
 -(3-(4-methoxyphenoxy)propyl)-2-(2-oxoethyl)decahydro-1*H*-cyclopenta[*cd*]indole-
 1-carboxylate (**49**)**

To a solution of **29** (820 mg, 1.34 mmol) in 1,4-dioxane (70.0 mL) and H₂O (23.0 mL) was added 2,6-lutidine (1.37 mL, 11.8 mmol), NaIO₄ (5.73 g, 26.8 mmol) and OsO₄ (33.6 mL, 1.34 mmol, 0.04 M in *t*-BuOH) at 0 °C. The mixture was stirred for 3 h at room temperature. After cooling to 0 °C, CH₂Cl₂ and aqueous NaHCO₃ were added. The resulting solution was extracted with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 2:1) to give **49** (730 mg, 89%) as a white amorphous solid. $[\alpha]_D^{27}$ 30.6° (*c* = 1.00, CHCl₃); IR (film) 2932, 2857, 2238, 1725, 1697, 1508, 1471, 1393, 1366, 1231, 1172, 1127, 1043, 836, 776 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 9.70 (t, *J*=2.5 Hz, 1H), 6.83 (s, 4H), 4.80 (m, 1H), 4.09 (ddd, *J*=10.5, 6.9, 6.9 Hz, 1H), 3.92 (t, *J*=5.7 Hz, 2H), 3.77 (s, 3H), 3.69 (m, 1H), 2.74-2.64 (m, 2H), 2.54 (ddd, *J*=14.6, 5.5, 2.5 Hz, 1H), 2.04-1.93 (m, 3H), 1.88-1.58 (m, 9H), 1.47 (s, 9H), 0.89 (s, 9H), 0.06 (s, 3H), 0.05 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 200.2 (CH), 154.0 (C), 153.1 (C), 152.8 (C), 122.7 (C), 115.5 (CH), 114.7 (CH), 80.8 (C), 72.6 (CH), 68.4 (CH₂), 60.4 (CH), 56.1 (CH), 55.7 (CH₃), 53.3 (CH), 50.2 (C), 49.0 (CH₂), 39.1 (CH₂), 36.9 (CH), 35.5 (CH₂), 29.4 (CH₂), 29.4 (CH₂), 28.4 (CH₃), 25.7 (CH₂), 25.7 (CH₃), 19.2 (CH), 17.9 (C), -4.6 (CH₃), -5.0 (CH₃); HRMS (ESI⁺) 635.3509 (calcd for C₃₄H₅₂N₂NaO₆Si 635.3492).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-*tert*-Butyl-3-((*tert*-butyldimethylsilyl)oxy)-6-cyano-2-(2-hydroxyethyl)-2*a*¹-(3-(4-methoxyphenoxy)propyl)decahydro-1*H*-cyclopenta[*cd*]indole-1-carboxylate (50**)**

To a solution of **49** (685 mg, 1.12 mmol) in MeOH (20.0 mL) was added NaBH₄ (96.0 mg, 2.46 mmol) at 0 °C. The solution was stirred for 30 min at the same temperature. To this solution was added CH₂Cl₂ and aqueous NaHCO₃. The resulting solution was extracted with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:1) to give **50** (510 mg, 74%) as a white amorphous solid. $[\alpha]_D^{26}$ 7.71° (*c* = 1.00, CHCl₃); IR (film) 3481, 2932, 2857, 2238, 1692, 1508, 1472, 1393, 1366, 1313, 1231, 1173, 1128, 1044, 836, 776 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 6.83 (s, 4H), 4.61 (br s, 1H), 4.07 (ddd, *J*=6.4, 6.4, 10.6 Hz, 1H), 3.92 (t, *J*=6.0 Hz, 2H), 3.77 (s, 3H), 3.68-3.45 (m, 4H), 2.70 (dddd, *J*=12.4, 12.4, 2.5, 2.5 Hz, 1H), 2.10-1.91 (m, 3H), 1.89-1.49 (m, 11H), 1.46 (s, 9H), 0.91 (s, 9H), 0.07 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 153.9 (C), 153.6 (C), 152.8 (C), 122.7 (C), 115.4 (CH), 114.7 (CH), 80.9 (C), 72.9 (CH), 68.4 (CH₂), 59.6 (CH), 59.5 (CH), 55.7 (CH), 55.3 (CH₃), 54.6 (C), 50.6 (CH₂), 38.9 (CH₂), 38.0 (CH₂), 36.8 (CH), 34.3 (CH₂), 28.8 (CH₂), 28.7 (CH₂), 28.4 (CH₃), 25.8 (CH₂), 25.8 (CH₃), 19.5 (CH), 18.0 (C), -4.5 (CH₃), -4.8 (CH₃); HRMS (ESI+) 637.3658 (calcd for C₃₄H₅₄N₂NaO₆Si 637.3649).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-*tert*-Butyl 3-((*tert*-butyldimethylsilyl)oxy)-6-cyano-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)decahydro-1*H*-cyclopenta[*cd*]indole-1-carboxylate (30**)**

To a solution of **50** (470 mg, 0.764 mmol) in toluene (11.0 mL) was added *N*-methyl-2-nitrobenzenesulfonamide (181 mg, 0.840 mmol), triphenylphosphine (260 mg, 0.993 mmol) and diethyl azodicarboxylate (451 μL , 0.993 mmol, 2.2 M in toluene) at 0 °C. The solution was stirred for 1 h at the room temperature. To this solution was added CH_2Cl_2 and aqueous NaHCO_3 . The resulting solution was extracted with CH_2Cl_2 . The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 5:1~2:1) to give **30** (600 mg, 97%) as a white amorphous solid. $[\alpha]_D^{26}$ 13.5° (c = 1.00, CHCl_3); IR (film) 2932, 2857, 2238, 1692, 1545, 1508, 1471, 1367, 1231, 1169, 1126, 1042, 835, 776 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.96 (m, 1H), 7.68-7.59 (m, 2H), 7.58 (m, 1H), 6.84 (s, 4H), 4.30 (m, 1H), 4.07 (ddd, J =10.5, 6.4, 6.4 Hz, 1H), 3.94 (t, J =5.7 Hz, 2H), 3.76 (s, 3H), 3.58 (m, 1H), 3.23 (ddd, J =13.1, 13.1, 4.1 Hz, 1H), 3.07 (ddd, J =13.1, 12.8, 5.3 Hz, 1H), 2.89 (s, 3H), 2.68 (dddd, J =13.0, 13.0, 2.8, 2.8 Hz, 1H), 2.03-1.92 (m, 4H), 1.87-1.61 (m, 7H), 1.61-1.51 (m, 3H), 1.46 (s, 9H), 0.85 (s, 9H), 0.04 (s, 3H), 0.02 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 153.9 (C), 153.3 (C), 152.9 (C), 148.1 (C), 133.5 (CH), 132.6 (C), 131.6 (CH), 130.7 (CH), 124.1 (CH), 122.9 (C), 115.5 (CH), 114.7 (CH), 80.1 (C), 72.6 (CH), 68.5 (CH_2), 60.1 (CH), 55.7 (CH_3), 55.3 (CH), 54.9 (CH), 50.5 (C), 47.3 (CH_2), 39.1 (CH_2), 36.9 (CH), 34.6 (CH_3), 34.6 (CH_2), 34.6 (CH_2), 32.9 (CH_2), 29.1 (CH_2), 28.4 (CH_3), 25.8 (CH_3), 25.7 (CH_2), 19.3 (CH), 17.9 (C), -4.5 (CH_3), -4.9 (CH_3); HRMS (ESI+) 835.3735 (calcd for $\text{C}_{41}\text{H}_{60}\text{N}_4\text{NaO}_9\text{SSi}$ 835.3748).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-tert-Butyl 3-((*tert*-butyldimethylsilyl)oxy)-6-formyl-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)decahydro-1*H*-cyclopenta[*cd*]indole-1-carboxylate (51**)**

To a solution of **30** (544 mg, 0.669 mmol) in toluene (14.0 mL) was added DIBAL (1.0 M in toluene, 1.34 mL, 1.34 mmol) at -78°C . The solution was stirred for 1 h at the same temperature. To this solution was added H_2O carefully. The resulting solution was extracted with CH_2Cl_2 . The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 2:1~5:1) to give **51** (510 mg, 93%) as a pale yellow amorphous solid. $[\alpha]_D^{26} 13.9^\circ$ ($c = 0.600$, CHCl_3); IR (film) 2930, 2857, 2390, 1722, 1693, 1546, 1508, 1470, 1366, 1231, 1169, 1126, 1043, 835, 776 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 9.69 (s, 1H), 7.97 (m, 1H), 7.70-7.60 (m, 2H), 7.59 (m, 1H), 6.83 (s, 4H), 4.33 (m, 1H), 4.08 (ddd, $J=10.5, 6.4, 6.4$ Hz, 1H), 3.93 (t, $J=5.7$ Hz, 2H), 3.77 (s, 3H), 3.64 (m, 1H), 3.24 (ddd, $J=13.1, 13.1, 4.1$ Hz, 1H), 3.09 (ddd, $J=13.1, 12.7, 4.6$ Hz, 1H), 2.90 (s, 3H), 2.50 (m, 1H), 2.05-1.70 (m, 7H), 1.70-1.50 (m, 6H), 1.44 (s, 9H), 1.32 (ddd, $J=13.3, 13.3, 2.7$ Hz, 1H), 0.86 (s, 9H), 0.05 (s, 3H), 0.02 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 204.7 (CH), 153.9 (C), 153.3 (C), 153.0 (C), 148.1 (C), 133.4 (CH), 132.7 (C), 131.5 (CH), 130.7 (CH), 124.1 (CH), 115.5 (CH), 114.7 (CH), 79.6 (C), 72.8 (CH), 68.7 (CH_2), 60.9 (CH), 55.7 (CH_3), 55.5 (CH), 55.2 (CH), 51.3 (C), 47.3 (CH_2), 40.2 (CH), 39.5 (CH_2), 37.2 (CH), 34.9 (CH_2), 34.6 (CH_3), 33.7 (CH_2), 28.4 (CH_2), 28.4 (CH_3), 25.8 (CH_3), 25.8 (CH_2), 25.0 (CH_2), 17.9 (C), -4.5 (CH_3), -4.9 (CH_3); HRMS (ESI+) 838.3752 (calcd for $\text{C}_{41}\text{H}_{61}\text{N}_3\text{NaO}_{10}\text{SSi}$ 838.3745).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-*tert*-Butyl 3-((*tert*-butyldimethylsilyl)oxy)-6-(hydroxymethyl)-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)decahydro-1*H*-cyclopenta[*cd*]indole-1-carboxylate (31**)**

To a solution of **51** (1.41 g, 1.73 mmol) in THF (10.0 mL) and MeOH (10.0 mL) was added NaBH₄ (130 mg, 3.46 mmol) at 0 °C. The solution was stirred for 1 h at the same temperature. To this solution were added CH₂Cl₂ and aqueous NH₄Cl. The resulting solution was extracted with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:1) to give **31** (1.37 g, 97%) as a pale yellow amorphous solid. [α]_D²⁷ 20.9° (*c* = 1.70, CHCl₃); IR (film) 3464, 2930, 2857, 1681, 1545, 1508, 1471, 1366, 1231, 1169, 1125, 1042, 971, 898, 835, 776 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.97 (m, 1H), 7.70-7.60 (m, 2H), 7.58 (m, 1H), 6.84 (s, 4H), 4.32 (m, 1H), 4.07 (ddd, *J*=9.6, 7.3, 7.3 Hz, 1H), 3.93 (t, *J*=6.2 Hz, 2H), 3.76 (s, 3H), 3.57 (m, 1H), 3.53-3.40 (m, 2H), 3.22 (ddd, *J*=12.8, 12.8, 4.3 Hz, 1H), 3.18-3.00 (m, 2H), 2.90 (s, 3H), 2.00-1.84 (m, 3H), 1.84-1.51 (m, 10H), 1.44 (s, 9H), 1.27 (ddd, *J*=13.3, 13.3, 6.4 Hz, 1H), 1.08 (ddd, *J*=14.2, 11.9, 2.7 Hz, 1H), 0.86 (s, 9H), 0.04 (s, 3H), 0.01 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 153.9 (C), 153.3 (C), 153.1 (C), 148.1 (C), 133.4 (CH), 132.8 (C), 131.5 (CH), 130.7 (CH), 124.1 (CH), 115.5 (CH), 114.7 (CH), 79.3 (C), 73.0 (CH), 68.9 (CH₂), 68.1 (CH₂), 61.8 (CH), 55.7 (CH₃), 55.4 (CH), 55.4 (CH), 51.2 (C), 47.5 (CH₂), 39.7 (CH₂), 37.9 (CH), 35.1 (CH₂), 34.6 (CH₃), 33.5 (CH₂), 29.4 (CH), 28.8 (2CH₂), 28.5 (CH₃), 26.0 (CH₂), 25.8 (CH₃), 18.0 (C), -4.4 (CH₃), -4.8 (CH₃); HRMS (ESI+) 840.3923 (calcd for C₄₁H₆₃N₃NaO₁₀SSi 840.3901).

(2*S*,2*aR*,2*a*¹*S*,3*R*,4*aS*,6*R*,7*aS*)-tert-Butyl 3-hydroxy-6-(hydroxymethyl)-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)decahydro-1*H*-cyclopenta[*cd*]indole-1-carboxylate (52**)**

To a solution of **31** (650 mg, 0.795 mmol) in *N,N*-dimethylformamide (4.50 mL) was added tris(dimethylamino)sulfonium difluorotrimethylsilicate (1.20 g, 4.36 mmol) in *N,N*-dimethylformamide (3.00 mL) at room temperature. The solution was stirred for 3 h at the same temperature. To this solution was added CH₂Cl₂ and H₂O. The resulting solution was extracted with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 0:1) to afford **52** (540 mg, 97%) as a pale yellow amorphous solid. $[\alpha]_D^{27}$ 13.6° (*c* = 1.00, CHCl₃); IR (film) 3444, 2932, 1671, 1543, 1508, 1456, 1405, 1367, 1347, 1231, 1165, 1137, 1041, 826, 758 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.96 (m, 1H), 7.70-7.65 (m, 2H), 7.59 (m, 1H), 6.83 (s, 4H), 4.38 (m, 1H), 4.21 (ddd, *J*=8.0, 8.0, 8.0 Hz, 1H), 3.92 (t, *J*=6.2 Hz, 2H), 3.77 (s, 3H), 3.60 (m, 1H), 3.52-3.42 (m, 2H), 3.36 (ddd, *J*=13.3, 8.2, 8.2 Hz, 1H), 3.19 (ddd, *J*=13.3, 8.2, 4.1 Hz, 1H), 2.92 (s, 3H), 2.49 (br s, 1H), 2.10 (dd, *J*=7.3, 1.8 Hz, 1H), 2.05-1.93 (m, 2H), 1.85-1.51 (m, 11H), 1.46 (s, 9H), 1.28 (m, 1H), 1.10 (ddd, *J*=14.2, 11.9, 2.8 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 153.8 (C), 153.5 (C), 153.0 (C), 148.2 (C), 133.6 (CH), 131.8 (C), 131.6 (CH), 130.8 (CH), 124.1 (CH), 115.5 (CH), 114.7 (CH), 79.6 (C), 72.3 (CH), 68.9 (CH₂), 68.1 (CH₂), 61.7 (CH), 55.7 (CH₃), 54.9 (CH), 54.1 (CH), 52.4 (C), 47.1 (CH₂), 38.8 (CH₂), 37.9 (CH), 35.3 (CH₂), 34.4 (CH₃), 31.8 (CH₂), 29.2 (CH), 28.9 (2CH₂), 28.6 (CH₃), 26.0 (CH₂); HRMS (ESI+) 726.3032 (calcd for C₃₅H₄₉N₃NaO₁₀S 726.3036).

(2*S*,2*aR*,2*a*¹*S*,4*aS*,6*R*,7*aS*)-*tert*-Butyl 6-formyl-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)-3-oxodecahydro-1*H*-cyclopenta[*cd*]indole-1-carboxylate (32**)**

To a solution of **52** (510 mg, 0.725 mmol) in CH₂Cl₂ (25.0 mL) was added sodium hydrogen carbonate (1.90 g, 22.6 mmol), Dess-Martine periodinane (677 mg, 1.56 mmol) at 0 °C and the solution was stirred for 3 h at the same temperature. To this solution was added CH₂Cl₂ and H₂O and then the resulting solution was extracted with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:2) to afford **32** (497 mg, 98%) as a white amorphous solid. $[\alpha]_D^{27}$ 16.3° (*c* = 1.00, CHCl₃); IR (film) 2933, 2360, 1737, 1687, 1545, 1508, 1468, 1368, 1232, 1166, 1125, 1038, 982, 826, 760 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 9.70 (s, 1H), 7.96 (m, 1H), 7.70-7.64 (m, 2H), 7.58 (m, 1H), 6.82 (s, 4H), 4.00-3.91 (m, 2H), 3.86 (m, 1H), 3.76 (s, 3H), 3.60 (br s, 1H), 3.38 (m, 1H), 3.14 (ddd, *J*=15.6, 10.1, 5.5 Hz, 1H), 2.94 (s, 3H), 2.77-2.61 (m, 2H), 2.60-2.20 (m, 3H), 2.13 (dd, *J*=18.5, 6.0 Hz, 1H), 2.18-1.53 (m, 9H), 1.46 (s, 9H); ¹³C NMR (100 MHz, CDCl₃; one sp³ carbon signal missing, possibly due to broadening) δ 217.1 (C), 202.3 (CH), 153.9 (C), 153.5 (C), 152.9 (C), 148.2 (C), 133.5 (CH), 132.1 (C), 131.6 (CH), 130.7 (CH), 124.1 (CH), 115.5 (CH), 114.7 (CH), 80.4 (C), 68.3 (CH₂), 60.2 (CH), 58.3 (CH), 58.2 (CH), 55.7 (CH₃), 52.1 (C), 47.1 (CH₂), 44.0 (CH₂), 41.1 (CH), 35.1 (CH₃), 34.6 (CH₂), 33.6 (CH₂), 32.3 (CH₂), 28.4 (CH₃), 24.8 (CH₂), 22.6 (CH); HRMS (ESI+) 722.2705 (calcd for C₃₅H₄₅N₃NaO₁₀S 722.2723).

(2*S*,2*aR*,2*a*¹*S*,4*S*,4*aS*,6*S*,7*aS*,8*S*)-tert-Butyl 8-hydroxy-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)-3-oxodecahydro-1*H*-4,6-methanocyclopenta[*cd*]indole-1-carboxylate (33**)**

To a solution of **32** (497 mg, 0.710 mmol) in MeOH (27.0 mL) was added KOH (8.70 mL, 15.5 mmol, 10% w/w in MeOH) at 0 °C and the solution was stirred for 1.5 h at room temperature. To this solution was added CH₂Cl₂ and aqueous NH₄Cl at 0 °C and then the resulting solution was extracted with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:3) to afford **33** (490 mg, 98%) as a white amorphous solid. $[\alpha]_D^{27}$ 40.1° (*c* = 1.00, CHCl₃); IR (film) 3466, 2946, 2391, 1731, 1683, 1544, 1508, 1455, 1370, 1348, 1231, 1164, 1132, 1033, 826, 760 cm⁻¹; ¹H NMR (400 MHz, CDCl₃; this material was observed as a 3:2 mixture of rotamers) δ 8.00-7.91 (m, 1H), 7.68-7.60 (m, 2H), 7.58-7.52 (m, 1H), 6.84 (s, 4H), 4.05-3.94 (m, 3H), 3.85 (m, (3/5)1H), 3.80-3.73 (m, 3H+(2/5)1H), 3.63 (t, *J*=9.1 Hz, (2/5)1H), 3.47 (t, *J*=9.1 Hz, (3/5)1H), 3.41-3.25 (m, 1H), 3.22-3.08 (m, 1H), 2.94 (s, (3/5)3H), 2.92 (s, (2/5)3H), 2.84-2.72 (m, 3H), 2.69-2.36 (m, 3H), 2.17-2.10 (m, 1H+(3/5)1H), 2.6-1.84 (m, 4H+(3/5)1H), 1.78-1.43 (m, 1H), 1.70 (br s, (3/5)1H), 1.67 (br s, (2/5)1H), 1.44 (s, (2/5)9H), 1.42 (s, (3/5)9H), 0.38 (m, 1H); ¹³C NMR (100 MHz, CDCl₃; this material was observed as a mixture of rotamers) δ 218.0 (C), 217.6 (C), 154.5 (C), 154.1 (C), 153.9 (2C), 153.0 (2C), 148.3 (C), 148.1 (C), 133.6 (CH), 133.4 (CH), 132.2 (C), 131.6 (CH), 130.7 (CH), 130.6 (CH), 124.2 (CH), 124.1 (CH), 115.5 (CH), 114.7 (CH), 84.5 (CH), 84.4 (CH), 80.3 (C), 79.9 (C), 68.5 (2CH₂), 60.9 (CH), 60.8 (CH), 60.5 (CH), 59.2 (CH), 58.2 (CH), 58.1 (CH), 57.4 (CH), 57.0 (CH), 55.8 (CH₃), 54.0 (C), 53.0 (C), 47.8 (2CH₂), 39.3 (CH), 39.2 (CH), 38.9 (CH), 38.7 (CH), 35.0 (2CH₃), 34.0 (2CH₂), 32.9 (CH₂), 31.8 (CH₂), 30.4 (CH₂), 29.6 (CH₂), 28.5 (2CH₃), 27.8 (CH₂), 27.7 (CH₂), 24.6 (CH₂), 24.4 (CH₂); HRMS (ESI+) 722.2701 (calcd for C₃₅H₄₅N₃NaO₁₀S 722.2723).

(2*S*,2*aR*,2*a*¹*S*,4*S*,4*aS*,6*S*,7*aS*,8*S*)-tert-Butyl 8-(benzoyloxy)-2*a*¹-(3-(4-methoxyphenoxy)propyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)-3-oxodecahydro-1*H*-4,6-methanocyclopenta[*cd*]indole-1-carboxylate (53**)**

To a solution of **33** (460 mg, 0.657 mmol) in pyridine (8.0 mL) was added benzoyl chloride (152 μ L, 1.31 mmol) at 0 °C and the solution was stirred for 1 h at room temperature. To this solution was added CH_2Cl_2 and aqueous $NaHCO_3$ at 0 °C and then the resulting solution was extracted three times with CH_2Cl_2 . The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO_2 ; *n*-hexane:EtOAc = 4:1~1:2) to afford **53** (486 mg, 92%) as a white amorphous solid. $[\alpha]_D^{26}$ 88.1° ($c = 1.50$, $CHCl_3$); IR (film) 3471, 2952, 1736, 1716, 1687, 1545, 1508, 1453, 1369, 1273, 1231, 1168, 1110, 1069, 1026, 953, 826, 760 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$; this material was observed as a 3:2 mixture of rotamers) δ 8.10-7.92 (m, 3H), 7.70-7.52 (m, 4H), 7.50-7.27 (m, 2H), 6.92-6.80 (m, 4H), 5.10-5.00 (m, 1H), 4.20-3.70 (m, 3H), 3.75 (s, 3H), 3.67 (t, $J=9.2$ Hz, (2/5)1H), 3.50 (t, $J=9.2$ Hz, (3/5)1H), 3.45-3.25 (m, 1H), 3.22-2.84 (m, 3H), 2.94 (s, (3/5)3H), 2.92 (s, (2/5)3H), 2.78-2.55 (m, 1+(3/5)2H), 2.55-2.40 (m, 1+(2/5)1H), 2.15-1.35 (m, 7+(2/5)1H), 1.45 (s, 3.8H), 1.43 (s, 5.2H), 0.60 (m, 1H); ^{13}C NMR (100 MHz, $CDCl_3$; this material was observed as a mixture of rotamers) δ 216.5 (C), 216.2 (C), 165.5 (2C), 154.3 (C), 154.0 (C), 153.8 (2C), 153.0 (2C), 148.3 (2C), 133.5 (CH), 133.4 (CH), 133.1 (2CH), 132.1 (2C), 131.6 (2CH), 130.6 (CH), 130.5 (CH), 130.1 (2C), 130.0 (2CH), 128.4 (2CH), 124.1 (CH), 124.0 (CH), 115.5 (2CH), 114.7 (2CH), 85.8 (2CH), 80.3 (C), 80.0 (C), 68.4 (2CH₂), 60.4 (CH), 59.2 (CH), 57.9 (CH), 57.8 (CH), 57.6 (2CH), 57.3 (CH), 57.1 (CH), 55.7 (2CH₃), 54.2 (C), 53.1 (C), 47.9 (2CH₂), 39.4 (CH), 39.3 (CH), 36.7 (CH), 36.5 (CH), 35.2 (CH₃), 35.1 (CH₃), 33.9 (2CH₂), 33.1 (CH₂), 32.1 (CH₂), 30.3 (CH₂), 29.4 (CH₂), 28.4 (2CH₃), 28.3 (CH₂), 28.0 (CH₂), 24.5 (CH₂), 24.3 (CH₂); HRMS (ESI+) 826.2996 (calcd for $C_{42}H_{49}N_3NaO_{11}S$ 826.2986).

(2*S*,2*aR*,2*a*¹*S*,4*S*,4*aS*,6*S*,7*aS*,8*S*)-tert-Butyl 8-(benzoyloxy)-2*a*¹-(3-hydroxypropyl)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)-3-oxodecahydro-1*H*-4,6-methanocyclopenta[*cd*]indole-1-carboxylate (54**)**

To a solution of **53** (760 mg, 0.946 mmol) in CH₃CN (20.0 mL) and H₂O (5.0 mL) was added cerium(IV) ammonium nitrate (570 mg, 1.04 mmol) at 0 °C and the solution was stirred for 2.5 h at the same temperature. To this solution was added CH₂Cl₂ and aqueous NaHCO₃ at 0 °C and then the resulting solution was extracted three times with CH₂Cl₂. The organic phase was washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:1~1:3) to afford **54** (590 mg, 89%) as a white amorphous solid. $[\alpha]_D^{27}$ 67.7° (*c* = 1.00, CHCl₃); IR (film) 3503, 2944, 1735, 1716, 1683, 1543, 1454, 1369, 1274, 1166, 1112, 1068, 985, 852, 760 cm⁻¹; ¹H NMR (400 MHz, CDCl₃; this material was observed as a 2:1 mixture of rotamers) δ 8.00-7.91 (m, 3H), 7.74-7.52 (m, 4H), 7.47-7.40 (m, 2H), 5.08-5.02 (m, 1H), 3.92 (d, *J*=9.2 Hz, (2/3)1H), 3.84-3.75 (m, 1H+(1/3)1H), 3.73-3.60 (m, 1H+(1/3)1H), 3.50-3.30 (m, 1H+(2/3)1H), 3.15-2.84 (m, 3H), 2.95 (s, (2/3)3H), 2.94 (s, (1/3)3H), 2.75-2.62 (m, 2H+(2/3)1H), 2.55-2.38 (m, 1H+(1/3)1H), 1.98-1.55 (m, 7H), 1.46 (s, (1/3)9H), 1.43 (s, (2/3)9H), 0.66-0.53 (m, 1H); ¹³C NMR (100 MHz, CDCl₃; this material was observed as a mixture of rotamers) δ 216.7 (C), 216.4 (C), 165.5 (2C), 154.2 (C), 153.9 (C), 148.2 (2C), 133.7 (CH), 133.6 (CH), 133.1 (2CH), 131.9 (C), 131.8 (C), 131.7 (2CH), 130.6 (CH), 130.4 (CH), 130.1 (C), 129.5 (2CH), 128.4 (2CH), 124.1 (CH), 124.1 (CH), 85.8 (CH), 85.7 (CH), 80.2 (C), 80.0 (C), 62.6 (2CH₂), 59.4 (CH), 58.4 (CH), 58.2 (CH), 58.0 (CH), 57.5 (CH), 57.5 (CH), 57.3 (CH), 57.0 (CH), 54.4 (C), 53.3 (C), 48.0 (2CH₂), 39.2 (CH), 39.0 (CH), 36.7 (CH), 36.5 (CH), 35.5 (CH₃), 35.3 (CH₃), 33.1 (CH₂), 33.0 (CH₂), 32.9 (CH₂), 31.9 (CH₂), 30.5 (CH₂), 29.6 (CH₂), 28.4 (2CH₃), 28.1 (CH₂), 27.8 (CH₂), 27.5 (CH₂), 27.3 (CH₂); HRMS (ESI+) 720.2587 (calcd for C₃₅H₄₃N₃NaO₁₀S 720.2567).

(2*S*,2*aR*,2*a*¹*S*,4*S*,4*aS*,6*S*,7*aS*,8*S*)-tert-Butyl 8-(benzoyloxy)-2-(2-(*N*-methyl-2-nitrophenylsulfonamido)ethyl)-3-oxo-2*a*¹-(3-oxopropyl)decahydro-1*H*-4,6-methanocyclopenta[*cd*]indole-1-carboxylate (34**)**

To a solution of **54** (550 mg, 0.788 mmol) in CH₂Cl₂ (600 mL) was added sodium hydrogen carbonate (1.10 g, 13.1 mmol), Dess-Martine periodinane (401 mg, 0.946 mmol) at 0 °C. After stirring for 1 h at room temperature, water was added at the same temperature. The resulting solution was extracted three times with CH₂Cl₂. The combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; *n*-hexane:EtOAc = 1:2) to give **34** (540 mg, 98%) as a colorless oil. $[\alpha]_D^{27}$ 76.7° (*c* = 1.75, CHCl₃); IR (film) 3854, 2968, 1718, 1685, 1544, 1453, 1369, 1273, 1167, 1111, 1069, 985, 852, 759 cm⁻¹; ¹H NMR (400 MHz, CDCl₃; this material was observed as a 2:1 mixture of rotamers) δ 9.90 (s, (2/3)1H), 9.88 (s, (1/3)1H), 8.00-7.88 (m, 3H), 7.75-7.65 (m, 2H), 7.65-7.52 (m, 2H), 7.47-7.40 (m, 2H), 5.04 (s, 1H), 3.91 (d, *J*=10.0 Hz, (2/3)1H), 3.81 (d, *J*=10.0 Hz, (1/3)1H), 3.67 (dd, *J*=8.6, 8.6 Hz, (1/3)1H), 3.50 (dd, *J*=9.2, 9.2 Hz, (2/3)1H), 3.46-3.30 (m, 1H), 3.15-3.00 (m, 2H), 2.95 (s, (2/3)3H), 2.94 (s, (1/3)3H), 2.90-2.58 (m, 5H+(2/3)1H), 2.55-2.41 (m, 1H+(1/3)1H), 2.22-2.17 (m, 1H), 2.02-1.92 (m, 2H), 1.75-1.49 (m, 3H), 1.45 (s, (1/3)9H), 1.43 (s, (2/3)9H), 0.65-0.50 (m, 1H); ¹³C NMR (100 MHz, CDCl₃; this material was observed as a mixture of rotamers) δ 216.1 (C), 215.7 (C), 201.2 (CH), 200.9 (CH), 165.5 (2C), 154.2 (C), 153.9 (C), 148.3 (C), 148.2 (C), 133.7 (CH), 133.6 (CH), 133.2 (2CH), 132.0 (2C), 131.7 (2CH), 130.6 (CH), 130.4 (CH), 130.0 (2C), 129.6 (2CH), 128.4 (2CH), 124.2 (CH), 124.1 (CH), 85.7 (CH), 85.6 (CH), 80.5 (C), 80.2 (C), 60.3 (CH), 58.6 (CH), 58.3 (CH), 57.9 (CH), 57.7 (CH), 57.5 (CH), 57.3 (CH), 57.1 (CH), 53.7 (C), 52.7 (C), 47.9 (2CH₂), 39.4 (CH), 39.2 (CH), 39.0 (CH₂), 38.8 (CH₂), 36.7 (CH), 36.5 (CH), 35.6 (CH₃), 35.4 (CH₃), 33.3 (CH₂), 32.3 (CH₂), 30.1 (CH₂), 29.3 (CH₂), 28.7 (CH₂), 28.5

(2CH₃), 28.3 (CH₂), 28.2 (CH₂), 27.8 (CH₂); HRMS (ESI+) 718.2388 (calcd for C₃₅H₄₁N₃NaO₁₀S 718.2410).

(2*S*,2*aS*,2*a*¹*S*,4*S*,5*aS*,6*S*,7*S*,10*aS*,10*bR*,13*S*)-8-Methyl-1-oxododecahydro-1*H*-2*a*¹,7-ethano-2,4-methanocyclopenta[*cd*]pyrimido[1,6-*a*]indol-13-yl benzoate (35**)**

To a solution of **34** (67.0 mg, 0.144 mmol) in CH₂Cl₂ (3.0 mL) was added TFA (2.0 mL) at -78°C . After stirring for 15 min at room temperature, CH₂Cl₂ and aqueous NaHCO₃ were added at the 0°C . The resulting solution was extracted twice with CH₂Cl₂. The combined organic phases were washed with brine, dried over sodium sulfate and filtered. The filtrate was concentrated *in vacuo*. The residual oil was dissolved in CH₃CN (40.0 mL) and potassium phosphate (200 mg, 0.941 mmol), thiophenol (230 μL , 2.26 mmol) was added at room temperature. After stirring for 2 h at room temperature, acetic acid (320 μL) was added and then the reaction mixture was heated at 50°C for 18 h. After cooling to room temperature, the solution was concentrated *in vacuo* and the residue was purified by flash column chromatography (SiO₂; CHCl₃:MeOH = 5:1) to give **35** (49.0 mg, 89% for 2 steps) as a white solid. mp $212.2\text{--}213.0^\circ\text{C}$; $[\alpha]_D^{27} 156^\circ$ ($c = 1.75$, CHCl₃); IR (film) 3421, 2943, 1722, 1451, 1274, 1177, 1110, 1068, 984, 855 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.98 (dd, $J=7.6, 1.4$ Hz, 2H), 7.55 (dd, $J=7.6, 7.6$ Hz, 1H), 7.42 (dd, $J=7.6, 7.6$ Hz, 2H), 5.02 (s, 1H), 3.70 (t, $J=8.2$ Hz, 1H), 3.51–3.43 (m, 1H), 3.00–2.90 (m, 2H), 2.71–2.57 (m, 2H), 2.56–2.46 (m, 2H), 2.45 (s, 3H), 2.41 (d, $J=6.4$ Hz, 1H), 2.38–2.30 (m, 1H), 2.22 (ddd, $J=15.6, 9.9, 9.9$ Hz, 1H), 2.10–1.93 (m, 2H), 1.80–1.70 (m, 3H), 0.99 (dd, $J=15.6, 10.1$ Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 219.9 (C), 165.5 (C), 133.0 (CH), 130.2 (C), 129.6 (CH), 128.4 (CH), 86.4 (CH), 76.5 (CH), 65.1 (CH), 63.2 (CH), 61.2 (CH), 60.8 (CH), 54.4 (C), 42.7 (CH₃), 42.6 (CH₂), 42.0 (CH), 38.5 (CH), 35.8 (CH₂), 28.1 (CH₂), 26.6 (CH₂), 24.5 (CH₂), 23.4 (CH₂); HRMS (ESI+) 393.2180 (calcd for C₂₄H₂₉N₂O₃ 393.2178).

(+)-Lycopalhine A (1)

To a solution of **35** (20.0 mg, 0.0509 mmol) in MeOH (1.0 mL) was added potassium carbonate (20 mg 0.145 mmol) at 0 °C. After stirring for 3 h at room temperature, the solution was concentrated *in vacuo* and the residue was purified by PTLC (NH₂ silica gel; CHCl₃:MeOH = 10:1) to give (+)-lycopalhine A (**1**) and *epi*-lycopalhine A (**36**) (13.0 mg, 89%) as a colorless gum. $[\alpha]_D^{25}$ 105° (c = 0.200, MeOH); lit.¹ $[\alpha]_D^{15}$ 89.1° (c = 0.17, MeOH); IR (film) 3411, 2940, 1714, 1659, 1454, 1367, 1300, 1177, 1048, 1026, 854 cm⁻¹; ¹H NMR (400 MHz, pyridine-*d*₅) δ 4.46 (dd, J =11.0, 5.5 Hz, 0.2H, minor isomer), 4.29 (s, 1H), 3.95-3.89 (m, 0.2H, minor isomer), 3.81 (dd, J =10.1, 7.4 Hz, 0.2H, minor isomer), 3.65 (dd, J =10.1, 6.4 Hz, 1H), 3.59-3.54 (m, 1H), 3.05 (d, J =8.2 Hz, 1H), 2.98 (ddd, J =12.8, 9.2, 3.2 Hz, 1H), 2.84 (ddd, J =10.6, 7.8, 2.5 Hz, 0.2H, minor isomer), 2.63-2.55 (m, 0.2H, minor isomer), 2.57 (dd, J =9.9, 9.9 Hz, 1H), 2.47-2.22 (m, 6H), 2.45 (s, 3H), 2.16 (ddd, J =11.5, 4.6, 4.6 Hz, 1H), 2.07 (ddd, J =15.6, 9.6, 9.6 Hz, 1H), 1.87 (ddd, J =11.9, 11.9, 7.4 Hz, 1H), 1.75-1.60 (m, 4H), 1.55 (dd, J =12.5, 7.8 Hz, 1H), 1.35-1.25 (m, 0.2H, minor isomer), 0.93 (dd, J =15.1, 10.3 Hz, 1H); ¹³C NMR (100 MHz, pyridine-*d*₅) major isomer: δ 221.6 (C), 85.6 (CH), 77.0 (CH), 65.6 (CH), 65.3 (CH), 63.4 (CH), 61.1 (CH), 54.3 (C), 43.0 (CH₂), 42.8 (CH₃), 42.3 (CH), 41.4 (CH), 36.0 (CH₂), 27.9 (CH₂), 27.3 (CH₂), 25.4 (CH₂), 24.2 (CH₂); minor isomer: δ 219.1 (C), 76.3 (CH), 75.5 (CH), 64.3 (CH), 62.1 (CH), 58.7 (CH), 58.2 (CH), 55.6 (C), 43.8 (CH), 42.1 (CH₃), 41.0 (CH₂), 37.6 (CH), 36.7 (CH₂), 27.0 (CH₂), 23.5 (CH₂), 23.0 (CH₂), 22.8 (CH₂); HRMS (ESI+) 289.1902 (calcd for C₁₇H₂₅N₂O₂ 289.1916).

¹ Dong, L.-B.; Yang, J.; He, J.; Luo, H.-R.; Wu, X.-D.; Deng, X.; Peng, L.-Y.; Cheng, X.; Zhao, Q.-S. *Chem. Commun.* **2012**, 48, 9038.

Table S1. ^1H and ^{13}C NMR data for (+)-lycopalhine A (1) in pyridine- d_5

(+)-lycopalhine A (1)

Synthetic			Natural ^a	
no.	δ_{C} (100 MHz)	δ_{H} (400 MHz)	δ_{C} (125 MHz)	δ_{H} (500 MHz)
1a	42.99	2.98 (ddd, 12.8, 9.2, 3.2 Hz, 1H)	43.06	2.98 (ddd, 13.5, 10.5, 4.5 Hz, 1H)
1b		2.34-2.31 (m, 1H)		2.34 (m, 1H)
2a	24.22	2.29-2.21 (m, 1H)	24.32	2.27 (m, 1H)
2b		1.75-1.73 (m, 1H)		1.74 (m, 1H)
3	61.10	3.58-3.54 (m, 1H)	61.17	3.56 (m, 1H)
4	63.45	2.37-2.35 (m, 1H)	63.57	2.36 (m, 1H)
5	221.64		221.56	
6	65.55	3.05 (d, 8.2 Hz, 1H)	65.58	3.05 (d, 7.5 Hz, 1H)
7	42.26	2.43-2.40 (m, 1H)	42.31	2.43 (m, 1H)
8a	27.93	2.16 (ddd, 11.5, 4.6, 4.6 Hz)	27.95	2.16 (dt, 12.0, 4.5 Hz, 1H)
8b		1.72-1.70 (m, 1H)		1.70 (overlap, 1H)
9	77.00	3.65 (dd, 10.1, 6.4 Hz)	77.12	3.64 (dd, 10.5, 7.5 Hz, 1H)
10a	25.41	1.75-1.73 (m, 1H)	25.48	1.70 (overlap, 1H)
10b		1.65-1.60 (m, 1H)		1.63 (m, 1H)
11a	35.99	1.87 (ddd, 11.9, 11.9, 7.4 Hz)	36.07	1.85 (td, 12.0, 7.5 Hz, 1H)
11b		1.55 (dd, 12.5, 7.8 Hz)		1.55 (dd, 12.0, 7.5 Hz, 1H)
12	54.33		54.38	
13	65.27	2.57 (dd, 9.9, 9.9 Hz, 1H)	65.35	2.56 (t, 10.0 Hz, 1H)
14a	27.28	2.07 (ddd, 15.6, 9.6, 9.6 Hz, 1H)	27.34	2.07 (dt, 15.5, 10.0 Hz, 1H)
14b		0.93 (dd, 15.1, 10.3 Hz, 1H)		0.92 (dd, 15.5, 10.0 Hz, 1H)
15	41.35	2.42-2.40 (m, 1H)	41.41	2.40 (m, 1H)
16	85.62	4.29 (s, 1H)	85.68	4.29 (brs, 1H)
17	42.83	2.45 (s, 3H)	42.86	2.45 (s, 3H)

^a Dong, L.-B.; Yang, J.; He, J.; Luo, H.-R.; Wu, X.-D.; Deng, X.; Peng, L.-Y.; Cheng, X.; Zhao, Q.-S. *Chem. Commun.* **2012**, 48, 9038.

12

X : parts per Million : Proton

12

X : parts per Million : Carbon13

X : parts per Million : Proton

X : parts per Million : Carbon13

```

Experiment      = single pulse.jxp
Sample_id      = YO-5-095
Solvent        = CHLOROFORM-D
Creation Time   = 22-JUN-2015 15:31:52

Field Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 2.18365952[s]
X_Domain       = 1H
X_Freq         = 399.78219838[MHz]
X_Offset       = 5[ppm]
X_Points       = 16384
X_Prescans     = 1
X_Resolution   = 0.45794685[Hz]
X_Sweep        = 7.5030012[kHz]
X_Sweep_Clippped = 6.00240096[kHz]
Irr_Domain     = Proton
Irr_Freq       = 399.78219838[MHz]
Irr_Offset     = 5[ppm]
Tri_Domain     = Proton
Tri_Freq       = 399.78219838[MHz]
Tri_Offset     = 5[ppm]
Clipped        = FALSE
Mod_Return     = 1
Probe_Recovery = 5[us]
Scans          = 8
Total_Scans    = 8

Relaxation_Delay = 5[s]
Recvr_Gain       = 24
Temp_Get         = 24.9[degC]
X_90_Width      = 9.8[us]
X_Acq_Time      = 2.18365952[s]
X_Angle         = 45[deg]
X_Atn           = 1[db]
X_Pulse         = 4.9[us]
Irr_Mode        = Off
Tri_Mode        = Off
Dante_Presat    = FALSE
Dead_Time       = 0[us]
Delay           = 0[us]
Initial_Wait    = 1[s]
Repetition_Time = 7.18365952[s]

```

```

Experiment      = single pulse dec.jxp
Sample_id      = YO-5-095 overNight
Solvent        = CHLOROFORM-D
Creation Time   = 23-JUN-2015 00:02:30

Field Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 1.04333312[s]
X_Domain       = 13C
X_Freq         = 100.52530333[MHz]
X_Offset       = 100[ppm]
X_Points       = 32768
X_Prescans     = 4
X_Resolution   = 0.95846665[Hz]
X_Sweep        = 31.40703518[kHz]
X_Sweep_Clippped = 25.12562814[kHz]
Irr_Domain     = Proton
Irr_Freq       = 399.78219838[MHz]
Irr_Offset     = 5[ppm]
Clipped        = FALSE
Mod_Return     = 1
Probe_Recovery = 20[us]
Scans          = 2048
Total_Scans    = 2048

Relaxation_Delay = 1.5[s]
Recvr_Gain       = 50
Temp_Get         = 25.1[degC]
X_90_Width      = 8.6[us]
X_Acq_Time      = 1.04333312[s]
X_Angle         = 30[deg]
X_Atn           = 4[db]
X_Pulse         = 2.86666667[us]
Irr_Atn_Dec     = 22.569[db]
Irr_Atn_No     = 22.569[db]
Irr_Noise       = WALTZ
Irr_Pwidth      = 0.115[ms]
Dead_Time       = 0[us]
Decoupling      = TRUE
Delay           = 0[us]
Initial_Wait    = 1[s]
Noe             = TRUE
Noe_Time        = 1.5[s]
Repetition_Time = 2.54333312[s]

```

X : parts per Million : Proton

X₂₀₇: parts per Million : Carbon13

Experiment = single pulse.jxp
Sample_id = YO-5-108
Solvent = CHLOROFORM-D
Creation_Time = 30-JUN-2015 21:05:56
Field_Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 2.18365952[s]
X_Domain = 1H
X_Freq = 399.78219838[MHz]
X_Offset = 5[ppm]
X_Points = 16384
X_Prescans = 1
X_Resolution = 0.45794685[Hz]
X_Sweep = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Tri_Domain = Proton
Tri_Freq = 399.78219838[MHz]
Tri_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 5[us]
Scans = 8
Total_Scans = 8
Relaxation_Delay = 5[s]
Recvr_Gain = 30
Temp_Get = 25[dc]
X_90_Width = 9.8[us]
X_Acq_Time = 2.18365952[s]
X_Angle = 45[deg]
X_Atn = 1[db]
X_Pulse = 4.9[us]
Irr_Mode = Off
Tri_Mode = Off
Dante_Presat = FALSE
Dead_Time = 0[us]
Delay = 0[us]
Initial_Wait = 1[s]
Repetition_Time = 7.18365952[s]

Experiment = single pulse_dec.jxp
Sample_id = YO-5-108 overnight
Solvent = CHLOROFORM-D
Creation_Time = 3-JUL-2015 00:03:15
Field_Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 1.04333312[s]
X_Domain = 13C
X_Freq = 100.52530333[MHz]
X_Offset = 100[ppm]
X_Points = 32768
X_Prescans = 4
X_Resolution = 0.95846665[Hz]
X_Sweep = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 20[us]
Scans = 2048
Total_Scans = 2048
Relaxation_Delay = 1.5[s]
Recvr_Gain = 50
Temp_Get = 25.2[dc]
X_90_Width = 8.6[us]
X_Acq_Time = 1.04333312[s]
X_Angle = 30[deg]
X_Atn = 4[db]
X_Pulse = 2.86666667[us]
Irr_Atn_Dec = 22.569[db]
Irr_Atn_No2 = 22.569[db]
Irr_Noise = WALTZ
Irr_Pwidth = 0.115[ms]
Dead_Time = 0[us]
Decoupling = TRUE
Delay = 0[us]
Initial_Wait = 1[s]
Noe = TRUE
Noe_Time = 1.5[s]
Repetition_Time = 2.54333312[s]

Experiment = single pulse.jxp
 Sample_id = YO-5-109
 Solvent = CHLOROFORM-D
 Creation Time = 1-JUL-2015 21:31:11
 Field Strength = 9.389766[T] (400[MHz])
 X_Acq_Duration = 2.18365952[s]
 X_Domain = 1H
 X_Freq = 399.78219838 [MHz]
 X_Offset = 5 [ppm]
 X_Points = 16384
 X_Prescans = 1
 X_Resolution = 0.45794685 [Hz]
 X_Sweep = 7.5030012 [kHz]
 X_Sweep_Clippped = 6.00240096 [kHz]
 Irr_Domain = Proton
 Irr_Freq = 399.78219838 [MHz]
 Irr_Offset = 5 [ppm]
 Tri_Domain = Proton
 Tri_Freq = 399.78219838 [MHz]
 Tri_Offset = 5 [ppm]
 Clipped = FALSE
 Mod_Return = 1
 Probe_Recovery = 5 [us]
 Scans = 8
 Total_Scans = 8
 Relaxation_Delay = 5 [s]
 Recvr_Gain = 34
 Temp_Get = 25 [dc]
 X_90_Width = 9.8 [us]
 X_Acq_Time = 2.18365952 [s]
 X_Angle = 45 [deg]
 X_Atn = 1 [dB]
 X_Pulse = 4.9 [us]
 Irr_Mode = Off
 Tri_Mode = Off
 Dante_Presat = FALSE
 Dead_Time = 0 [us]
 Delay = 0 [us]
 Initial_Wait = 1 [s]
 Repetition_Time = 7.18365952 [s]

R = $-(CH_2)_3OPMP$
 Ar = $-C_6H_4-p-CF_3$

Experiment = single pulse dec.jxp
 Sample_id = YO-5-109 overNight
 Solvent = CHLOROFORM-D
 Creation Time = 5-JUL-2015 00:03:16
 Field Strength = 9.389766[T] (400[MHz])
 X_Acq_Duration = 1.04333312 [s]
 X_Domain = 13C
 X_Freq = 100.52530333 [MHz]
 X_Offset = 100 [ppm]
 X_Points = 32768
 X_Prescans = 4
 X_Resolution = 0.95846665 [Hz]
 X_Sweep = 31.40703518 [kHz]
 X_Sweep_Clippped = 25.12562814 [kHz]
 Irr_Domain = Proton
 Irr_Freq = 399.78219838 [MHz]
 Irr_Offset = 5 [ppm]
 Clipped = FALSE
 Mod_Return = 1
 Probe_Recovery = 20 [us]
 Scans = 2048
 Total_Scans = 2048
 Relaxation_Delay = 1.5 [s]
 Recvr_Gain = 50
 Temp_Get = 25.1 [dc]
 X_90_Width = 8.6 [us]
 X_Acq_Time = 1.04333312 [s]
 X_Angle = 30 [deg]
 X_Atn = 4 [dB]
 X_Pulse = 2.86666667 [us]
 Irr_Atn_Dec = 22.569 [dB]
 Irr_Atn_No2 = 22.569 [dB]
 Irr_Noise = WALTZ
 Irr_Pwidth = 0.115 [ms]
 Dead_Time = 0 [us]
 Decoupling = TRUE
 Delay = 0 [us]
 Initial_Wait = 1 [s]
 Noe = TRUE
 Noe_Time = 1.5 [s]
 Repetition_Time = 2.54333312 [s]

R = $-(CH_2)_3OPMP$
 Ar = $-C_6H_4-p-CF_3$

X : parts per Million : Proton

```

Experiment      = single pulse.jxp
Sample Id       = YO-5-118
Solvent         = CHLOROFORM-D
Creation Time    = 6-JUL-2015 18:56:11

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 2.18365952[s]
X_Domain        = 1H
X_Freq          = 399.78219838[MHz]
X_Offset        = 5[ppm]
X_Points        = 16384
X_Prescans      = 1
X_Resolution    = 0.45794685[Hz]
X_Sweep         = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Tri_Domain      = Proton
Tri_Freq        = 399.78219838[MHz]
Tri_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 5[us]
Scans           = 8
Total_Scans     = 8

Relaxation_Delay = 5[s]
Recvr_Gain       = 30
Temp_Get         = 25[dc]
X_90_Width      = 9.8[us]
X_Acq_Time      = 2.18365952[s]
X_Angle         = 45[deg]
X_Atn           = 1[db]
X_Pulse         = 4.9[us]
Irr_Mode        = Off
Tri_Mode        = Off
Dante_Presat    = FALSE
Dead_Time       = 0[us]
Delay           = 0[us]
Initial_Wait    = 1[s]
Repetition_Time = 7.18365952[s]

```

R = $-(CH_2)_3OPMP$

Ar = $-C_6H_4-p-CF_3$

X : parts per Million : Carbon13

```

Experiment      = single pulse dec.jxp
Sample Id       = YO-5-118 overnight
Solvent         = CHLOROFORM-D
Creation Time    = 7-JUL-2015 00:02:08

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 1.04333312[s]
X_Domain        = 13C
X_Freq          = 100.52530333[MHz]
X_Offset        = 100[ppm]
X_Points        = 32768
X_Prescans      = 4
X_Resolution    = 0.95846665[Hz]
X_Sweep         = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 20[us]
Scans           = 2048
Total_Scans     = 2048

Relaxation_Delay = 1.5[s]
Recvr_Gain       = 50
Temp_Get         = 25[dc]
X_90_Width      = 8.6[us]
X_Acq_Time      = 1.04333312[s]
X_Angle         = 30[deg]
X_Atn           = 4[db]
X_Pulse         = 2.86666667[us]
Irr_Atn_Dec     = 22.569[db]
Irr_Atn_Noise   = 22.569[db]
Irr_Noise       = WALTZ
Irr_Pwidth      = 0.115[ms]
Dead_Time       = 0[us]
Decoupling      = TRUE
Delay           = 0[us]
Initial_Wait    = 1[s]
Noe              = TRUE
Noe_Time        = 1.5[s]
Repetition_Time = 2.54333312[s]

```

R = $-(CH_2)_3OPMP$

Ar = $-C_6H_4-p-CF_3$

Experiment = single pulse.jxp
 Sample Id = YO-5-126
 Solvent = CHLOROFORM-D
 Creation Time = 9-JUL-2015 20:54:39
 Field Strength = 9.389766[T] (400[MHz])
 X_Acq_Duration = 2.18365952[s]
 X_Domain = 1H
 X_Freq = 399.78219838[MHz]
 X_Offset = 5[ppm]
 X_Points = 16384
 X_Prescans = 1
 X_Resolution = 0.45794685[Hz]
 X_Sweep = 7.5030012[kHz]
 X_Sweep_Clippped = 6.00240096[kHz]
 Irr_Domain = Proton
 Irr_Freq = 399.78219838[MHz]
 Irr_Offset = 5[ppm]
 Tri_Domain = Proton
 Tri_Freq = 399.78219838[MHz]
 Tri_Offset = 5[ppm]
 Clipped = FALSE
 Mod_Return = 1
 Probe_Recovery = 5[us]
 Scans = 8
 Total_Scans = 8
 Relaxation_Delay = 5[s]
 Recvr_Gain = 30
 Temp_Get = 25.1[dc]
 X_90_Width = 9.8[us]
 X_Acq_Time = 2.18365952[s]
 X_Angle = 45[deg]
 X_Atn = 1[db]
 X_Pulse = 4.9[us]
 Irr_Mode = Off
 Tri_Mode = Off
 Dante_Presat = FALSE
 Dead_Time = 0[us]
 Delay = 0[us]
 Initial_Wait = 1[s]
 Repetition_Time = 7.18365952[s]

27
 R = -(CH₂)₃OPMP
 Ar = -C₆H₄-p-CF₃

Experiment = single pulse dec.jxp
 Sample Id = YO-5-126 overnight
 Solvent = CHLOROFORM-D
 Creation Time = 11-JUL-2015 00:03:19
 Field Strength = 9.389766[T] (400[MHz])
 X_Acq_Duration = 1.04333312[s]
 X_Domain = 13C
 X_Freq = 100.52530333[MHz]
 X_Offset = 100[ppm]
 X_Points = 32768
 X_Prescans = 4
 X_Resolution = 0.95846665[Hz]
 X_Sweep = 31.40703518[kHz]
 X_Sweep_Clippped = 25.12562814[kHz]
 Irr_Domain = Proton
 Irr_Freq = 399.78219838[MHz]
 Irr_Offset = 5[ppm]
 Clipped = FALSE
 Mod_Return = 1
 Probe_Recovery = 20[us]
 Scans = 2048
 Total_Scans = 2048
 Relaxation_Delay = 1.5[s]
 Recvr_Gain = 50
 Temp_Get = 25.2[dc]
 X_90_Width = 8.6[us]
 X_Acq_Time = 1.04333312[s]
 X_Angle = 30[deg]
 X_Atn = 4[db]
 X_Pulse = 2.86666667[us]
 Irr_Atn_Dec = 22.569[db]
 Irr_Atn_Noise = 22.569[db]
 Irr_Noise = WALTZ
 Irr_Pwidth = 0.115[ms]
 Dead_Time = 0[us]
 Decoupling = TRUE
 Delay = 0[us]
 Initial_Wait = 1[s]
 Noe = TRUE
 Noe_Time = 1.5[s]
 Repetition_Time = 2.54333312[s]

27
 R = -(CH₂)₃OPMP
 Ar = -C₆H₄-p-CF₃

```

Experiment      = single pulse.jxp
Sample Id       = YO-5-130
Solvent         = CHLOROFORM-D
Creation Time    = 13-JUL-2015 22:23:14

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 2.18365952[s]
X_Domain        = 1H
X_Freq          = 399.78219838[MHz]
X_Offset        = 5[ppm]
X_Points        = 16384
X_Prescans      = 1
X_Resolution    = 0.45794685[Hz]
X_Sweep         = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Tri_Domain      = Proton
Tri_Freq        = 399.78219838[MHz]
Tri_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 5[us]
Scans           = 8
Total_Scans     = 8

Relaxation_Delay = 5[s]
Recvr_Gain       = 30
Temp_Get         = 25.5[dc]
X_90_Width       = 9.8[us]
X_Acq_Time       = 2.18365952[s]
X_Angle          = 45[deg]
X_Atn            = 1[db]
X_Pulse          = 4.9[us]
Irr_Mode         = Off
Tri_Mode         = Off
Dante_Presat     = FALSE
Dead_Time        = 0[us]
Delay            = 0[us]
Initial_Wait     = 1[s]
Repetition_Time  = 7.18365952[s]

```

```

Experiment      = single pulse dec.jxp
Sample Id       = YO-5-130 overNight
Solvent         = CHLOROFORM-D
Creation Time    = 14-JUL-2015 00:01:57

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 1.04333312[s]
X_Domain        = 13C
X_Freq          = 100.52530333[MHz]
X_Offset        = 100[ppm]
X_Points        = 32768
X_Prescans      = 4
X_Resolution    = 0.95846665[Hz]
X_Sweep         = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Tri_Domain      = Proton
Tri_Freq        = 399.78219838[MHz]
Tri_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 20[us]
Scans           = 2048
Total_Scans     = 2048

Relaxation_Delay = 1.5[s]
Recvr_Gain       = 50
Temp_Get         = 25.5[dc]
X_90_Width       = 8.6[us]
X_Acq_Time       = 1.04333312[s]
X_Angle          = 30[deg]
X_Atn            = 4[db]
X_Pulse          = 2.86666667[us]
Irr_Atn_Dec      = 22.569[db]
Irr_Atn_Noise   = 22.569[db]
Irr_Noise       = WALTZ
Irr_Pwidth       = 0.115[ms]
Dead_Time        = 0[us]
Decoupling       = TRUE
Delay            = 0[us]
Initial_Wait     = 1[s]
Noe              = TRUE
Noe_Time         = 1.5[s]
Repetition_Time  = 2.54333312[s]

```

X : parts per Million : Proton

```

Experiment      = single pulse.jxp
Sample_id       = YO-5-139
Solvent         = CHLOROFORM-D
Creation Time   = 21-JUL-2015 19:23:52

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration = 2.18365952[s]
X_Domain        = 1H
X_Freq          = 399.78219838[MHz]
X_Offset        = 5[ppm]
X_Points        = 16384
X_Prescans      = 1
X_Resolution    = 0.45794685[Hz]
X_Sweep         = 7.5030012[kHz]
X_Sweep_Clippped = 6.00240096[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Tri_Domain      = Proton
Tri_Freq        = 399.78219838[MHz]
Tri_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 5[us]
Scans           = 8
Total_Scans     = 8

Relaxation_Delay = 5[s]
Recvr_Gain       = 30
Temp_Get         = 25.5[dC]
X_90_Width      = 9.8[us]
X_Acq_Time       = 2.18365952[s]
X_Angle         = 45[deg]
X_Atn           = 1[dB]
X_Pulse         = 4.9[us]
Irr_Mode        = Off
Tri_Mode        = Off
Dante_Presat    = FALSE
Dead_Time       = 0[us]
Delay           = 0[us]
Initial_Wait     = 1[s]
Repetition_Time  = 7.18365952[s]

```

```

Experiment      = single pulse_dec.jxp
Sample_id       = YO-5-139 overnigh
Solvent         = CHLOROFORM-D
Creation Time   = 22-JUL-2015 00:02:54

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration = 1.04333312[s]
X_Domain        = 13C
X_Freq          = 100.52530333[MHz]
X_Offset        = 100[ppm]
X_Points        = 32768
X_Prescans      = 4
X_Resolution    = 0.95846665[Hz]
X_Sweep         = 31.40703518[kHz]
X_Sweep_Clippped = 25.12562814[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 20[us]
Scans           = 2048
Total_Scans     = 2048

Relaxation_Delay = 1.5[s]
Recvr_Gain       = 50
Temp_Get         = 25.6[dC]
X_90_Width      = 8.6[us]
X_Acq_Time       = 1.04333312[s]
X_Angle         = 30[deg]
X_Atn           = 4[dB]
X_Pulse         = 2.86666666[us]
Irr_Atn_Dec     = 22.569[dB]
Irr_Atn_Noise   = 22.569[dB]
Irr_Noise       = WALTZ
Irr_Pwidth      = 0.115[ms]
Dead_Time       = 0[us]
Decoupling      = TRUE
Delay           = 0[us]
Initial_Wait     = 1[s]
Noe              = TRUE
Noe_Time        = 1.5[s]
Repetition_Time  = 2.54333312[s]

```

```

Experiment      = single pulse.jxp
Sample_id       = YO-5-153
Solvent         = CHLOROFORM-D
Creation Time   = 29-JUL-2015 13:45:50

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration = 2.18365952[s]
X_Domain       = 1H
X_Freq         = 399.78219838[MHz]
X_Offset       = 5[ppm]
X_Points       = 16384
X_Prescans     = 1
X_Resolution   = 0.45794685[Hz]
X_Sweep        = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain     = Proton
Irr_Freq       = 399.78219838[MHz]
Irr_Offset     = 5[ppm]
Tri_Domain     = Proton
Tri_Freq       = 399.78219838[MHz]
Tri_Offset     = 5[ppm]
Clipped        = FALSE
Mod_Return     = 1
Probe_Recovery = 5[us]
Scans          = 8
Total_Scans    = 8

Relaxation_Delay = 5[s]
Recvr_Gain       = 30
Temp_Get         = 26[dc]
X_90_Width      = 9.8[us]
X_Acq_Time      = 2.18365952[s]
X_Angle         = 45[deg]
X_Atn           = 1[db]
X_Pulse         = 4.9[us]
Irr_Mode        = Off
Tri_Mode        = Off
Dante_Presat    = FALSE
Dead_Time       = 0[us]
Delay           = 0[us]
Initial_Wait    = 1[s]
Repetition_Time = 7.18365952[s]

```

```

Experiment      = single pulse dec.jxp
Sample_id       = YO-5-153 overNight
Solvent         = CHLOROFORM-D
Creation Time   = 31-JUL-2015 00:02:57

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration = 1.04333312[s]
X_Domain       = 13C
X_Freq         = 100.52530333[MHz]
X_Offset       = 100[ppm]
X_Points       = 32768
X_Prescans     = 4
X_Resolution   = 0.95846665[Hz]
X_Sweep        = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain     = Proton
Irr_Freq       = 399.78219838[MHz]
Irr_Offset     = 5[ppm]
Clipped        = FALSE
Mod_Return     = 1
Probe_Recovery = 20[us]
Scans          = 4096
Total_Scans    = 4096

Relaxation_Delay = 1.5[s]
Recvr_Gain       = 50
Temp_Get         = 26[dc]
X_90_Width      = 8.6[us]
X_Acq_Time      = 1.04333312[s]
X_Angle         = 30[deg]
X_Atn           = 4[db]
X_Pulse         = 2.86666667[us]
Irr_Atn_Dec     = 22.569[db]
Irr_Atn_Noise  = 22.569[db]
Irr_Noise      = WALTZ
Irr_Pwidth     = 0.115[ms]
Dead_Time       = 0[us]
Decoupling      = TRUE
Delay           = 0[us]
Initial_Wait    = 1[s]
Noe             = TRUE
Noe_Time        = 1.5[s]
Repetition_Time = 2.54333312[s]

```

Experiment = single pulse.jxp
 Sample_id = YO-5-154
 Solvent = CHLOROFORM-D
 Creation Time = 30-JUL-2015 20:19:45
 Field Strength = 9.389766[T] (400[MHz])
 X_Acq_Duration = 2.18365952[s]
 X_Domain = 1H
 X_Freq = 399.78219838[MHz]
 X_Offset = 5[ppm]
 X_Points = 16384
 X_Prescans = 1
 X_Resolution = 0.45794665[Hz]
 X_Sweep = 7.5030012[kHz]
 X_Sweep_Clippped = 6.00240096[kHz]
 Irr_Domain = Proton
 Irr_Freq = 399.78219838[MHz]
 Irr_Offset = 5[ppm]
 Tri_Domain = Proton
 Tri_Freq = 399.78219838[MHz]
 Tri_Offset = 5[ppm]
 Clipped = FALSE
 Mod_Return = 1
 Probe_Recovery = 5[us]
 Scans = 8
 Total_Scans = 8
 Relaxation_Delay = 5[s]
 Recvr_Gain = 30
 Temp_Get = 25.9[dc]
 X_90_Width = 9.8[us]
 X_Acq_Time = 2.18365952[s]
 X_Angle = 45[deg]
 X_Atn = 1[db]
 X_Pulse = 4.9[us]
 Irr_Mode = Off
 Tri_Mode = Off
 Dante_Presat = FALSE
 Dead_Time = 0[us]
 Delay = 0[us]
 Initial_Wait = 1[s]
 Repetition_Time = 7.18365952[s]

Experiment = single pulse dec.jxp
 Sample_id = YO-5-154 overNight
 Solvent = CHLOROFORM-D
 Creation Time = 1-AUG-2015 00:03:38
 Field Strength = 9.389766[T] (400[MHz])
 X_Acq_Duration = 1.04333312[s]
 X_Domain = 13C
 X_Freq = 100.52530333[MHz]
 X_Offset = 100[ppm]
 X_Points = 32768
 X_Prescans = 4
 X_Resolution = 0.95846665[Hz]
 X_Sweep = 31.40703518[kHz]
 X_Sweep_Clippped = 25.12562814[kHz]
 Irr_Domain = Proton
 Irr_Freq = 399.78219838[MHz]
 Irr_Offset = 5[ppm]
 Clipped = FALSE
 Mod_Return = 1
 Probe_Recovery = 20[us]
 Scans = 4096
 Total_Scans = 4096
 Relaxation_Delay = 1.5[s]
 Recvr_Gain = 50
 Temp_Get = 26[dc]
 X_90_Width = 8.6[us]
 X_Acq_Time = 1.04333312[s]
 X_Angle = 30[deg]
 X_Atn = 4[db]
 X_Pulse = 2.86666667[us]
 Irr_Atn_Dec = 22.569[db]
 Irr_Atn_No = 22.569[db]
 Irr_Noise = WALTZ
 Irr_Pwidth = 0.115[ms]
 Dead_Time = 0[us]
 Decoupling = TRUE
 Delay = 0[us]
 Initial_Wait = 1[s]
 Noe = TRUE
 Noe_Time = 1.5[s]
 Repetition_Time = 2.54333312[s]

Experiment = single pulse.jxp
Sample_id = YO-5-156
Solvent = CHLOROFORM-D
Creation Time = 31-JUL-2015 16:42:26

Field Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 2.18365952[s]
X_Domain = 1H
X_Freq = 399.78219838[MHz]
X_Offset = 5[ppm]
X_Points = 16384
X_Prescans = 1
X_Resolution = 0.45794685[Hz]
X_Sweep = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Tri_Domain = Proton
Tri_Freq = 399.78219838[MHz]
Tri_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 5[us]
Scans = 8
Total_Scans = 8

Relaxation_Delay = 5[s]
Recvr_Gain = 32
Temp_Get = 26.2[dC]
X_90_Width = 9.8[us]
X_Acq_Time = 2.18365952[s]
X_Angle = 45[deg]
X_Atn = 1[dB]
X_Pulse = 4.9[us]
Irr_Mode = Off
Tri_Mode = Off
Dante_Presat = FALSE
Dead_Time = 0[us]
Delay = 0[us]
Initial_Wait = 1[s]
Repetition_Time = 7.18365952[s]

Experiment = single pulse dec.jxp
Sample_id = YO-5-156 overNight
Solvent = CHLOROFORM-D
Creation Time = 2-AUG-2015 00:02:58

Field Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 1.04333312[s]
X_Domain = 13C
X_Freq = 100.52530333[MHz]
X_Offset = 100[ppm]
X_Points = 32768
X_Prescans = 4
X_Resolution = 0.95846665[Hz]
X_Sweep = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 20[us]
Scans = 4096
Total_Scans = 4096

Relaxation_Delay = 1.5[s]
Recvr_Gain = 50
Temp_Get = 26.1[dC]
X_90_Width = 8.6[us]
X_Acq_Time = 1.04333312[s]
X_Angle = 30[deg]
X_Atn = 4[dB]
X_Pulse = 2.86666667[us]
Irr_Atn_Dec = 22.569[dB]
Irr_Atn_No2 = 22.569[dB]
Irr_Noise = WALTZ
Irr_Pwidth = 0.115[ms]
Dead_Time = 0[us]
Decoupling = TRUE
Delay = 0[us]
Initial_Wait = 1[s]
Noe = TRUE
Noe_Time = 1.5[s]
Repetition_Time = 2.54333312[s]

```

Experiment      = single pulse.jxp
Sample_id       = YO-5-160
Solvent         = CHLOROFORM-D
Creation Time   = 3-AUG-2015 19:12:45

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 2.18365952[s]
X_Domain        = 1H
X_Freq          = 399.78219838[MHz]
X_Offset        = 5[ppm]
X_Points        = 16384
X_Prescans      = 1
X_Resolution    = 0.45794665[Hz]
X_Sweep         = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Tri_Domain      = Proton
Tri_Freq        = 399.78219838[MHz]
Tri_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 5[us]
Scans           = 8
Total_Scans     = 8

Relaxation_Delay = 5[s]
Recvr_Gain       = 30
Temp_Get         = 26.1[dc]
X_90_Width      = 9.8[us]
X_Acq_Time       = 2.18365952[s]
X_Angle         = 45[deg]
X_Atn           = 1[db]
X_Pulse         = 4.9[us]
Irr_Mode        = Off
Tri_Mode        = Off
Dante_Presat    = FALSE
Dead_Time       = 0[us]
Delay           = 0[us]
Initial_Wait    = 1[s]
Repetition_Time = 7.18365952[s]

```

53

```

Experiment      = single pulse dec.jxp
Sample_id       = YO-5-160 overnight
Solvent         = CHLOROFORM-D
Creation Time   = 6-AUG-2015 00:04:10

Field Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 1.04333312[s]
X_Domain        = 13C
X_Freq          = 100.52530333[MHz]
X_Offset        = 100[ppm]
X_Points        = 32768
X_Prescans      = 4
X_Resolution    = 0.95846665[Hz]
X_Sweep         = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 20[us]
Scans           = 4096
Total_Scans     = 4096

Relaxation_Delay = 1.5[s]
Recvr_Gain       = 50
Temp_Get         = 26.1[dc]
X_90_Width      = 8.6[us]
X_Acq_Time       = 1.04333312[s]
X_Angle         = 30[deg]
X_Atn           = 4[db]
X_Pulse         = 2.866666667[us]
Irr_Atn_Dec     = 22.569[db]
Irr_Atn_Noise   = 22.569[db]
Irr_Noise       = WALTZ
Irr_Pwidth      = 0.115[ms]
Dead_Time       = 0[us]
Decoupling      = TRUE
Delay           = 0[us]
Initial_Wait    = 1[s]
Noe              = TRUE
Noe_Time        = 1.5[s]
Repetition_Time = 2.54333312[s]

```

53

Experiment = single pulse.jxp
Sample_id = YO-5-172
Solvent = CHLOROFORM-D
Creation Time = 19-AUG-2015 13:01:20
Field Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 2.18365952[s]
X_Domain = 1H
X_Freq = 399.78219838[MHz]
X_Offset = 5[ppm]
X_Points = 16384
X_Prescans = 1
X_Resolution = 0.45794685[Hz]
X_Sweep = 7.5030012[kHz]
X_Sweep_Clippped = 6.00240096[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Tri_Domain = Proton
Tri_Freq = 399.78219838[MHz]
Tri_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 5[us]
Scans = 8
Total_Scans = 8
Relaxation_Delay = 5[s]
Recvr_Gain = 34
Temp_Get = 25.6[dC]
X_90_Width = 9.8[us]
X_Acq_Time = 2.18365952[s]
X_Angle = 45[deg]
X_Atn = 1[dB]
X_Pulse = 4.9[us]
Irr_Mode = Off
Tri_Mode = Off
Dante_Presat = FALSE
Dead Time = 0[us]
Delay = 0[us]
Initial_Wait = 1[s]
Repetition_Time = 7.18365952[s]

Experiment = single pulse_dec.jxp
Sample_id = YO-5-172 overNight
Solvent = CHLOROFORM-D
Creation Time = 20-AUG-2015 00:02:18
Field Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 1.04333312[s]
X_Domain = 13C
X_Freq = 100.52530333[MHz]
X_Offset = 100[ppm]
X_Points = 32768
X_Prescans = 4
X_Resolution = 0.95846665[Hz]
X_Sweep = 31.40703518[kHz]
X_Sweep_Clippped = 25.12562814[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 20[us]
Scans = 4096
Total_Scans = 4096
Relaxation_Delay = 1.5[s]
Recvr_Gain = 50
Temp_Get = 25.7[dC]
X_90_Width = 8.6[us]
X_Acq_Time = 1.04333312[s]
X_Angle = 30[deg]
X_Atn = 4[dB]
X_Pulse = 2.86666667[us]
Irr_Atn_Dec = 22.569[dB]
Irr_Atn_Noe = 22.569[dB]
Irr_Noise = WALTZ
Irr_Pwidth = 0.115[ms]
Dead Time = 0[us]
Decoupling = TRUE
Delay = 0[us]
Initial_Wait = 1[s]
Noe = TRUE
Noe Time = 1.5[s]
Repetition_Time = 2.54333312[s]

Experiment = single pulse.jxp
Sample_id = YO-5-189
Solvent = CHLOROFORM-D
Creation_Time = 24-AUG-2015 19:02:00
Field_Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 2.18365952[s]
X_Domain = 1H
X_Freq = 399.78219838[MHz]
X_Offset = 5[ppm]
X_Points = 16384
X_Prescans = 1
X_Resolution = 0.45794685[Hz]
X_Sweep = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Tri_Domain = Proton
Tri_Freq = 399.78219838[MHz]
Tri_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 5[us]
Scans = 8
Total_Scans = 8
Relaxation_Delay = 5[s]
Recvr_Gain = 36
Temp_Get = 25.6[dC]
X_90_Width = 9.8[us]
X_Acq_Time = 2.18365952[s]
X_Angle = 45[deg]
X_Atn = 1[dB]
X_Pulse = 4.9[us]
Irr_Mode = Off
Tri_Mode = Off
Dante_Presat = FALSE
Dead_Time = 0[us]
Delay = 0[us]
Initial_Wait = 1[s]
Repetition_Time = 7.18365952[s]

35

Experiment = single pulse dec.jxp
Sample_id = YO-5-189 overNight
Solvent = CHLOROFORM-D
Creation_Time = 25-AUG-2015 00:03:02
Field_Strength = 9.389766[T] (400[MHz])
X_Acq_Duration = 1.04333312[s]
X_Domain = 13C
X_Freq = 100.52530333[MHz]
X_Offset = 100[ppm]
X_Points = 32768
X_Prescans = 4
X_Resolution = 0.95846665[Hz]
X_Sweep = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain = Proton
Irr_Freq = 399.78219838[MHz]
Irr_Offset = 5[ppm]
Clipped = FALSE
Mod_Return = 1
Probe_Recovery = 20[us]
Scans = 4096
Total_Scans = 4096
Relaxation_Delay = 1.5[s]
Recvr_Gain = 50
Temp_Get = 25.5[dC]
X_90_Width = 8.6[us]
X_Acq_Time = 1.04333312[s]
X_Angle = 30[deg]
X_Atn = 4[dB]
X_Pulse = 2.86666667[us]
Irr_Atn_Dec = 22.569[dB]
Irr_Atn_No2 = 22.569[dB]
Irr_Noise = WALTZ
Irr_Pwidth = 0.115[ms]
Dead_Time = 0[us]
Decoupling = TRUE
Delay = 0[us]
Initial_Wait = 1[s]
Noe = TRUE
Noe_Time = 1.5[s]
Repetition_Time = 2.54333312[s]

35

```

Experiment      = single pulse.jxp
Sample_id       = YO-5-19b py
Solvent         = PYRIDINE-D5
Creation_Time    = 25-AUG-2015 16:36:50

Field_Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 2.18365952[s]
X_Domain        = 1H
X_Freq          = 399.78219838[MHz]
X_Offset        = 5[ppm]
X_Points        = 16384
X_Prescans      = 1
X_Resolution    = 0.457946685[Hz]
X_Sweep         = 7.5030012[kHz]
X_Sweep_Clipped = 6.00240096[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Tri_Domain      = Proton
Tri_Freq        = 399.78219838[MHz]
Tri_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 5[us]
Scans           = 8
Total_Scans     = 8

Relaxation_Delay = 5[s]
Recvr_Gain       = 38
Temp_Get         = 25.3[dC]
X_90_Width      = 9.8[us]
X_Acq_Time       = 2.18365952[s]
X_Angle         = 45[deg]
X_Atn           = 1[db]
X_Pulse         = 4.9[us]
Irr_Mode        = Off
Tri_Mode        = Off
Dante_Presat    = FALSE
Dead_Time       = 0[us]
Delay           = 0[us]
Initial_Wait    = 1[s]
Repetition_Time  = 7.18365952[s]
  
```

```

Experiment      = single pulse_dec.jxp
Sample_id       = YO-5-19b py Overnight
Solvent         = PYRIDINE-D5
Creation_Time    = 28-AUG-2015 22:37:11

Field_Strength  = 9.389766[T] (400[MHz])
X_Acq_Duration  = 1.04333312[s]
X_Domain        = 13C
X_Freq          = 100.52530333[MHz]
X_Offset        = 100[ppm]
X_Points        = 32768
X_Prescans      = 4
X_Resolution    = 0.95846665[Hz]
X_Sweep         = 31.40703518[kHz]
X_Sweep_Clipped = 25.12562814[kHz]
Irr_Domain      = Proton
Irr_Freq        = 399.78219838[MHz]
Irr_Offset      = 5[ppm]
Clipped         = FALSE
Mod_Return      = 1
Probe_Recovery  = 20[us]
Scans           = 8192
Total_Scans     = 8192

Relaxation_Delay = 1.5[s]
Recvr_Gain       = 50
Temp_Get         = 25.3[dC]
X_90_Width      = 8.6[us]
X_Acq_Time       = 1.04333312[s]
X_Angle         = 30[deg]
X_Atn           = 4[db]
X_Pulse         = 2.86666667[us]
Irr_Atn_Dec     = 22.569[db]
Irr_Atn_Noise   = 22.569[db]
Irr_Noise       = WALTZ
Irr_Pwidth      = 0.115[ms]
Dead_Time       = 0[us]
Decoupling      = TRUE
Delay           = 0[us]
Initial_Wait    = 1[s]
Noe              = TRUE
Noe_Time        = 1.5[s]
Repetition_Time  = 2.54333312[s]
  
```