Total Synthesis of Topopyrones B and D

Jason Tan, Marco A. Ciufolini*

Department of Chemistry, University of British Columbia, 2036 Main Mall, Vancouver, BC V6T 1Z1, Canada

Supporting Information

Table of Contents

<i>Subject</i>	<i>page</i>
Experimental Protocols	3
Synthesis of intermediates 6, 7, 8, 22, 24	3
preparation of 2-carbethoxymethyl-2-methyl-dioxolane	3
preparation of 2-(2-hydroxyethyl)-2-methyl-dioxolane	3
preparation of aldehyde 10	3
preparation of benzamide 6	4
preparation of 3-chloro-2,4-dihydroxybenzaldehyde	4
preparation of 3-chloro-2,4-dimethoxybenzaldehyde	4
preparation of 3-chloro-2,4-dimethoxybenzoic acid	4
preparation of benzamide 7	5
preparation of 4-bromo-3,5-dimethoxybenzyl alcohol	5
preparation of TBS ether 9	5
preparation of compound 11	5
preparation of compound 12	6
preparation of compound 13	6
preparation of compound 14	6
preparation of aldehyde 8	7
preparation of aldehyde 21	7
preparation of anthraquinone 23	7
Synthesis of topopyrone D	7
preparation of anthraquinone 17	7
deprotection of anthraquinone 17	8
preparation of anthraquinone 24	8
topopyrone D 4	8
topopyrone D triacetate 26	9
Synthesis of topopyrone B	9
preparation of anthraquinone 18	9
deprotection of anthraquinone 18	9
preparation of anthraquinone 24	10
topopyrone B trimethyl ether 27	10
Index of hardcopy ¹ H and ¹³ C NMR spectra	11
spectra	12

Experimental Protocols.

Unless otherwise stated, proton and ^{13}C NMR spectra were obtained from CDCl_3 solutions. Chemical shifts are reported in parts per million (ppm) on the δ scale and coupling constants, J , are in hertz (Hz). Multiplicities are reported as “s” (singlet), “d” (doublet), “t” (triplet), “q” (quartet), “dd” (doublet of doublets), “td” (triplet of doublets), “m” (multiplet), “c” (complex), “br” (broad). FT-IR spectra (cm^{-1}) were from thin films deposited on NaCl plates. Low- and high-resolution mass spectra (m/z) were obtained in the electrospray (ESI) mode. Melting points are uncorrected. All reagents and solvents were commercial products and used without further purification except THF (freshly distilled from Na/benzophenone under Ar) and CH_2Cl_2 (freshly distilled from CaH_2 under Ar). Commercial $n\text{-BuLi}$ was titrated against N -benzylbenzamide in THF at -78°C until persistence of a light blue color. Flash chromatography was performed on Silicycle 230 – 400 mesh silica gel. All reactions were performed under dry Ar in flame or over dried flasks equipped with TeflonTM stirbars. All flasks were fitted with rubber septa for the introduction of substrates, reagents, and solvents via syringe.

Synthesis of intermediates 6, 7, 8, 22, 24.

2-carbethoxymethyl-2-methyl-dioxolane. A solution of ethyl acetoacetate (100.0 g, 0.77 mol), ethylene glycol (143.0 g, 2.30 mol), and $p\text{-TsOH}\cdot\text{H}_2\text{O}$ (6.5 g, 34.2 mmol) in benzene (1000 mL; **CAUTION:** cancer suspect agent) was refluxed for 5 h with continuous azeotropic H_2O separation (Dean-Stark

trap). The mixture was then cooled to RT, concentrated to 500 mL *in vacuo*, washed sequentially with sat. aq. NaHCO_3 (2 x 100 mL) and brine (100 mL), dried (Na_2SO_4), filtered and concentrated to afford the title ketal ester (120.5 g, 90%) as a colorless oil. This material was of excellent quality and required no further purification. ^1H : 4.06 (q, 2H, $J = 6.6$), 3.88 (s, 4H), 2.57 (s, 2H), 1.41 (s, 3H), 1.17 (t, 3H, $J = 6.6$). ^{13}C : 169.2, 107.4, 64.5, 60.2, 44.0, 24.2, 13.9. **IR**: 1747. **HRMS** calcd for $\text{C}_8\text{H}_{14}\text{O}_4$ [$\text{M} + \text{Na}$] $^+$ = 197.0790, found 197.0791.

2-(2-hydroxyethyl)-2-methyl-dioxolane. A solution of the above ketal ester (9.0 g, 51.7 mmol) in dry THF (20 mL) was added dropwise to a cold (-5°C ; ice-salt bath) suspension of LAH (1.97 g, 51.7 mmol) in dry THF (80 mL). The reaction mixture was stirred at 0°C for 1.5 h then quenched with aq. sat.

Na / K tartrate soln. (100 mL. **CAUTION:** vigorous evolution of highly flammable H_2 gas). The organic layer was separated and the aqueous layer was extracted with Et_2O (2 x 50 mL). The combined organic layers were washed with brine (100 mL), dried (Na_2SO_4) and concentrated to give pure title alcohol (6.2 g, 91%) as a colorless oil. ^1H : 3.99 (s, 4H), 3.76 (broad s, 2H), 2.76 (broad s, 1H), 1.95 (t, 2H, $J = 5.2$), 1.36 (s, 3H). ^{13}C : 110.3, 64.6, 58.8, 40.5, 23.9; **IR**: 3440. **HRMS** calcd for $\text{C}_6\text{H}_{12}\text{O}_3$ [$\text{M} + \text{Na}$] $^+$ = 155.0684, found 155.0683.

Aldehyde 10 (cf. Langer, P.; Freifeld, I. *Synlett*. **2001**, 4, 523). Dry DMSO (2.79 mL, 39.3 mmol) was added dropwise to a cold (-78°C) solution of oxalyl chloride (3.38 mL, 39.3 mmol) in dry DCM (130 mL; **CAUTION:** a vigorous exotherm ensues that produces poisonous CO gas). The mixture was stirred at -78°C for 15 min, then solution of the above alcohol (4.0 g, 30.3 mmol) in dry DCM (20 mL) was added dropwise. The resultant solution was stirred at -78°C for 30 min, then Et_3N (16.9 mL, 121 mmol) was added. The mixture was stirred at -78°C for 5 min then it was allowed to warm to RT over 40 min, with continued stirring. The reaction mixture was washed successively with aq. sat. NH_4Cl (3 x 50 mL), dried (Na_2SO_4) and concentrated to

give pure **10** (3.6 g, 91%) as a pale yellow oil. ^1H : 9.75 (t, 1H, $J = 2.8$), 4.01 (m, 4H), 2.71 (d, 2H, $J = 2.8$), 1.43 (s, 3H). ^{13}C : 200.3, 107.6, 64.8, 52.2, 24.9. IR: 1724. HRMS calcd for $\text{C}_6\text{H}_{10}\text{O}_3$ $[\text{M} + \text{Na}]^+ = 153.0528$, found 153.0527.

***N,N*-diethyl-2,4-dimethoxybenzamide 6** (cf. Kamila, S.; Mukherjee, C.; Mondal, S.; De, A. *Tetrahedron*. **2003**, 59, 1339). Diethylamine (7.12 mL, 68.5 mmol) was carefully added to a cold (0 °C) solution of 2,4-dimethoxybenzoyl chloride (4.58 g, 22.8 mmol) in dry toluene (70 mL), and the mixture was stirred at 0 °C for 1 h then warmed to RT and stirred

for another 5 h. The solution was concentrated and the residue was partitioned between ethyl acetate (50 mL) and aq. sat. NaHCO_3 (50 mL). The organic layer was washed successively with 1M HCl (50 mL), H_2O (50 mL), and brine (50 mL), dried (Na_2SO_4) and concentrated to give pure amide **6** (4.8 g, 89%) as a pale yellow oil. ^1H : 7.05 (d, 1H, $J = 8.4$), 6.43 (dd, 1H, $J = 8.4, 2.0$), 6.39 (d, 1H, $J = 2.0$), 3.74 (s, 3H), 3.73 (s, 3H), 3.49 (broad q, 2H), 3.09 (q, 2H, $J = 7.2$), 1.16 (t, 3H, $J = 7.6$), 0.97 (t, 3H, $J = 7.2$). ^{13}C : 168.6, 161.0, 156.3, 128.1, 119.6, 104.4, 98.4, 55.3, 55.2, 42.6, 38.7, 13.8, 12.7. IR: intense band with fine structure between 1680 and 1615. HRMS calcd for $\text{C}_{13}\text{H}_{18}^{35}\text{ClNO}_3$ $[\text{M} + \text{Na}]^+ = 294.0873$, found 294.0870.

3-Chloro-2,4-dihydroxybenzaldehyde (cf. Bui, E.; Bayle, J. P.; Perez, F.; Liebert, L.; Courtieu, J. *Liq. Cryst.* **1990**, 8, 513). A suspension of 2,4-dihydroxybenzaldehyde (2.0 g, 14.5 mmol, Fluka) in H_2O (10 mL) was sequentially treated at RT with a solution of KOH (2.0 g, 35.6 mmol) in H_2O (15 mL), followed by commercial bleach containing 6% NaOCl (21

mL, 16 mmol). The reaction mixture was stirred at RT for 3 h then it was acidified with 6M HCl (20 mL) and extracted with EtOAc (2 x 50 mL). The combined organic extracts were dried (Na_2SO_4) and concentrated. Purification by flash chromatography (EtOAc:hexanes, 1:1) gave pure aldehyde 3-chloro-2,4-dihydroxybenzaldehyde (2.4 g, quantitative) as a white solid, m.p. 146-148 °C (no m.p. was recorded in the literature). ^1H (acetone- d_6): 9.80 (s, 1H), 7.60 (d, 1H, $J = 8.8$), 6.75 (d, 1H, $J = 8.8$). ^{13}C : 197.1, 162.5, 161.6, 135.5, 117.1, 110.5, 109.1. IR: 1634. HRMS calcd for $\text{C}_7\text{H}_5^{35}\text{ClO}_3$ $[\text{M} + \text{H}]^+ = 173.0005$, found 173.0007.

3-Chloro-2,4-dimethoxybenzaldehyde. Neat MeI (7.9mL, 127 mmol. **CAUTION:** toxic, volatile, cancer suspect agent) was added at RT to a solution of the above aldehyde (2.2 g, 12.7 mmol) in acetone (50 mL) containing suspended K_2CO_3 (17.6 g, 127 mmol). The mixture was refluxed for 5 h, then it was filtered and concentrated, and the residue was

purified by flash chromatography (EtOAc : hexanes, 1:4) to give pure 3-chloro-2,4-dimethoxybenzaldehyde (2.2 g, 85%) as a white solid, m.p. 105-107 °C (lit. 109-111 °C: Plattner, J. J.; Fung, A. K. L.; Parks, J. A.; Pariza, R. J.; Crowley, S. R.; Pernet, A. G.; Bunnell, P. R.; Dodge, P. W. *J. Med. Chem.* **1984**, 27, 1016). ^1H : 10.19 (s, 1H), 7.75 (d, 1H, $J = 9.1$), 6.81 (d, 1H, $J = 9.1$), 3.96 (s, 3H), 3.95 (s, 3H). ^{13}C : 187.9, 161.1, 160.5, 127.8, 123.8, 116.6, 107.7, 62.9, 56.6. IR: 1681. HRMS calcd for $\text{C}_9\text{H}_9^{35}\text{ClO}_3$ $[\text{M} + \text{H}]^+ = 201.0318$, found 201.0317;

3-Chloro-2,4-dimethoxybenzoic acid. A solution of NaClO_2 (80%, 1.4 g, 12.3 mmol) in a NaH_2PO_4 buffer (pH 3.5, 50 mL) was carefully added to a solution of the above aldehyde (2.06 g, 10.3 mmol) in *t*-butanol (50 mL) containing 2-methyl-2-butene (2.0M in THF, 6.2 mL, 12.4 mmol). The mixture was stirred at RT overnight, then it was basified with 1M NaOH

(10 mL) and the aqueous layer was washed with EtOAc (2 x 25 mL). The aqueous layer was reacidified with 6M HCl (5 mL) and extracted with EtOAc (2 x 40 mL). The combined extracts were dried (Na₂SO₄) and concentrated to give pure title acid (2.0 g, 90%) as a white solid, m.p. 168-170 °C. ¹H: 8.07 (d, 1H, *J* = 9.1), 6.88 (d, 1H, *J* = 9.1), 4.08 (s, 3H), 3.99 (s, 3H). ¹³C (acetone-*d*₆): 165.8, 160.4, 158.5, 131.9, 119.0, 118.1, 108.2, 62.0, 57.0. **IR**: 1695. **HRMS** calcd for C₉H₉³⁵ClO₄ [M + Na]⁺ = 239.0087, found 239.0089.

***N,N*-diethyl-3-chloro-2,4-dimethoxybenzamide 7.** Thionyl chloride (1.0 mL, 13.8 mmol; **CAUTION**: corrosive, toxic) was carefully added to a solution of the above acid (2.0 g, 9.2 mmol) in dry benzene (65 mL; **CAUTION**: cancer suspect agent) and the mixture was refluxed for 3 h

(**CAUTION**: evolution of corrosive HCl gas). The mixture was then cooled to 0 °C prior to dropwise addition of a solution of diethylamine (7.65 mL, 73.6 mmol) in dry benzene (10 mL **CAUTION**: cancer suspect agent). The mixture was stirred overnight at RT, then it was washed with 1M aq. NaOH (2 x 50 mL), dried (Na₂SO₄) and concentrated. The residue of crude amide was recrystallized (EtOAc:hexanes, 1:1) to afford pure **7** (1.9 g, 76%) as tan crystals, m.p. 70-72 °C. ¹H: 7.11 (d, 1H, *J* = 8.7), 6.73 (d, 1H, *J* = 8.7), 3.91 (s, 3H), 3.87 (s, 3H), 3.74, 3.36 (br s, 2H), 3.15 (q, 2H, *J* = 7.3), 1.24 (t, 3H, *J* = 7.3), 1.02 (t, 3H, *J* = 7.3). ¹³C: 167.8, 156.8, 153.4, 125.8, 125.6, 116.8, 107.7, 61.9, 56.6, 43.2, 39.2, 14.1, 12.9. **IR**: 1634. **HRMS** calcd for C₁₃H₁₈³⁵ClNO₃ [M + Na]⁺ = 294.0873, found 294.0870.

4-Bromo-3,5-dimethoxybenzyl alcohol. Commercial BH₃•SMe₂ solution (36.3 mL, 0.38 mol) was added dropwise to a solution of 3,5-dimethoxy-4-bromobenzoic acid (50.0 g, 0.19 mol) in dry THF (600 mL). The mixture was stirred at 40 °C overnight, then it was quenched with 1M aq. HCl (200 mL. **CAUTION**: evolution of highly flammable H₂ gas) and the

volatiles were removed *in vacuo*. The aqueous residue was extracted with EtOAc (3 x 200 mL), and the combined extracts were dried (Na₂SO₄) and concentrated to give pure title alcohol (47 g, quantitative) as a white solid, m.p. 100 – 102 °C (lit. 100-102 °C: Inoue, I. WO 2003072536 A1 2003). ¹H: 6.51 (s, 2H), 4.59 (s, 2H), 3.84 (s, 6H), 2.41 (s, 1H). ¹³C: 156.8, 141.7, 102.8, 99.3, 64.8, 56.3. **IR**: 3310. **HRMS** calcd for C₉H₁₁⁷⁹BrO₃ [M + Na]⁺ = 268.9789, found 268.9797.

TBS ether 9. A solution of the above alcohol (47.0 g, 0.19 mol), imidazole (25.9 g, 0.38 mol), 4-DMAP (1.8 g, 15 mmol), and TBSCl (30.1 g, 0.20 mol) in dry CH₂Cl₂ (400 mL) was stirred at RT overnight, then it was concentrated. The residue was taken up with sat. NH₄Cl aq. (200 mL) and extracted with EtOAc (3 x 150 mL). The combined extracts

were dried (Na₂SO₄) and concentrated, and the residue was purified by flash chromatography (hexanes → EtOAc:hexanes, 1:1) to afford silyl ether **9** (68 g, quantitative) as a waxy solid, m.p. 34 – 36 °C. ¹H: 6.57 (s, 2H), 4.71 (s, 2H), 3.89 (s, 6H), 0.96 (s, 9H), 0.11 (s, 6H). ¹³C: 157.1, 142.7, 102.3, 64.8, 56.5, 26.1, 18.6, -5.1. **IR**: 2856, 1589, 1233. **HRMS** calcd for C₁₅H₂₅⁷⁹BrO₃Si [M + Na]⁺ = 383.0654, found 383.0655.

Compound 11. A pentane solution of *t*-BuLi (1.45M 10.5 mL, 15.2 mmol) was added dropwise to a cold (-78 °C) solution of **9** (2.5 g, 6.9 mmol) in dry THF (25 mL), followed by a solution of aldehyde **10** (0.90 g, 6.9 mmol) in dry THF (5 mL). The mixture was stirred at -78 °C for 1 h then it was allowed to warm to RT overnight. The reaction was quenched

with sat. NH_4Cl aq. (30 mL) and the aqueous layer was extracted with EtOAc (2 x 30 mL). The combined extracts were dried (Na_2SO_4) and concentrated. Chromatographic purification of the residue (EtOAc:hexanes, 1:1) gave **11** (1.9 g, 67%) as a pale orange oil. ^1H : 6.53 (s, 2H), 5.43 (ddd, 1H, $J = 9.6, 9.6, 3.9$), 4.70 (s, 2H), 3.97 (m, 4H), 3.82 (s, 6H), 3.54 (d, 1H, $J = 9.6$), 2.43 (dd, 1H, $J = 15.0, 9.6$), 1.97 (dd, 1H, $J = 15.0, 3.9$), 1.45 (s, 3H), 0.95 (s, 9H), 0.10 (s, 6H). ^{13}C : 157.7, 142.4, 118.8, 109.9, 101.9, 65.1, 64.7, 64.3, 63.8, 55.8, 45.2, 26.3, 24.3, 18.5, -4.8. **IR**: 3565. **HRMS** calcd for $\text{C}_{21}\text{H}_{36}\text{O}_6\text{Si}$ $[\text{M} + \text{Na}]^+ = 435.2179$, found 435.2178.

Compound 12. Neat TIPS-OTf (1.85 mL, 6.9 mmol) was carefully added to a solution of **11** (2.7 g, 6.6 mmol), imidazole (0.9 g, 13.2 mmol), and 4-DMAP (64.0 mg, 0.53 mmol) in dry DMF (13 mL), and the mixture was stirred at RT overnight. The reaction was quenched with aq. sat. NaHCO_3 (50 mL) and extracted with EtOAc (2 x 50 mL). The combined extracts were washed with brine (50 mL), dried (Na_2SO_4) and concentrated. The residue was purified by flash chromatography (hexanes $\rightarrow$ EtOAc:hexanes, 1:4) to afford silyl ether **12** (3.5 g, 94%) as colorless oil. ^1H : 6.50 (br s, 1H), 6.46 (br s, 1H), 5.57 (dd, 1H, $J = 9.2, 4.4$), 4.70 (br s, 2H), 3.85-3.78 (c, 4H), 3.78 (s, 3H), 3.77 (s, 3H), 2.76 (dd, 1H, $J = 14.4, 9.2$), 2.23 (dd, 1H, $J = 14.4, 4.4$), 1.24 (s, 3H), 1.00-0.98 (m, 12H), 0.94 (s, 9H), 0.89-0.85 (m, 9H), 0.08 (s, 6H). ^{13}C : 160.1, 157.0, 142.1, 120.0, 109.5, 102.8, 101.1, 65.3, 64.4, 64.3, 63.1, 55.6, 55.6, 45.1, 26.0, 24.5, 18.2, 18.0, 17.9, 12.5, -5.0. **IR**: 2865, 1610, 1228. **HRMS** calcd for $\text{C}_{30}\text{H}_{56}\text{O}_6\text{Si}_2$ $[\text{M} + \text{Na}]^+ = 591.3513$, found 591.3518.

Compound 13. A solution of **12** (3.5 g, 6.2 mmol) and NBS (1.1 g, 6.2 mmol) in CH_3CN (20 mL) was stirred at RT for 1.5 h, then it was diluted with aq. sat. NaHCO_3 aq. (30 mL) and extracted with EtOAc (2 x 30 mL). The combined extracts were washed with brine (50 mL), dried (Na_2SO_4) and concentrated to give pure **13** (4.0 g, quantitative) as a colorless oil. Proton and ^{13}C NMR spectra indicated that this material exists as a ca. 4 :1 mixture of atropisomers. Major atropisomer: ^1H : 6.94 (s, 1H), 5.57 (dd, 1H, $J = 10.5, 3.6$), 4.70 (br AB-type system, 2H, $J = 15.1$), 3.93 (s, 3H), 3.80 (s, 3H), 3.81-3.69 (c, 4H), 3.14 (dd, 1H, $J = 14.2, 10.5$), 2.12 (dd, 1H, $J = 14.2, 3.6$), 1.18 (s, 3H), 1.11-0.84 (c, 21H), 0.97 (s, 9H), 0.13 (s, 6H). ^{13}C : 157.2, 156.4, 141.1, 126.5, 109.0, 108.2, 105.4, 64.9, 64.5, 64.4, 63.2, 62.1, 55.6, 44.7, 26.1, 24.4, 18.4, 18.2, 12.5, -5.1. **IR**: 2866. **HRMS** calcd for $\text{C}_{30}\text{H}_{55}\text{O}_6^{79}\text{BrSi}_2$ $[\text{M} + \text{Na}]^+ = 669.2618$, found 669.2614.

Compound 14. A solution of **13** (4.0 g, 6.2 mmol) and TBAF (1.0M in THF, 5.9 mL, 6.2 mmol) in THF (20 mL) was stirred at RT for 20 min, then it was quenched with aq. sat. NaHCO_3 (30 mL) and extracted with EtOAc (2 x 30 mL). The combined extracts were washed with brine (50 mL), dried (Na_2SO_4) and concentrated to give **14** (3.2 g, quantitative) as a colorless oil. Proton and ^{13}C NMR spectra indicated that this material exists as a ca. 4:1 mixture of atropisomers. Major atropisomer: ^1H : 6.84 (s, 1H), 5.58 (dd, 1H, $J = 10.5, 3.5$), 4.71 (br AB-type system, 2H, $J = 14.0$), 3.93 (s, 3H), 3.82 (s, 3H), 3.78-3.68 (c, 4H), 3.13 (dd, 1H, $J = 14.0, 10.5$), 2.26 (br, 1H, OH), 2.13 (dd, 1H, $J = 14.0, 3.5$), 1.40-0.80 (m, 21H), 1.17 (s, 3H). ^{13}C : 157.4, 156.4, 140.9, 127.0, 109.4, 108.9, 106.0, 64.3, 64.2, 63.0, 62.0, 55.7, 53.7, 44.5, 28.5, 24.2, 20.9, 18.2, 18.1, 12.4. **IR**: 3430. **HRMS** calcd for $\text{C}_{24}\text{H}_{41}^{79}\text{BrO}_6\text{Si}$ $[\text{M} + \text{Na}]^+ = 555.1753$, found 555.1755.

Aldehyde 8. Dry DMSO (0.35 mL, 4.8 mmol) was cautiously added to a cold (-78 °C) solution of oxalyl chloride (0.42 mL, 4.8 mmol) in dry DCM (15 mL). **CAUTION:** a vigorous exotherm ensues that produces poisonous CO gas). The mixture was stirred at -78 °C for 15 min, then a solution of **14** (2.0 g, 3.7 mmol) in dry DCM (5 mL) was added and the reaction was stirred at -78 °C for 1 h. Triethylamine (2.1 mL, 14.8 mmol) was injected and the mixture was stirred at -78 °C for 5 min, then it was allowed to warm to RT with continued stirring. The reaction mixture was washed successively with aq. sat. NH₄Cl aq. (3 x 50 mL), dried (Na₂SO₄) and concentrated. Chromatography of the residue (EtOAc:hexanes, 1:2) gave pure aldehyde **8** (1.6 g, 81%) as a colorless semisolid. Proton and ¹³C NMR spectra indicated that this material exists as a ca. 6 : 1 mixture of atropisomers. Major atropisomer: ¹H: 10.36 (s, 1H), 7.21 (s, 1H), 5.61 (dd, 1H, *J* = 10.5, 3.2), 3.99 (s, 3H), 3.90-3.60 (cm, 4H), 3.85 (s, 3H), 3.17 (dd, 1H, *J* = 14.2, 10.5), 2.17 (dd, 1H, *J* = 14.2, 3.2), 1.20 (s, 3H), 1.15-0.84 (cm, 21H). ¹³C: 191.9, 158.2, 156.6, 135.4, 133.6, 116.3, 108.7, 106.3, 64.4, 64.3, 63.1, 62.5, 56.0, 44.5, 24.3, 18.2, 18.1, 12.4. **IR:** 1694. **HRMS** calcd for C₂₄H₃₉⁷⁹BrO₆Si [M + Na]⁺ = 553.1597, found 553.1596.

Aldehyde 21 (cf. Broering, T. J.; Morrow, Gary W. *Synth. Commun.* **1999**, 29, 1135). A solution of bromine (3.9 mL, 12.2 g, 76 mmol. **CAUTION:** highly corrosive) in acetic acid (50 mL) was cautiously added over 1 h to a solution of 3,5-dimethoxybenzaldehyde (12.6 g, 76 mmol) in acetic acid (350 mL) at RT. The mixture was stirred overnight, then it was diluted with water (500 mL), whereupon the product precipitated. Filtration afforded 16.4 g (89%) of **21**, as a white solid, m.p. 102-104 °C (lit. 105-106 °C), which required no further purification. ¹H: 10.42 (s, 1H), 7.04 (d, 1H, *J* = 2.7), 6.71 (d, 1H, *J* = 2.7), 3.91 (s, 3H), 3.85 (s, 3H).

Anthraquinone 23. A solution of benzamide **6** (10.3 g, 43 mmol) in dry THF (50 mL) was added to a cold (-78 °C) solution of *s*-BuLi (33 mL, 1.05 eq., 45 mmol) and TMEDA (8.5 mL, 56 mmol) in dry THF (75 mL), followed, after 1 h, by a solution of **21** (11.2 g, 46 mmol) in dry THF (50 mL). The mixture was stirred for 45 min, then *t*-BuLi (58 mL of 1.7M pentane solution, 96 mmol). The reaction was stirred at -78 °C for 20 min and allowed to warm to RT overnight. The mixture was quenched with H₂O (10 mL) and stirred for 1 h while bubbling air through the solution, then it was concentrated, taken up with aq. sat. NaHCO₃ (250 mL) and extracted with CH₂Cl₂ (3 x 200 mL). The combined extracts were dried (Na₂SO₄) and concentrated. The yellow residue was recrystallized from EtOAc to give 5.4 g (40%) of pure **23** as yellow solid, m.p. 216-218 °C (lit. 217-218 °C: Castonguay, A. *Synth. Commun.* **1975**, 5, 377). ¹H: 7.31 (d, 2H, *J* = 2.4), 6.76 (d, 2H, *J* = 2.4), 3.96 (s, 6H), 3.95 (s, 6H).

Synthesis of topopyrone D.

Anthraquinone 17. A solution of benzamide **6** (200 mg, 0.86 mmol) in dry THF (2 mL) was added to a cold (-78 °C) solution of *s*-BuLi (0.68 mL, 0.90 mmol) and TMEDA (0.14 mL, 0.95 mmol) in dry THF (4 mL), followed, after 3 h, by a solution of **8** (460 mg, 0.86 mmol) in dry THF (2 mL). The mixture was stirred for 1 h, then *t*-BuLi (1.32 mL of 1.45M pentane solution 1.9 mmol) was added. The reaction was stirred at -78 °C for 2 h

and allowed to warm to RT overnight. The mixture was quenched with H₂O (1 mL) and stirred for 1 h while bubbling air through the solution, then it was diluted with aq. sat. NaHCO₃ (20 mL) and extracted with EtOAc (2 x 30 mL). The combined extracts were washed with brine (50 mL), dried (Na₂SO₄) and concentrated. The yellow residue was purified by flash chromatography (EtOAc:hexanes, 1:1) to give **17** (80 mg, 17%) as a bright yellow solid. Proton and ¹³C NMR spectra indicated that this material exists as a ca. 2 : 1 mixture of atropisomers. Major atropisomer: ¹H: 7.48 (s, 1H), 7.33 (d, 1H, *J* = 2.7), 6.78 (d, 1H, *J* = 2.7), 5.63 (dd, 1H, *J* = 10.5, 3.2), 4.01 (s, 3H), 3.96 (br s, 6H), 3.95 (s, 3H), 3.90-3.70 (m, 4H), 3.27 (dd, 1H, *J* = 14.2, 10.5), 2.15 (dd, 1H, *J* = 14.2, 3.2), 1.23 (s, 3H), 1.05-0.85 (m, 21 H). ¹³C: 184.1, 181.2, 163.8, 162.7, 162.0, 160.2, 136.6, 134.8, 134.5, 122.5, 118.5, 108.8, 105.5, 104.1, 102.3, 64.5, 64.4, 63.7, 63.1, 56.8, 56.1, 44.4, 24.3, 18.3, 18.2, 12.5. **IR**: 1670. **HRMS** calcd for C₃₃H₄₆O₉Si [M + Na]⁺ = 637.2809, found 637.2807.

Deprotection of anthraquinone 17. A solution of **17** (45 mg, 0.073 mmol) and TBAF (1.0M in THF, 0.15 mL, 0.15 mmol) in THF (0.5 mL) was stirred at RT overnight, then it was diluted with aq. sat. NaHCO₃ (30 mL) and extracted with EtOAc (2 x 20 mL). The combined extracts were washed with brine (50 mL), dried (Na₂SO₄) and concentrated. Chromatographic purification of the residue (EtOAc ; hexanes, 1:1) gave the free alcohol (30 mg, quantitative) as a yellow solid. Unlike the case of **17**, proton and ¹³C NMR spectra of the present compound indicated the existence of a single species in solution. ¹H: 7.54 (s, 1H), 7.33 (d, 1H, *J* = 2.8), 6.78 (d, 1H, *J* = 2.0), 5.53 (ddd, 1H, *J* = 8.4, 8.4, 3.2), 4.02-3.96 (m, 4H), 4.02 (s, 3H), 3.99 (s, 1H), 3.97 (s, 3H), 3.96 (s, 3H), 3.47 (d, 1H, *J* = 10.0), 2.55 (dd, 1H, *J* = 14.4, 9.2), 2.02 (dd, 1H, *J* = 14.4, 3.0), 1.46 (s, 3H). ¹³C: 183.8, 181.1, 164.1, 162.1, 161.5, 159.6, 136.5, 134.9, 133.3, 122.4, 118.3, 109.9, 105.6, 105.0, 102.5, 64.9, 64.6, 64.3, 63.5, 44.9, 24.5. **IR**: 3500, 1669. **HRMS** calcd for C₂₄H₂₆O₉ [M + Na]⁺ = 481.1475, found 481.1478.

Anthraquinone 24. A mixture of the above alcohol (7.0 mg, 0.015 mmol) and IBX (13 mg, 0.045 mmol) in CH₃CN (1 mL) was refluxed for 45 min, then it was cooled to RT, filtered and concentrated. Purification of the residue by preparative TLC (EtOAc) provided **24** (6.0 mg, 88%) as a yellow solid. Proton and ¹³C NMR spectra of **24** indicated the existence of a single species in solution. ¹H (7.55 (s, 1H), 7.35 (d, 1H, *J* = 2.8), 6.80 (d, 1H, *J* = 2.8), 3.98 (s, 3H), 3.97 (s, 3H), 3.97 (s, 3H), 3.96-3.91 (m, 4H), 3.94 (s, 3H), 3.18 (s, 2H), 1.54 (s, 3H). ¹³C: 199.8, 183.5, 180.4, 164.3, 162.3, 159.5, 158.8, 136.5, 136.3, 132.3, 122.3, 117.9, 107.9, 105.7, 105.2, 102.8, 64.8, 64.2, 56.8, 56.6, 56.2, 52.8, 24.6. **IR**: 1718, 1670. **HRMS** calcd for C₂₄H₂₄O₉ [M + Na]⁺ = 479.1318, found 479.1320.

Topopyrone D, 4. A solution of **24** (6.0 mg, 0.013 mmol) in AcOH (1.5 mL) and aq. 48% HBr (1.0 mL) was refluxed overnight. The mixture was diluted with H₂O (10 mL) and extracted with EtOAc (2 x 40 mL). The combined extracts were washed with brine (50 mL), dried (Na₂SO₄) and concentrated to give fully synthetic topopyrone D (5 mg, quantitative) as an orange solid. ¹H (DMSO-*d*₆): 7.55 (s, 1H), 7.07 (d, 1H, *J* = 2.3), 6.60 (d, 1H, *J* = 2.3), 6.52 (s, 1H), 2.47 (s, 3H). ¹³C (DMSO-*d*₆): 185.4, 182.1, 180.8, 169.9, 164.6, 164.45, 164.41, 164.0, 159.1, 138.1, 134.0, 113.5, 110.4, 110.0, 108.7, 107.6, 106.6, 20.0.

Topopyrone D triacetate 26. Synthetic **4** was acetylated as described in the literature (ref. 7 in the paper) to afford the corresponding triacetate, which was purified by preparative TLC (EtOAc). This material produced ^1H and ^{13}C NMR spectra in complete accord with the literature (values in brackets, ref. 7 in the paper). ^1H : 8.19 [8.19] (s, 1H), 7.97 [7.97] (d, 1H, $J = 2.1$), 7.28 [7.28] (d, 1H, $J = 2.1$), 6.13 [6.13] (s, 1H), 2.52 [2.52] (s, 3H), 2.45 [2.45] (s, 3H), 2.40 [2.40] (s, 3H), 2.36 [2.36] (s, 3H). ^{13}C : 180.4 [180.2], 178.5 [178.4], 175.9 [175.7], 169.3 (2 overlapping resonances) [169.0], 168.2 [169.0], 165.9 [165.6], 159.7 [159.4], 154.9 (2 overlapping resonances) [154.7], 151.8 [151.6], 136.5 [136.3], 135.5 [135.3], 124.3 [124.1], 124.2 [124.1], 122.4 [122.3], 121.4 [121.2], 118.7 [118.4], 115.6 [115.3], 113.1 [112.8], 21.34 [21.1], 21.27 [21.0], 21.2 [20.9], 20.4 [20.1]. **HRMS** calcd for $\text{C}_{24}\text{H}_{16}\text{O}_{10}$ [$\text{M} + \text{Na}$] $^+$ = 487.0641, found 487.0645.

Synthesis of topopyrone B.

Anthraquinone 18. A solution of **7** (0.20 g, 0.74 mmol) in dry THF (1 mL) was added to a cold ($-78\text{ }^\circ\text{C}$) solution of $t\text{-BuLi}$ (1.5M pentane solution, 0.49 mL, 0.74 mmol) and TMEDA (0.11 mL, 0.78 mmol) in dry THF (3 mL). After 3 h, a solution of **8** (0.39 g, 0.74 mmol) in dry THF (1 mL) was added, and after stirring for 1 h, $t\text{-BuLi}$ (1.5M pentane solution, 0.97 mL, 1.48 mmol) was injected. The solution was stirred at $-78\text{ }^\circ\text{C}$, then it was allowed to warm to RT overnight, and finally it was quenched with H_2O (1mL) and stirred under a stream of air for 2 h. The mixture was diluted with aq. sat. NaHCO_3 (30 mL) and extracted with EtOAc (2 x 30 mL). The combined extracts were washed with brine (50 mL), dried (Na_2SO_4) and concentrated. Chromatographic purification of the residue (EtOAc:hexanes, 1:4) gave **18** (97 mg, 20%) as an orange oil. Proton and ^{13}C NMR spectra indicated that this material exists as a ca. 2:1 mixture of atropisomers. Major atropisomer: ^1H : 7.57 (s, 1H), 7.50 (s, 1H), 5.63 (dd, 0.5H, $J = 10.5, 3.1$), 4.06 (s, 3H), 4.04 (s, 3H), 3.98 (s, 3H), 3.96 (s, 3H), 3.90-3.70 (m, 4H), 3.25 (dd, 1H, $J = 14.0, 10.5\text{ Hz}$), 2.19 (dd, 1H, $J = 14.4, 3.1$), 1.23 (s, 3H), 1.10-0.81 (c, 21H). ^{13}C : 183.0, 180.5, 162.8, 160.6, 159.3, 134.8, 134.7, 133.4, 125.7, 123.3, 122.0, 108.9, 105.1, 104.3, 64.5, 64.4, 63.4, 63.0, 62.2, 57.1, 56.2, 44.7, 24.3, 18.3, 18.2, 12.5. Minor atropisomer: ^1H : 7.57 (s, 1H), 7.51 (s, 3H), 5.76 (dd, 0.5H, $J = 7.4, 5.2$), 4.06 (s, 3H), 4.04 (s, 3H), 3.98 (s, 3H), 3.92 (s, 3H), 2.58 (dd, 1H, $J = 14.4, 7.4$), 2.41 (dd, 1H, $J = 14.4, 5.2$), 1.32 (s, 3H), 1.10-0.81 (c, 21H). ^{13}C : 182.96, 180.7, 163.4, 159.2, 157.9, 157.8, 134.8, 134.6, 134.3, 125.6, 123.4, 120.2, 109.1, 105.2, 64.5, 64.3, 63.9, 62.8, 62.2, 56.1, 56.0, 45.2, 24.6, 18.2, 18.1, 12.6. **IR**: 1672. **HRMS** calcd for $\text{C}_{33}\text{H}_{45}^{35}\text{ClO}_9\text{Si}$ [$\text{M} + \text{Na}$] $^+$ = 671.2419, found 671.2422.

Deprotection of 18. A solution of **18** (84 mg, 0.13 mmol) and TBAF (1.0M in THF, 0.26 mL, 0.26 mmol) in THF (1 mL) was stirred at RT overnight, then it was diluted with aq. sat. NaHCO_3 (30 mL) and extracted with EtOAc (2 x 30 mL). The combined extracts were washed with brine (50 mL), dried (Na_2SO_4) and concentrated. Chromatographic purification of the residue (EtOAc : hexanes, 1:1) gave the free alcohol (58 mg, 91%) as a yellow solid. Unlike the case of **18**, proton and ^{13}C NMR spectra of the present compound indicated the existence of a single species in solution. ^1H : 7.58 (s, 1H), 7.55 (s, 1H), 5.55 (td, 1H, $J = 9.2, 3.1\text{ Hz}$), 4.07 (s, 3H), 4.04 (s, 3H), 4.00 (s, 3H), 4.02-3.98 (m, 4H), 3.97 (s, 3H), 3.48 (c, 1H), 2.56 (dd, 1H, $J = 14.8, 9.2$), 2.00 (dd, 1H, $J = 14.8, 3.5$), 1.46 (s, 3H). ^{13}C : 182.7, 180.4, 161.8, 160.1, 159.4, 158.0, 134.9 (2 overlapping resonances), 133.3, 125.9, 123.0, 121.7, 109.9,* 105.2,* 105.1, 64.9, 64.6, 64.1,

63.4, 62.2, 57.1, 56.6, 44.8, 24.5. **IR**: 3520, 1671. **HRMS** calcd for $C_{24}H_{25}^{35}ClO_9$ $[M + Na]^+$ = 515.1085, found 515.1087.

Anthraquinone 25. A solution of the above alcohol (100 mg, 0.20 mmol) IBX (114 mg, 0.40 mmol) in EtOAc (2 mL) was refluxed for 3 h, then it was cooled to RT, diluted with more EtOAc (30 mL) and washed with 1M NaOH (2 x 50 mL). The organic layer was dried (Na_2SO_4) and concentrated. Purification of the residue by preparative TLC afforded **25** (90 mg, 90%) as a yellow solid. **¹H**: 7.59 (s, 1H), 7.55 (s, 1H), 4.08 (s, 3H), 4.02 (s, 3H), 3.98 (s, 3H), 3.94 (c, 4H), 3.91 (s, 3H), 3.18 (s, 2H), 1.54 (s, 3H). **¹³C**: 199.6, 182.5, 179.8, 160.0, 159.6, 159.0, 158.2, 136.2, 133.2, 132.9, 126.2, 122.6, 121.6, 107.9, 105.4 (two overlapping resonances), 64.8, 64.0, 62.1, 57.2, 56.7, 52.9, 29.9. **IR**: 1718, 1672. **HRMS** calcd for $C_{24}H_{23}^{35}ClO_9$ $[M + H]^+$ = 491.1109, found 491.1106.

Topopyrone B trimethyl ether 27. A solution of ketone **25** (17 mg, 0.035 mmol) in conc. HBr (48%, 1 mL) and acetic acid (1.5 mL) was refluxed overnight., then it was concentrated to give crude topopyrone B (13 mg, quantitative), which without further purification was suspended in dry CH_2Cl_2 (0.8 mL) and treated with solid Meerwein's salt (30 mg, 0.2 mmol) while stirring at RT for 2h. The mixture was partitioned between EtOAc (10 mL) and sat. $NaHCO_3$ aq. (10 mL). The organic layer was dried (Na_2SO_4) and concentrated. Purification of the residue by preparative TLC (4% MeOH in DCM containing 0.5% triethylamine) gave pure **27** (3.0 mg, 30%) as a yellow solid. This material produced **¹H** and **¹³C** NMR spectra in complete accord with the literature (values in brackets, ref. 7 in the paper). **¹H**: 8.00 [8.00] (s, 1H), 7.59 [7.59] (s, 1H), 6.16 [6.16] (s, 1H), 4.13 [4.13] (s, 3H), 4.08 [4.08] (s, 3H), 4.04 [4.05] (s, 3H), 2.38 [2.38] (s, 3H). **¹³C**: 181.9 [181.7], 179.7 [179.5], 176.4 [176.2], 165.0 [164.8], 162.5 [162.3], 160.2 [160.0], 159.7 [159.5], 158.2 [158.0], 136.9 [136.7], 133.2 [133.0], 126.3 [126.1], 124.8 [124.6], 123.7 [123.5], 122.9 [122.7], 113.2 [113.0], 112.9 [112.6], 105.1 [104.9], 64.1 [63.9], 62.3 [62.1], 57.2 [57.0], 20.2 [20.0]. **IR**: 1734, 1673. **HRMS** calcd for $C_{21}H_{15}^{35}ClO_7$ $[M + H]^+$ = 415.0585, found 415.0587.

Index of ^1H and ^{13}C Spectra

<i>Subject</i>	<i>page</i>
Spectra of various Intermediates	
Spectra of 2-carbethoxymethyl-2-methyl-dioxolane	12
Spectra of 2-(2-hydroxyethyl)-2-methyl-dioxolane	14
Spectra of aldehyde 10	16
Spectra of benzamide 6	18
Spectra of 3-chloro-2,4-dihydroxybenzaldehyde	20
Spectra of 3-chloro-2,4-dimethoxybenzaldehyde	22
Spectra of 3-chloro-2,4-dimethoxybenzoic acid	24
Spectra of benzamide 7	26
Spectra of 4-bromo-3,5-dimethoxybenzyl alcohol	28
Spectra of TBS ether 9	30
Spectra of compound 11	32
Spectra of compound 12	34
Spectra of compound 13	36
Spectra of compound 14	38
Spectra of aldehyde 8	40
Synthesis of topopyrone D	42
Spectra of anthraquinone 17	42
Spectra of deprotected anthraquinone 17	44
Spectra of anthraquinone 24	46
Spectra of topopyrone D 4	48
Spectra of topopyrone D triacetate 26	52
Synthesis of topopyrone B	54
Spectra of anthraquinone 18	54
Spectra of deprotected anthraquinone 18	56
Spectra of anthraquinone 25	58
Spectra of topopyrone B trimethyl ether 26	60

11

12

25