Supporting Information

Total Synthesis of the Proposed Structures of Indole Alkaloids Lyaline and Lyadine

*M.-Lluïsa Bennasar, * Tomàs Roca, and Manuel Monerri*

Laboratory of Organic Chemistry, Faculty of Pharmacy, University of Barcelona,
08028-Barcelona, Spain

bennasar@ub.edu

- 1 Y = CH=CH₂ (Lyaline ?)
2 Y = CH(OH)CH₃ (Lyadine ?)

Contents:

- Pages S1 to S5: General experimental protocols and experimental procedures for the preparation of **12a-c**, **18a**, and **18d**.
- Pages S6 to S49: NMR spectra of all compounds.
- Pages S50 and S51: ESI-MS spectra of **1** and **2**.

General Procedures.

All nonaqueous reactions were performed under an argon atmosphere. All solvents were dried by standard methods. Reaction courses and product mixtures were routinely monitored by TLC on silica gel (precoated F₂₅₄ Merck plates). Drying of organic extracts during the workup of reactions was performed over anhydrous Na₂SO₄. Evaporation of the solvents was accomplished under reduced pressure with a rotatory evaporator. Flash chromatography was carried out on SiO₂ (silica gel 60, SDS, 0.04–0.06 mm). Melting points are uncorrected. Chemical shifts of NMR spectra are reported in ppm downfield (δ) from Me₄Si. *N*-methylharman (**6**) was prepared from commercially available harman following a reported procedure.¹ Pyridine **8a** was prepared by esterification of commercially available 3,5-pyridinedicarboxylic acid. Pyridines **8b**,² **8c**,^{2,3} and **8d**² were prepared following reported procedures.

Organometallic Derivative 9. *n*-BuLi (1.6 M in hexane, 0.35 mL, 0.56 mmol) was slowly added under Ar to a cooled (0°C) solution of harmane **6** (0.10 g, 0.51 mmol) in dry THF (10 mL), and the resulting red solution was stirred at 0°C for 30 min. After the solution was cooled to –78°C, a suspension of CuCN (46 mg, 0.51 mmol) in dry THF (5 mL) was added. The resulting mixture was stirred until the reaction temperature rose to 0°C, and then cooled again to –78°C. ZnCl₂ (1M in Et₂O, 0.51 mL, 0.51 mmol) was added and the mixture was stirred for 10 min. Then was used in the next reaction.

Dimethyl 1-Acetyl-4-[(9-methyl- β -carbolin-1-yl)methyl]-1,4-dihydropyridine-3,5-dicarboxylate (12a). Acetyl chloride (36 μ L, 0.51 mmol) was added to a solution of pyridine **8a** (0.10 g, 0.51 mmol) in anhydrous THF (6 mL) cooled at –10°C. The resulting mixture was stirred at –10°C for 15 min and then cooled to –78°C. Organometallic **9** (0.51 mmol) was added by cannula, and the mixture was allowed to slowly warm to rt for 12 h. The reaction mixture was poured into a 1:1 mixture of 20% NH₄OH and saturated aqueous NH₄Cl (25 mL), and extracted with CH₂Cl₂ (3 x 30 mL). The organic extracts were concentrated and the resulting residue was chromatographed (95:5 AcOEt-MeOH) to give 1,2-dihydropyridine **14a** (20 mg, 9%) and 1,4-dihydropyridine **12a** (90 mg, 41%). **14a**: ¹H NMR (CDCl₃, 400 MHz, assignments aided by HSQC) δ 2.11 (br s, 3H, COMe), 3.30 (t, *J* = 10.7 Hz, 1H, CH₂), 3.64 (s, 3H, OMe), 3.65 (dd, *J* = 5.2, 13 Hz, 1H, CH₂), 3.86 (s, 3H, OMe), 4.31 (s, 3H,

NMe), 6.20 (br s, 1H, pyr 2-H), 7.26 (t, $J = 7.2$ Hz, 1H, 6-H), 7.49 (d, $J = 8.4$ Hz, 1H, 8-H), 7.59 (t, $J = 7.2$ Hz, 1H, 7-H), 7.67 (s, 1H, pyr 4-H), 7.84 (d, $J = 5.2$ Hz, 1H, 4-H), 7.98 (br s, 1H, pyr 6-H), 8.08 (d, $J = 7.6$ Hz, 1H, 5-H), 8.24 (br s, 1H, 3-H); ^{13}C NMR (CDCl_3 , 100.6 MHz, assignments aided by HSQC) δ 21.0 (COMe), 32.2 (NMe), 38.3 (CH_2), 50.0 (pyr C-2), 51.9 (OMe), 52.0 (OMe), 109.7 (C-8), 113.6 (C-4), 119.6 (C-6), 121.0 (C-4b), 121.4 (C-5), 121.8 (pyr C-5), 128.3 (C-7), 129.5 (C-4a), 130.4 (pyr C-4), 136.2 (C-9a), 137.4 (C-3), 138.9 (pyr C-6), 139.6 (pyr C-3), 142.4 (C-1, C-8a), 165.1 (CO), 165.3 (CO), 169.7 (CO). **12a**: ^1H NMR (CDCl_3 , 300 MHz, assignments aided by HSQC and HMBC) δ 2.41 (s, 3H, COMe), 3.2-3.7 (br s, 8H, OMe, CH_2), 4.30 (s, 3H, NMe), 4.35 (t, $J = 7.5$ Hz, 1H, pyr 4-H), 7.26 (ddd, $J = 1.2, 6.9, 8.0$ Hz, 1H, 6-H), 7.4-8.2 (br s, 2H, pyr 2-H and 6-H), 7.48 (d, $J = 8.1$ Hz, 1H, 8-H), 7.59 (ddd, $J = 1.2, 6.9, 8.1$ Hz, 1H, 7-H), 7.80 (d, $J = 5.1$ Hz, 1H, 4-H), 8.09 (d, $J = 8.0$ Hz, 1H, 5-H), 8.22 (d, $J = 5.1$ Hz, 1H, 3-H); ^1H NMR (CDCl_3 , 300 MHz, 55 °C) δ 3.42 (s, 6H, OMe), 3.55 (d, $J = 7.5$ Hz, 2H, CH_2), 7.97 (s, 2H, pyr 2-H and 6-H); ^{13}C NMR (CDCl_3 , 75.4 MHz, assignments aided by HSQC and HMBC) δ 21.8 (COMe), 32.1 (NMe), 33.7 (pyr C-4), 43.4 (CH_2), 51.6 (OMe), 109.5 (C-8), 113.1 (C-4), 114.2 (pyr C-3 and C-5), 119.4 (C-6), 120.9 (C-4b), 121.2 (C-5), 128.1 (C-7), 129.2 (C-4a), 131.1 (pyr C-2 and C-6), 136.1 (C-9a), 137.2 (C-3), 141.4 (C-1), 142.2 (C-8a), 166.4 (CO), 167.2 (CO); HRMS calcd for $\text{C}_{24}\text{H}_{23}\text{N}_3\text{O}_5$ 433.1638, found 433.1639.

Methyl 1-Acetyl-5-ethyl-4-[(9-methyl- β -carbolin-1-yl)methyl]-1,4-dihydropyridine-3-carboxylate (12b). Acetyl chloride (36 μL , 0.51 mmol) was added to a solution of pyridine **8b** (84 mg, 0.51 mmol) in anhydrous THF (6 mL) cooled at -25°C . The resulting mixture was stirred at -25°C for 30 min, cooled to -78°C and then treated with **9** (0.51 mmol) as above. After the extractive workup and chromatography of the residue (AcOEt) a 4:1:1 mixture of dihydropyridines **12b**, **14b** and **16b** (83 mg, 40%) was obtained. An additional chromatography allowed the isolation of pure **12b** (2:1 mixture of rotamers): ^1H NMR (CDCl_3 , 300 MHz, assignment aided by HSQC and HMBC) δ 0.85 and 0.95 (2 br m, 3H, CH_2CH_3), 1.4-1.8 (3 br m, 2H, CH_2CH_3), 2.34 (s, 3H, COMe), 3.37 and 3.55 (2 br s, 3H, OMe), 3.4-3.7 (br m, 2H, CH_2), 3.90 (br m, 1H, pyr 4-H), 4.17 (br s, 3H, NMe), 6.40 and 7.0 (2 br s, 1H, pyr 6-H), 7.28 (t, $J = 8.2$ Hz, 1H, 6-H), 7.48 (d, $J = 8.4$, 1H, 8-H), 7.61 (t, $J = 7.6$ Hz, 1H, 7-H), 7.75 and 8.30 (2 br s, 1H, pyr 2-H), 7.85 (d, $J = 5.2$ Hz, 1H, 4-H), 8.12 (d, $J = 8$ Hz, 1H, 5-H), 8.34 (d, $J = 5.2$

Hz, 1H, 3-H); ^{13}C NMR (CDCl_3 , 75.4 MHz, assignment aided by HSQC and HMBC) δ 12.0 (CH_2CH_3), 21.7 and 21.8 (COMe), 25.9 (CH_2CH_3), 32.4 (NMe), 37.4 (pyr C-4), 42.4 (CH_2), 51.5 (OMe), 109.5 (C-8), 112.2 (pyr C-3), 113.0 (C-4), 116.6 and 118.2 (pyr C-6), 119.6 (C-6), 121.0 (C-4b), 121.3 (C-5), 127.1 (pyr C-5), 128.2 (C-7), 129.5 (C-4a), 131.9 and 133.3 (pyr C-2), 135.8 (C-9a), 137.9 (C-3), 142.3 (C-1), 142.4 (C-8a), 166.9 (CO), 167.2 (CO); HRMS calcd for $\text{C}_{24}\text{H}_{25}\text{N}_3\text{O}_3$ 403.1896, found 403.1890.

Methyl 1,5-Diacetyl-4-[(9-methyl- β -carbolin-1-yl)methyl]-1,4-dihydropyridine-3-carboxylate (12c). Acetyl chloride (36 μL , 0.51 mmol) was added to a solution of pyridine **8c** (91 mg, 0.51 mmol) in anhydrous CH_2Cl_2 (5 mL) cooled at -10°C , and the mixture was stirred at -10°C for 5 min. Trimethylsilyl triflate (92 μL , 0.51 mmol) was added and the mixture was stirred at -10°C for 1 h. The resulting pyridinium triflate (**11c**) was allowed to react as above with **9** (0.51 mmol). After the extractive workup and chromatography of the residue (96:4 AcOEt-MeOH) 1,4-dihydropyridine **12c** was obtained: 85 mg (40%); ^1H NMR (CDCl_3 , 300 MHz) δ 2.26 (br s, 3H, COMe), 2.44 (s, 3H, COMe), 3.24 (br s, 3H, OMe), 3.3 and 3.5 (2 br m, 2H, CH_2), 4.32 (s, 3H, NMe), 4.47 (t, $J = 7.2$ Hz, 1H, pyr 4-H), 7.26 (t, $J = 7.5$ Hz, 1H, 6-H), 7.49 (d, $J = 8.1$ Hz, 1H, 8-H), 7.59 (t, $J = 7.2$ Hz, 1H, 7-H), 7.80 (d, $J = 5.1$ Hz, 1H, 4-H), 7.85 (br s, 1H, pyr 2-H), 8.08 (d, $J = 7.5$ Hz, 1H, 5-H), 8.18 (d, $J = 5.1$ Hz, 1H, 3-H), 8.20 (br s, 1H, pyr 6-H); ^{13}C NMR (CDCl_3 , 75.4 MHz) δ 21.7 (COMe), 25.2 (COMe), 32.0 (NMe), 32.3 (pyr C-4), 42.6 (CH_2), 51.4 (OMe), 109.5 (C-8), 113.1 (C-4), 114.7 (pyr C-3), 119.3 (C-6), 120.8 (C-4b), 121.1 (C-5), 123.4 (pyr C-5), 128.1 (C-7), 129.2 (C-4a), 131.5 (pyr C-2), 136.0 (C-9a), 136.9 (C-3), 141.4 (C-1), 142.2 (C-8a), 166.0 (CO), 167.4 (CO), 196.2 (CO).

General Procedure for the N-deprotection of 12a and 12c. Dihydropyridines **12a,c** (0.1 mmol) were added to a solution of KOH in MeOH (1%, 3 mL) and stirring was maintained at rt for 15 min. The reaction mixture was poured into 5% aqueous Na_2CO_3 (10 mL) and extracted with CH_2Cl_2 (6x20 mL). The organic extracts were dried and concentrated. Pure **18a,c** were obtained after flash chromatography. Eluent and yields are given below.

Dimethyl 4-[(9-methyl- β -carbolin-1-yl)methyl]-1,4-dihydropyridine-3,5-dicarboxylate (18a). 98:2 AcOEt-MeOH; yield 85%; ^1H NMR ($\text{DMSO}-d_6$, 300 MHz,

assignment aided by HSQC and HMBC) δ 3.21 (s, 6H, OMe), 3.24 (d, J = 7.5 Hz, 2H, CH₂), 4.21 (t, J = 7.5 Hz, 1H, pyr 4-H), 4.25 (s, 3H, NMe), 7.24 (ddd, J = 0.9, 7.0, 7.8 Hz, 1H, 6-H), 7.30 (dd, J = 0.9, 5.4 Hz, 2H, pyr 2-H and 6-H), 7.59 (ddd, J = 1.2, 7.0, 8.4 Hz, 1H, 7-H), 7.73 (d, J = 8.4 Hz, 1H, 8-H), 7.92 (d, J = 5.1 Hz, 1H, 4-H), 8.10 (d, J = 5.1 Hz, 1H, 3-H), 8.20 (d, J = 7.8 Hz, 1H, 5-H), 9.24 (t, J = 5.7 Hz, 1H, NH); ¹³C NMR (DMSO-d₆, 75.4 MHz, assignment aided by HSQC and HMBC) δ 31.9 (NMe), 32.5 (pyr C-4), 42.9 (CH₂), 50.7 (OMe), 104.3 (pyr C-3 and C-5), 110.2 (C-8), 112.8 (C-4), 119.3 (C-6), 120.6 (C-4b), 121.3 (C-5), 128.0 (C-7), 128.1 (C-4a), 135.7 (C-9a), 136.4 (pyr C-2 and C-6), 136.9 (C-3), 141.8 (C-1), 142.3 (C-8a), 167.0 (CO).

Methyl 5-Acetyl-4-[(9-methyl- β -carbolin-1-yl)methyl]-1,4-dihydropyridine 3-carboxylate (18c). 9:1 AcOEt-MeOH; yield 80%; mp 134-135°C; 4:1 mixture of rotamers; ¹H NMR (DMSO-d₆, 300 MHz, major rotamer) δ 2.03 (s, 3H, COMe), 3.09 (dd, J = 8.1, 12.6 Hz, 1H, CH₂), 3.18 (dd, J = 6.6, 12.6 Hz, 1H, CH₂), 3.13 (s, 3H, OMe), 4.28 (s, 3H, NMe), 4.33 (dd, J = 8.1, 12.6 Hz, 1H, pyr 4-H), 7.24 (t, J = 7.8 Hz, 1H, 6-H), 7.28 (d, J = 5.1 Hz, 1H, pyr 2-H), 7.48 (d, J = 5.4, 1H, pyr 6-H), 7.59 (ddd, J = 0.9, 7.8, 8.7 Hz, 1H, 7-H), 7.73 (d, J = 8.7 Hz, 1H, 8-H), 7.90 (d, J = 5.0, 1H, 4-H), 8.08 (d, J = 5.0 Hz, 1H, 3-H), 8.19 (d, J = 7.8, 1H, 5-H), 9.36 (t, J = 5.1 Hz, NH); ¹³C NMR (DMSO-d₆, 100.6 MHz, major rotamer) δ 24.8 (COMe), 31.1 (pyr C-4), 31.9 (NMe), 42.3 (CH₂), 50.6 (OMe), 105.4 (pyr C-3), 110.3 (C-8), 112.8 (C-4), 115.2 (pyr C-5), 119.3 (C-6), 120.7 (C-4b), 121.3 (C-5), 128.0 (C-7), 128.2 (C-4a), 135.8 (C-9a), 136.0, 136.8 (pyr C-2, C-3), 138.3 (pyr C-6), 141.8 (C-1), 142.5 (C-8a), 166.9 (CO), 194.8 (CO).

References

1. (a) Karrer, P.; Müller, H. *J. Org. Chem.* **1957**, 22, 1433-1434. (b) Seki, H.; Hashimoto, A.; Hino, T. *Chem. Pharm. Bull.* **1993**, 41, 1169-1172.
2. Bracher, F.; Papke, T. *Monatsh. Chem.* **1995**, 126, 805-809.
3. Bennasar, M.-L.; Zulaica, E.; Roca, T. Alonso, Y.; Moneris, M. *Tetrahedron Lett.* **2003**, 44, 4711-4744.